



**THE UNITED
KINGDOM
ACID
WATERS
MONITORING
NETWORK**

**DATA
REPORT FOR
2003 – 2004
(YEAR 16)**



**THE UNITED KINGDOM ACID WATERS
MONITORING NETWORK
DATA REPORT FOR 2003 – 2004 (YEAR 16)**

Report to the Department for Environment, Food and Rural Affairs
(Contract EPG 1/3/160)
and the Department of the Environment Northern Ireland
(Contract CON 4/4 (38))

2004

Editors

E. M. Shilland¹
D. T. Monteith¹
M. Bonjean²
W. R. C. Beaumont³

¹ENSIS Ltd
²CEH Wallingford
³CEH Wareham

TABLE OF CONTENTS

TABLE OF CONTENTS	3
INTRODUCTION	10
THE MONITORING NETWORK.....	10
DATA FORMAT.....	11
REFERENCES	13
LOCATION OF UKAWMN SITES	14
1. LOCH COIRE NAN ARR	15
1.1. Spot sampled chemistry data	15
1.2. Macroinvertebrate data	16
1.2.1. Percentage abundance summary, Loch Coire nan Arr.....	16
1.2.2. Summary statistics, Loch Coire nan Arr	17
1.3. Fish data (for outflow stream).....	18
1.3.1. Summary of mean Trout density (numbers 100m ⁻²), Loch Coire nan Arr	18
1.4. Epilithic diatom data.....	19
1.4.1. Percentage abundance summary, Loch Coire nan Arr.....	19
1.4.2. Summary statistics, Loch Coire nan Arr	20
1.5. Aquatic macrophyte data, Loch Coire nan Arr.....	21
1.6. Sediment trap data, Loch Coire nan Arr	22
2. ALLT A'MHARCAIDH	23
2.1. Spot sampled chemistry data	23
2.2. Macroinvertebrate data	24
2.2.1. Percentage abundance summary, Allt a'Mharcaidh	24
2.2.2. Summary statistics, Allt a'Mharcaidh	25
2.3. Fish data	26
2.3.1. Summary of mean Salmon density (total numbers 100m ⁻²), Allt a'Mharcaidh.....	26
2.3.2. Summary of mean Trout density (numbers 100m ⁻²), Allt a'Mharcaidh...	26
2.4. Epilithic diatom data.....	27
2.4.1. Percentage abundance summary, Allt a'Mharcaidh	27
2.4.2. Summary statistics, Allt a'Mharcaidh	28
2.5. Aquatic macrophyte data, Allt a'Mharcaidh.....	29
3. ALLT NA COIRE NAN CON.....	30
3.1. Spot sampled chemistry data	30
3.2. Macroinvertebrate data	31

3.2.1. Percentage abundance summary, Allt na Coire nan Con.....	31
3.2.2. Summary statistics, Allt na Coire nan Con	32
3.3. Fish data	33
3.3.1. Summary of mean Salmon density (total numbers 100m ⁻²), Allt na Coire nan Con.....	33
3.3.2. Summary of mean Trout density (numbers 100m ⁻²), Allt na Coire nan Con.....	33
3.4. Epilithic diatom data.....	34
3.4.1. Percentage abundance summary, Allt na Coire nan Con.....	34
3.4.2. Summary statistics, Allt na Coire nan Con	35
3.5. Aquatic macrophyte data, Allt na Coire nan Con.....	36
4. LOCHNAGAR.....	37
4.1. Spot sampled chemistry data	37
4.2. Macroinvertebrate data	38
4.2.1. Percentage abundance summary, Lochnagar.....	38
4.2.2. Summary statistics, Lochnagar	39
4.3. Fish data (for outflow stream).....	40
4.3.1. Summary of mean Trout density (numbers 100m ⁻²), Lochnagar	40
4.4. Epilithic diatom data.....	41
4.4.1. Percentage abundance summary, Lochnagar.....	41
4.4.2. Summary statistics, Lochnagar	42
4.5. Aquatic macrophyte data, Lochnagar	43
4.6. Sediment trap data, Lochnagar	44
5. LOCH CHON	45
5.1. Spot sampled chemistry data	45
5.2. Macroinvertebrate data	46
5.2.1. Percentage abundance summary, Loch Chon	46
5.2.2. Summary statistics, Loch Chon	47
5.3. Fish data (for outflow stream).....	48
5.3.1. Summary of mean Trout density (numbers 100m ⁻²), Loch Chon	48
5.4. Epilithic diatom data.....	49
5.4.1. Percentage abundance summary, Loch Chon	49
5.4.2. Summary statistics, Loch Chon	50
5.5. Aquatic macrophyte data, Loch Chon	51
5.6. Sediment trap data, Loch Chon	52
6. LOCH TINKER	53
6.1. Spot sampled chemistry data	53
6.2. Macroinvertebrate data	54
6.2.1. Percentage abundance summary, Loch Tinker	54
6.2.2. Summary statistics, Loch Tinker.....	55
6.3. Fish data (for outflow stream).....	56
6.3.1. Summary of mean Trout density (numbers 100m ⁻²), Loch Tinker	56
6.4. Epilithic diatom data.....	57
6.4.1. Percentage abundance summary, Loch Tinker	57

6.4.2. Summary statistics, Loch Tinker.....	58
6.5. Aquatic macrophyte data, Loch Tinker	59
6.6. Sediment trap data, Loch Tinker	60
7. ROUND LOCH OF GLENHEAD.....	61
7.1. Spot sampled chemistry data	61
7.2. Macroinvertebrate data	62
7.2.1. Percentage abundance summary, Round Loch of Glenhead.....	62
7.2.2. Summary statistics, Round Loch of Glenhead.....	63
7.3. Fish data (for outflow stream).....	65
7.3.1. Summary of mean Trout density (numbers 100m ⁻²), Round Loch of Glenhead.....	65
7.4. Epilithic diatom data	66
7.4.1. Percentage abundance summary, Round Loch of Glenhead.....	66
7.4.2. Summary statistics, Round Loch of Glenhead.....	67
7.5. Aquatic macrophyte data, Round Loch of Glenhead.....	68
7.6. Sediment trap data, Round Loch of Glenhead	69
8. LOCH GRANNOCH.....	70
8.1. Spot sampled chemistry data	70
8.2. Macroinvertebrate data	71
8.2.1. Percentage abundance summary, Loch Grannoch	71
8.2.2. Summary statistics, Loch Grannoch.....	72
8.3. Fish data (for outflow stream).....	73
8.3.1. Summary of mean Trout density (numbers 100m ⁻²), Loch Grannoch....	73
8.4. Epilithic diatom data.....	74
8.4.1. Percentage abundance summary, Loch Grannoch	74
8.4.2. Summary statistics, Loch Grannoch.....	75
8.5. Aquatic macrophyte data, Loch Grannoch.....	76
8.6. Sediment trap data, Loch Grannoch	77
9. DARGALL LANE.....	78
9.1. Spot sampled chemistry data	78
9.2. Macroinvertebrate data	79
9.2.1. Percentage abundance summary, Dargall Lane	79
9.2.2. Summary statistics, Dargall Lane	80
9.3. Fish data	81
9.3.1. Summary of mean Trout density (numbers 100m ⁻²), Dargall Lane.....	81
9.4. Epilithic diatom data.....	82
9.4.1. Percentage abundance summary, Dargall Lane	82
9.4.2. Summary statistics, Dargall Lane.....	83
9.5. Aquatic macrophyte data, Dargall Lane.....	84
10. SCOAT TARN	85
10.1. Spot sampled chemistry data	85
10.2. Macroinvertebrate data	86
10.2.1. Percentage abundance summary, Scoat Tarn	86

10.2.2. Summary statistics, Scoat Tarn.....	87
10.3. Fish data (for outflow stream).....	88
10.3.1. Summary of mean Trout density (numbers 100m ⁻²), Scoat Tarn.....	88
10.4. Epilithic diatom data.....	89
10.4.1. Percentage abundance summary, Scoat Tarn	89
10.4.2. Summary statistics, Scoat Tarn.....	90
10.5. Aquatic macrophyte data, Scoat Tarn.....	91
10.6. Sediment trap data, Scoat Tarn	92
11. BURNMOOR TARN	93
11.1. Spot sampled chemistry data	93
11.2. Macroinvertebrate data	94
11.2.1. Percentage abundance summary, Burnmoor Tarn.....	94
11.2.2. Summary statistics, Burnmoor Tarn	95
11.3. Fish data (for outflow stream).....	96
11.3.1. Summary of mean Trout density (numbers 100m ⁻²), Burnmoor Tarn ..	96
11.4. Epilithic diatom data.....	97
11.4.1. Percentage abundance summary, Burnmoor Tarn.....	97
11.4.2. Summary statistics, Burnmoor Tarn	98
11.5. Aquatic macrophyte data, Burnmoor Tarn	99
11.6. Sediment trap data, Burnmoor Tarn.....	100
12. RIVER ETHEROW.....	101
12.1. Spot sampled chemistry data	101
12.2. Macroinvertebrate data	102
12.2.1. Percentage abundance summary, River Etherow	102
12.2.2. Summary statistics, River Etherow	103
12.3. Fish data	104
12.4. Epilithic diatom data.....	104
12.4.1. Percentage abundance summary, River Etherow	104
12.4.2. Summary statistics, River Etherow	105
12.5. Aquatic macrophyte data, River Etherow	106
13. OLD LODGE.....	107
13.1. Spot sampled chemistry data	107
13.2. Macroinvertebrate data	108
13.2.1. Percentage abundance summary, Old Lodge	108
13.2.2. Summary statistics, Old Lodge.....	110
13.3. Fish data	111
13.3.1. Summary of mean Trout density (numbers 100m ⁻²), Old Lodge.....	111
13.4. Epilithic diatom data.....	112
13.4.1. Percentage abundance summary, Old Lodge	112
13.4.2. Summary statistics, Old Lodge.....	113
13.5. Aquatic macrophyte data, Old Lodge	114
14. NARRATOR BROOK.....	115
14.1. Spot sampled chemistry data	115

14.2. Macroinvertebrate data	116
14.2.1. Percentage abundance summary, Narrator Brook	116
14.2.2. Summary statistics, Narrator Brook	117
14.3. Fish data	118
14.3.1. Summary of mean Trout density (numbers 100m ⁻²), Narrator Brook	118
14.4. Epilithic diatom data	119
14.4.1. Percentage abundance summary, Narrator Brook	119
14.4.2. Summary statistics, Narrator Brook	120
14.5. Aquatic macrophyte data, Narrator Brook	121
15. LLYN LLAGI	122
15.1. Spot sampled chemistry data	122
15.2. Macroinvertebrate data	123
15.2.1. Percentage abundance summary, Llyn Llagi	123
15.2.2. Summary statistics, Llyn Llagi	124
15.3. Fish data (for outflow stream)	125
15.3.1. Summary of mean Trout density (numbers 100m ⁻²), Llyn Llagi	125
15.4. Epilithic diatom data	126
15.4.1. Percentage abundance summary, Llyn Llagi	126
15.4.2. Summary statistics, Llyn Llagi	127
15.5. Aquatic macrophyte data, Llyn Llagi	128
15.6. Sediment trap data, Llyn Llagi	129
16. LLYN CWM MYNACH	130
16.1. Spot sampled chemistry data	130
16.2. Macroinvertebrate data	131
16.2.1. Percentage abundance summary, Llyn Cwm Mynach	131
16.2.2. Summary statistics, Llyn Cwm Mynach	132
16.3. Fish data (for outflow stream)	133
16.3.1. Summary of mean Trout density (numbers 100m ⁻²), Llyn Cwm Mynach	133
16.4. Epilithic diatom data	134
16.4.1. Percentage abundance summary, Llyn Cwm Mynach	134
16.4.2. Summary statistics, Llyn Cwm Mynach	135
16.5. Aquatic macrophyte data, Llyn Cwm Mynach	136
16.6. Sediment trap data, Llyn Cwm Mynach	137
17. AFON HAFREN	138
17.1. Spot sampled chemistry data	138
17.2. Macroinvertebrate data	139
17.2.1. Percentage abundance summary, Afon Hafren	139
17.2.2. Summary statistics, Afon Hafren	140
17.3. Fish data	141
17.3.1. Summary of mean Trout density (numbers 100m ⁻²), Afon Hafren	141
17.4. Epilithic diatom data	142
17.4.1. Percentage abundance summary, Afon Hafren	142
17.4.2. Summary statistics, Afon Hafren	143

17.5. Aquatic macrophyte data, Afon Hafren.....	144
18. AFON GWY	145
18.1. Spot sampled chemistry data	145
18.2. Macroinvertebrate data	146
18.2.1. Percentage abundance summary, Afon Gwy	146
18.2.2. Summary statistics, Afon Gwy	147
18.3. Fish data	148
18.3.1. Summary of mean Trout density (numbers 100m ⁻²), Afon Gwy.....	148
18.4. Epilithic diatom data	149
18.4.1. Percentage abundance summary, Afon Gwy	149
18.4.2. Summary statistics, Afon Gwy	150
18.5. Aquatic macrophyte data, Afon Gwy	151
19. BEAGHS BURN	152
19.1. Spot sampled chemistry data	152
19.2. Macroinvertebrate data	153
19.2.1. Percentage abundance summary, Beaghs Burn	153
19.2.2. Summary statistics, Beaghs Burn.....	154
19.3. Fish data	155
19.3.1. Summary of mean Trout density (numbers 100m ⁻²), Beaghs Burn ...	155
19.4. Epilithic diatom data	156
19.4.1. Percentage abundance summary, Beaghs Burn	156
19.4.2. Summary statistics, Beaghs Burn.....	157
19.5. Aquatic macrophyte data, Beaghs Burn.....	158
20. BENCROM RIVER	159
20.1. Spot sampled chemistry data	159
20.2. Macroinvertebrate data	160
20.2.1. Percentage abundance summary, Bencrom River	160
20.2.2. Summary statistics, Bencrom River.....	161
20.3. Fish data	162
20.3.1. Summary of mean Trout density (numbers 100m ⁻²), Bencrom River	162
20.4. Epilithic diatom data	163
20.4.1. Percentage abundance summary, Bencrom River	163
20.4.2. Summary statistics, Bencrom River.....	164
20.5. Aquatic macrophyte data, Bencrom River	165
21. BLUE LOUGH	166
21.1. Spot sampled chemistry data	166
21.2. Macroinvertebrate data	168
21.2.1. Percentage abundance summary, Blue Lough.....	168
21.2.2. Summary statistics, Blue Lough	169
21.3. Fish data (for outflow stream).....	170
21.3.1. Summary of mean Trout density (numbers 100m ⁻²), Blue Lough	170
21.4. Epilithic diatom data	171
21.4.1. Percentage abundance summary, Blue Lough.....	171

21.4.2. Summary statistics, Blue Lough	172
21.5. Aquatic macrophyte data, Blue Lough.....	173
21.6. Sediment trap data, Blue Lough	174
22. CONEYGLEN BURN.....	175
22.1. Spot sampled chemistry data	175
22.2. Macroinvertebrate data	176
22.2.1. Percentage abundance summary, Coneyglen Burn	176
22.2.2. Summary statistics, Coneyglen Burn.....	177
22.3. Fish data	178
22.3.1. Summary of mean Trout density (numbers 100m ⁻²), Coneyglen Burn	178
22.4. Epilithic diatom data	179
22.4.1. Percentage abundance summary, Coneyglen Burn	179
22.4.2. Summary statistics, Coneyglen Burn.....	180
22.5. Aquatic macrophyte data, Coneyglen Burn.....	181
23. LOCH COIRE FIONNARAICH	182
23.1. Spot sampled chemistry data	182
23.2. Macroinvertebrate data	183
23.2.1. Percentage abundance summary, Loch Coire Flonnaraich.....	183
23.2.2. Summary statistics, Loch Coire Flonnaraich	184
23.3. Fish data (for outflow stream).....	185
23.3.1. Summary of mean Trout density (numbers 100m ⁻²), Loch Coire Flonnaraich	185
23.4. Epilithic diatom data.....	186
23.4.1. Percentage abundance summary, Loch Coire Flonnaraich.....	186
23.4.2. Summary statistics, Loch Coire Flonnaraich	187
23.5. Aquatic macrophyte data, Loch Coire Flonnaraich	188
23.6. Sediment trap data, Loch Coire Flonnaraich.....	189

Cover photographs: *Sparganium angustifolium*, sediment trap buoy on Loch Grannoch and macrophyte surveying at Dargall Lane. All © Ewan Shilland.

INTRODUCTION

The UK Acid Waters Monitoring Network (UKAWMN) has been in continuous operation since 1988. For the first ten years biological and chemical data were summarised in an annual series of printed reports and these were followed by a detailed analysis of data in an interpretative report (Monteith and Evans, 2000), which is available on the [UKAWMN](#) web page. From the year 2000 annual data reports have been available from the [UKAWMN](#) web page. These are of a similar format to earlier annual reports but focus on graphical representations of time trends in raw data and diagnostic statistics (e.g. species richness and diversity indices). An analytical report, providing an interpretation of the first 15 years of data will be completed and made available on the project web page in the near future. A full description of sampling methods and analytical procedures, together with site descriptions, is presented in Patrick *et al.* (1991).

THE MONITORING NETWORK

The UKAWMN, funded by the UK Department of the Environment, originally consisted of 10 stream sites and 10 lakes, situated in those parts of the country most susceptible to acidification (see map, page 14). In 1990, two additional sites, Blue Lough and Coneyglen Burn, were added to the Network with funding from the Department of Environment (Northern Ireland). In January 1991 site 18, the Nant y Gronwen, was withdrawn from the Network at the request of the landowner and was replaced by a nearby moorland stream, the Afon Gwy. Due to water abstraction and damming by a local fish farm at site 1, Coire nan Arr, a nearby replacement control site was chosen, site 23, named Loch Coire Fionnaraich. Since 2001 the entire network has been funded by the UK Department of the Environment Food and Rural Affairs ([DEFRA](#)).

All sites are monitored chemically and biologically according to methodologies described by Patrick *et al.* (1991). Water samples are collected monthly at stream sites and quarterly at lake sites. Epilithic diatoms and benthic invertebrates are sampled annually. Aquatic macrophytes are surveyed bi-annually between June and September. Stream sites and the outflow streams of lake sites are electro-fished annually in the autumn.

In addition to the annual surveys, sediment cores were taken from all lake sites during the first five years of monitoring. These were radiometrically dated and analysed for diatoms, carbonaceous particles (derived from the combustion of fossil fuels) (Rose *et al.* 1995) and trace metals. Results of this work are presented in Patrick *et al.* (1995). Sediment traps have now been installed in all lakes and are emptied annually. The contents are analysed for diatom species composition and the flux of carbonaceous particles, allowing direct comparisons to be made with the historical (sediment core) record.

Water chemistry and macroinvertebrate sampling was prevented at several sites in the spring of 2001 by foot-and-mouth related access restrictions. Sampling was resumed across the Network in June 2001.

All chemical, physical and biological data are stored in a database managed by the Centre for Ecology and Hydrology and ENSIS. Summary data are available to scientific and other interested organisations on request. Further information on the UKAWMN, including site descriptions and photographs, is available via the Internet at the address: <http://www.ukawmn.ucl.ac.uk>

DATA FORMAT

The chemical and biological data are presented in a series of sections, summarised below, on a site-by-site basis.

Section 1: Time series graphs of key spot sampled chemical determinands for individual samples.

Summary table for key chemical determinands including: the mean over the 1988-1993 baseline period; the mean for the current year (2003-2004), the standard deviation for the current year; the Seasonal Kendall (Hirsch *et al.*, 1982) slope estimate for the period 1988-2004; and, the Seasonal Kendall trend significance level (p) (we consider values of less than 0.05 as evidence for a significant temporal trend). The normal number of observations per year is 4 for lakes and 12 for streams.

Section 2: Macroinvertebrates. Time series of macroinvertebrate taxon % abundance in annual aggregated samples (5 kick samples from lake littoral habitats or from riffle areas in streams), and annual total number of individual animals. Some species occurring at less than 1% relative abundance are omitted.

Macroinvertebrate summary statistic time series:

- 1) total number of individuals;
- 2) number of individuals identified at Genus level only (excludes some ubiquitous groups such as the chironomids and oligochaetes);
- 3) total number of taxa;
- 4) Diversity Indices. Although we have observed a general between-site relationship between acidity and the total number of macroinvertebrate species found, it is difficult to predict how chemical recovery might influence measures of diversity at specific sites. However, trends in the diversity scores described below should provide an indication of directional changes in community structure.

a) Hill's N_1 , the exponent of Shannon's Index and a measure of the number of abundant species in a sample (Hill, 1973).

b) Hill's N_2 , the reciprocal of Simpson's Index and a measure of the number of very abundant species in a sample (Hill, 1973).

c) E_5 , a measure of evenness based on the ratio $(N_2-1):(N_1-1)$. As a single species becomes more and more dominant, E_5 tends to zero.

Section 3: Salmonids. Summary histogram of mean density of trout and salmon, if present, in three 50m reaches (number of individuals caught per 100m² survey area) for each year of the monitoring period. (0+ = new recruits, >0+ = all fish over one year of age).

- Section 4: Epilithic diatoms. Time series of annual mean percentage frequency (from 3-4 replicate samples) of taxa occurring at greater than 2 % abundance in any one sample.
Epilithic diatom summary statistic time series. Mean, maximum and minimum for:
- a) Hill's N_1 (see above)
 - b) Hill's N_2 (see above)
 - c) E_5 (see above)
 - d) Diatom inferred pH (Di pH), based on the weighted average of species pH optima in the surface sediments of the 167 lake Surface Water Acidification Project dataset (Stevenson *et al.* 1991).
pH reconstructions are intended only for application to sedimentary diatoms but directional trends in inferred pH of epilithic assemblages should provide an indication of the direction of a response to changing acidity.
- Section 5: Aquatic macrophytes. For lakes relative species abundance determined on a five point scale (comparable to the DAFOR scoring system, Palmer *et al.* 1992) following shoreline survey, shore transects and deep water grapnel trawls, as follows:
1. rare/infrequent
 2. occasional but not abundant
 3. widespread but not abundant
 4. locally abundant
 5. widespread and abundant
- For streams, total macrophyte cover estimated for 5m sections of a 50m survey stretch and each then partitioned into proportional species abundance to provide percentage cover for each species. Data analysed for this report are the mean species cover estimates for the 50m stretches.
- Section 6: For lake sites only. Histogram of diatom species composition and carbonaceous particle flux estimated from annually retrieved sediment traps. Species occurring at less than 1% abundance in all years are omitted. Carbonaceous particle flux data presented in units of number of particles accumulated per trap (uniform trap size for all sites) per day.

REFERENCES

Hill, M. O. 1973 Diversity and evenness: a unifying notation and its consequences. *Ecology*, **54**, 427-31.

Hirsch, R. M., Slack, J. R. & Smith, R. A. (1982) A nonparametric trend test for seasonal data with serial dependence. *Water Resources Research*, **18**, **1**, 107-121.

Monteith, D. T. & Evans, C. D. (Eds.) 2000 *UK Acid Waters Monitoring Network: 10 Year Report. Analysis and Interpretation of Results, April 1988-March 1998*. ENSIS Ltd, London.

Palmer, M. A., Bell, S. L. & Butterfield, I. 1992 A botanical classification of standing waters in Britain: applications for conservation and monitoring. *Aquatic conservation: marine and freshwater ecosystems*, **2**, 125-143.


Patrick, S. T., Waters, D., Juggins, S. & Jenkins, A. (Eds.) 1991 *The United Kingdom Acid Waters Monitoring Network. Site descriptions and methodology report*. ENSIS Ltd, London.

Patrick, S. T., Monteith, D. T. & Jenkins, A. 1995 *UK Acid Waters Monitoring Network: The First Five Years. Analysis and interpretation of results, April 1988 - March 1993*. ENSIS Ltd, London.

Rose, N. L., Harlock, S., Appleby, P. G. & Battarbee, R. W. 1995 Dating of recent lake sediments in the United Kingdom and Ireland using spheroidal carbonaceous particle (SCP) concentration profiles. *The Holocene*, **5**, **3**, 328-335.



Stevenson, A. C., Juggins, S., Birks, H. J. B., Anderson, N. J., Battarbee, R. W., Berge, F., Davis, R. B., Flower, R. J., Haworth, E. Y., Jones, V. J., Kingston, J. C., Kreiser, A. M., Line, J. M., Munro, M. A. R. & Renberg, I. 1991 *The surface waters acidification project palaeolimnology programme: Modern diatom/lake-water chemistry data-set*. ENSIS Ltd, London.

LOCATION OF UKAWMN SITES



1. Loch Coire nan Arr

1.1. Spot sampled chemistry data



Determinand statistics


	mean 4/1988-3/1993	mean 4/2003-3/2004	std.dev. 4/2003-3/2004	SK*	p*
pH	6.39	6.39	0.20	0.00	0.93
ANC	42.66	54.15	19.15	0.59	0.13
Ca	42.53	42.75	11.18	0.00	1.00
Mg	63.67	61.25	17.85	0.00	0.86
Na	239.6	225.0	60.39	-0.02	0.52
K	9.51	9.04	2.47	0.00	0.14
Sol.Al	0.46	0.57	0.26	0.18	0.49

	mean 4/1988-3/1993	mean 4/2003-3/2004	std.dev. 4/2003-3/2004	SK*	p*
Sol.lab.Al	0.11	0.07	0.00	0.00	0.37
Cl	273.9	247.9	88.39	-0.02	0.74
SO_4	41.15	35.94	7.09	-0.03	0.01
XSO_4	12.38	9.91	4.53	-0.02	0.04
NO_3	2.80	2.21	0.00	0.00	0.27
Si	33.21	35.36	11.45	0.00	0.27
DOC	124.6	297.9	104.6	0.15	0.00


* Seasonal Kendall trend analysis: slope estimate (SK) and significance level (p)
Most units $\mu\text{eq l}^{-1}$, except Sol.Al, Sol.lab.Al and DOC ($\mu\text{mol l}^{-1}$)

1.2. Macroinvertebrate data

1.2.1. Percentage abundance summary, Loch Coire nan Arr




1.2.2. Summary statistics, Loch Coire nan Arr



1.3. Fish data (for outflow stream)


1.3.1. Summary of mean Trout density (numbers 100m⁻²), Loch Coire nan Arr




NF = Not fished

1.4. Epilithic diatom data

1.4.1. Percentage abundance summary, Loch Coire nan Arr




1.4.2. Summary statistics, Loch Coire nan Arr



1.5. Aquatic macrophyte data, Loch Coire nan Arr


Species Scores (1-5)



Aquatic macrophytes no longer surveyed after 1999.

1.6. Sediment trap data, Loch Coire nan Arr


Relative percentage frequency of diatom taxa and carbonaceous particle flux (no. trap⁻¹ day⁻¹).




Sediment trap samples no longer collected after 2002.

2. Allt a'Mharcaidh

2.1. Spot sampled chemistry data



_____ mean for first 5 years



_____ mean for first 5 years

Determinand statistics


	mean 4/1988-3/1993	mean 4/2003-3/2004	std.dev. 4/2003-3/2004	SK* 4/1988-3/2004	p* 4/1988-3/2004
pH	6.45	6.63	0.36	0.00	0.60
ANC	51.57	69.07	22.79	0.88	0.02
Ca	42.41	47.54	8.76	0.00	0.51
Mg	30.39	30.62	2.89	0.00	0.14
Na	132.6	134.1	14.81	0.00	0.46
K	6.75	6.20	0.58	0.00	0.02
Sol.AI	1.31	1.22	1.10	-0.20	0.56

	mean 4/1988-3/1993	mean 4/2003-3/2004	std.dev. 4/2003-3/2004	SK* 4/1988-3/2004	p* 4/1988-3/2004
Sol.lab.AI	0.23	0.17	0.14	-0.09	0.09
Cl	108.4	96.48	6.70	-0.02	0.27
SO_4	44.41	41.32	3.86	-0.02	0.01
X SO_4	33.02	31.19	4.37	-0.01	0.01
NO_3	1.50	2.21	0.00	0.00	0.07
Si	170.8	211.9	49.00	0.02	0.13
DOC	167.7	215.3	147.9	0.07	0.01


* Seasonal Kendall trend analysis: slope estimate (SK) and significance level (p)
Most units $\mu\text{eq l}^{-1}$, except Sol.AI, Sol.lab.AI and DOC ($\mu\text{mol l}^{-1}$)

2.2. Macroinvertebrate data

2.2.1. Percentage abundance summary, Allt a'Mharcaidh




2.2.2. Summary statistics, Allt a'Mharcaidh




2.3. Fish data

2.3.1. Summary of mean Salmon density (total numbers 100m⁻²), Allt a'Mharcaidh




2.3.2. Summary of mean Trout density (numbers 100m⁻²), Allt a'Mharcaidh




2.4. Epilithic diatom data

2.4.1. Percentage abundance summary, Allt a'Mharcaidh




2.4.2. Summary statistics, Allt a'Mharcaidh



2.5. Aquatic macrophyte data, Allt a'Mharcaidh



Percentage Species Cover



+ Represents <0.1% abundance

3. Allt na Coire nan Con

3.1. Spot sampled chemistry data



Determinand statistics


	mean 4/1988-3/1993	mean 4/2003-3/2004	std.dev. 4/2003-3/2004	SK* 4/1988-3/2004	p* 4/1988-3/2004
pH	5.81	6.03	0.43	0.01	0.17
ANC	32.53	62.50	30.76	1.63	0.01
Ca	58.97	58.46	19.23	0.00	0.83
Mg	70.89	63.06	15.60	0.00	0.90
Na	272.0	238.8	48.11	-0.02	0.62
K	9.14	6.15	1.63	-0.01	0.02
Sol.Al	2.40	2.52	1.06	0.20	0.75

	mean 4/1988-3/1993	mean 4/2003-3/2004	std.dev. 4/2003-3/2004	SK* 4/1988-3/2004	p* 4/1988-3/2004
Sol.lab.Al	0.80	0.19	0.09	-0.42	0.05
Cl	321.5	249.8	58.28	-0.07	0.60
SO_4	61.95	52.43	11.16	-0.04	0.01
XSO_4	28.19	26.20	6.84	-0.02	0.18
NO_3	4.91	2.21	0.00	0.00	0.26
Si	65.54	83.99	36.39	0.02	0.03
DOC	266.7	576.4	242.6	0.21	0.00


* Seasonal Kendall trend analysis: slope estimate (SK) and significance level (p)
Most units $\mu\text{eq l}^{-1}$, except Sol.Al, Sol.lab.Al and DOC ($\mu\text{mol l}^{-1}$)

3.2. Macroinvertebrate data

3.2.1. Percentage abundance summary, Allt na Coire nan Con




3.2.2. Summary statistics, Allt na Coire nan Con




3.3. Fish data

3.3.1. Summary of mean Salmon density (total numbers 100m⁻²), Allt na Coire nan Con




3.3.2. Summary of mean Trout density (numbers 100m⁻²), Allt na Coire nan Con




3.4. Epilithic diatom data

3.4.1. Percentage abundance summary, Allt na Coire nan Con




3.4.2. Summary statistics, Allt na Coire nan Con



3.5. Aquatic macrophyte data, Allt na Coire nan Con



Percentage Species Cover



+ Represents <0.1% abundance

4. Lochnagar

4.1. Spot sampled chemistry data



Determinand statistics


	mean 4/1988-3/1993	mean 4/2003-3/2004	std.dev. 4/2003-3/2004	SK* 4/1988-3/2004	p* 4/1988-3/2004
pH	5.40	5.51	0.09	0.01	0.31
ANC	2.84	5.87	1.23	0.25	0.07
Ca	29.06	29.12	1.89	-0.01	0.04
Mg	30.57	33.33	0.68	0.00	0.11
Na	92.22	84.78	4.35	-0.02	0.06
K	7.54	5.26	0.44	-0.01	0.00
Sol.AI	1.19	0.59	0.25	-0.91	0.20

	mean 4/1988-3/1993	mean 4/2003-3/2004	std.dev. 4/2003-3/2004	SK* 4/1988-3/2004	p* 4/1988-3/2004
Sol.lab.AI	0.66	0.31	0.09	-0.33	0.32
Cl	86.88	78.87	6.09	-0.03	0.15
SO_4	60.64	46.35	1.04	-0.05	0.00
XSO_4	51.51	38.07	0.51	-0.05	0.00
NO_3	10.77	22.14	2.74	0.01	0.04
Si	74.21	74.64	11.96	0.00	0.92
DOC	72.37	97.92	7.98	0.05	0.00


* Seasonal Kendall trend analysis: slope estimate (SK) and significance level (p)
Most units $\mu\text{eq l}^{-1}$, except Sol.AI, Sol.lab.AI and DOC ($\mu\text{mol l}^{-1}$)

4.2. Macroinvertebrate data

4.2.1. Percentage abundance summary, Lochnagar




4.2.2. Summary statistics, Lochnagar



4.3. Fish data (for outflow stream)


4.3.1. Summary of mean Trout density (numbers 100m⁻²), Lochnagar




NF = Not fished

4.4. Epilithic diatom data

4.4.1. Percentage abundance summary, Lochnagar




4.4.2. Summary statistics, Lochnagar




4.5. Aquatic macrophyte data, Lochnagar

Species Scores (1-5)





4.6. Sediment trap data, Lochnagar

Relative percentage frequency of diatom taxa and carbonaceous particle flux (no. cm⁻² yr⁻¹).



5. Loch Chon

5.1. Spot sampled chemistry data



Determinand statistics


	mean 4/1988-3/1993	mean 4/2003-3/2004	std.dev. 4/2003-3/2004	SK* 4/1988-3/2004	p* 4/1988-3/2004
pH	5.47	6.11	0.16	0.04	0.00
ANC	14.84	54.75	10.99	2.83	0.00
Ca	75.74	86.00	6.26	0.00	0.95
Mg	47.50	51.04	2.58	0.00	0.76
Na	186.3	146.7	10.87	-0.07	0.11
K	5.65	8.33	0.65	0.00	0.48
Sol.Al	2.43	1.60	0.34	-1.25	0.00

	mean 4/1988-3/1993	mean 4/2003-3/2004	std.dev. 4/2003-3/2004	SK* 4/1988-3/2004	p* 4/1988-3/2004
Sol.lab.Al	0.97	0.17	0.11	-1.23	0.00
Cl	223.4	156.3	6.71	-0.14	0.15
SO_4	72.15	57.29	2.08	-0.07	0.00
XSO_4	48.69	40.87	1.40	-0.05	0.02
NO_3	9.88	19.64	2.77	0.01	0.02
Si	31.58	32.50	13.36	0.00	0.44
DOC	225.4	433.3	32.63	0.21	0.00


* Seasonal Kendall trend analysis: slope estimate (SK) and significance level (p)
Most units $\mu\text{eq l}^{-1}$, except Sol.Al, Sol.lab.Al and DOC ($\mu\text{mol l}^{-1}$)

5.2. Macroinvertebrate data

5.2.1. Percentage abundance summary, Loch Chon




5.2.2. Summary statistics, Loch Chon



5.3. Fish data (for outflow stream)


5.3.1. Summary of mean Trout density (numbers 100m⁻²), Loch Chon




NF = Not fished

5.4. Epilithic diatom data

5.4.1. Percentage abundance summary, Loch Chon




5.4.2. Summary statistics, Loch Chon




5.5. Aquatic macrophyte data, Loch Chon

Species Scores (1-5)





5.6. Sediment trap data, Loch Chon

Relative percentage frequency of diatom taxa and carbonaceous particle flux (no. cm⁻² yr⁻¹).



6. Loch Tinker

6.1. Spot sampled chemistry data



Determinand statistics


	mean	mean	std.dev.	SK*	p*
	4/1988-3/1993	4/2003-3/2004	4/2003-3/2004	4/1988-3/2004	4/1988-3/2004
pH	6.10	6.40	0.17	0.01	0.05
ANC	55.51	76.93	19.26	1.48	0.03
Ca	86.00	88.50	13.31	-0.01	0.57
Mg	46.45	42.29	4.78	-0.01	0.18
Na	140.0	101.1	8.96	-0.06	0.07
K	7.22	6.15	2.07	0.00	0.20
Sol.Al	0.72	0.47	0.18	-0.09	0.65

	mean	mean	std.dev.	SK*	p*
	4/1988-3/1993	4/2003-3/2004	4/2003-3/2004	4/1988-3/2004	4/1988-3/2004
Sol.lab.Al	0.12	0.07	0.00	0.00	0.75
Cl	162.6	93.66	12.68	-0.10	0.10
SO ₄	52.96	44.79	5.24	-0.06	0.02
XSO ₄	35.88	34.95	4.91	-0.05	0.06
NO ₃	2.03	2.21	0.00	0.00	0.03
Si	24.06	18.39	16.37	0.00	0.66
DOC	345.6	477.1	95.59	0.19	0.00

* Seasonal Kendall trend analysis: slope estimate (SK) and significance level (p)
Most units $\mu\text{eq l}^{-1}$, except Sol.Al, Sol.lab.Al and DOC ($\mu\text{mol l}^{-1}$)


6.2. Macroinvertebrate data

6.2.1. Percentage abundance summary, Loch Tinker



No sampling in 2001 due to Foot and Mouth restrictions.


6.2.2. Summary statistics, Loch Tinker



No sampling in 2001 due to Foot and Mouth restrictions.

6.3. Fish data (for outflow stream)


6.3.1. Summary of mean Trout density (numbers 100m⁻²), Loch Tinker




NF = Not fished

6.4. Epilithic diatom data


6.4.1. Percentage abundance summary, Loch Tinker



6.4.2. Summary statistics, Loch Tinker




6.5. Aquatic macrophyte data, Loch Tinker




6.6. Sediment trap data, Loch Tinker

Relative percentage frequency of diatom taxa and carbonaceous particle flux (no. cm⁻² yr⁻¹).




7. Round Loch of Glenhead

7.1. Spot sampled chemistry data



— mean for first 5 years



— mean for first 5 years

Determinand statistics


	mean 4/1988-3/1993	mean 4/2003-3/2004	std.dev. 4/2003-3/2004	SK* 4/1988-3/2004	p* 4/1988-3/2004
pH	4.90	5.29	0.18	0.02	0.00
ANC	-7.13	12.58	4.76	1.57	0.00
Ca	34.92	31.00	2.74	-0.01	0.01
Mg	48.55	38.54	2.75	-0.01	0.01
Na	193.8	135.9	5.47	-0.08	0.00
K	9.00	7.12	1.05	-0.01	0.07
Sol.Al	3.65	2.65	0.12	-2.00	0.01

	mean 4/1988-3/1993	mean 4/2003-3/2004	std.dev. 4/2003-3/2004	SK* 4/1988-3/2004	p* 4/1988-3/2004
Sol.lab.Al	2.54	0.96	0.44	-3.60	0.00
Cl	223.6	147.9	7.45	-0.16	0.02
SO_4	68.42	43.75	1.70	-0.11	0.00
XSO_4	44.94	28.22	1.63	-0.08	0.01
NO_3	4.81	7.86	2.86	0.00	0.11
Si	26.32	16.43	13.64	0.00	0.31
DOC	233.3	311.1	31.55	0.13	0.00

* Seasonal Kendall trend analysis: slope estimate (SK) and significance level (p)
Most units $\mu\text{eq l}^{-1}$, except Sol.Al, Sol.lab.Al and DOC ($\mu\text{mol l}^{-1}$)


7.2. Macroinvertebrate data

7.2.1. Percentage abundance summary, Round Loch of Glenhead



No sampling in 2001 due to Foot and Mouth restrictions.


7.2.2. Summary statistics, Round Loch of Glenhead



No sampling in 2001 due to Foot and Mouth restrictions.

7.3. Fish data (for outflow stream)


7.3.1. Summary of mean Trout density (numbers 100m⁻²), Round Loch of Glenhead




NF = Not fished

7.4. Epilithic diatom data


7.4.1. Percentage abundance summary, Round Loch of Glenhead



7.4.2. Summary statistics, Round Loch of Glenhead




7.5. Aquatic macrophyte data, Round Loch of Glenhead





7.6. Sediment trap data, Round Loch of Glenhead

Relative percentage frequency of diatom taxa and carbonaceous particle flux (no. cm⁻² yr⁻¹).



8. Loch Grannoch

8.1. Spot sampled chemistry data



Determinand statistics


	mean 4/1988-3/1993	mean 4/2003-3/2004	std.dev. 4/2003-3/2004	SK* 4/1988-3/2004	p* 4/1988-3/2004
pH	4.65	4.86	0.15	0.01	0.19
ANC	-34.68	4.04	6.34	2.50	0.00
Ca	51.47	40.67	1.04	-0.03	0.00
Mg	56.54	46.11	3.15	-0.01	0.01
Na	239.1	179.7	9.05	-0.10	0.03
K	4.73	6.07	1.16	0.00	0.15
Sol.Al	11.72	5.81	0.67	-6.40	0.02

	mean 4/1988-3/1993	mean 4/2003-3/2004	std.dev. 4/2003-3/2004	SK* 4/1988-3/2004	p* 4/1988-3/2004
Sol.lab.Al	9.12	2.53	0.77	-9.55	0.00
Cl	283.8	197.2	14.08	-0.18	0.05
SO_4	98.14	60.42	2.08	-0.16	0.00
XSO_4	68.34	39.71	1.86	-0.12	0.01
NO_3	13.89	11.61	4.89	0.00	0.55
Si	59.02	22.38	21.46	-0.02	0.02
DOC	319.3	479.2	43.83	0.17	0.00

* Seasonal Kendall trend analysis: slope estimate (SK) and significance level (p)
Most units $\mu\text{eq l}^{-1}$, except Sol.Al, Sol.lab.Al and DOC ($\mu\text{mol l}^{-1}$)


8.2. Macroinvertebrate data

8.2.1. Percentage abundance summary, Loch Grannoch



No sampling in 2001 due to Foot and Mouth restrictions.


8.2.2. Summary statistics, Loch Grannoch



No sampling in 2001 due to Foot and Mouth restrictions.

8.3. Fish data (for outflow stream)


8.3.1. Summary of mean Trout density (numbers 100m⁻²), Loch Grannoch




NF = Not fished

8.4. Epilithic diatom data

8.4.1. Percentage abundance summary, Loch Grannoch




8.4.2. Summary statistics, Loch Grannoch



8.5. Aquatic macrophyte data, Loch Grannoch


Species Scores (1-5)



No aquatic macrophyte survey in 2003.



8.6. Sediment trap data, Loch Grannoch

Relative percentage frequency of diatom taxa and carbonaceous particle flux (no. cm⁻² yr⁻¹).



9. Dargall Lane

9.1. Spot sampled chemistry data



Determinand statistics


	mean 4/1988-3/1993	mean 4/2003-3/2004	std.dev. 4/2003-3/2004	SK* 4/1988-3/2004	p* 4/1988-3/2004
pH	5.44	6.10	0.42	0.03	0.00
ANC	3.39	26.98	15.55	1.45	0.00
Ca	51.58	49.88	13.14	-0.01	0.04
Mg	56.62	52.36	9.09	-0.01	0.03
Na	182.2	146.7	22.02	-0.07	0.01
K	9.06	8.42	3.32	0.00	0.06
Sol.AI	2.06	0.68	0.56	-1.80	0.00

	mean 4/1988-3/1993	mean 4/2003-3/2004	std.dev. 4/2003-3/2004	SK* 4/1988-3/2004	p* 4/1988-3/2004
Sol.lab.AI	1.48	0.21	0.19	-1.50	0.00
Cl	207.9	152.6	30.67	-0.13	0.02
SO₄	82.02	68.23	10.48	-0.08	0.00
XSO₄	60.18	52.21	9.31	-0.06	0.01
NO₃	10.70	9.08	6.76	0.00	0.19
Si	70.68	90.83	60.62	0.00	0.23
DOC	117.8	152.1	66.11	0.07	0.00

* Seasonal Kendall trend analysis: slope estimate (SK) and significance level (p)
Most units $\mu\text{eq l}^{-1}$, except Sol.AI, Sol.lab.AI and DOC ($\mu\text{mol l}^{-1}$)


9.2. Macroinvertebrate data

9.2.1. Percentage abundance summary, Dargall Lane



No sampling in 2001 due to Foot and Mouth restrictions.


9.2.2. Summary statistics, Dargall Lane



No sampling in 2001 due to Foot and Mouth restrictions.


9.3. Fish data

9.3.1. Summary of mean Trout density (numbers 100m⁻²), Dargall Lane




9.4. Epilithic diatom data

9.4.1. Percentage abundance summary, Dargall Lane




9.4.2. Summary statistics, Dargall Lane



9.5. Aquatic macrophyte data, Dargall Lane

Percentage Species Cover





+ Represents <0.1% abundance

2003 data pending.

10. Scoat Tarn

10.1. Spot sampled chemistry data



mean for first 5 years

mean for first 5 years

Determinand statistics


	mean 4/1988-3/1993	mean 4/2003-3/2004	std.dev. 4/2003-3/2004	SK* 4/1988-3/2004	p* 4/1988-3/2004
pH	5.00	5.19	0.11	0.01	0.00
ANC	-19.95	-1.63	5.42	1.33	0.00
Ca	35.75	27.00	1.29	-0.02	0.00
Mg	51.50	39.58	3.76	-0.01	0.00
Na	178.3	123.9	12.55	-0.07	0.00
K	8.27	5.77	0.99	-0.01	0.00
Sol.AI	5.36	1.44	0.80	-7.86	0.00

	mean 4/1988-3/1993	mean 4/2003-3/2004	std.dev. 4/2003-3/2004	SK* 4/1988-3/2004	p* 4/1988-3/2004
Sol.lab.AI	4.91	0.88	0.81	-7.67	0.00
Cl	206.3	128.2	10.91	-0.13	0.00
SO_4	60.73	50.52	1.99	-0.05	0.00
XSO_4	39.06	37.06	2.59	-0.02	0.10
NO_3	21.21	14.29	3.25	0.00	0.18
Si	42.50	39.64	16.29	0.00	0.94
DOC	44.50	102.1	23.94	0.07	0.00

* Seasonal Kendall trend analysis: slope estimate (SK) and significance level (p)
Most units $\mu\text{eq l}^{-1}$, except Sol.AI, Sol.lab.AI and DOC ($\mu\text{mol l}^{-1}$)


10.2. Macroinvertebrate data

10.2.1. Percentage abundance summary, Scoat Tarn



No sampling in 2001 due to Foot and Mouth restrictions.


10.2.2. Summary statistics, Scoat Tarn



No sampling in 2001 due to Foot and Mouth restrictions.

10.3. Fish data (for outflow stream)


10.3.1. Summary of mean Trout density (numbers 100m⁻²), Scoat Tarn




NF = Not fished

10.4. Epilithic diatom data


10.4.1. Percentage abundance summary, Scoat Tarn



10.4.2. Summary statistics, Scoat Tarn




10.5. Aquatic macrophyte data, Scoat Tarn





10.6. Sediment trap data, Scoat Tarn

Relative percentage frequency of diatom taxa and carbonaceous particle flux (no. cm⁻² yr⁻¹).



11. Burnmoor Tarn

11.1. Spot sampled chemistry data



Determinand statistics


	mean 4/1988-3/1993	mean 4/2003-3/2004	std.dev. 4/2003-3/2004	SK* 4/1988-3/2004	p* 4/1988-3/2004
pH	6.50	6.78	0.16	0.01	0.31
ANC	54.11	88.06	29.09	1.65	0.02
Ca	95.97	91.12	18.40	-0.03	0.02
Mg	68.50	62.50	9.67	-0.01	0.00
Na	208.5	155.4	10.87	-0.10	0.00
K	9.24	6.92	0.66	-0.01	0.00
Sol.AI	0.28	0.10	0.06	0.00	0.89

	mean 4/1988-3/1993	mean 4/2003-3/2004	std.dev. 4/2003-3/2004	SK* 4/1988-3/2004	p* 4/1988-3/2004
Sol.lab.AI	0.08	0.07	0.00	0.00	0.44
Cl	233.5	149.3	2.30	-0.20	0.00
SO₄	81.46	63.02	5.21	-0.10	0.00
XSO₄	56.93	47.34	5.01	-0.06	0.01
NO₃	5.63	3.62	2.82	0.00	0.37
Si	53.21	49.82	17.12	0.00	0.98
DOC	126.5	200.0	26.35	0.10	0.00

* Seasonal Kendall trend analysis: slope estimate (SK) and significance level (p)
Most units $\mu\text{eq l}^{-1}$, except Sol.AI, Sol.lab.AI and DOC ($\mu\text{mol l}^{-1}$)


11.2. Macroinvertebrate data

11.2.1. Percentage abundance summary, Burnmoor Tarn



No sampling in 2001 due to Foot and Mouth restrictions.


11.2.2. Summary statistics, Burnmoor Tarn



No sampling in 2001 due to Foot and Mouth restrictions.

11.3. Fish data (for outflow stream)


11.3.1. Summary of mean Trout density (numbers 100m⁻²), Burnmoor Tarn




NF = Not fished

11.4. Epilithic diatom data


11.4.1. Percentage abundance summary, Burnmoor Tarn



11.4.2. Summary statistics, Burnmoor Tarn




11.5. Aquatic macrophyte data, Burnmoor Tarn





11.6. Sediment trap data, Burnmoor Tarn

Relative percentage frequency of diatom taxa and carbonaceous particle flux (no. cm⁻² yr⁻¹).



12. River Etherow

12.1. Spot sampled chemistry data



Determinand statistics


	mean 4/1989-3/1994	mean 4/2003-3/2004	std.dev. 4/2003-3/2004	SK* 4/1989-3/2004	p* 4/1989-3/2004
pH	5.53	5.92	0.97	0.02	0.16
ANC	7.12	74.67	58.50	6.50	0.01
Ca	178.3	164.9	20.84	-0.05	0.02
Mg	172.1	158.6	23.53	-0.03	0.01
Na	300.5	288.4	51.95	-0.03	0.25
K	19.83	20.06	2.80	-0.01	0.05
Sol.Al	5.24	3.27	3.34	-1.29	0.37

	mean 4/1989-3/1994	mean 4/2003-3/2004	std.dev. 4/2003-3/2004	SK* 4/1989-3/2004	p* 4/1989-3/2004
Sol.lab.Al	2.64	0.93	1.16	-1.00	0.02
Cl	316.4	269.2	45.97	-0.11	0.04
SO ₄	288.7	211.5	20.44	-0.40	0.00
XSO ₄	255.4	183.2	18.84	-0.39	0.00
NO ₃	44.47	45.18	12.07	0.00	0.57
Si	234.0	268.2	61.08	-0.01	0.36
DOC	377.3	553.5	455.9	0.31	0.01


* Seasonal Kendall trend analysis: slope estimate (SK) and significance level (p)
Most units μeq l⁻¹, except Sol.Al, Sol.lab.Al and DOC (μmol l⁻¹)

12.2. Macroinvertebrate data

12.2.1. Percentage abundance summary, River Etherow



12.2.2. Summary statistics, River Etherow




12.3. Fish data


No fish are present in this reach of the river.

12.4. Epilithic diatom data

12.4.1. Percentage abundance summary, River Etherow




12.4.2. Summary statistics, River Etherow



12.5. Aquatic macrophyte data, River Etherow



Percentage Species Cover



+ Represents <0.1% abundance

13. Old Lodge

13.1. Spot sampled chemistry data



Determinand statistics


	mean 4/1988-3/1993	mean 4/2003-3/2004	std.dev. 4/2003-3/2004	SK* 4/1988-3/2004	p* 4/1988-3/2004
pH	4.55	5.05	0.44	0.05	0.00
ANC	-41.74	15.53	43.18	6.25	0.00
Ca	167.6	163.5	36.26	-0.03	0.06
Mg	158.2	144.5	36.68	-0.02	0.03
Na	490.8	459.4	106.6	-0.11	0.02
K	22.18	24.40	11.60	0.00	0.43
Sol.AI	9.97	4.28	1.90	-10.17	0.00

	mean 4/1988-3/1993	mean 4/2003-3/2004	std.dev. 4/2003-3/2004	SK* 4/1988-3/2004	p* 4/1988-3/2004
Sol.lab.AI	8.26	2.76	1.49	-12.80	0.00
Cl	614.2	524.6	164.9	-0.32	0.00
SO_4	289.2	220.7	100.1	-0.41	0.00
XSO₄	224.7	165.6	106.6	-0.40	0.00
NO_3	7.39	7.72	7.63	0.00	0.67
Si	133.5	139.0	30.59	0.00	0.79
DOC	288.8	456.2	245.0	0.35	0.00


* Seasonal Kendall trend analysis: slope estimate (SK) and significance level (p)
Most units $\mu\text{eq l}^{-1}$, except Sol.AI, Sol.lab.AI and DOC ($\mu\text{mol l}^{-1}$)

13.2. Macroinvertebrate data

13.2.1. Percentage abundance summary, Old Lodge




13.2.2. Summary statistics, Old Lodge



13.3. Fish data


13.3.1. Summary of mean Trout density (numbers 100m⁻²), Old Lodge




NF = Not fished

13.4. Epilithic diatom data

13.4.1. Percentage abundance summary, Old Lodge




13.4.2. Summary statistics, Old Lodge



13.5. Aquatic macrophyte data, Old Lodge



Percentage Species Cover



+ Represents <0.1% abundance

14. Narrator Brook

14.1. Spot sampled chemistry data



Determinand statistics


	mean 4/1991-3/1996	mean 4/2003-3/2004	std.dev. 4/2003-3/2004	SK* 4/1991-3/2004	p* 4/1991-3/2004
pH	5.71	6.05	0.27	0.03	0.03
ANC	18.50	8.37	5.13	0.62	0.32
Ca	33.77	33.83	2.38	0.00	0.35
Mg	64.52	65.62	3.88	0.00	0.28
Na	254.5	267.4	91.72	-0.04	0.00
K	19.52	20.88	2.54	0.00	0.76
Sol.Al	2.09	2.31	1.25	1.50	0.00

	mean 4/1991-3/1996	mean 4/2003-3/2004	std.dev. 4/2003-3/2004	SK* 4/1991-3/2004	p* 4/1991-3/2004
Sol.lab.Al	0.93	0.93	0.48	0.50	0.16
Cl	275.7	248.1	12.04	-0.10	0.00
SO₄	73.59	73.09	4.12	0.00	0.37
XSO₄	44.63	47.03	4.01	0.02	0.04
NO₃	6.42	6.55	3.32	0.00	0.08
Si	151.3	187.0	17.80	0.02	0.04
DOC	125.3	164.2	82.88	0.06	0.00

* Seasonal Kendall trend analysis: slope estimate (SK) and significance level (p)
Most units μeq l⁻¹, except Sol.Al, Sol.lab.Al and DOC (μmol l⁻¹)


14.2. Macroinvertebrate data

14.2.1. Percentage abundance summary, Narrator Brook



No sampling in 2001 due to Foot and Mouth restrictions.


14.2.2. Summary statistics, Narrator Brook



No sampling in 2001 due to Foot and Mouth restrictions.

14.3. Fish data


14.3.1. Summary of mean Trout density (numbers 100m⁻²), Narrator Brook




NF = Not fished

14.4. Epilithic diatom data

14.4.1. Percentage abundance summary, Narrator Brook




14.4.2. Summary statistics, Narrator Brook



14.5. Aquatic macrophyte data, Narrator Brook



Percentage Species Cover



+ Represents <0.1% abundance

15. Llyn Llago

15.1. Spot sampled chemistry data



mean for first 5 years

mean for first 5 years

Determinand statistics


	mean 4/1988-3/1993	mean 4/2003-3/2004	std.dev. 4/2003-3/2004	SK* 4/1988-3/2004	p* 4/1988-3/2004
pH	5.23	5.79	0.21	0.05	0.00
ANC	5.86	17.75	3.92	1.52	0.00
Ca	57.06	41.12	7.27	-0.02	0.02
Mg	50.53	36.88	6.14	-0.01	0.07
Na	187.4	135.9	17.53	-0.06	0.10
K	3.60	2.82	1.09	0.00	0.37
Sol.Al	2.81	2.04	0.48	-0.16	0.67

	mean 4/1988-3/1993	mean 4/2003-3/2004	std.dev. 4/2003-3/2004	SK* 4/1988-3/2004	p* 4/1988-3/2004
Sol.lab.Al	1.58	0.24	0.13	-1.83	0.01
Cl	222.4	139.4	24.61	-0.13	0.08
SO_4	62.17	40.10	3.56	-0.10	0.00
XSO_4	38.82	25.46	3.17	-0.07	0.00
NO_3	10.42	4.05	3.34	0.00	0.05
Si	34.96	15.36	14.05	0.00	0.16
DOC	177.9	289.4	54.55	0.10	0.00


* Seasonal Kendall trend analysis: slope estimate (SK) and significance level (p)
Most units $\mu\text{eq l}^{-1}$, except Sol.Al, Sol.lab.Al and DOC ($\mu\text{mol l}^{-1}$)

15.2. Macroinvertebrate data

15.2.1. Percentage abundance summary, Llyn Llagi




15.2.2. Summary statistics, Llyn Llgi



15.3. Fish data (for outflow stream)


15.3.1. Summary of mean Trout density (numbers 100m⁻²), Llyn Llagi




NF = Not fished

15.4. Epilithic diatom data


15.4.1. Percentage abundance summary, Llyn Llago



15.4.2. Summary statistics, Llyn Llgi




15.5. Aquatic macrophyte data, Llyn Llagi





15.6. Sediment trap data, Llyn Llgi

Relative percentage frequency of diatom taxa and carbonaceous particle flux (no. cm⁻² yr⁻¹).



16. Llyn Cwm Mynach

16.1. Spot sampled chemistry data



Determinand statistics


	mean 4/1988-3/1993	mean 4/2003-3/2004	std.dev. 4/2003-3/2004	SK* 4/1988-3/2004	p* 4/1988-3/2004
pH	5.38	5.67	0.29	0.01	0.23
ANC	9.18	11.36	5.31	0.77	0.07
Ca	79.13	74.38	18.89	-0.02	0.29
Mg	68.86	64.38	11.71	-0.01	0.50
Na	294.5	265.2	54.88	-0.04	0.38
K	3.41	4.10	1.59	0.00	0.01
Sol.Al	3.94	2.71	1.99	1.25	0.28

	mean 4/1988-3/1993	mean 4/2003-3/2004	std.dev. 4/2003-3/2004	SK* 4/1988-3/2004	p* 4/1988-3/2004
Sol.lab.Al	2.42	0.55	0.58	-1.62	0.14
Cl	342.6	293.7	65.96	-0.10	0.39
SO_4	88.16	73.44	5.74	-0.07	0.01
XSO_4	52.17	42.60	7.14	-0.04	0.11
NO_3	9.59	10.12	10.65	0.00	0.35
Si	56.77	28.93	25.90	-0.01	0.01
DOC	208.8	200.0	65.68	0.05	0.09


* Seasonal Kendall trend analysis: slope estimate (SK) and significance level (p)
 Most units $\mu\text{eq l}^{-1}$, except Sol.Al, Sol.lab.Al and DOC ($\mu\text{mol l}^{-1}$)

16.2. Macroinvertebrate data

16.2.1. Percentage abundance summary, Llyn Cwm Mynach




16.2.2. Summary statistics, Llyn Cwm Mynach



16.3. Fish data (for outflow stream)


16.3.1. Summary of mean Trout density (numbers 100m⁻²), Llyn Cwm Mynach




NF = Not fished

16.4. Epilithic diatom data

16.4.1. Percentage abundance summary, Llyn Cwm Mynach




16.4.2. Summary statistics, Llyn Cwm Mynach




16.5. Aquatic macrophyte data, Llyn Cwm Mynach

Species Scores (1-5)





16.6. Sediment trap data, Llyn Cwm Mynach

Relative percentage frequency of diatom taxa and carbonaceous particle flux (no. cm⁻² yr⁻¹).



17. Afon Hafren

17.1. Spot sampled chemistry data



Determinand statistics


	mean 4/1988-3/1993	mean 4/2003-3/2004	std.dev. 4/2003-3/2004	SK* 4/1988-3/2004	p* 4/1988-3/2004
pH	5.27	5.36	0.49	0.01	0.47
ANC	-2.57	7.50	6.25	0.93	0.17
Ca	48.08	43.88	10.47	-0.01	0.02
Mg	67.32	68.40	9.73	0.00	0.36
Na	201.0	195.3	17.63	-0.02	0.12
K	3.20	4.94	2.48	0.00	0.00
Sol.AI	6.38	5.83	3.44	1.59	0.52

	mean 4/1988-3/1993	mean 4/2003-3/2004	std.dev. 4/2003-3/2004	SK* 4/1988-3/2004	p* 4/1988-3/2004
Sol.lab.AI	3.83	2.21	2.00	-1.00	0.47
Cl	221.8	199.5	16.64	-0.06	0.10
SO_4	82.70	66.84	6.79	-0.06	0.00
XSO_4	59.41	45.88	5.90	-0.06	0.00
NO_3	20.45	25.17	8.68	0.00	0.10
Si	113.8	106.0	29.48	-0.02	0.02
DOC	149.3	223.0	78.20	0.10	0.00

* Seasonal Kendall trend analysis: slope estimate (SK) and significance level (p)
Most units $\mu\text{eq l}^{-1}$, except Sol.AI, Sol.lab.AI and DOC ($\mu\text{mol l}^{-1}$)


17.2. Macroinvertebrate data

17.2.1. Percentage abundance summary, Afon Hafren



No sampling in 2001 due to Foot and Mouth restrictions.


17.2.2. Summary statistics, Afon Hafren



No sampling in 2001 due to Foot and Mouth restrictions.

17.3. Fish data


17.3.1. Summary of mean Trout density (numbers 100m⁻²), Afon Hafren




NF = Not fished

17.4. Epilithic diatom data

17.4.1. Percentage abundance summary, Afon Hafren




17.4.2. Summary statistics, Afon Hafren



17.5. Aquatic macrophyte data, Afon Hafren



Percentage Species Cover



+ Represents <0.1% abundance

18. Afon Gwy

18.1. Spot sampled chemistry data



Determinand statistics


	mean 4/1991-3/1996	mean 4/2003-3/2004	std.dev. 4/2003-3/2004	SK* 4/1991-3/2004	p* 4/1991-3/2004
pH	5.51	5.69	0.43	0.02	0.06
ANC	14.00	10.32	4.25	0.96	0.09
Ca	40.57	39.96	11.57	-0.01	0.29
Mg	54.08	56.60	10.62	0.00	0.63
Na	147.5	155.4	19.10	0.00	0.94
K	3.24	3.38	2.11	0.00	0.38
Sol.Al	3.93	3.10	1.64	-1.00	0.50

	mean 4/1991-3/1996	mean 4/2003-3/2004	std.dev. 4/2003-3/2004	SK* 4/1991-3/2004	p* 4/1991-3/2004
Sol.lab.Al	1.98	0.90	0.85	-2.00	0.04
Cl	159.7	158.5	22.22	-0.03	0.42
SO_4	65.90	56.60	9.52	-0.08	0.00
XSO_4	49.13	39.96	8.16	-0.07	0.00
NO_3	8.87	6.46	5.08	0.00	0.46
Si	69.88	66.61	22.11	0.00	0.45
DOC	163.5	220.5	68.44	0.06	0.01

* Seasonal Kendall trend analysis: slope estimate (SK) and significance level (p)
Most units $\mu\text{eq l}^{-1}$, except Sol.Al, Sol.lab.Al and DOC ($\mu\text{mol l}^{-1}$)


18.2. Macroinvertebrate data

18.2.1. Percentage abundance summary, Afon Gwy



No sampling in 2001 due to Foot and Mouth restrictions.


18.2.2. Summary statistics, Afon Gwy



No sampling in 2001 due to Foot and Mouth restrictions.

18.3. Fish data


18.3.1. Summary of mean Trout density (numbers 100m⁻²), Afon Gwy




NF = Not fished

18.4. Epilithic diatom data

18.4.1. Percentage abundance summary, Afon Gwy




18.4.2. Summary statistics, Afon Gwy





18.5. Aquatic macrophyte data, Afon Gwy

Percentage Species Cover



19. Beaghs Burn

19.1. Spot sampled chemistry data



Determinand statistics


	mean 4/1988-3/1993	mean 4/2003-3/2004	std.dev. 4/2003-3/2004	SK* 4/1988-3/2004	p* 4/1988-3/2004
pH	5.76	5.86	0.88	0.00	1.00
ANC	86.98	131.8	104.1	2.48	0.05
Ca	103.2	117.3	55.00	-0.02	0.36
Mg	113.9	119.4	38.49	-0.01	0.20
Na	306.7	308.7	54.84	-0.05	0.11
K	11.31	12.26	3.29	0.00	0.62
Sol.Al	2.14	1.31	0.82	-1.00	0.10

	mean 4/1988-3/1993	mean 4/2003-3/2004	std.dev. 4/2003-3/2004	SK* 4/1988-3/2004	p* 4/1988-3/2004
Sol.lab.Al	0.44	0.25	0.26	0.00	0.53
Cl	350.5	336.6	68.99	-0.10	0.10
SO_4	73.68	55.38	30.30	-0.10	0.00
XSO_4	37.39	20.03	27.60	-0.08	0.01
NO_3	3.11	2.21	0.00	0.00	0.78
Si	68.92	67.26	39.41	0.00	0.41
DOC	783.8	1182.6	533.4	0.42	0.00


* Seasonal Kendall trend analysis: slope estimate (SK) and significance level (p)
Most units $\mu\text{eq l}^{-1}$, except Sol.Al, Sol.lab.Al and DOC ($\mu\text{mol l}^{-1}$)

19.2. Macroinvertebrate data

19.2.1. Percentage abundance summary, Beaghs Burn




19.2.2. Summary statistics, Beaghs Burn




19.3. Fish data

19.3.1. Summary of mean Trout density (numbers 100m⁻²), Beaghs Burn




19.4. Epilithic diatom data

19.4.1. Percentage abundance summary, Beaghs Burn




19.4.2. Summary statistics, Beaghs Burn



19.5. Aquatic macrophyte data, Beaghs Burn



Percentage Species Cover



+ Represents <math><0.1\%</math> abundance

20. Bencrom River

20.1. Spot sampled chemistry data



Determinand statistics


	mean 4/1988-3/1993	mean 4/2003-3/2004	std.dev. 4/2003-3/2004	SK* 4/1988-3/2004	p* 4/1988-3/2004
pH	5.20	5.73	0.82	0.03	0.02
ANC	-3.90	7.44	80.42	2.39	0.00
Ca	52.21	75.12	15.49	0.00	0.97
Mg	62.05	57.64	6.61	-0.01	0.08
Na	259.7	241.3	22.36	-0.04	0.11
K	11.66	11.88	2.05	0.00	0.16
Sol.AI	7.40	4.75	2.30	-6.12	0.02

	mean 4/1988-3/1993	mean 4/2003-3/2004	std.dev. 4/2003-3/2004	SK* 4/1988-3/2004	p* 4/1988-3/2004
Sol.lab.AI	4.43	2.91	2.09	-4.18	0.01
Cl	252.3	211.0	24.95	-0.07	0.11
SO₄	94.96	76.91	10.34	-0.10	0.00
XSO₄	68.45	54.75	9.57	-0.08	0.00
NO₃	26.86	68.21	79.75	0.00	0.66
Si	214.7	223.8	56.68	-0.01	0.51
DOC	287.7	261.1	82.75	0.06	0.12


* Seasonal Kendall trend analysis: slope estimate (SK) and significance level (p)
Most units $\mu\text{eq l}^{-1}$, except Sol.AI, Sol.lab.AI and DOC ($\mu\text{mol l}^{-1}$)

20.2. Macroinvertebrate data

20.2.1. Percentage abundance summary, Bencrom River




20.2.2. Summary statistics, Bencrom River




20.3. Fish data

20.3.1. Summary of mean Trout density (numbers 100m⁻²), Bencrom River




20.4. Epilithic diatom data

20.4.1. Percentage abundance summary, Bencrom River




20.4.2. Summary statistics, Bencrom River



20.5. Aquatic macrophyte data, Bencrom River

Percentage Species Cover





+ Represents $<0.1\%$ abundance

No survey undertaken in 2002 due to spate conditions.

21. Blue Lough

21.1. Spot sampled chemistry data



Determinand statistics


	mean 4/1990-3/1995	mean 4/2003-3/2004	std.dev. 4/2003-3/2004	SK* 4/1990-3/2004	p* 4/1990-3/2004
pH	4.66	4.89	0.15	0.01	0.00
ANC	-44.87	-12.74	6.69	2.88	0.00
Ca	40.74	30.38	2.66	-0.02	0.01
Mg	57.42	43.75	4.17	-0.01	0.05
Na	245.0	216.3	22.83	-0.05	0.20
K	11.38	10.06	1.83	0.00	0.80
Sol.AI	14.13	9.31	0.96	-12.14	0.00

	mean 4/1990-3/1995	mean 4/2003-3/2004	std.dev. 4/2003-3/2004	SK* 4/1990-3/2004	p* 4/1990-3/2004
Sol.lab.AI	11.63	6.19	0.38	-12.38	0.00
Cl	265.2	214.1	25.61	-0.10	0.34
SO_4	95.62	63.02	5.98	-0.17	0.00
XSO_4	67.77	40.53	3.99	-0.15	0.00
NO_3	24.69	21.25	7.52	0.00	0.56
Si	71.25	55.36	36.60	0.00	0.95
DOC	265.8	343.8	12.50	0.12	0.00


* Seasonal Kendall trend analysis: slope estimate (SK) and significance level (p)
Most units $\mu\text{eq l}^{-1}$, except Sol.AI, Sol.lab.AI and DOC ($\mu\text{mol l}^{-1}$)

21.2. Macroinvertebrate data

21.2.1. Percentage abundance summary, Blue Lough




21.2.2. Summary statistics, Blue Lough



21.3. Fish data (for outflow stream)


21.3.1. Summary of mean Trout density (numbers 100m⁻²), Blue Lough




NF = Not fished

21.4. Epilithic diatom data

21.4.1. Percentage abundance summary, Blue Lough




21.4.2. Summary statistics, Blue Lough




21.5. Aquatic macrophyte data, Blue Lough

Species Scores (1-5)





21.6. Sediment trap data, Blue Lough

Relative percentage frequency of diatom taxa and carbonaceous particle flux (no. cm⁻² yr⁻¹).



22. Coneyglen Burn

22.1. Spot sampled chemistry data



Determinand statistics


	mean 4/1990-3/1995	mean 4/2003-3/2004	std.dev. 4/2003-3/2004	SK* 4/1990-3/2004	p* 4/1990-3/2004
pH	6.53	6.29	0.79	-0.02	0.08
ANC	193.9	192.3	150.2	-1.07	0.56
Ca	146.8	144.0	76.99	-0.02	0.32
Mg	118.7	116.0	42.86	-0.01	0.29
Na	241.1	217.8	37.51	-0.07	0.02
K	8.91	8.14	3.01	0.00	0.97
Sol.AI	1.35	1.12	0.53	0.05	0.91

	mean 4/1990-3/1995	mean 4/2003-3/2004	std.dev. 4/2003-3/2004	SK* 4/1990-3/2004	p* 4/1990-3/2004
Sol.lab.AI	0.22	0.21	0.29	0.00	0.21
Cl	261.5	231.5	68.52	-0.12	0.06
SO_4	52.05	46.53	13.60	-0.06	0.03
XSO_4	24.58	22.22	13.21	-0.04	0.08
NO_3	1.56	7.74	18.28	0.00	0.04
Si	89.29	83.33	59.71	-0.02	0.12
DOC	597.8	946.5	457.9	0.46	0.01

* Seasonal Kendall trend analysis: slope estimate (SK) and significance level (p)
 Most units $\mu\text{eq l}^{-1}$, except Sol.AI, Sol.lab.AI and DOC ($\mu\text{mol l}^{-1}$)


22.2. Macroinvertebrate data

22.2.1. Percentage abundance summary, Coneyglen Burn



No sampling in 2001 due to Foot and Mouth restrictions.


22.2.2. Summary statistics, Coneyglen Burn



No sampling in 2001 due to Foot and Mouth restrictions.

22.3. Fish data


22.3.1. Summary of mean Trout density (numbers 100m⁻²), Coneyglen Burn




NF = Not fished

22.4. Epilithic diatom data

22.4.1. Percentage abundance summary, Coneyglen Burn




22.4.2. Summary statistics, Coneyglen Burn



22.5. Aquatic macrophyte data, Coneyglen Burn


Percentage Species Cover



+ Represents <0.1% abundance

23. Loch Coire Fionnaraich

23.1. Spot sampled chemistry data




Determinand statistics

	mean 4/2003-3/2004	std.dev. 4/2003-3/2004		mean 4/2003-3/2004	std.dev. 4/2003-3/2004
pH	5.89	0.43	Sol.lab.AI	0.10	0.06
ANC	39.47	25.47	Cl	190.14	67.08
Ca	30.81	12.68	SO ₄	28.63	8.04
Mg	43.70	14.76	XSO ₄	8.68	4.48
Na	178.34	46.93	NO ₃	2.24	0.00
K	6.65	1.60	Si	24.21	10.02
Sol.AI	2.06	1.22	DOC	456.67	302.71


Most units $\mu\text{eq l}^{-1}$, except Sol.AI, Sol.lab.AI and DOC ($\mu\text{mol l}^{-1}$)

23.2. Macroinvertebrate data

23.2.1. Percentage abundance summary, Loch Coire Fionnaraich




23.2.2. Summary statistics, Loch Coire Fionnaraich



23.3. Fish data (for outflow stream)


23.3.1. Summary of mean Trout density (numbers 100m⁻²), Loch Coire Fionnaraich




NF = Not fished

23.4. Epilithic diatom data

23.4.1. Percentage abundance summary, Loch Coire Fionnaraich



23.4.2. Summary statistics, Loch Coire Fionnaraich



23.5. Aquatic macrophyte data, Loch Coire Fionnaraich

Species Scores (1-5)

Species	2003 Score
<i>Sphagnum auriculatum</i>	2
<i>Scapania undulata</i>	2
<i>Fontinalis antipyretica</i>	2
<i>Isoetes lacustris</i>	4
<i>Lobelia dortmanna</i>	4
<i>Littorella uniflora</i>	4
<i>Myriophyllum alterniflorum</i>	2
<i>Callitriche hamulata</i>	2
<i>Subularia aquatica</i>	3
<i>Juncus effusus</i>	2
<i>Carex nigra</i>	3
<i>Ranunculus flammula</i>	3
<i>Juncus bulbosus</i>	3
<i>Potamogeton polygonifolius</i>	1
<i>Sparganium angustifolium</i>	2
<i>Nitella flexilis</i> agg.	2
<i>Batrachospermum</i> sp.	3
Filamentous algae	2

23.6. Sediment trap data, Loch Coire Fionnaraich

Relative percentage frequency of diatom taxa and carbonaceous particle flux
(no. cm⁻² yr⁻¹).

