

Annexes

ANNEX 1: Key Categories	577
A 1.1 GENERAL APPROACH USED TO IDENTIFY KEY CATEGORIES	578
A 1.2 QUALITATIVE ANALYSIS USED TO IDENTIFY KEY CATEGORIES	578
A 1.3 QUANTITATIVE APPROACH 1 KCA FOLLOWING IPCC 2006 GUIDELINES	579
A 1.4 QUANTITATIVE APPROACH 2 KCA FOLLOWING IPCC 2006 GUIDELINES	616
A 1.5 KEY CATEGORY ANALYSIS (KCA) RANKING SYSTEM	639
A 1.6 APPROACH USED TO IDENTIFY KP-LULUCF KEY CATEGORIES	641
A 1.7 USING THE UNCERTAINTY ANALYSIS TO PLAN IMPROVEMENTS IN THE PREPARATION OF THE INVENTORY	641
A 1.8 TABLE NIR 3, AS CONTAINED IN THE ANNEX TO DECISION 6/CMP.3	642
ANNEX 2: Assessment of Uncertainty	645
A 2.1 ESTIMATION OF UNCERTAINTIES USING AN ERROR PROPAGATION APPROACH (APPROACH 1)	645
A 2.1.1 Previous Improvements to the Error Propagation Model	646
A 2.1.2 Review of Changes Made to the Error Propagation Model since the last NIR	646
A 2.1.3 Key Categories	646
A 2.1.4 Tables of uncertainty estimates from the error propagation approach	648
A 2.2 ESTIMATION OF UNCERTAINTY BY SIMULATION (APPROACH 2)	656
A 2.2.1 Overview of the Method	656
A 2.2.2 Summary of Recent Improvements to the Monte Carlo Model	657
A 2.2.3 Review of changes made to the Monte Carlo model since the last NIR	657
A 2.2.4 Methodological details of the Monte Carlo model	658
A 2.2.5 Quality Control Checks on the Monte Carlo Model Output	660
A 2.3 UNCERTAINTIES ACCORDING TO GAS	660
A 2.3.1 Carbon Dioxide Emission Uncertainties	660
A 2.3.2 Methane Emission Uncertainties	676
A 2.3.3 Nitrous Oxide Emission Uncertainties	677

	A 2.3.4 Halocarbons, NF ₃ and SF ₆	678
A 2.4	UNCERTAINTIES IN GWP WEIGHTED EMISSIONS	678
	A 2.4.1 Uncertainty in the emissions	678
	A 2.4.2 Uncertainty in the Trend	678
A 2.5	SECTORAL UNCERTAINTIES	680
	A 2.5.1 Overview of the Method	680
	A 2.5.2 Review of Changes made to the Monte Carlo Model since the last NIR	680
A 2.6	COMPARISON OF UNCERTAINTIES FROM THE ERROR PROPAGATION AND MONTE CARLO ANALYSES	684

ANNEX 3: Other Detailed Methodological Descriptions for Individual Source or Sink Categories, Including for KP-LULUCF Activities. 687

A 3.1	ENERGY 687	
	A 3.1.1 Emission factors	688
	A 3.1.2 Commonly used references	746
	A 3.1.3 Feedstocks and NEU of fuels	753
A 3.2	INDUSTRIAL PROCESSES (CRF SECTOR 2)	764
	A 3.2.1 Potential Emissions of Halocarbons and SF ₆	764
A 3.3	AGRICULTURE (CRF SECTOR 3)	764
	A 3.3.1 Enteric Fermentation (3A)	764
	A 3.3.2 Manure Management (3B)	774
	A 3.3.3 Agricultural Soils (3D)	780
	A 3.3.4 Field Burning of Agricultural Residues (3F)	787
A 3.4	LAND USE, LAND USE CHANGE AND FORESTRY (CRF SECTOR 4)	788
	A 3.4.1 Carbon stock changes due to afforestation and forest management (4A)	788
	A 3.4.2 Land Use Change and Soils (4B, 4C, 4E)	798
	A 3.4.3 Changes in stocks of carbon in non-forest biomass due to land use change (4B2, 4C2, 4E2)	808
	A 3.4.4 Carbon stock changes and biomass burning emissions due to Deforestation (4B, 4C, 4E, 4G)	811
	A 3.4.5 Biomass Burning – Forest and Non-Forest Wildfires (4A, 4B, 4C)	817

Contents of Annexes

A 3.4.6	Emissions from drainage (organic soils) (4B1, 4C1)	820
A 3.4.7	Changes in Stocks of Carbon in Non-Forest Biomass due to Yield Improvements (4B1)	821
A 3.4.8	Emissions of N ₂ O due to disturbance associated with land use conversion and land management changes (4(III))	821
A 3.4.9	On-site and off-site emissions from peat extraction (4D)	822
A 3.4.10	Flooded Lands (4D)	826
A 3.4.11	Harvested Wood Products (4G)	827
A 3.4.12	Methods for the Overseas Territories (OTs) and Crown Dependencies (CDs)	828
A 3.4.13	Uncertainty analysis of the LULUCF sector	832
A 3.5	WASTE (CRF SECTOR 5)	834
A 3.5.1	Solid Waste Disposal on Land (5A)	834
A 3.5.2	Waste Incineration (5C)	841
A 3.5.3	Wastewater Handling (5D)	844
A 3.6	DATA FOR THE UK'S CROWN DEPENDENCIES AND OVERSEAS TERRITORIES	848
ANNEX 4: National Energy Balance for the Most Recent Inventory Year		853
A 4.1	UK ENERGY BALANCE	853
A 4.2	FUELS DATA	859
A 4.2.1	Reallocations of energy data and differences from UK energy statistics	860
ANNEX 5: Additional Information to be Considered as Part of the Annual Inventory Submission and the Supplementary Information Required Under Article 7, paragraph 1, of the Kyoto Protocol Other Useful Reference Information.		873
A 5.1	ANNUAL INVENTORY SUBMISSION	873
A 5.2	SUPPLEMENTARY INFORMATION UNDER ARTICLE 7, PARAGRAPH 1	873
ANNEX 6: Verification		875
A 6.1	MODELLING APPROACH USED FOR THE VERIFICATION OF THE UK GHGI	875
A 6.2	METHANE	875
A 6.3	NITROUS OXIDE	877

A 6.4	HYDROFLUOROCARBONS	878
	A 6.4.1 HFC-134a	878
	A 6.4.2 HFC-152a	880
	A 6.4.3 HFC-125	881
	A 6.4.4 HFC-143a	882
	A 6.4.5 HFC-23	883
	A 6.4.6 HFC-32	884
	A 6.4.7 HFC-43-10mee	885
	A 6.4.8 HFC-227ea	886
	A 6.4.9 HFC-365mfc	887
	A 6.4.10 HFC-245fa	888
A 6.5	PERFLUOROCARBONS	889
	A 6.5.1 PFC-14	889
	A 6.5.2 PFC-116	890
	A 6.5.3 PFC-218	891
	A 6.5.4 PFC-318	892
A 6.6	SULPHUR HEXAFLUORIDE	893
A 6.7	NITROGEN TRIFLUORIDE	894
A 6.8	CARBON DIOXIDE	895
ANNEX 7: Analysis of EU ETS Data		899
A 7.1	INTRODUCTION	899
A 7.2	BACKGROUND	902
	A 7.2.1 EU ETS Data and GHG Inventories	902
	A 7.2.2 Scope of the UK EU ETS and Implications for the GHG Inventory	903
	A 7.2.3 Limitations of EU ETS Data Integration with GHG Inventory: Autogeneration	905
A 7.3	DATA PROCESSING	906
A 7.4	EU ETS DATA COVERAGE	907
A 7.5	EU ETS DATA USE IN THE UK GHGI	909
	A 7.5.1 Activity and Emissions Data	910

	A 7.5.2 Implied Emission Factors	914
ANNEX 8: UK Domestic Emissions Reporting Requirements		925
A 8.1	NATIONAL STATISTICS	925
A 8.2	UK ONLY EMISSIONS STATISTICS	926
ANNEX 9: End User Emissions		929
A 9.1	INTRODUCTION	929
A 9.2	DEFINITION OF FINAL USERS	929
A 9.3	OVERVIEW OF THE FINAL USER CALCULATIONS	930
A 9.4	EXAMPLE FINAL USER CALCULATION	932
A 9.5	FINAL USER CALCULATION METHODOLOGY FOR THE UK GREENHOUSE GAS INVENTORY	936
A 9.6	DETAILED EMISSIONS ACCORDING TO FINAL USER CATEGORIES	964
 Table of Tables:		
Table A 1.2.1	Qualitative Key Category Analysis.....	579
Table A 1.3.1	Approach 1 Key Category Analysis for the base year based on level of emissions (including LULUCF)	580
Table A 1.3.2	Approach 1 Key Category Analysis for the base year based on level of emissions (excluding LULUCF)	586
Table A 1.3.3	Approach 1 Key Category Analysis for the latest reported year based on level of emissions (including LULUCF)	591
Table A 1.3.4	Approach 1 Key Category Analysis for the latest reported year based on level of emissions (excluding LULUCF).....	597
Table A 1.3.5	Approach 1 Key Category Analysis based on trend in emissions (from base year to latest reported year, including LULUCF)	603
Table A 1.3.6	Approach 1 Key Category Analysis based on trend in emissions (from base year to latest reported year, excluding LULUCF)	610
Table A 1.4.1	Approach 2 Level Assessment for Base year (including LULUCF) with Key Categories Shaded in Grey	617
Table A 1.4.2	Approach 2 Level Assessment for 2013 (including LULUCF) with Key Categories Shaded in Grey	621

Contents of Annexes

Table A 1.4.3	Approach 2 Level Assessment for Base year (not including LULUCF) with Key Categories Shaded in Grey.....	625
Table A 1.4.4	Approach 2 Level Assessment for 2013 (not including LULUCF) with Key Categories Shaded in Grey.....	628
Table A 1.4.5	Approach 2 Assessment for Trend (including LULUCF) with Key Categories Shaded in Grey.....	631
Table A 1.4.6	Approach 2 Assessment for Trend (not including LULUCF) with Key Categories Shaded in Grey.....	635
Table A 1.5.1	KCA Ranking.....	639
Table A 1.8.1	Table NIR 3. Summary overview for Key Categories for Land Use, Land-Use Change and Forestry Activities under the Kyoto Protocol.....	642
Table A 2.1.1	Summary of error propagation uncertainty estimates including LULUCF, base year to the latest reported year.....	648
Table A 2.3.1	Uncertainties in the activity data and emission factors for fuels used in the carbon dioxide (CO ₂) inventory.....	662
Table A 2.3.2	Estimated uncertainties in the activity data and emission factors used in the methane (CH ₄) inventory.....	668
Table A 2.3.3	Estimated uncertainties in the activity data and emission factors used in the nitrous oxide (N ₂ O) inventory.....	671
Table A 2.3.4	Estimated uncertainties in the activity data and emission factors used in the F-gas inventory.....	674
Table A 2.4.1	Summary of Monte Carlo Uncertainty Estimates.....	679
Table A 2.5.1	Sectoral Uncertainty Estimates.....	681
Table A 2.6.1	Comparison of the central estimates and trends in emissions from the error propagation (Approach 1) and Monte Carlo (Approach 2) uncertainty analyses.....	684
Table A 3.1.1	Emission factors used for the 2015 submission (sector 1A).....	688
Table A 3.1.2	Emission factors used for the 2015 submission (sector 1B).....	741
Table A 3.1.3	Approaches of estimating activity and emission factors.....	750
Table A 3.1.4	Summary of Emission Sources for UK Fuels Allocated as Non Energy Use in UK Energy Statistics.....	754
Table A 3.1.5	Activity data and methane leakage estimates for Gas leakage at Point of Use, including cooking appliances, gas fires and boilers.....	760
Table A 3.1.6	Petroleum Coke Carbon Emissions and Carbon Stored across all UK Supply 1990-2012.....	763

Contents of Annexes

Table A 3.3.1	Livestock Population Data for 2013 by Animal Type ^a	764
Table A 3.3.2	Trends in Livestock Numbers ('000s) 1990-2013.....	765
Table A 3.3.3	Methane Emission Factors for Livestock Emissions for 2013	766
Table A 3.3.4	Dairy Cows Tier 2 Methane Emission Factors ^a	767
Table A 3.3.5	Energy and digestibility coefficients of ruminant feeds (dairy cows) (MAFF, 1990)	768
Table A 3.3.6	Calculation of digestibility (dairy cows) ^a	770
Table A 3.3.7	Parameters used in the calculations of gross energy for dairy cows Tier 2 emission factors	771
Table A 3.3.8	Beef Cows Tier 2 Methane Emission Factors ^a	772
Table A 3.3.9	Parameters used in the calculation of gross energy for beef cows Tier 2 emission factors	773
Table A 3.3.10	Methane conversion factors for Manure Management Systems in the UK	774
Table A 3.3.11	Nitrogen Excretion Factors, kg N hd ⁻¹ year ⁻¹ for livestock in the UK (1990-2013) ^a	776
Table A 3.3.12	Distribution of Animal Waste Management Systems used for Different Animal types, 2013 ^a	777
Table A 3.3.13	Nitrous Oxide Emission Factors for Animal Waste Handling Systems ^a	780
Table A 3.3.14	Areas of UK Crops and quantities of fertiliser applied for 2013 ^a	780
Table A 3.3.15	Trends in area grown ('000 ha) and N fertiliser applied (kg/ha) for the major UK crops, 1990-2013 ^a	782
Table A 3.3.16	Nitrous oxide emissions from sewage sludge (kt N ₂ O/yr) ^a	784
Table A 3.3.17	Dry Mass Content and Residue Fraction of UK Crops for 2013	784
Table A 3.3.18	Production of UK Crops for 2013 ^a	785
Table A 3.3.19	Mineralised N from soils	786
Table A 3.3.20	Constants for crop residue calculations (Tables 2.5, 2.6 & p.11.14 2006GL)	787
Table A 3.3.21	CO ₂ emissions from the application of lime.....	787
Table A 3.3.22	CO ₂ emissions from the application of urea	788
Table A 3.4.1	Main parameters for forest carbon flow model used to estimate carbon uptake by planting of forests of Sitka spruce (<i>P. Sitchensis</i>), yield class 12.	789
Table A 3.4.2	Forest area by management prescription in the UK (from 1990-2012 inventory)	792
Table A 3.4.3	Forest area by rotation length for forests managed for production (from 1990- 2012 inventory).....	792

Contents of Annexes

Table A 3.4.4	Afforestation rate of conifers and broadleaves in the United Kingdom since 1500 (using pre-1920 estimates from woodland age and post-1921 planting data)	794
Table A 3.4.5	Grouping of MLC land cover types for soil carbon change modelling	798
Table A 3.4.6	Grouping of Countryside Survey Broad Habitat types (Jackson, 2000) for soil carbon change modelling.....	799
Table A 3.4.7	Sources of land use change data in Great Britain for different periods in estimation of changes in soil carbon.....	800
Table A 3.4.8	Sources of land use change data in Northern Ireland for different periods in estimation of changes in soil carbon.	800
Table A 3.4.9	Soil carbon stock (TgC = MtC) for depths to 1 m in different land types in the UK	802
Table A 3.4.10	Weighted average change in equilibrium soil carbon density ($t\ ha^{-1}$) to 1 m deep for changes between different land types in England	803
Table A 3.4.11	Weighted average change in equilibrium soil carbon density ($t\ ha^{-1}$) to 1 m deep for changes between different land types in Scotland	803
Table A 3.4.12	Weighted average change in equilibrium soil carbon density ($t\ ha^{-1}$) to 1 m deep for changes between different land types in Wales.....	804
Table A 3.4.13	Weighted average change in equilibrium soil carbon density ($t\ ha^{-1}$) to 1 m deep for changes between different land types in Northern Ireland.....	804
Table A 3.4.14	Rates of change of soil carbon for land use change transitions.	804
Table A 3.4.15	Range of times for soil carbon to reach 99% of a new value after a change in land use in England (E), Scotland (S) and Wales (W)	805
Table A 3.4.16	Equilibrium biomass carbon density ($kg\ m^{-2}$) for different land types	809
Table A 3.4.17	Weighted average change in equilibrium biomass carbon density ($kg\ m^{-2}$) for changes between different land types in England.....	809
Table A 3.4.18	Weighted average change in equilibrium biomass carbon density ($kg\ m^{-2}$) for changes between different land types in Scotland.....	810
Table A 3.4.19	Weighted average change in equilibrium biomass carbon density ($kg\ m^{-2}$) for changes between different land types in Wales.....	810
Table A 3.4.20	Weighted average change in equilibrium biomass carbon density ($kg\ m^{-2}$) for changes between different land types in Northern Ireland.	810
Table A 3.4.21	Countryside Survey data for Forest conversion.....	813
Table A 3.4.22	Corrected Forest conversion rates.....	813
Table A 3.4.23	IRS database property type descriptions by LULUCF sub-category.....	818

Contents of Annexes

Table A 3.4.24	Activity data for peat extraction sites in Northern Ireland.....	822
Table A 3.4.25	Activity data for peat extraction sites in England, Scotland and Wales	824
Table A 3.4.26	Annual peat production, m ³ for England and Scotland (from Annual Minerals Raised Inquiry/Mineral Extraction in Great Britain reports)	824
Table A 3.4.27	Information sources for estimating LULUCF emissions from the Overseas Territories and Crown Dependencies	828
Table A 3.4.28	Assumptions used in applying the Tier 1 methodology to the Overseas Territories and Crown Dependencies	829
Table A 3.4.29	Tier 1 factors used for estimating LULUCF emissions from Overseas Territories and Crown Dependencies	832
Table A 3.4.30	Combined uncertainties for the LULUCF sector 1990-2013 (rounded to 5%).....	833
Table A 3.5.1	Waste composition data	836
Table A 3.5.2	Amount of waste landfilled and methane generated, captured, utilised, flared, oxidised and emitted	837
Table A 3.5.3	Parameters used in landfill emission estimates for overseas territories and crown dependencies.....	839
Table A 3.5.4	Activity Data: UK Waste Incineration 1990-2013	841
Table A 3.5.5	Emissions Data: UK Waste Incineration 1990-2013.....	842
Table A 3.5.6	UK Domestic and Commercial Waste Water Treatment (5D1) Activity Data.....	844
Table A 3.5.7	UK Domestic and Commercial Waste Water Treatment (5D1) Implied Emission Factors.....	845
Table A 3.5.8	UK Domestic and Commercial Waste Water Treatment (5D1) Emission Estimates (kt CH ₄)	846
Table A 3.5.9	UK Industrial Waste Water Treatment Activity Data (5D2) (1990-2013)	846
Table A 3.6.1	Isle of Man, Guernsey and Jersey – Emissions of Direct GHGs (Mt CO ₂ equivalent)	848
Table A 3.6.2	Isle of Man, Guernsey and Jersey – Fuel use data	848
Table A 3.6.3	Isle of Man, Guernsey and Jersey – Animal numbers.....	848
Table A 3.6.4	Isle of Man, Guernsey and Jersey – Total emissions from Agricultural Soils (kg N ₂ O-N).....	849
Table A 3.6.5	Isle of Man, Guernsey and Jersey – Amount of synthetic fertilizer applied	849
Table A 3.6.6	Cayman Islands, Falklands Islands, Bermuda and Montserrat – Emissions of Direct GHGs (Mt CO ₂ equivalent)	849

Contents of Annexes

Table A 3.6.7	Cayman Islands, Falklands Islands, Bermuda and Montserrat – Fuel use data	850
Table A 3.6.8	Cayman Islands, Falklands Islands, Bermuda and Montserrat – Animal numbers	850
Table A 3.6.9	Cayman Islands, Falklands Islands, Bermuda and Montserrat – Total emissions from Agricultural Soils (kg N ₂ O-N)	850
Table A 3.6.10	Cayman Islands, Falklands Islands, Bermuda and Montserrat - Amount of synthetic fertilizer applied	851
Table A 3.6.11	Cayman Islands, Falklands Islands, Bermuda and Montserrat – Production of non-N-fixing crops (tonnes).....	851
Table A 3.6.12	Montserrat – Production of N-fixing crops (tonnes)	851
Table A 3.6.13	Gibraltar – Emissions of Direct GHGs (Mt CO ₂ equivalent).....	851
Table A 3.6.14	Gibraltar – Fuel use data	851
Table A 4.1.1	UK Energy Balance (thousand tonnes of oil equivalent, gross energy basis)	854
Table A 4.2.1	Mapping of fuels used in IPCC and the NAEI	859
Table A 4.2.2	Fuel reconciliation - coal use in the latest year (Mtonnes)	861
Table A 4.2.3	Fuel reconciliation – natural gas use in 2013 (Mtherms)	863
Table A 4.2.4	Fuel reconciliation – Fuel oils use in 2013 (Mtonnes)	865
Table A 4.2.5	Fuel reconciliation – Gas oil use in 2013 (Mtonnes).....	867
Table A 4.2.6	Fuel reconciliation – Use of petroleum gases in 2013 (Mtherms).....	870
Table A 7.1.1	Summary of the use of EU ETS data in the UK inventory	901
Table A 7.4.1	Numbers of installations included in the EU ETS data.....	908
Table A 7.5.1	Refinery Emissions Data Comparison and Revision to OPG Activity	911
Table A 7.5.2	EU ETS data for Fuels used at Power Stations and Autogenerators (Emission Factors in kt / Mt for Coal & Fuel Oil, kt / Mth for Gases).....	914
Table A 7.5.3	Refinery EU ETS Data for Fuel Oil, OPG and Natural Gas (Emission Factors in kt / Mt for Fuel Oil and kt / Mth for OPG and Natural Gas).....	917
Table A 7.5.4	EU ETS data for Fuels used at Integrated Steelworks and Coke Ovens (Emission Factors in kt / Mt for Solid & Liquid Fuels, kt / Mth for Gases)	918
Table A 7.5.5	EU ETS data for Fuels used at Cement Kilns (kt / Mt).....	920
Table A 7.5.6	Comparison of Cement Sector Carbon Dioxide Emissions* within the UK GHGI and the EU ETS for 2008-2012	921
Table A 7.5.7	EU ETS data for Fuels used at Lime Kilns (Emission Factors in kt / Mt for Solid Fuels and kt / Mth for Gases).....	921

Contents of Annexes

Table A 7.5.8	EU ETS emission factor data for production of lime (kt / Mt lime produced)	922
Table A 7.5.9	EU ETS data for Coal, Fuel Oil and Natural Gas used by Industrial Combustion Plant (Emission Factors in kt / Mt for Coal & Fuel Oil, kt / Mth for Natural Gas)	923
Table A 8.1.1	Summary table of GHG (Including net emissions/removals from LULUCF) emissions by NC Category (Mt CO ₂ eq) – National Statistics coverage (UK and Crown Dependencies)	925
Table A 8.1.2	Summary table of GHG emissions by Gas (Mt CO ₂ eq) – National Statistics coverage	925
Table A 8.2.1	Summary table of GHG emissions by NC Category (Mt CO ₂ eq) – UK only	926
Table A 8.2.2	Summary table of GHG emissions by Gas (Mt CO ₂ eq) – UK Only	927
Table A 9.4.1	Example of the outputs during a final user calculation	934
Table A 9.5.1	Sources reallocated to final users and the fuels used	936
Table A 9.5.2	Final user category, IPCC sectors, and NAEI source names and activity names used in the emission calculation	939
Table A 9.6.1	Final user emissions from all National Communication categories, MtCO ₂ equivalent	965
Table A 9.6.2	Final user CO ₂ emissions from all National Communication categories, MtCO ₂ equivalent	965
Table A 9.6.3	Final user CH ₄ emissions from all National Communication categories, MtCO ₂ equivalent	966
Table A 9.6.4	Final user N ₂ O emissions from all National Communication categories, MtCO ₂ equivalent	967

Table of Figures:

Figure A 3.4.1	Proportional land use contribution to forest conversion in the UK in the previous 20 years, 1990-2013.....	796
Figure A 3.4.2	Area of forest on different drained soil types in the UK 1990-2013	797
Figure A 3.4.3	Stratification of environments across the UK with areas 1 to 6 based on the underlying Land Classification (45 classes).	801
Figure A 3.4.4	Decision tree for assessing the effects of Cropland Management on soil carbon stocks.	807
Figure A 3.4.5	Comparison of estimated areas of deforestation in the UK between the 1990-2012 and 1990-2013 inventories	815

Contents of Annexes

Figure A 3.4.6	Country-specific biomass densities for biomass burning, 1990-2013. Living densities (top) and dead organic matter – litter and dead wood (bottom).	816
Figure A 3.4.7	Annual area of FIRMS thermal anomalies for GB for 2001-2013 (thermal anomalies were filtered to exclude those recorded over urban/industrial areas).	819
Figure A 3.4.8	Time series of wildfire burnt areas in the UK 1990-2013	820
Figure A 3.4.9	Peat extraction site visible on Google Earth imagery	823
Figure A 3.4.10	The largest uncertainties in the LULUCF inventory, in terms of standard deviation in the output distributions	833
Figure A 6.2.1	Monthly Northern Hemisphere trend in methane estimated from Mace Head observations (blue line). Red line denotes the de-seasonalised long-term trend. Data under grey shading are not yet ratified.....	876
Figure A 6.3.1	Monthly Northern Hemisphere trend in nitrous oxide estimated from Mace Head observations (blue line). Red line denotes the de-seasonalised long-term trend. Data under grey shading are not yet ratified.....	878
Figure A 6.3.2	Verification of the UK emission inventory estimates for nitrous oxide in Gg yr ⁻¹ from 1990. GHGI estimates are shown in orange (uncertainty in red). InTEM (MHD-only, 3-year) estimates are shown in blue (uncertainty in light blue). InTEM (all data, 1-year) estimates are shown in black (uncertainty in grey).	878
Figure A 6.4.1	Monthly Northern Hemisphere trend in HFC-134a estimated from Mace Head observations (blue line). Red line denotes the de-seasonalised long-term trend. Data under grey shading are not yet ratified.....	879
Figure A 6.4.2	Verification of the UK emission inventory estimates for HFC-134a in Gg yr ⁻¹ from 1990. GHGI estimates are shown in orange (uncertainty in red). InTEM (MHD-only, 3-year) estimates are shown in blue (uncertainty in light blue). InTEM (all data, 1-year) estimates are shown in black (uncertainty in grey).	879
Figure A 6.4.3	Monthly Northern Hemisphere trend in HFC-152a estimated from Mace Head observations (blue line). Red line denotes the de-seasonalised long-term trend. Data under grey shading are not yet ratified.....	880
Figure A 6.4.4	Verification of the UK emission inventory estimates for HFC-152a in Gg yr ⁻¹ from 1990. GHGI estimates are shown in orange (uncertainty in red). InTEM (MHD-only, 3-year) estimates are shown in blue (uncertainty in light blue). InTEM (all data, 1-year) estimates are shown in black (uncertainty in grey).	881
Figure A 6.4.5	Monthly Northern Hemisphere trend in HFC-125 estimated from Mace Head observations (blue line). Red line denotes the de-seasonalised long-term trend. Data under grey shading are not yet ratified.....	881
Figure A 6.4.6	Verification of the UK emission inventory estimates for HFC-125 in Gg yr ⁻¹ from 1990. GHGI estimates are shown in orange (uncertainty in red). InTEM (MHD-only, 3-year) estimates are shown in blue (uncertainty in light blue).	882

Figure A 6.4.7	Monthly Northern Hemisphere trend in HFC-143a estimated from Mace Head observations (blue line). Red line denotes the de-seasonalised long-term trend. Data under grey shading are not yet ratified.....	882
Figure A 6.4.8	Verification of the UK emission inventory estimates for HFC-143a in Gg yr ⁻¹ from 1990. GHGI estimates are shown in orange (uncertainty in red). InTEM (MHD-only, 3-year) estimates are shown in blue (uncertainty in light blue). InTEM (all data, 1-year) estimates are shown in black (uncertainty in grey).....	883
Figure A 6.4.9	Monthly Northern Hemisphere trend in HFC-23 estimated from Mace Head observations (blue line). Red line denotes the de-seasonalised long-term trend. Data under grey shading are not yet ratified.....	883
Figure A 6.4.10	Verification of the UK emission inventory estimates for HFC-23 in Gg yr ⁻¹ from 1990. GHGI estimates are shown in orange (uncertainty in red). InTEM (MHD-only, 3-year) estimates are shown in blue (uncertainty in light blue).....	884
Figure A 6.4.11	Verification of the UK emission inventory estimates for HFC-23 in Gg yr ⁻¹ zoomed in for 2004-2013. GHGI estimates are shown in orange (uncertainty in red). InTEM (MHD-only, 3-year) estimates are shown in blue (uncertainty in light blue).	884
Figure A 6.4.12	Monthly Northern Hemisphere trend in HFC-32 estimated from Mace Head observations (blue line). Red line denotes the de-seasonalised long-term trend. Data under grey shading are not yet ratified.....	885
Figure A 6.4.13	Verification of the UK emission inventory estimates for HFC-32 in Gg yr ⁻¹ from 1990. GHGI estimates are shown in orange (uncertainty in red). InTEM (MHD-only, 3-year) estimates are shown in blue (uncertainty in light blue).....	885
Figure A 6.4.14	Monthly Northern Hemisphere trend in HFC-43-10mee estimated from Mace Head observations (blue line). Red line denotes the de-seasonalised long-term trend. Data under grey shading are not yet ratified.....	886
Figure A 6.4.15	Verification of the UK emission inventory estimates for HFC-43-10mee in Gg yr ⁻¹ from 1990. GHGI estimates are shown in orange (uncertainty in red). InTEM (MHD-only, 3-year) estimates are shown in blue (uncertainty in light blue). InTEM (all data, 1-year) estimates are shown in black (uncertainty in grey).	886
Figure A 6.4.16	Monthly Northern Hemisphere trend in HFC-227ea estimated from Mace Head observations (blue line). Red line denotes the de-seasonalised long-term trend. Data under grey shading are not yet ratified.....	887
Figure A 6.4.17	Verification of the UK emission inventory estimates for HFC-227ea in Gg yr ⁻¹ from 1990. GHGI estimates are shown in orange (uncertainty in red). InTEM (MHD-only, 3-year) estimates are shown in blue (uncertainty in light blue). InTEM (all data, 1-year) estimates are shown in black (uncertainty in grey).....	887
Figure A 6.4.18	Monthly Northern Hemisphere trend in HFC-365mfc estimated from Mace Head observations (blue line). Red line denotes the de-seasonalised long-term trend. Data under grey shading are not yet ratified.....	888

Figure A 6.4.19	Verification of the UK emission inventory estimates for HFC-365mfc in Gg yr ⁻¹ from 1990. GHGI estimates are shown in orange (uncertainty in red). InTEM (MHD-only, 3-year) estimates are shown in blue (uncertainty in light blue). InTEM (all data, 1-year) estimates are shown in black (uncertainty in grey).....	888
Figure A 6.4.20	Monthly Northern Hemisphere trend in HFC-245fa estimated from Mace Head observations (blue line). Red line denotes the de-seasonalised long-term trend. Data under grey shading are not yet ratified.....	889
Figure A 6.4.21	Verification of the UK emission inventory estimates for HFC-245fa in Gg yr ⁻¹ from 1990. GHGI estimates are shown in orange (uncertainty in red). InTEM (MHD-only, 3-year) estimates are shown in blue (uncertainty in light blue).....	889
Figure A 6.5.1	Monthly Northern Hemisphere trend in PFC-14 estimated from Mace Head observations (blue line). Red line denotes the de-seasonalised long-term trend. Data under grey shading are not yet ratified.....	890
Figure A 6.5.2	Verification of the UK emission inventory estimates for PFC-14 in Gg yr ⁻¹ from 1990. GHGI estimates are shown in orange (uncertainty in red). InTEM (MHD-only, 3-year) estimates are shown in blue (uncertainty in light blue). InTEM (all data, 1-year) estimates are shown in black (uncertainty in grey).....	890
Figure A 6.5.3	Monthly Northern Hemisphere trend in PFC-116 estimated from Mace Head observations (blue line). Red line denotes the de-seasonalised long-term trend. Data under grey shading are not yet ratified.....	891
Figure A 6.5.4	Verification of the UK emission inventory estimates for PFC-116 in Gg yr ⁻¹ from 1990. GHGI estimates are shown in orange (uncertainty in red). InTEM (MHD-only, 3-year) estimates are shown in blue (uncertainty in light blue). InTEM (all data, 1-year) estimates are shown in black (uncertainty in grey).....	891
Figure A 6.5.5	Monthly Northern Hemisphere trend in PFC-218 estimated from Mace Head observations (blue line). Red line denotes the de-seasonalised long-term trend. Data under grey shading are not yet ratified.....	892
Figure A 6.5.6	Verification of the UK emission inventory estimates for PFC-218 in Gg yr ⁻¹ from 1990. GHGI estimates are shown in orange (uncertainty in red). InTEM (MHD-only, 3-year) estimates are shown in blue (uncertainty in light blue). InTEM (all data, 1-year) estimates are shown in black (uncertainty in grey).....	892
Figure A 6.5.7	Monthly Northern Hemisphere trend in PFC-318 estimated from Mace Head observations (blue line). Red line denotes the de-seasonalised long-term trend. Data under grey shading are not yet ratified.....	893
Figure A 6.5.8	Verification of the UK emission inventory estimates for PFC-318 in Gg yr ⁻¹ from 1990. GHGI estimates are shown in orange (uncertainty in red). InTEM (MHD-only, 3-year) estimates are shown in blue (uncertainty in light blue). InTEM (all data, 1-year) estimates are shown in black (uncertainty in grey).....	893

Figure A 6.6.1	Monthly Northern Hemisphere trend in SF ₆ estimated from Mace Head observations (blue line). Red line denotes the de-seasonalised long-term trend. Data under grey shading are not yet ratified.....	894
Figure A 6.6.2	Verification of the UK emission inventory estimates for SF ₆ in Gg yr ⁻¹ from 1990. GHGI estimates are shown in orange (uncertainty in red). InTEM (MHD-only, 3-year) estimates are shown in blue (uncertainty in light blue). InTEM (all data, 1-year) estimates are shown in black (uncertainty in grey).....	894
Figure A 6.7.1	Monthly Northern Hemisphere trend in NF ₃ estimated from Mace Head observations (blue line). Red line denotes the de-seasonalised long-term trend. Data under grey shading are not yet ratified.....	895
Figure A 6.8.1	Monthly Northern Hemisphere trend in CO ₂ estimated from Mace Head observations (blue line). Red line denotes the de-seasonalised long-term trend. Data under grey shading are not yet ratified.....	895
Figure A 6.8.2	Verification of the UK emission inventory estimates for CO ₂ in Gg yr ⁻¹ for 1990-2011. GHGI estimates are shown in orange (uncertainty in red). InTEM (MHD-only, 3-year, using CO ratio) estimates are shown in blue (uncertainty in light blue). InTEM (all network, winter & daytime data only) estimate shown in black (uncertainty in black).	896
Figure A 9.3.1	Simplified fuel flows for a final user calculation.....	931
Figure A 9.4.1	Fuel use in the example calculation.....	933
Figure A 9.4.2	Comparison of 'direct' and final user emissions of carbon according the sectors considered in the final user example.....	935

ANNEX 1: Key Categories

This annex contains the key category analysis for the latest GHG inventory¹. It contains:

- A description of the methodology used for identifying key categories
- Information on the level of disaggregation
- Information to fulfil the reporting requirements of Tables 4.2 and 4.3 of volume 1 of the 2006 IPCC Guidelines, including and excluding land use, land-use change and forestry (LULUCF).

The annex also contains information relevant to the requirements of reporting under the Kyoto Protocol. The table below contains the additional KP information that Annex 1 needs to contain, and the locations of this information in the Annex².

Requirements	Locations of the relevant information in this Annex
Description of methodology used for identifying key categories, including KP-LULUCF	See sections immediately below including “ <i>General approach used to identify Key Categories</i> ” and “ <i>Approach used to identify KP-LULUCF Key Categories</i> ”.
Reference to the key category tables in the CRF	This Annex of the NIR presents detailed tables of information of the data derived from the key category analysis. These data are used to create the key category tables (Table 7) in the CRF.
Reference to the key category tables in the CRF, including in the KP-LULUCF CRF tables	This Annex of the NIR presents detailed tables of information of the data derived from the key category analysis. These data are used to create the key category KP-LULUCF tables (Table NIR 3) in the CRF.
Information on the level of disaggregation	The tables in this Annex contain information on the level of disaggregation used. The level of disaggregation follows IPCC 2006 Guidelines.
Tables 4.2 to 4.4 of Volume 4 the 2006 IPCC guidelines	The data requested in the 2006 Guidelines tables, including and excluding LULUCF, are provided in Table A 1.3.1 to Table A 1.4.6 and Table 1.7 to Table 1.10 .
<i>Table NIR.3, as contained in the annex to decision 6/CMP.3</i>	A facsimile of Table NIR 3, provided in the CRF, is given in Table A 1.8.1 .

¹ Following the requirements to report information about uncertainties as set out in FCCC/CP/2013/10/Add.3. Report of the Conference of the Parties on its nineteenth session, held in Warsaw from 11 to 23 November 2013. Addendum Part two: Action taken by the Conference of the Parties at its nineteenth session.

² The information in this table has been taken directly from the UNFCCC document “Annotated outline of the National Inventory Report including reporting elements under the Kyoto Protocol”.

A 1.1 GENERAL APPROACH USED TO IDENTIFY KEY CATEGORIES

In the UK inventory, certain source categories are particularly significant in terms of their contribution to the overall uncertainty of the inventory. These key source categories have been identified so that the resources available for inventory preparation may be prioritised, and the best possible estimates prepared for the most significant source categories.

The UK completes both quantitative and qualitative Key Category Analyses (KCAs).

The UK has used the method set out in Section 4.3.1 and Section 4.3.2 of the 2006 IPCC Guidelines Volume 1 General Guidance and Reporting (*Approach 1 to identify key categories*, and *Approach 2 to identify key categories* respectively) to quantitatively determine the key source categories.

The method used in the qualitative KCA is described below, and further descriptions of the methods the UK uses to quantitatively determine key categories are given later in this section.

A 1.2 QUALITATIVE ANALYSIS USED TO IDENTIFY KEY CATEGORIES

Following IPCC good practice, a qualitative analysis of the inventory has been made to identify any additional key source categories, which may not have been identified using the quantitative analysis. The approach set out in Section 4.3.3 of the IPCC 2006 GLs has been applied, using the four criteria set out in the guidance, to judge whether a category is a key category. The criteria are:

1. (Use of) mitigation techniques and technologies;
2. Emissions growth (increase or decrease);
3. No quantitative assessment of uncertainties performed;
4. Completeness (examine qualitatively potential key categories that are not yet estimated quantitatively by applying the qualitative considerations above).

In addition, additional criteria have also been taken in account

5. High uncertainty (links to point 3 above);
6. Unexpectedly low or high emissions;
7. External recommendation has also been used as an additional criterion to identify key categories.

The results of this qualitative analysis are summarised in **Table A 1.2.1**. This is kept under review.

Table A 1.2.1 Qualitative Key Category Analysis

Source category	2006 IPCC GLs qualitative key category criteria				Other criteria
	(Use of) mitigation techniques and technologies	Emissions growth (increase or decrease)	No quantitative assessment of uncertainties performed	Completeness	
Cement Production (2A1)					X ^a

Notes

- a Following UNFCCC Expert Review Team recommendation from the 2010 Centralised Review (FCCC/ARR/2010/GBR) to include this source category as a key category: "...excluding uncertainties, this category is by far the most significant category within the industrial processes sector. The ERT recommends therefore, based on this quantitative and qualitative criterion, that the United Kingdom consider this category as key." Note that cement production is now being identified by the UK's approach 1 key category analysis, so this doesn't present an additional key category.

A 1.3 QUANTITATIVE APPROACH 1 KCA FOLLOWING IPCC 2006 GUIDELINES

A key category analysis has been completed for both level and trend. This KCA has been created using the 2006 IPCC GLs Approach 1 methodology. The factors that make a source a key category are:

- A high contribution to the level of emissions;
- A high contribution to the trend;

The results of the key category analysis with and without LULUCF, for the base year and the latest reported year and for both Approaches 1 and 2 KCA, are summarised by sector and gas in **Section 1.5.1**. The tables indicate whether a key category arises from the level (L1) assessment or the trend (T1) assessment.

For example, transport fuel (1A3b) is a key category for carbon dioxide because it is a large source of emissions and nitric acid production (2B2) because it shows a significant trend.

The results of the **level assessment** (based on Approach 1) with and without LULUCF for the base year and the latest reported year are shown in **Table A 1.3.1** to **Table A 1.3.4**. The key source categories are highlighted by the shaded cells in the table. The source categories (i.e. rows of the table) were sorted in descending order of magnitude based on the results of the "Level Parameter", and then the cumulative total was included in the final column of the table. The key source categories are those whose contributions add up to 95% of the sum of the level parameters in the final column after this sorting process, which according to the 2006 IPCC guidelines, should account for 90% of the uncertainty in level.

The results of the **trend assessment** (based on Approach 1) with and without LULUCF for the base year to the latest reported year are shown in **Table A 1.3.5** and **Table A 1.3.6**. The key source categories are highlighted by the shaded cells in the table. The trend parameter was calculated using the absolute value of the result; an absolute function is used since Land Use,

Land Use Change and Forestry contains negative sources (sinks) and the absolute function is necessary to produce positive uncertainty contributions for these sinks. The source categories (i.e. rows of the table) were sorted in descending order of the “Trend parameter”, and then the cumulative total was included in the final column of the table. The key source categories are those whose contributions add up to 95% of the sum of the trend parameters in the final column after this sorting process, which according to the 2006 IPCC guidelines, should account for 90% of the uncertainty in trend.

Table A 1.3.1 Approach 1 Key Category Analysis for the base year based on level of emissions (including LULUCF)

IPCC Code	IPCC Category	Greenhouse Gas	Base year emissions (GgCO ₂ e)	Absolute value of Base year emissions (GgCO ₂ e)	Level Assessment	Cumulative Total
1A1	Energy industries: solid fuels	CO ₂	185,494.19	185,494.19	0.2171	0.2171
1A3b	Road transportation: liquid fuels	CO ₂	108,570.51	108,570.51	0.1271	0.3441
1A4	Other sectors: gaseous fuels	CO ₂	70,384.62	70,384.62	0.0824	0.4265
5A	Solid waste disposal	CH ₄	62,749.72	62,749.72	0.0734	0.4999
1A1	Energy industries: liquid fuels	CO ₂	40,878.89	40,878.89	0.0478	0.5478
1A2	Manufacturing industries and construction: solid fuels	CO ₂	38,947.00	38,947.00	0.0456	0.5933
1A2	Manufacturing industries and construction: liquid fuels	CO ₂	29,119.63	29,119.63	0.0341	0.6274
1A2	Manufacturing industries and construction: gaseous fuels	CO ₂	28,170.48	28,170.48	0.0330	0.6604
3D	Agricultural soils	N ₂ O	23,153.96	23,153.96	0.0271	0.6875
3A1	Enteric fermentation from Cattle	CH ₄	21,921.41	21,921.41	0.0257	0.7131
1B1	Coal mining and handling	CH ₄	21,788.17	21,788.17	0.0255	0.7386
2B3	Adipic acid production	N ₂ O	19,934.61	19,934.61	0.0233	0.7620
1A4	Other sectors: solid fuels	CO ₂	19,876.40	19,876.40	0.0233	0.7852
1A4	Other sectors: liquid fuels	CO ₂	18,559.20	18,559.20	0.0217	0.8069
2B9	Fluorochemical production	HFCs, PFCs, SF ₆ and NF ₃	17,793.94	17,793.94	0.0208	0.8278
4A	Forest land	CO ₂	-16,016.07	16,016.07	0.0187	0.8465
4B	Cropland	CO ₂	15,130.79	15,130.79	0.0177	0.8642
1B2	Oil and gas extraction	CH ₄	12,332.94	12,332.94	0.0144	0.8786
1A1	Energy industries: gaseous fuels	CO ₂	9,179.46	9,179.46	0.0107	0.8894
2A1	Cement production	CO ₂	7,295.26	7,295.26	0.0085	0.8979

IPCC Code	IPCC Category	Greenhouse Gas	Base year emissions (GgCO ₂ e)	Absolute value of Base year emissions (GgCO ₂ e)	Level Assessment	Cumulative Total
4E	Settlements	CO ₂	6,919.10	6,919.10	0.0081	0.9060
1B2	Oil and gas extraction	CO ₂	5,777.92	5,777.92	0.0068	0.9128
3A2	Enteric fermentation from Sheep	CH ₄	5,727.52	5,727.52	0.0067	0.9195
2C1	Iron and steel production	CO ₂	5,582.62	5,582.62	0.0065	0.9260
1A5	Other: liquid fuels	CO ₂	5,284.82	5,284.82	0.0062	0.9322
5D	Wastewater treatment and discharge	CH ₄	4,169.64	4,169.64	0.0049	0.9371
2B8	Petrochemical and carbon black production	CO ₂	4,036.08	4,036.08	0.0047	0.9418
2B2	Nitric acid production	N ₂ O	3,860.26	3,860.26	0.0045	0.9463
4C	Grassland	CO ₂	-3,677.20	3,677.20	0.0043	0.9506
3B1	Manure management from Cattle	CH ₄	3,072.25	3,072.25	0.0036	0.9542
1A3d	Domestic Navigation: liquid fuels	CO ₂	2,168.02	2,168.02	0.0025	0.9568
2B1	Ammonia production	CO ₂	2,004.50	2,004.50	0.0023	0.9591
1A3a	Domestic aviation: liquid fuels	CO ₂	1,859.57	1,859.57	0.0022	0.9613
1B1	Coal mining and handling solid fuels	CO ₂	1,698.56	1,698.56	0.0020	0.9633
3G	Liming	CO ₂	1,579.17	1,579.17	0.0018	0.9651
1A1	Energy industries: solid fuels	N ₂ O	1,553.80	1,553.80	0.0018	0.9669
1A4	Other sectors: solid fuels	CH ₄	1,477.49	1,477.49	0.0017	0.9687
2A2	Lime production	CO ₂	1,462.05	1,462.05	0.0017	0.9704
1A3c	Railways: liquid fuels	CO ₂	1,455.18	1,455.18	0.0017	0.9721
2C6	Zinc production	CO ₂	1,358.83	1,358.83	0.0016	0.9737
5C	Incineration and open burning of waste	CO ₂	1,304.38	1,304.38	0.0015	0.9752
2D	Non-energy products from fuels and solvent use	CO ₂	1,232.62	1,232.62	0.0014	0.9766
1A3b	Road transportation: liquid fuels	N ₂ O	1,142.41	1,142.41	0.0013	0.9780
5D	Wastewater treatment and discharge	N ₂ O	1,111.13	1,111.13	0.0013	0.9793
3B1	Manure management from Cattle	N ₂ O	1,107.19	1,107.19	0.0013	0.9806
3B3	Manure management from Swine	CH ₄	1,091.66	1,091.66	0.0013	0.9818

IPCC Code	IPCC Category	Greenhouse Gas	Base year emissions (GgCO ₂ e)	Absolute value of Base year emissions (GgCO ₂ e)	Level Assessment	Cumulative Total
3B4	Manure management from Other livestock	N ₂ O	1,053.80	1,053.80	0.0012	0.9831
1A2	Manufacturing industries and construction: liquid fuels	N ₂ O	910.70	910.70	0.0011	0.9841
2F1	Refrigeration and air conditioning	HFCs, PFCs, SF ₆ and NF ₃	855.96	855.96	0.0010	0.9851
1A3b	Road transportation: liquid fuels	CH ₄	758.31	758.31	0.0009	0.9860
2G1	Electrical equipment	HFCs, PFCs, SF ₆ and NF ₃	747.79	747.79	0.0009	0.9869
2F4	Aerosols	HFCs, PFCs, SF ₆ and NF ₃	663.14	663.14	0.0008	0.9877
4B	Cropland	N ₂ O	649.15	649.15	0.0008	0.9884
2A4	Other process uses of carbonates	CO ₂	646.76	646.76	0.0008	0.9892
1A4	Other sectors: liquid fuels	N ₂ O	595.99	595.99	0.0007	0.9899
4D	Wetlands	CO ₂	481.73	481.73	0.0006	0.9905
1A4	Other sectors: peat	CO ₂	475.59	475.59	0.0006	0.9910
2C3	Aluminium production	CO ₂	450.32	450.32	0.0005	0.9915
2A3	Glass production	CO ₂	408.14	408.14	0.0005	0.9920
1A2	Manufacturing industries and construction: solid fuels	N ₂ O	387.27	387.27	0.0005	0.9925
2C4	Magnesium production	HFCs, PFCs, SF ₆ and NF ₃	387.17	387.17	0.0005	0.9929
3H	Urea application to land	CO ₂	385.46	385.46	0.0005	0.9934
4E	Settlements	N ₂ O	369.67	369.67	0.0004	0.9938
2C3	Aluminium production	HFCs, PFCs, SF ₆ and NF ₃	333.43	333.43	0.0004	0.9942
3A4	Enteric fermentation from Other livestock	CH ₄	295.24	295.24	0.0003	0.9945
3A3	Enteric fermentation from Swine	CH ₄	283.32	283.32	0.0003	0.9949
2G2	SF ₆ and PFCs from other product use	HFCs, PFCs, SF ₆ and NF ₃	278.58	278.58	0.0003	0.9952

IPCC Code	IPCC Category	Greenhouse Gas	Base year emissions (GgCO ₂ e)	Absolute value of Base year emissions (GgCO ₂ e)	Level Assessment	Cumulative Total
1A4	Other sectors: solid fuels	N ₂ O	277.04	277.04	0.0003	0.9955
2B7	Soda ash production	CO ₂	231.55	231.55	0.0003	0.9958
1A1	Energy industries: other fuels	CO ₂	226.99	226.99	0.0003	0.9961
1A3e	Other transportation: liquid fuels	CO ₂	224.74	224.74	0.0003	0.9963
3B2	Manure management from Sheep	CH ₄	222.48	222.48	0.0003	0.9966
3F	Field burning of agricultural residues	CH ₄	205.48	205.48	0.0002	0.9968
1A1	Energy industries: gaseous fuels	N ₂ O	198.68	198.68	0.0002	0.9971
3B3	Manure management from Swine	N ₂ O	196.57	196.57	0.0002	0.9973
2B10	Other Chemical Industry	CH ₄	185.65	185.65	0.0002	0.9975
2F2	Foam blowing agents	HFCs, PFCs, SF ₆ and NF ₃	184.55	184.55	0.0002	0.9977
2F6	Other product uses as substitutes for ODS	HFCs, PFCs, SF ₆ and NF ₃	165.54	165.54	0.0002	0.9979
1A4	Other sectors: gaseous fuels	CH ₄	157.20	157.20	0.0002	0.9981
1A1	Energy industries: liquid fuels	N ₂ O	156.12	156.12	0.0002	0.9983
5C	Incineration and open burning of waste	CH ₄	137.22	137.22	0.0002	0.9984
2B6	Titanium dioxide production	CO ₂	104.63	104.63	0.0001	0.9986
1A1	Energy industries: gaseous fuels	CH ₄	92.19	92.19	0.0001	0.9987
3B4	Manure management from Other livestock	CH ₄	86.40	86.40	0.0001	0.9988
1A4	Other sectors: biomass	CH ₄	75.42	75.42	0.0001	0.9989
3F	Field burning of agricultural residues	N ₂ O	63.50	63.50	0.0001	0.9989
2C1	Iron and steel production	N ₂ O	62.46	62.46	0.0001	0.9990
1A4	Other sectors: liquid fuels	CH ₄	56.73	56.73	0.0001	0.9991
4A	Forest land	N ₂ O	48.26	48.26	0.0001	0.9991
1A5	Other: liquid fuels	N ₂ O	47.24	47.24	0.0001	0.9992
3B2	Manure management from Sheep	N ₂ O	45.91	45.91	0.0001	0.9992
1A1	Energy industries: liquid fuels	CH ₄	43.71	43.71	0.0001	0.9993

IPCC Code	IPCC Category	Greenhouse Gas	Base year emissions (GgCO ₂ e)	Absolute value of Base year emissions (GgCO ₂ e)	Level Assessment	Cumulative Total
1A2	Manufacturing industries and construction: liquid fuels	CH ₄	43.69	43.69	0.0001	0.9993
1A1	Energy industries: solid fuels	CH ₄	41.90	41.90	0.0000	0.9994
1B2	Oil and gas extraction	N ₂ O	40.75	40.75	0.0000	0.9994
4G	Harvested wood products	CO ₂	40.72	40.72	0.0000	0.9995
1A4	Other sectors: gaseous fuels	N ₂ O	37.48	37.48	0.0000	0.9995
2C1	Iron and steel production	CH ₄	36.89	36.89	0.0000	0.9996
1A4	Other sectors: peat	CH ₄	33.65	33.65	0.0000	0.9996
2A4	Other process uses of carbonates	CH ₄	31.27	31.27	0.0000	0.9996
5C	Incineration and open burning of waste	N ₂ O	29.85	29.85	0.0000	0.9997
1A3e	Other transportation: liquid fuels	N ₂ O	27.86	27.86	0.0000	0.9997
2B8	Petrochemical and carbon black production	CH ₄	27.79	27.79	0.0000	0.9997
1A2	Manufacturing industries and construction: solid fuels	CH ₄	21.01	21.01	0.0000	0.9998
1A3a	Domestic aviation: liquid fuels	N ₂ O	17.60	17.60	0.0000	0.9998
1A3d	Domestic Navigation: liquid fuels	N ₂ O	16.26	16.26	0.0000	0.9998
1A2	Manufacturing industries and construction: gaseous fuels	N ₂ O	14.99	14.99	0.0000	0.9998
1A1	Energy industries: other fuels	CH ₄	13.57	13.57	0.0000	0.9998
1A2	Manufacturing industries and construction: gaseous fuels	CH ₄	12.57	12.57	0.0000	0.9999
4C	Grassland	N ₂ O	12.10	12.10	0.0000	0.9999
4C	Grassland	CH ₄	11.29	11.29	0.0000	0.9999
1A4	Other sectors: biomass	N ₂ O	11.20	11.20	0.0000	0.9999
1A2	Manufacturing industries and construction: biomass	N ₂ O	10.44	10.44	0.0000	0.9999
2E1	Integrated circuit or semiconductor	HFCs, PFCs, SF ₆ and NF ₃	9.56	9.56	0.0000	0.9999
1A2	Manufacturing industries and construction: biomass	CH ₄	6.57	6.57	0.0000	0.9999
1A1	Energy industries: other fuels	N ₂ O	5.96	5.96	0.0000	0.9999
4E	Settlements	CH ₄	5.89	5.89	0.0000	0.9999

IPCC Code	IPCC Category	Greenhouse Gas	Base year emissions (GgCO ₂ e)	Absolute value of Base year emissions (GgCO ₂ e)	Level Assessment	Cumulative Total
5B	Biological treatment of solid waste	CH ₄	5.48	5.48	0.0000	1.0000
5B	Biological treatment of solid waste	N ₂ O	4.90	4.90	0.0000	1.0000
1A3a	Domestic aviation: liquid fuels	CH ₄	4.17	4.17	0.0000	1.0000
4D	Wetlands	N ₂ O	3.83	3.83	0.0000	1.0000
1A5	Other: liquid fuels	CH ₄	3.73	3.73	0.0000	1.0000
4A	Forest land	CH ₄	3.72	3.72	0.0000	1.0000
1A3c	Railways: liquid fuels	N ₂ O	3.27	3.27	0.0000	1.0000
1A3c	Railways: liquid fuels	CH ₄	2.46	2.46	0.0000	1.0000
1A3d	Domestic Navigation: liquid fuels	CH ₄	2.22	2.22	0.0000	1.0000
1A4	Other sectors: peat	N ₂ O	1.87	1.87	0.0000	1.0000
2B8	Petrochemical and carbon black production	N ₂ O	1.79	1.79	0.0000	1.0000
2D	Non-energy products from fuels and solvent use	N ₂ O	1.64	1.64	0.0000	1.0000
2F3	Fire protection	HFCs, PFCs, SF ₆ and NF ₃	1.42	1.42	0.0000	1.0000
1A2	Manufacturing industries and construction: other fuels	CO ₂	1.07	1.07	0.0000	1.0000
2D	Non-energy products from fuels and solvent use	CH ₄	0.69	0.69	0.0000	1.0000
1A1	Energy industries: biomass	CH ₄	0.47	0.47	0.0000	1.0000
2G3	N ₂ O from product uses	N ₂ O	0.44	0.44	0.0000	1.0000
2B1	Ammonia production	N ₂ O	0.31	0.31	0.0000	1.0000
1A3e	Other transportation: liquid fuels	CH ₄	0.29	0.29	0.0000	1.0000
2B1	Ammonia production	CH ₄	0.26	0.26	0.0000	1.0000
1A1	Energy industries: biomass	N ₂ O	0.25	0.25	0.0000	1.0000
4B	Cropland	CH ₄	0.11	0.11	0.0000	1.0000
1B1	Coal mining and handling biomass	CH ₄	0.10	0.10	0.0000	1.0000
1B1	Coal mining and handling solid fuels	N ₂ O	0.09	0.09	0.0000	1.0000
1B1	Coal mining and handling solid fuels	CH ₄	0.08	0.08	0.0000	1.0000

IPCC Code	IPCC Category	Greenhouse Gas	Base year emissions (GgCO ₂ e)	Absolute value of Base year emissions (GgCO ₂ e)	Level Assessment	Cumulative Total
1A2	Manufacturing industries and construction: other fuels	N ₂ O	0.02	0.02	0.0000	1.0000
1A2	Manufacturing industries and construction: other fuels	CH ₄	0.01	0.01	0.0000	1.0000
Total			815,143.98	854,530.52	1.0000	

Table A 1.3.2 Approach 1 Key Category Analysis for the base year based on level of emissions (excluding LULUCF)

IPCC Code	IPCC Category	Greenhouse Gas	Base year emissions (Gg CO ₂ e)	Absolute value of Base year emissions (Gg CO ₂ e)	Level Assessment	Cumulative Total
1A1	Energy industries: solid fuels	CO ₂	185,494.19	185,494.19	0.2287	0.2287
1A3b	Road transportation: liquid fuels	CO ₂	108,570.51	108,570.51	0.1338	0.3625
1A4	Other sectors: gaseous fuels	CO ₂	70,384.62	70,384.62	0.0868	0.4493
5A	Solid waste disposal	CH ₄	62,749.72	62,749.72	0.0774	0.5267
1A1	Energy industries: liquid fuels	CO ₂	40,878.89	40,878.89	0.0504	0.5770
1A2	Manufacturing industries and construction: solid fuels	CO ₂	38,947.00	38,947.00	0.0480	0.6251
1A2	Manufacturing industries and construction: liquid fuels	CO ₂	29,119.63	29,119.63	0.0359	0.6610
1A2	Manufacturing industries and construction: gaseous fuels	CO ₂	28,170.48	28,170.48	0.0347	0.6957
3D	Agricultural soils	N ₂ O	23,153.96	23,153.96	0.0285	0.7242
3A1	Enteric fermentation from Cattle	CH ₄	21,921.41	21,921.41	0.0270	0.7513
1B1	Coal mining and handling	CH ₄	21,788.17	21,788.17	0.0269	0.7781
2B3	Adipic acid production	N ₂ O	19,934.61	19,934.61	0.0246	0.8027
1A4	Other sectors: solid fuels	CO ₂	19,876.40	19,876.40	0.0245	0.8272
1A4	Other sectors: liquid fuels	CO ₂	18,559.20	18,559.20	0.0229	0.8501
2B9	Fluorochemical production	HFCs, PFCs, SF ₆ and NF ₃	17,793.94	17,793.94	0.0219	0.8720
1B2	Oil and gas extraction	CH ₄	12,332.94	12,332.94	0.0152	0.8872
1A1	Energy industries: gaseous fuels	CO ₂	9,179.46	9,179.46	0.0113	0.8985

IPCC Code	IPCC Category	Greenhouse Gas	Base year emissions (Gg CO ₂ e)	Absolute value of Base year emissions (Gg CO ₂ e)	Level Assessment	Cumulative Total
2A1	Cement production	CO ₂	7,295.26	7,295.26	0.0090	0.9075
1B2	Oil and gas extraction	CO ₂	5,777.92	5,777.92	0.0071	0.9146
3A2	Enteric fermentation from Sheep	CH ₄	5,727.52	5,727.52	0.0071	0.9217
2C1	Iron and steel production	CO ₂	5,582.62	5,582.62	0.0069	0.9286
1A5	Other: liquid fuels	CO ₂	5,284.82	5,284.82	0.0065	0.9351
5D	Wastewater treatment and discharge	CH ₄	4,169.64	4,169.64	0.0051	0.9402
2B8	Petrochemical and carbon black production	CO ₂	4,036.08	4,036.08	0.0050	0.9452
2B2	Nitric acid production	N ₂ O	3,860.26	3,860.26	0.0048	0.9500
3B1	Manure management from Cattle	CH ₄	3,072.25	3,072.25	0.0038	0.9538
1A3d	Domestic Navigation: liquid fuels	CO ₂	2,168.02	2,168.02	0.0027	0.9564
2B1	Ammonia production	CO ₂	2,004.50	2,004.50	0.0025	0.9589
1A3a	Domestic aviation: liquid fuels	CO ₂	1,859.57	1,859.57	0.0023	0.9612
1B1	Coal mining and handling solid fuels	CO ₂	1,698.56	1,698.56	0.0021	0.9633
3G	Liming	CO ₂	1,579.17	1,579.17	0.0019	0.9652
1A1	Energy industries: solid fuels	N ₂ O	1,553.80	1,553.80	0.0019	0.9672
1A4	Other sectors: solid fuels	CH ₄	1,477.49	1,477.49	0.0018	0.9690
2A2	Lime production	CO ₂	1,462.05	1,462.05	0.0018	0.9708
1A3c	Railways: liquid fuels	CO ₂	1,455.18	1,455.18	0.0018	0.9726
2C6	Zinc production	CO ₂	1,358.83	1,358.83	0.0017	0.9743
5C	Incineration and open burning of waste	CO ₂	1,304.38	1,304.38	0.0016	0.9759
2D	Non-energy products from fuels and solvent use	CO ₂	1,232.62	1,232.62	0.0015	0.9774
1A3b	Road transportation: liquid fuels	N ₂ O	1,142.41	1,142.41	0.0014	0.9788
5D	Wastewater treatment and discharge	N ₂ O	1,111.13	1,111.13	0.0014	0.9802
3B1	Manure management from Cattle	N ₂ O	1,107.19	1,107.19	0.0014	0.9815
3B3	Manure management from Swine	CH ₄	1,091.66	1,091.66	0.0013	0.9829
3B4	Manure management from Other livestock	N ₂ O	1,053.80	1,053.80	0.0013	0.9842

IPCC Code	IPCC Category	Greenhouse Gas	Base year emissions (Gg CO ₂ e)	Absolute value of Base year emissions (Gg CO ₂ e)	Level Assessment	Cumulative Total
1A2	Manufacturing industries and construction: liquid fuels	N ₂ O	910.70	910.70	0.0011	0.9853
2F1	Refrigeration and air conditioning	HFCs, PFCs, SF ₆ and NF ₃	855.96	855.96	0.0011	0.9863
1A3b	Road transportation: liquid fuels	CH ₄	758.31	758.31	0.0009	0.9873
2G1	Electrical equipment	HFCs, PFCs, SF ₆ and NF ₃	747.79	747.79	0.0009	0.9882
2F4	Aerosols	HFCs, PFCs, SF ₆ and NF ₃	663.14	663.14	0.0008	0.9890
2A4	Other process uses of carbonates	CO ₂	646.76	646.76	0.0008	0.9898
1A4	Other sectors: liquid fuels	N ₂ O	595.99	595.99	0.0007	0.9906
1A4	Other sectors: peat	CO ₂	475.59	475.59	0.0006	0.9911
2C3	Aluminium production	CO ₂	450.32	450.32	0.0006	0.9917
2A3	Glass production	CO ₂	408.14	408.14	0.0005	0.9922
1A2	Manufacturing industries and construction: solid fuels	N ₂ O	387.27	387.27	0.0005	0.9927
2C4	Magnesium production	HFCs, PFCs, SF ₆ and NF ₃	387.17	387.17	0.0005	0.9932
3H	Urea application to land	CO ₂	385.46	385.46	0.0005	0.9936
2C3	Aluminium production	HFCs, PFCs, SF ₆ and NF ₃	333.43	333.43	0.0004	0.9940
3A4	Enteric fermentation from Other livestock	CH ₄	295.24	295.24	0.0004	0.9944
3A3	Enteric fermentation from Swine	CH ₄	283.32	283.32	0.0003	0.9948
2G2	SF ₆ and PFCs from other product use	HFCs, PFCs, SF ₆ and NF ₃	278.58	278.58	0.0003	0.9951
1A4	Other sectors: solid fuels	N ₂ O	277.04	277.04	0.0003	0.9954
2B7	Soda ash production	CO ₂	231.55	231.55	0.0003	0.9957
1A1	Energy industries: other fuels	CO ₂	226.99	226.99	0.0003	0.9960
1A3e	Other transportation: liquid fuels	CO ₂	224.74	224.74	0.0003	0.9963
3B2	Manure management from Sheep	CH ₄	222.48	222.48	0.0003	0.9966

IPCC Code	IPCC Category	Greenhouse Gas	Base year emissions (Gg CO ₂ e)	Absolute value of Base year emissions (Gg CO ₂ e)	Level Assessment	Cumulative Total
3F	Field burning of agricultural residues	CH ₄	205.48	205.48	0.0003	0.9968
1A1	Energy industries: gaseous fuels	N ₂ O	198.68	198.68	0.0002	0.9971
3B3	Manure management from Swine	N ₂ O	196.57	196.57	0.0002	0.9973
2B10	Other Chemical Industry	CH ₄	185.65	185.65	0.0002	0.9975
2F2	Foam blowing agents	HFCs, PFCs, SF ₆ and NF ₃	184.55	184.55	0.0002	0.9978
2F6	Other product uses as substitutes for ODS	HFCs, PFCs, SF ₆ and NF ₃	165.54	165.54	0.0002	0.9980
1A4	Other sectors: gaseous fuels	CH ₄	157.20	157.20	0.0002	0.9982
1A1	Energy industries: liquid fuels	N ₂ O	156.12	156.12	0.0002	0.9983
5C	Incineration and open burning of waste	CH ₄	137.22	137.22	0.0002	0.9985
2B6	Titanium dioxide production	CO ₂	104.63	104.63	0.0001	0.9986
1A1	Energy industries: gaseous fuels	CH ₄	92.19	92.19	0.0001	0.9988
3B4	Manure management from Other livestock	CH ₄	86.40	86.40	0.0001	0.9989
1A4	Other sectors: biomass	CH ₄	75.42	75.42	0.0001	0.9990
3F	Field burning of agricultural residues	N ₂ O	63.50	63.50	0.0001	0.9990
2C1	Iron and steel production	N ₂ O	62.46	62.46	0.0001	0.9991
1A4	Other sectors: liquid fuels	CH ₄	56.73	56.73	0.0001	0.9992
1A5	Other: liquid fuels	N ₂ O	47.24	47.24	0.0001	0.9992
3B2	Manure management from Sheep	N ₂ O	45.91	45.91	0.0001	0.9993
1A1	Energy industries: liquid fuels	CH ₄	43.71	43.71	0.0001	0.9993
1A2	Manufacturing industries and construction: liquid fuels	CH ₄	43.69	43.69	0.0001	0.9994
1A1	Energy industries: solid fuels	CH ₄	41.90	41.90	0.0001	0.9995
1B2	Oil and gas extraction	N ₂ O	40.75	40.75	0.0001	0.9995
1A4	Other sectors: gaseous fuels	N ₂ O	37.48	37.48	0.0000	0.9996
2C1	Iron and steel production	CH ₄	36.89	36.89	0.0000	0.9996
1A4	Other sectors: peat	CH ₄	33.65	33.65	0.0000	0.9996

IPCC Code	IPCC Category	Greenhouse Gas	Base year emissions (Gg CO ₂ e)	Absolute value of Base year emissions (Gg CO ₂ e)	Level Assessment	Cumulative Total
2A4	Other process uses of carbonates	CH ₄	31.27	31.27	0.0000	0.9997
5C	Incineration and open burning of waste	N ₂ O	29.85	29.85	0.0000	0.9997
1A3e	Other transportation: liquid fuels	N ₂ O	27.86	27.86	0.0000	0.9997
2B8	Petrochemical and carbon black production	CH ₄	27.79	27.79	0.0000	0.9998
1A2	Manufacturing industries and construction: solid fuels	CH ₄	21.01	21.01	0.0000	0.9998
1A3a	Domestic aviation: liquid fuels	N ₂ O	17.60	17.60	0.0000	0.9998
1A3d	Domestic Navigation: liquid fuels	N ₂ O	16.26	16.26	0.0000	0.9998
1A2	Manufacturing industries and construction: gaseous fuels	N ₂ O	14.99	14.99	0.0000	0.9999
1A1	Energy industries: other fuels	CH ₄	13.57	13.57	0.0000	0.9999
1A2	Manufacturing industries and construction: gaseous fuels	CH ₄	12.57	12.57	0.0000	0.9999
1A4	Other sectors: biomass	N ₂ O	11.20	11.20	0.0000	0.9999
1A2	Manufacturing industries and construction: biomass	N ₂ O	10.44	10.44	0.0000	0.9999
2E1	Integrated circuit or semiconductor	HFCs, PFCs, SF ₆ and NF ₃	9.56	9.56	0.0000	0.9999
1A2	Manufacturing industries and construction: biomass	CH ₄	6.57	6.57	0.0000	0.9999
1A1	Energy industries: other fuels	N ₂ O	5.96	5.96	0.0000	1.0000
5B	Biological treatment of solid waste	CH ₄	5.48	5.48	0.0000	1.0000
5B	Biological treatment of solid waste	N ₂ O	4.90	4.90	0.0000	1.0000
1A3a	Domestic aviation: liquid fuels	CH ₄	4.17	4.17	0.0000	1.0000
1A5	Other: liquid fuels	CH ₄	3.73	3.73	0.0000	1.0000
1A3c	Railways: liquid fuels	N ₂ O	3.27	3.27	0.0000	1.0000
1A3c	Railways: liquid fuels	CH ₄	2.46	2.46	0.0000	1.0000
1A3d	Domestic Navigation: liquid fuels	CH ₄	2.22	2.22	0.0000	1.0000
1A4	Other sectors: peat	N ₂ O	1.87	1.87	0.0000	1.0000
2B8	Petrochemical and carbon black production	N ₂ O	1.79	1.79	0.0000	1.0000

IPCC Code	IPCC Category	Greenhouse Gas	Base year emissions (Gg CO ₂ e)	Absolute value of Base year emissions (Gg CO ₂ e)	Level Assessment	Cumulative Total
2D	Non-energy products from fuels and solvent use	N ₂ O	1.64	1.64	0.0000	1.0000
2F3	Fire protection	HFCs, PFCs, SF ₆ and NF ₃	1.42	1.42	0.0000	1.0000
1A2	Manufacturing industries and construction: other fuels	CO ₂	1.07	1.07	0.0000	1.0000
2D	Non-energy products from fuels and solvent use	CH ₄	0.69	0.69	0.0000	1.0000
1A1	Energy industries: biomass	CH ₄	0.47	0.47	0.0000	1.0000
2G3	N ₂ O from product uses	N ₂ O	0.44	0.44	0.0000	1.0000
2B1	Ammonia production	N ₂ O	0.31	0.31	0.0000	1.0000
1A3e	Other transportation: liquid fuels	CH ₄	0.29	0.29	0.0000	1.0000
2B1	Ammonia production	CH ₄	0.26	0.26	0.0000	1.0000
1A1	Energy industries: biomass	N ₂ O	0.25	0.25	0.0000	1.0000
1B1	Coal mining and handling biomass	CH ₄	0.10	0.10	0.0000	1.0000
1B1	Coal mining and handling solid fuels	N ₂ O	0.09	0.09	0.0000	1.0000
1B1	Coal mining and handling solid fuels	CH ₄	0.08	0.08	0.0000	1.0000
1A2	Manufacturing industries and construction: other fuels	N ₂ O	0.02	0.02	0.0000	1.0000
1A2	Manufacturing industries and construction: other fuels	CH ₄	0.01	0.01	0.0000	1.0000
Total			811,160.90	811,160.90	1.0000	

Table A 1.3.3 Approach 1 Key Category Analysis for the latest reported year based on level of emissions (including LULUCF)

IPCC Code	IPCC Category	Greenhouse Gas	2013 emissions (Gg CO ₂ e)	Absolute value of 2013 emissions (Gg CO ₂ e)	Level Assessment	Cumulative Total
1A1	Energy industries: solid fuels	CO ₂	113,205.74	113,205.74	0.1829	0.1829
1A3b	Road transportation: liquid fuels	CO ₂	107,048.58	107,048.58	0.1729	0.3558
1A4	Other sectors: gaseous fuels	CO ₂	87,203.06	87,203.06	0.1409	0.4966

IPCC Code	IPCC Category	Greenhouse Gas	2013 emissions (Gg CO ₂ e)	Absolute value of 2013 emissions (Gg CO ₂ e)	Level Assessment	Cumulative Total
1A1	Energy industries: gaseous fuels	CO ₂	44,618.23	44,618.23	0.0721	0.5687
1A2	Manufacturing industries and construction: gaseous fuels	CO ₂	22,842.35	22,842.35	0.0369	0.6056
1A2	Manufacturing industries and construction: solid fuels	CO ₂	20,641.24	20,641.24	0.0333	0.6389
3D	Agricultural soils	N ₂ O	19,513.90	19,513.90	0.0315	0.6705
3A1	Enteric fermentation from Cattle	CH ₄	18,703.95	18,703.95	0.0302	0.7007
1A1	Energy industries: liquid fuels	CO ₂	17,608.72	17,608.72	0.0284	0.7291
4A	Forest land	CO ₂	-17,297.63	17,297.63	0.0279	0.7571
5A	Solid waste disposal	CH ₄	16,681.83	16,681.83	0.0269	0.7840
2F1	Refrigeration and air conditioning	HFCs, PFCs, SF ₆ and NF ₃	13,144.55	13,144.55	0.0212	0.8052
1A2	Manufacturing industries and construction: liquid fuels	CO ₂	12,554.23	12,554.23	0.0203	0.8255
1A4	Other sectors: liquid fuels	CO ₂	12,541.14	12,541.14	0.0203	0.8458
4B	Cropland	CO ₂	12,150.39	12,150.39	0.0196	0.8654
4C	Grassland	CO ₂	-5,897.14	5,897.14	0.0095	0.8749
4E	Settlements	CO ₂	5,876.98	5,876.98	0.0095	0.8844
1B2	Oil and gas extraction	CH ₄	5,687.26	5,687.26	0.0092	0.8936
2C1	Iron and steel production	CO ₂	4,945.17	4,945.17	0.0080	0.9016
3A2	Enteric fermentation from Sheep	CH ₄	4,243.16	4,243.16	0.0069	0.9084
2A1	Cement production	CO ₂	4,029.11	4,029.11	0.0065	0.9150
1B2	Oil and gas extraction	CO ₂	3,789.72	3,789.72	0.0061	0.9211
5D	Wastewater treatment and discharge	CH ₄	3,359.33	3,359.33	0.0054	0.9265
1A4	Other sectors: solid fuels	CO ₂	3,147.53	3,147.53	0.0051	0.9316
2B8	Petrochemical and carbon black production	CO ₂	2,948.24	2,948.24	0.0048	0.9364
3B1	Manure management from Cattle	CH ₄	2,549.91	2,549.91	0.0041	0.9405
1A5	Other: liquid fuels	CO ₂	2,285.42	2,285.42	0.0037	0.9442
1A3d	Domestic Navigation: liquid fuels	CO ₂	2,165.99	2,165.99	0.0035	0.9477
2F4	Aerosols	HFCs, PFCs, SF ₆ and NF ₃	2,158.68	2,158.68	0.0035	0.9511

IPCC Code	IPCC Category	Greenhouse Gas	2013 emissions (Gg CO ₂ e)	Absolute value of 2013 emissions (Gg CO ₂ e)	Level Assessment	Cumulative Total
1A1	Energy industries: other fuels	CO ₂	2,040.24	2,040.24	0.0033	0.9544
1A3a	Domestic aviation: liquid fuels	CO ₂	1,961.73	1,961.73	0.0032	0.9576
1A3c	Railways: liquid fuels	CO ₂	1,925.85	1,925.85	0.0031	0.9607
1B1	Coal mining and handling	CH ₄	1,682.41	1,682.41	0.0027	0.9634
2B1	Ammonia production	CO ₂	1,383.38	1,383.38	0.0022	0.9657
2A2	Lime production	CO ₂	1,239.15	1,239.15	0.0020	0.9677
4G	Harvested wood products	CO ₂	-1,121.03	1,121.03	0.0018	0.9695
5D	Wastewater treatment and discharge	N ₂ O	1,063.03	1,063.03	0.0017	0.9712
2D	Non-energy products from fuels and solvent use	CO ₂	1,032.92	1,032.92	0.0017	0.9729
1A1	Energy industries: solid fuels	N ₂ O	935.10	935.10	0.0015	0.9744
1A3b	Road transportation: liquid fuels	N ₂ O	909.33	909.33	0.0015	0.9759
3B1	Manure management from Cattle	N ₂ O	897.98	897.98	0.0015	0.9773
3B4	Manure management from Other livestock	N ₂ O	837.18	837.18	0.0014	0.9787
3G	Liming	CO ₂	774.69	774.69	0.0013	0.9799
2A4	Other process uses of carbonates	CO ₂	771.83	771.83	0.0012	0.9812
5B	Biological treatment of solid waste	CH ₄	715.72	715.72	0.0012	0.9823
3B3	Manure management from Swine	CH ₄	633.80	633.80	0.0010	0.9833
5B	Biological treatment of solid waste	N ₂ O	608.32	608.32	0.0010	0.9843
1A2	Manufacturing industries and construction: liquid fuels	N ₂ O	538.92	538.92	0.0009	0.9852
2F2	Foam blowing agents	HFCs, PFCs, SF ₆ and NF ₃	495.71	495.71	0.0008	0.9860
3A4	Enteric fermentation from Other livestock	CH ₄	491.61	491.61	0.0008	0.9868
1A4	Other sectors: liquid fuels	N ₂ O	475.43	475.43	0.0008	0.9875
1A3e	Other transportation: liquid fuels	CO ₂	458.95	458.95	0.0007	0.9883
2A3	Glass production	CO ₂	389.18	389.18	0.0006	0.9889
1A2	Manufacturing industries and construction: other fuels	CO ₂	386.12	386.12	0.0006	0.9895
4B	Cropland	N ₂ O	329.94	329.94	0.0005	0.9901

IPCC Code	IPCC Category	Greenhouse Gas	2013 emissions (Gg CO ₂ e)	Absolute value of 2013 emissions (Gg CO ₂ e)	Level Assessment	Cumulative Total
2G1	Electrical equipment	HFCs, PFCs, SF ₆ and NF ₃	318.54	318.54	0.0005	0.9906
4E	Settlements	N ₂ O	318.07	318.07	0.0005	0.9911
4D	Wetlands	CO ₂	300.49	300.49	0.0005	0.9916
2B7	Soda ash production	CO ₂	285.27	285.27	0.0005	0.9920
2G2	SF ₆ and PFCs from other product use	HFCs, PFCs, SF ₆ and NF ₃	271.13	271.13	0.0004	0.9925
2F3	Fire protection	HFCs, PFCs, SF ₆ and NF ₃	269.86	269.86	0.0004	0.9929
5C	Incineration and open burning of waste	CO ₂	264.30	264.30	0.0004	0.9934
3H	Urea application to land	CO ₂	245.22	245.22	0.0004	0.9937
1A4	Other sectors: biomass	CH ₄	238.17	238.17	0.0004	0.9941
1A1	Energy industries: gaseous fuels	N ₂ O	221.69	221.69	0.0004	0.9945
1A4	Other sectors: solid fuels	CH ₄	208.43	208.43	0.0003	0.9948
1A4	Other sectors: gaseous fuels	CH ₄	191.56	191.56	0.0003	0.9951
3A3	Enteric fermentation from Swine	CH ₄	183.08	183.08	0.0003	0.9954
1A2	Manufacturing industries and construction: solid fuels	N ₂ O	171.54	171.54	0.0003	0.9957
3B2	Manure management from Sheep	CH ₄	164.91	164.91	0.0003	0.9960
2B9	Fluorochemical production	HFCs, PFCs, SF ₆ and NF ₃	158.24	158.24	0.0003	0.9962
2C4	Magnesium production	HFCs, PFCs, SF ₆ and NF ₃	148.80	148.80	0.0002	0.9965
1B1	Coal mining and handling solid fuels	CO ₂	133.60	133.60	0.0002	0.9967
2B6	Titanium dioxide production	CO ₂	123.43	123.43	0.0002	0.9969
3B4	Manure management from Other livestock	CH ₄	122.88	122.88	0.0002	0.9971
3B3	Manure management from Swine	N ₂ O	121.88	121.88	0.0002	0.9973
2F6	Other product uses as substitutes for ODS	HFCs, PFCs, SF ₆ and NF ₃	110.00	110.00	0.0002	0.9975
1A1	Energy industries: gaseous fuels	CH ₄	95.30	95.30	0.0002	0.9976
1A1	Energy industries: liquid fuels	N ₂ O	93.43	93.43	0.0002	0.9978
2C3	Aluminium production	CO ₂	68.14	68.14	0.0001	0.9979

IPCC Code	IPCC Category	Greenhouse Gas	2013 emissions (Gg CO ₂ e)	Absolute value of 2013 emissions (Gg CO ₂ e)	Level Assessment	Cumulative Total
1A1	Energy industries: biomass	N ₂ O	66.71	66.71	0.0001	0.9980
2B10	Other Chemical Industry	CH ₄	66.42	66.42	0.0001	0.9981
1A3e	Other transportation: liquid fuels	N ₂ O	56.82	56.82	0.0001	0.9982
1A3b	Road transportation: liquid fuels	CH ₄	55.62	55.62	0.0001	0.9983
4A	Forest land	N ₂ O	50.29	50.29	0.0001	0.9984
1A1	Energy industries: other fuels	N ₂ O	48.24	48.24	0.0001	0.9984
1A4	Other sectors: peat	CO ₂	46.96	46.96	0.0001	0.9985
1A4	Other sectors: gaseous fuels	N ₂ O	45.67	45.67	0.0001	0.9986
2C1	Iron and steel production	N ₂ O	45.23	45.23	0.0001	0.9987
5C	Incineration and open burning of waste	N ₂ O	44.28	44.28	0.0001	0.9987
1A4	Other sectors: solid fuels	N ₂ O	43.36	43.36	0.0001	0.9988
2B2	Nitric acid production	N ₂ O	42.91	42.91	0.0001	0.9989
1A1	Energy industries: biomass	CH ₄	42.77	42.77	0.0001	0.9989
1B1	Coal mining and handling liquid fuels	CO ₂	41.68	41.68	0.0001	0.9990
1A2	Manufacturing industries and construction: biomass	N ₂ O	39.02	39.02	0.0001	0.9991
1A1	Energy industries: other fuels	CH ₄	37.57	37.57	0.0001	0.9991
2B8	Petrochemical and carbon black production	CH ₄	37.45	37.45	0.0001	0.9992
1A4	Other sectors: biomass	N ₂ O	36.24	36.24	0.0001	0.9992
3B2	Manure management from Sheep	N ₂ O	34.24	34.24	0.0001	0.9993
1A4	Other sectors: liquid fuels	CH ₄	33.31	33.31	0.0001	0.9994
1A3c	Railways: solid fuels	CO ₂	32.60	32.60	0.0001	0.9994
1B2	Oil and gas extraction	N ₂ O	30.79	30.79	0.0000	0.9995
4C	Grassland	CH ₄	28.46	28.46	0.0000	0.9995
1A2	Manufacturing industries and construction: liquid fuels	CH ₄	26.69	26.69	0.0000	0.9995
1A1	Energy industries: solid fuels	CH ₄	25.28	25.28	0.0000	0.9996
1A2	Manufacturing industries and construction: biomass	CH ₄	24.61	24.61	0.0000	0.9996

IPCC Code	IPCC Category	Greenhouse Gas	2013 emissions (Gg CO ₂ e)	Absolute value of 2013 emissions (Gg CO ₂ e)	Level Assessment	Cumulative Total
2F5	Solvents	HFCs, PFCs, SF ₆ and NF ₃	21.93	21.93	0.0000	0.9997
2C1	Iron and steel production	CH ₄	20.49	20.49	0.0000	0.9997
1A5	Other: liquid fuels	N ₂ O	20.30	20.30	0.0000	0.9997
1A3a	Domestic aviation: liquid fuels	N ₂ O	18.56	18.56	0.0000	0.9998
1A1	Energy industries: liquid fuels	CH ₄	17.70	17.70	0.0000	0.9998
1A3d	Domestic Navigation: liquid fuels	N ₂ O	16.22	16.22	0.0000	0.9998
2E1	Integrated circuit or semiconductor	HFCs, PFCs, SF ₆ and NF ₃	14.83	14.83	0.0000	0.9998
1A2	Manufacturing industries and construction: gaseous fuels	N ₂ O	11.96	11.96	0.0000	0.9999
1A2	Manufacturing industries and construction: gaseous fuels	CH ₄	10.04	10.04	0.0000	0.9999
5C	Incineration and open burning of waste	CH ₄	9.95	9.95	0.0000	0.9999
4C	Grassland	N ₂ O	9.15	9.15	0.0000	0.9999
2C3	Aluminium production	HFCs, PFCs, SF ₆ and NF ₃	6.95	6.95	0.0000	0.9999
1A2	Manufacturing industries and construction: other fuels	N ₂ O	6.55	6.55	0.0000	0.9999
1A2	Manufacturing industries and construction: solid fuels	CH ₄	5.94	5.94	0.0000	0.9999
4A	Forest land	CH ₄	5.04	5.04	0.0000	0.9999
1A3d	Domestic Navigation: liquid fuels	CH ₄	4.90	4.90	0.0000	0.9999
2A4	Other process uses of carbonates	CH ₄	4.52	4.52	0.0000	1.0000
1A3c	Railways: liquid fuels	N ₂ O	4.32	4.32	0.0000	1.0000
1A2	Manufacturing industries and construction: other fuels	CH ₄	4.12	4.12	0.0000	1.0000
1B1	Coal mining and handling biomass	CH ₄	3.82	3.82	0.0000	1.0000
1A4	Other sectors: peat	CH ₄	3.32	3.32	0.0000	1.0000
1A3c	Railways: liquid fuels	CH ₄	2.12	2.12	0.0000	1.0000
4E	Settlements	CH ₄	1.99	1.99	0.0000	1.0000
1A5	Other: liquid fuels	CH ₄	1.59	1.59	0.0000	1.0000
2B8	Petrochemical and carbon black production	N ₂ O	1.53	1.53	0.0000	1.0000

IPCC Code	IPCC Category	Greenhouse Gas	2013 emissions (Gg CO ₂ e)	Absolute value of 2013 emissions (Gg CO ₂ e)	Level Assessment	Cumulative Total
2D	Non-energy products from fuels and solvent use	N ₂ O	0.91	0.91	0.0000	1.0000
1A3a	Domestic aviation: liquid fuels	CH ₄	0.86	0.86	0.0000	1.0000
1A3c	Railways: solid fuels	CH ₄	0.86	0.86	0.0000	1.0000
4D	Wetlands	N ₂ O	0.50	0.50	0.0000	1.0000
2G3	N ₂ O from product uses	N ₂ O	0.49	0.49	0.0000	1.0000
1A3e	Other transportation: liquid fuels	CH ₄	0.48	0.48	0.0000	1.0000
2D	Non-energy products from fuels and solvent use	CH ₄	0.38	0.38	0.0000	1.0000
2B1	Ammonia production	N ₂ O	0.27	0.27	0.0000	1.0000
2B1	Ammonia production	CH ₄	0.23	0.23	0.0000	1.0000
1A4	Other sectors: peat	N ₂ O	0.18	0.18	0.0000	1.0000
4B	Cropland	CH ₄	0.14	0.14	0.0000	1.0000
1A3c	Railways: solid fuels	N ₂ O	0.08	0.08	0.0000	1.0000
1B1	Coal mining and handling solid fuels	N ₂ O	0.04	0.04	0.0000	1.0000
1B1	Coal mining and handling solid fuels	CH ₄	0.03	0.03	0.0000	1.0000
4F	Other land	CO ₂	0.02	0.02	0.0000	1.0000
Total			570,451.65	619,083.25	1.0000	

Table A 1.3.4 Approach 1 Key Category Analysis for the latest reported year based on level of emissions (excluding LULUCF)

IPCC Code	IPCC Category	Greenhouse Gas	2013 emissions (Gg CO ₂ e)	Absolute value of 2013 emissions (Gg CO ₂ e)	Level Assessment	Cumulative Total
1A1	Energy industries: solid fuels	CO ₂	113,205.74	113,205.74	0.1966	0.1966
1A3b	Road transportation: liquid fuels	CO ₂	107,048.58	107,048.58	0.1859	0.3826
1A4	Other sectors: gaseous fuels	CO ₂	87,203.06	87,203.06	0.1515	0.5341
1A1	Energy industries: gaseous fuels	CO ₂	44,618.23	44,618.23	0.0775	0.6116
1A2	Manufacturing industries and construction: gaseous fuels	CO ₂	22,842.35	22,842.35	0.0397	0.6512

IPCC Code	IPCC Category	Greenhouse Gas	2013 emissions (Gg CO ₂ e)	Absolute value of 2013 emissions (Gg CO ₂ e)	Level Assessment	Cumulative Total
1A2	Manufacturing industries and construction: solid fuels	CO ₂	20,641.24	20,641.24	0.0359	0.6871
3D	Agricultural soils	N ₂ O	19,513.90	19,513.90	0.0339	0.7210
3A1	Enteric fermentation from Cattle	CH ₄	18,703.95	18,703.95	0.0325	0.7535
1A1	Energy industries: liquid fuels	CO ₂	17,608.72	17,608.72	0.0306	0.7841
5A	Solid waste disposal	CH ₄	16,681.83	16,681.83	0.0290	0.8130
2F1	Refrigeration and air conditioning	HFCs, PFCs, SF ₆ and NF ₃	13,144.55	13,144.55	0.0228	0.8359
1A2	Manufacturing industries and construction: liquid fuels	CO ₂	12,554.23	12,554.23	0.0218	0.8577
1A4	Other sectors: liquid fuels	CO ₂	12,541.14	12,541.14	0.0218	0.8795
1B2	Oil and gas extraction	CH ₄	5,687.26	5,687.26	0.0099	0.8893
2C1	Iron and steel production	CO ₂	4,945.17	4,945.17	0.0086	0.8979
3A2	Enteric fermentation from Sheep	CH ₄	4,243.16	4,243.16	0.0074	0.9053
2A1	Cement production	CO ₂	4,029.11	4,029.11	0.0070	0.9123
1B2	Oil and gas extraction	CO ₂	3,789.72	3,789.72	0.0066	0.9189
5D	Wastewater treatment and discharge	CH ₄	3,359.33	3,359.33	0.0058	0.9247
1A4	Other sectors: solid fuels	CO ₂	3,147.53	3,147.53	0.0055	0.9302
2B8	Petrochemical and carbon black production	CO ₂	2,948.24	2,948.24	0.0051	0.9353
3B1	Manure management from Cattle	CH ₄	2,549.91	2,549.91	0.0044	0.9397
1A5	Other: liquid fuels	CO ₂	2,285.42	2,285.42	0.0040	0.9437
1A3d	Domestic Navigation: liquid fuels	CO ₂	2,165.99	2,165.99	0.0038	0.9475
2F4	Aerosols	HFCs, PFCs, SF ₆ and NF ₃	2,158.68	2,158.68	0.0037	0.9512
1A1	Energy industries: other fuels	CO ₂	2,040.24	2,040.24	0.0035	0.9548
1A3a	Domestic aviation: liquid fuels	CO ₂	1,961.73	1,961.73	0.0034	0.9582
1A3c	Railways: liquid fuels	CO ₂	1,925.85	1,925.85	0.0033	0.9615
1B1	Coal mining and handling	CH ₄	1,682.41	1,682.41	0.0029	0.9644
2B1	Ammonia production	CO ₂	1,383.38	1,383.38	0.0024	0.9668
2A2	Lime production	CO ₂	1,239.15	1,239.15	0.0022	0.9690

IPCC Code	IPCC Category	Greenhouse Gas	2013 emissions (Gg CO ₂ e)	Absolute value of 2013 emissions (Gg CO ₂ e)	Level Assessment	Cumulative Total
5D	Wastewater treatment and discharge	N ₂ O	1,063.03	1,063.03	0.0018	0.9708
2D	Non-energy products from fuels and solvent use	CO ₂	1,032.92	1,032.92	0.0018	0.9726
1A1	Energy industries: solid fuels	N ₂ O	935.10	935.10	0.0016	0.9743
1A3b	Road transportation: liquid fuels	N ₂ O	909.33	909.33	0.0016	0.9758
3B1	Manure management from Cattle	N ₂ O	897.98	897.98	0.0016	0.9774
3B4	Manure management from Other livestock	N ₂ O	837.18	837.18	0.0015	0.9789
3G	Liming	CO ₂	774.69	774.69	0.0013	0.9802
2A4	Other process uses of carbonates	CO ₂	771.83	771.83	0.0013	0.9815
5B	Biological treatment of solid waste	CH ₄	715.72	715.72	0.0012	0.9828
3B3	Manure management from Swine	CH ₄	633.80	633.80	0.0011	0.9839
5B	Biological treatment of solid waste	N ₂ O	608.32	608.32	0.0011	0.9849
1A2	Manufacturing industries and construction: liquid fuels	N ₂ O	538.92	538.92	0.0009	0.9859
2F2	Foam blowing agents	HFCs, PFCs, SF ₆ and NF ₃	495.71	495.71	0.0009	0.9867
3A4	Enteric fermentation from Other livestock	CH ₄	491.61	491.61	0.0009	0.9876
1A4	Other sectors: liquid fuels	N ₂ O	475.43	475.43	0.0008	0.9884
1A3e	Other transportation: liquid fuels	CO ₂	458.95	458.95	0.0008	0.9892
2A3	Glass production	CO ₂	389.18	389.18	0.0007	0.9899
1A2	Manufacturing industries and construction: other fuels	CO ₂	386.12	386.12	0.0007	0.9906
2G1	Electrical equipment	HFCs, PFCs, SF ₆ and NF ₃	318.54	318.54	0.0006	0.9911
2B7	Soda ash production	CO ₂	285.27	285.27	0.0005	0.9916
2G2	SF ₆ and PFCs from other product use	HFCs, PFCs, SF ₆ and NF ₃	271.13	271.13	0.0005	0.9921
2F3	Fire protection	HFCs, PFCs, SF ₆ and NF ₃	269.86	269.86	0.0005	0.9926
5C	Incineration and open burning of waste	CO ₂	264.30	264.30	0.0005	0.9930
3H	Urea application to land	CO ₂	245.22	245.22	0.0004	0.9934

IPCC Code	IPCC Category	Greenhouse Gas	2013 emissions (Gg CO ₂ e)	Absolute value of 2013 emissions (Gg CO ₂ e)	Level Assessment	Cumulative Total
1A4	Other sectors: biomass	CH ₄	238.17	238.17	0.0004	0.9939
1A1	Energy industries: gaseous fuels	N ₂ O	221.69	221.69	0.0004	0.9942
1A4	Other sectors: solid fuels	CH ₄	208.43	208.43	0.0004	0.9946
1A4	Other sectors: gaseous fuels	CH ₄	191.56	191.56	0.0003	0.9949
3A3	Enteric fermentation from Swine	CH ₄	183.08	183.08	0.0003	0.9953
1A2	Manufacturing industries and construction: solid fuels	N ₂ O	171.54	171.54	0.0003	0.9956
3B2	Manure management from Sheep	CH ₄	164.91	164.91	0.0003	0.9958
2B9	Fluorochemical production	HFCs, PFCs, SF ₆ and NF ₃	158.24	158.24	0.0003	0.9961
2C4	Magnesium production	HFCs, PFCs, SF ₆ and NF ₃	148.80	148.80	0.0003	0.9964
1B1	Coal mining and handling solid fuels	CO ₂	133.60	133.60	0.0002	0.9966
2B6	Titanium dioxide production	CO ₂	123.43	123.43	0.0002	0.9968
3B4	Manure management from Other livestock	CH ₄	122.88	122.88	0.0002	0.9970
3B3	Manure management from Swine	N ₂ O	121.88	121.88	0.0002	0.9972
2F6	Other product uses as substitutes for ODS	HFCs, PFCs, SF ₆ and NF ₃	110.00	110.00	0.0002	0.9974
1A1	Energy industries: gaseous fuels	CH ₄	95.30	95.30	0.0002	0.9976
1A1	Energy industries: liquid fuels	N ₂ O	93.43	93.43	0.0002	0.9978
2C3	Aluminium production	CO ₂	68.14	68.14	0.0001	0.9979
1A1	Energy industries: biomass	N ₂ O	66.71	66.71	0.0001	0.9980
2B10	Other Chemical Industry	CH ₄	66.42	66.42	0.0001	0.9981
1A3e	Other transportation: liquid fuels	N ₂ O	56.82	56.82	0.0001	0.9982
1A3b	Road transportation: liquid fuels	CH ₄	55.62	55.62	0.0001	0.9983
1A1	Energy industries: other fuels	N ₂ O	48.24	48.24	0.0001	0.9984
1A4	Other sectors: peat	CO ₂	46.96	46.96	0.0001	0.9985
1A4	Other sectors: gaseous fuels	N ₂ O	45.67	45.67	0.0001	0.9986
2C1	Iron and steel production	N ₂ O	45.23	45.23	0.0001	0.9986
5C	Incineration and open burning of waste	N ₂ O	44.28	44.28	0.0001	0.9987

IPCC Code	IPCC Category	Greenhouse Gas	2013 emissions (Gg CO ₂ e)	Absolute value of 2013 emissions (Gg CO ₂ e)	Level Assessment	Cumulative Total
1A4	Other sectors: solid fuels	N ₂ O	43.36	43.36	0.0001	0.9988
2B2	Nitric acid production	N ₂ O	42.91	42.91	0.0001	0.9989
1A1	Energy industries: biomass	CH ₄	42.77	42.77	0.0001	0.9989
1B1	Coal mining and handling liquid fuels	CO ₂	41.68	41.68	0.0001	0.9990
1A2	Manufacturing industries and construction: biomass	N ₂ O	39.02	39.02	0.0001	0.9991
1A1	Energy industries: other fuels	CH ₄	37.57	37.57	0.0001	0.9991
2B8	Petrochemical and carbon black production	CH ₄	37.45	37.45	0.0001	0.9992
1A4	Other sectors: biomass	N ₂ O	36.24	36.24	0.0001	0.9993
3B2	Manure management from Sheep	N ₂ O	34.24	34.24	0.0001	0.9993
1A4	Other sectors: liquid fuels	CH ₄	33.31	33.31	0.0001	0.9994
1A3c	Railways: solid fuels	CO ₂	32.60	32.60	0.0001	0.9994
1B2	Oil and gas extraction	N ₂ O	30.79	30.79	0.0001	0.9995
1A2	Manufacturing industries and construction: liquid fuels	CH ₄	26.69	26.69	0.0000	0.9995
1A1	Energy industries: solid fuels	CH ₄	25.28	25.28	0.0000	0.9996
1A2	Manufacturing industries and construction: biomass	CH ₄	24.61	24.61	0.0000	0.9996
2F5	Solvents	HFCs, PFCs, SF ₆ and NF ₃	21.93	21.93	0.0000	0.9997
2C1	Iron and steel production	CH ₄	20.49	20.49	0.0000	0.9997
1A5	Other: liquid fuels	N ₂ O	20.30	20.30	0.0000	0.9997
1A3a	Domestic aviation: liquid fuels	N ₂ O	18.56	18.56	0.0000	0.9998
1A1	Energy industries: liquid fuels	CH ₄	17.70	17.70	0.0000	0.9998
1A3d	Domestic Navigation: liquid fuels	N ₂ O	16.22	16.22	0.0000	0.9998
2E1	Integrated circuit or semiconductor	HFCs, PFCs, SF ₆ and NF ₃	14.83	14.83	0.0000	0.9998
1A2	Manufacturing industries and construction: gaseous fuels	N ₂ O	11.96	11.96	0.0000	0.9999
1A2	Manufacturing industries and construction: gaseous fuels	CH ₄	10.04	10.04	0.0000	0.9999
5C	Incineration and open burning of waste	CH ₄	9.95	9.95	0.0000	0.9999

IPCC Code	IPCC Category	Greenhouse Gas	2013 emissions (Gg CO ₂ e)	Absolute value of 2013 emissions (Gg CO ₂ e)	Level Assessment	Cumulative Total
2C3	Aluminium production	HFCs, PFCs, SF ₆ and NF ₃	6.95	6.95	0.0000	0.9999
1A2	Manufacturing industries and construction: other fuels	N ₂ O	6.55	6.55	0.0000	0.9999
1A2	Manufacturing industries and construction: solid fuels	CH ₄	5.94	5.94	0.0000	0.9999
1A3d	Domestic Navigation: liquid fuels	CH ₄	4.90	4.90	0.0000	0.9999
2A4	Other process uses of carbonates	CH ₄	4.52	4.52	0.0000	1.0000
1A3c	Railways: liquid fuels	N ₂ O	4.32	4.32	0.0000	1.0000
1A2	Manufacturing industries and construction: other fuels	CH ₄	4.12	4.12	0.0000	1.0000
1B1	Coal mining and handling biomass	CH ₄	3.82	3.82	0.0000	1.0000
1A4	Other sectors: peat	CH ₄	3.32	3.32	0.0000	1.0000
1A3c	Railways: liquid fuels	CH ₄	2.12	2.12	0.0000	1.0000
1A5	Other: liquid fuels	CH ₄	1.59	1.59	0.0000	1.0000
2B8	Petrochemical and carbon black production	N ₂ O	1.53	1.53	0.0000	1.0000
2D	Non-energy products from fuels and solvent use	N ₂ O	0.91	0.91	0.0000	1.0000
1A3a	Domestic aviation: liquid fuels	CH ₄	0.86	0.86	0.0000	1.0000
1A3c	Railways: solid fuels	CH ₄	0.86	0.86	0.0000	1.0000
2G3	N ₂ O from product uses	N ₂ O	0.49	0.49	0.0000	1.0000
1A3e	Other transportation: liquid fuels	CH ₄	0.48	0.48	0.0000	1.0000
2D	Non-energy products from fuels and solvent use	CH ₄	0.38	0.38	0.0000	1.0000
2B1	Ammonia production	N ₂ O	0.27	0.27	0.0000	1.0000
2B1	Ammonia production	CH ₄	0.23	0.23	0.0000	1.0000
1A4	Other sectors: peat	N ₂ O	0.18	0.18	0.0000	1.0000
1A3c	Railways: solid fuels	N ₂ O	0.08	0.08	0.0000	1.0000
1B1	Coal mining and handling solid fuels	N ₂ O	0.04	0.04	0.0000	1.0000
1B1	Coal mining and handling solid fuels	CH ₄	0.03	0.03	0.0000	1.0000
Total			575,695.98	575,695.98	1.0000	

Table A 1.3.5 Approach 1 Key Category Analysis based on trend in emissions (from base year to latest reported year, including LULUCF)

IPCC Code	IPCC Category	Greenhouse Gas	Base year emissions (Gg CO ₂ e)	2013 emissions (Gg CO ₂ e)	Trend Assessment	Contribution to Trend (%)	Cumulative Total
1A1	Energy industries: gaseous fuels	CO ₂	9,179.46	44,618.23	0.0447	0.1331	0.1331
1A4	Other sectors: gaseous fuels	CO ₂	70,384.62	87,203.06	0.0444	0.1322	0.2653
1A3b	Road transportation: liquid fuels	CO ₂	108,570.51	107,048.58	0.0364	0.1082	0.3735
5A	Solid waste disposal	CH ₄	62,749.72	16,681.83	0.0319	0.0949	0.4684
1A1	Energy industries: solid fuels	CO ₂	185,494.19	113,205.74	0.0194	0.0579	0.5262
2B3	Adipic acid production	N ₂ O	19,934.61	-	0.0163	0.0486	0.5748
1B1	Coal mining and handling	CH ₄	21,788.17	1,682.41	0.0159	0.0473	0.6221
2F1	Refrigeration and air conditioning	HFCs, PFCs, SF ₆ and NF ₃	855.96	13,144.55	0.0147	0.0437	0.6658
2B9	Fluorochemical production	HFCs, PFCs, SF ₆ and NF ₃	17,793.94	158.24	0.0144	0.0428	0.7086
1A1	Energy industries: liquid fuels	CO ₂	40,878.89	17,608.72	0.0129	0.0383	0.7470
1A4	Other sectors: solid fuels	CO ₂	19,876.40	3,147.53	0.0126	0.0375	0.7845
1A2	Manufacturing industries and construction: liquid fuels	CO ₂	29,119.63	12,554.23	0.0092	0.0273	0.8117
1A2	Manufacturing industries and construction: solid fuels	CO ₂	38,947.00	20,641.24	0.0077	0.0230	0.8348
4A	Forest land	CO ₂	-16,016.07	-17,297.63	0.0041	0.0123	0.8470
3A1	Enteric fermentation from Cattle	CH ₄	21,921.41	18,703.95	0.0039	0.0117	0.8588
3D	Agricultural soils	N ₂ O	23,153.96	19,513.90	0.0039	0.0115	0.8703
1A2	Manufacturing industries and construction: gaseous fuels	CO ₂	28,170.48	22,842.35	0.0037	0.0109	0.8812
1B2	Oil and gas extraction	CH ₄	12,332.94	5,687.26	0.0034	0.0103	0.8915
2B2	Nitric acid production	N ₂ O	3,860.26	42.91	0.0031	0.0093	0.9007
1A1	Energy industries: other fuels	CO ₂	226.99	2,040.24	0.0022	0.0066	0.9073

IPCC Code	IPCC Category	Greenhouse Gas	Base year emissions (Gg CO ₂ e)	2013 emissions (Gg CO ₂ e)	Trend Assessment	Contribution to Trend (%)	Cumulative Total
2F4	Aerosols	HFCs, PFCs, SF ₆ and NF ₃	663.14	2,158.68	0.0020	0.0059	0.9132
4B	Cropland	CO ₂	15,130.79	12,150.39	0.0018	0.0054	0.9186
1A5	Other: liquid fuels	CO ₂	5,284.82	2,285.42	0.0017	0.0049	0.9235
4G	Harvested wood products	CO ₂	40.72	-1,121.03	0.0013	0.0040	0.9275
4C	Grassland	CO ₂	-3,677.20	-5,897.14	0.0013	0.0039	0.9314
2A1	Cement production	CO ₂	7,295.26	4,029.11	0.0013	0.0037	0.9352
1B1	Coal mining and handling solid fuels	CO ₂	1,698.56	133.60	0.0012	0.0037	0.9389
2C1	Iron and steel production	CO ₂	5,582.62	4,945.17	0.0012	0.0036	0.9425
4E	Settlements	CO ₂	6,919.10	5,876.98	0.0012	0.0036	0.9461
2C6	Zinc production	CO ₂	1,358.83	-	0.0011	0.0033	0.9494
1A3c	Railways: liquid fuels	CO ₂	1,455.18	1,925.85	0.0011	0.0032	0.9526
1A4	Other sectors: solid fuels	CH ₄	1,477.49	208.43	0.0010	0.0029	0.9554
5B	Biological treatment of solid waste	CH ₄	5.48	715.72	0.0008	0.0025	0.9579
1A3a	Domestic aviation: liquid fuels	CO ₂	1,859.57	1,961.73	0.0008	0.0023	0.9602
1A3d	Domestic Navigation: liquid fuels	CO ₂	2,168.02	2,165.99	0.0008	0.0023	0.9625
5C	Incineration and open burning of waste	CO ₂	1,304.38	264.30	0.0008	0.0023	0.9647
5B	Biological treatment of solid waste	N ₂ O	4.90	608.32	0.0007	0.0021	0.9668
1A3b	Road transportation: liquid fuels	CH ₄	758.31	55.62	0.0006	0.0017	0.9685
1A4	Other sectors: liquid fuels	CO ₂	18,559.20	12,541.14	0.0005	0.0016	0.9700
5D	Wastewater treatment and discharge	CH ₄	4,169.64	3,359.33	0.0005	0.0015	0.9716
3B1	Manure management from Cattle	CH ₄	3,072.25	2,549.91	0.0005	0.0014	0.9730
1A2	Manufacturing industries and construction: other fuels	CO ₂	1.07	386.12	0.0005	0.0013	0.9743

IPCC Code	IPCC Category	Greenhouse Gas	Base year emissions (Gg CO ₂ e)	2013 emissions (Gg CO ₂ e)	Trend Assessment	Contribution to Trend (%)	Cumulative Total
2F2	Foam blowing agents	HFCs, PFCs, SF ₆ and NF ₃	184.55	495.71	0.0004	0.0013	0.9756
3G	Liming	CO ₂	1,579.17	774.69	0.0004	0.0012	0.9768
2A4	Other process uses of carbonates	CO ₂	646.76	771.83	0.0004	0.0011	0.9779
1A3e	Other transportation: liquid fuels	CO ₂	224.74	458.95	0.0004	0.0011	0.9789
1A4	Other sectors: peat	CO ₂	475.59	46.96	0.0003	0.0010	0.9799
5D	Wastewater treatment and discharge	N ₂ O	1,111.13	1,063.03	0.0003	0.0010	0.9809
3A4	Enteric fermentation from Other livestock	CH ₄	295.24	491.61	0.0003	0.0010	0.9819
2F3	Fire protection	HFCs, PFCs, SF ₆ and NF ₃	1.42	269.86	0.0003	0.0009	0.9828
1B2	Oil and gas extraction	CO ₂	5,777.92	3,789.72	0.0003	0.0009	0.9837
2C3	Aluminium production	CO ₂	450.32	68.14	0.0003	0.0009	0.9846
3A2	Enteric fermentation from Sheep	CH ₄	5,727.52	4,243.16	0.0003	0.0008	0.9854
2C3	Aluminium production	HFCs, PFCs, SF ₆ and NF ₃	333.43	6.95	0.0003	0.0008	0.9862
2A2	Lime production	CO ₂	1,462.05	1,239.15	0.0003	0.0008	0.9869
2G1	Electrical equipment	HFCs, PFCs, SF ₆ and NF ₃	747.79	318.54	0.0002	0.0007	0.9877
1A4	Other sectors: biomass	CH ₄	75.42	238.17	0.0002	0.0006	0.9883
2D	Non-energy products from fuels and solvent use	CO ₂	1,232.62	1,032.92	0.0002	0.0006	0.9889
1A1	Energy industries: solid fuels	N ₂ O	1,553.80	935.10	0.0002	0.0005	0.9894
1A4	Other sectors: solid fuels	N ₂ O	277.04	43.36	0.0002	0.0005	0.9899
3F	Field burning of agricultural residues	CH ₄	205.48	-	0.0002	0.0005	0.9904
3B3	Manure management from Swine	CH ₄	1,091.66	633.80	0.0002	0.0005	0.9909
4B	Cropland	N ₂ O	649.15	329.94	0.0001	0.0004	0.9913

IPCC Code	IPCC Category	Greenhouse Gas	Base year emissions (Gg CO ₂ e)	2013 emissions (Gg CO ₂ e)	Trend Assessment	Contribution to Trend (%)	Cumulative Total
2B8	Petrochemical and carbon black production	CO ₂	4,036.08	2,948.24	0.0001	0.0004	0.9918
2B7	Soda ash production	CO ₂	231.55	285.27	0.0001	0.0004	0.9922
3B1	Manure management from Cattle	N ₂ O	1,107.19	897.98	0.0001	0.0004	0.9926
2C4	Magnesium production	HFCs, PFCs, SF ₆ and NF ₃	387.17	148.80	0.0001	0.0004	0.9930
1A3b	Road transportation: liquid fuels	N ₂ O	1,142.41	909.33	0.0001	0.0004	0.9934
2A3	Glass production	CO ₂	408.14	389.18	0.0001	0.0004	0.9938
3B4	Manure management from Other livestock	N ₂ O	1,053.80	837.18	0.0001	0.0003	0.9941
1A2	Manufacturing industries and construction: solid fuels	N ₂ O	387.27	171.54	0.0001	0.0003	0.9945
1A2	Manufacturing industries and construction: liquid fuels	N ₂ O	910.70	538.92	0.0001	0.0003	0.9948
5C	Incineration and open burning of waste	CH ₄	137.22	9.95	0.0001	0.0003	0.9951
1A1	Energy industries: gaseous fuels	N ₂ O	198.68	221.69	0.0001	0.0003	0.9954
1A4	Other sectors: gaseous fuels	CH ₄	157.20	191.56	0.0001	0.0003	0.9957
2G2	SF ₆ and PFCs from other product use	HFCs, PFCs, SF ₆ and NF ₃	278.58	271.13	0.0001	0.0003	0.9960
1A1	Energy industries: biomass	N ₂ O	0.25	66.71	0.0001	0.0002	0.9962
2B10	Other Chemical Industry	CH ₄	185.65	66.42	0.0001	0.0002	0.9964
3B4	Manure management from Other livestock	CH ₄	86.40	122.88	0.0001	0.0002	0.9966
4E	Settlements	N ₂ O	369.67	318.07	0.0001	0.0002	0.9968
1A4	Other sectors: liquid fuels	N ₂ O	595.99	475.43	0.0001	0.0002	0.9970
2B6	Titanium dioxide production	CO ₂	104.63	123.43	0.0001	0.0002	0.9972
3F	Field burning of agricultural residues	N ₂ O	63.50	-	0.0001	0.0002	0.9974

IPCC Code	IPCC Category	Greenhouse Gas	Base year emissions (Gg CO ₂ e)	2013 emissions (Gg CO ₂ e)	Trend Assessment	Contribution to Trend (%)	Cumulative Total
1A1	Energy industries: other fuels	N ₂ O	5.96	48.24	0.0001	0.0002	0.9975
1A1	Energy industries: biomass	CH ₄	0.47	42.77	0.0000	0.0001	0.9977
1B1	Coal mining and handling liquid fuels	CO ₂	-	41.68	0.0000	0.0001	0.9978
1A3e	Other transportation: liquid fuels	N ₂ O	27.86	56.82	0.0000	0.0001	0.9980
4D	Wetlands	CO ₂	481.73	300.49	0.0000	0.0001	0.9981
1A3c	Railways: solid fuels	CO ₂	-	32.60	0.0000	0.0001	0.9982
1A2	Manufacturing industries and construction: biomass	N ₂ O	10.44	39.02	0.0000	0.0001	0.9983
1A1	Energy industries: gaseous fuels	CH ₄	92.19	95.30	0.0000	0.0001	0.9984
1A4	Other sectors: biomass	N ₂ O	11.20	36.24	0.0000	0.0001	0.9985
1A1	Energy industries: other fuels	CH ₄	13.57	37.57	0.0000	0.0001	0.9986
3H	Urea application to land	CO ₂	385.46	245.22	0.0000	0.0001	0.9987
5C	Incineration and open burning of waste	N ₂ O	29.85	44.28	0.0000	0.0001	0.9988
2F5	Solvents	HFCs, PFCs, SF ₆ and NF ₃	-	21.93	0.0000	0.0001	0.9989
4C	Grassland	CH ₄	11.29	28.46	0.0000	0.0001	0.9989
1A4	Other sectors: peat	CH ₄	33.65	3.32	0.0000	0.0001	0.9990
1A2	Manufacturing industries and construction: biomass	CH ₄	6.57	24.61	0.0000	0.0001	0.9991
1A4	Other sectors: gaseous fuels	N ₂ O	37.48	45.67	0.0000	0.0001	0.9991
2B1	Ammonia production	CO ₂	2,004.50	1,383.38	0.0000	0.0001	0.9992
2B8	Petrochemical and carbon black production	CH ₄	27.79	37.45	0.0000	0.0001	0.9993
2A4	Other process uses of carbonates	CH ₄	31.27	4.52	0.0000	0.0001	0.9993
4A	Forest land	N ₂ O	48.26	50.29	0.0000	0.0001	0.9994
1A1	Energy industries: liquid fuels	N ₂ O	156.12	93.43	0.0000	0.0001	0.9994

IPCC Code	IPCC Category	Greenhouse Gas	Base year emissions (Gg CO _{2e})	2013 emissions (Gg CO _{2e})	Trend Assessment	Contribution to Trend (%)	Cumulative Total
3B3	Manure management from Swine	N ₂ O	196.57	121.88	0.0000	0.0001	0.9995
3A3	Enteric fermentation from Swine	CH ₄	283.32	183.08	0.0000	0.0001	0.9995
1A1	Energy industries: liquid fuels	CH ₄	43.71	17.70	0.0000	0.0000	0.9996
1A5	Other: liquid fuels	N ₂ O	47.24	20.30	0.0000	0.0000	0.9996
3B2	Manure management from Sheep	CH ₄	222.48	164.91	0.0000	0.0000	0.9997
1A2	Manufacturing industries and construction: solid fuels	CH ₄	21.01	5.94	0.0000	0.0000	0.9997
2E1	Integrated circuit or semiconductor	HFCs, PFCs, SF ₆ and NF ₃	9.56	14.83	0.0000	0.0000	0.9997
1A2	Manufacturing industries and construction: other fuels	N ₂ O	0.02	6.55	0.0000	0.0000	0.9997
1A4	Other sectors: liquid fuels	CH ₄	56.73	33.31	0.0000	0.0000	0.9998
1A3a	Domestic aviation: liquid fuels	N ₂ O	17.60	18.56	0.0000	0.0000	0.9998
2F6	Other product uses as substitutes for ODS	HFCs, PFCs, SF ₆ and NF ₃	165.54	110.00	0.0000	0.0000	0.9998
2C1	Iron and steel production	CH ₄	36.89	20.49	0.0000	0.0000	0.9998
1A3d	Domestic Navigation: liquid fuels	N ₂ O	16.26	16.22	0.0000	0.0000	0.9998
1A2	Manufacturing industries and construction: other fuels	CH ₄	0.01	4.12	0.0000	0.0000	0.9999
1A1	Energy industries: solid fuels	CH ₄	41.90	25.28	0.0000	0.0000	0.9999
1A2	Manufacturing industries and construction: liquid fuels	CH ₄	43.69	26.69	0.0000	0.0000	0.9999
1B1	Coal mining and handling biomass	CH ₄	0.10	3.82	0.0000	0.0000	0.9999
1A3d	Domestic Navigation: liquid fuels	CH ₄	2.22	4.90	0.0000	0.0000	0.9999
4A	Forest land	CH ₄	3.72	5.04	0.0000	0.0000	0.9999

IPCC Code	IPCC Category	Greenhouse Gas	Base year emissions (Gg CO ₂ e)	2013 emissions (Gg CO ₂ e)	Trend Assessment	Contribution to Trend (%)	Cumulative Total
1B2	Oil and gas extraction	N ₂ O	40.75	30.79	0.0000	0.0000	0.9999
4D	Wetlands	N ₂ O	3.83	0.50	0.0000	0.0000	0.9999
4E	Settlements	CH ₄	5.89	1.99	0.0000	0.0000	0.9999
3B2	Manure management from Sheep	N ₂ O	45.91	34.24	0.0000	0.0000	1.0000
1A3a	Domestic aviation: liquid fuels	CH ₄	4.17	0.86	0.0000	0.0000	1.0000
1A3c	Railways: liquid fuels	N ₂ O	3.27	4.32	0.0000	0.0000	1.0000
2C1	Iron and steel production	N ₂ O	62.46	45.23	0.0000	0.0000	1.0000
1A2	Manufacturing industries and construction: gaseous fuels	N ₂ O	14.99	11.96	0.0000	0.0000	1.0000
1A2	Manufacturing industries and construction: gaseous fuels	CH ₄	12.57	10.04	0.0000	0.0000	1.0000
1A4	Other sectors: peat	N ₂ O	1.87	0.18	0.0000	0.0000	1.0000
1A5	Other: liquid fuels	CH ₄	3.73	1.59	0.0000	0.0000	1.0000
1A3c	Railways: solid fuels	CH ₄	-	0.86	0.0000	0.0000	1.0000
4C	Grassland	N ₂ O	12.10	9.15	0.0000	0.0000	1.0000
1A3c	Railways: liquid fuels	CH ₄	2.46	2.12	0.0000	0.0000	1.0000
1A3e	Other transportation: liquid fuels	CH ₄	0.29	0.48	0.0000	0.0000	1.0000
2B8	Petrochemical and carbon black production	N ₂ O	1.79	1.53	0.0000	0.0000	1.0000
2D	Non-energy products from fuels and solvent use	N ₂ O	1.64	0.91	0.0000	0.0000	1.0000
2G3	N ₂ O from product uses	N ₂ O	0.44	0.49	0.0000	0.0000	1.0000
2D	Non-energy products from fuels and solvent use	CH ₄	0.69	0.38	0.0000	0.0000	1.0000
1A3c	Railways: solid fuels	N ₂ O	-	0.08	0.0000	0.0000	1.0000
2B1	Ammonia production	N ₂ O	0.31	0.27	0.0000	0.0000	1.0000
4B	Cropland	CH ₄	0.11	0.14	0.0000	0.0000	1.0000
2B1	Ammonia production	CH ₄	0.26	0.23	0.0000	0.0000	1.0000
1B1	Coal mining and handling solid fuels	N ₂ O	0.09	0.04	0.0000	0.0000	1.0000
4F	Other land	CO ₂	-	0.02	0.0000	0.0000	1.0000

IPCC Code	IPCC Category	Greenhouse Gas	Base year emissions (Gg CO ₂ e)	2013 emissions (Gg CO ₂ e)	Trend Assessment	Contribution to Trend (%)	Cumulative Total
1B1	Coal mining and handling solid fuels	CH ₄	0.08	0.03	0.0000	0.0000	1.0000
Total			815,143.98	570,451.65	0.3359	1.0000	

Table A 1.3.6 Approach 1 Key Category Analysis based on trend in emissions (from base year to latest reported year, excluding LULUCF)

IPCC Code	IPCC Category	Greenhouse Gas	Base year emissions (Gg CO ₂ e)	2013 emissions (Gg CO ₂ e)	Trend Assessment	Contribution to Trend (%)	Cumulative Total
1A1	Energy industries: gaseous fuels	CO ₂	9,179.46	44,618.23	0.0447	0.1372	0.1372
1A4	Other sectors: gaseous fuels	CO ₂	70,384.62	87,203.06	0.0444	0.1363	0.2735
1A3b	Road transportation: liquid fuels	CO ₂	108,570.51	107,048.58	0.0364	0.1116	0.3851
5A	Solid waste disposal	CH ₄	62,749.72	16,681.83	0.0319	0.0978	0.4829
1A1	Energy industries: solid fuels	CO ₂	185,494.19	113,205.74	0.0194	0.0597	0.5426
2B3	Adipic acid production	N ₂ O	19,934.61	-	0.0163	0.0501	0.5927
1B1	Coal mining and handling	CH ₄	21,788.17	1,682.41	0.0159	0.0487	0.6414
2F1	Refrigeration and air conditioning	HFCs, PFCs, SF ₆ and NF ₃	855.96	13,144.55	0.0147	0.0451	0.6865
2B9	Fluorochemical production	HFCs, PFCs, SF ₆ and NF ₃	17,793.94	158.24	0.0144	0.0442	0.7307
1A1	Energy industries: liquid fuels	CO ₂	40,878.89	17,608.72	0.0129	0.0395	0.7702
1A4	Other sectors: solid fuels	CO ₂	19,876.40	3,147.53	0.0126	0.0387	0.8088
1A2	Manufacturing industries and construction: liquid fuels	CO ₂	29,119.63	12,554.23	0.0092	0.0281	0.8370
1A2	Manufacturing industries and construction: solid fuels	CO ₂	38,947.00	20,641.24	0.0077	0.0238	0.8607
3A1	Enteric fermentation from Cattle	CH ₄	21,921.41	18,703.95	0.0039	0.0121	0.8728

IPCC Code	IPCC Category	Greenhouse Gas	Base year emissions (Gg CO ₂ e)	2013 emissions (Gg CO ₂ e)	Trend Assessment	Contribution to Trend (%)	Cumulative Total
3D	Agricultural soils	N ₂ O	23,153.96	19,513.90	0.0039	0.0119	0.8847
1A2	Manufacturing industries and construction: gaseous fuels	CO ₂	28,170.48	22,842.35	0.0037	0.0112	0.8959
1B2	Oil and gas extraction	CH ₄	12,332.94	5,687.26	0.0034	0.0106	0.9065
2B2	Nitric acid production	N ₂ O	3,860.26	42.91	0.0031	0.0096	0.9160
1A1	Energy industries: other fuels	CO ₂	226.99	2,040.24	0.0022	0.0068	0.9228
2F4	Aerosols	HFCs, PFCs, SF ₆ and NF ₃	663.14	2,158.68	0.0020	0.0061	0.9289
1A5	Other: liquid fuels	CO ₂	5,284.82	2,285.42	0.0017	0.0051	0.9340
2A1	Cement production	CO ₂	7,295.26	4,029.11	0.0013	0.0039	0.9378
1B1	Coal mining and handling solid fuels	CO ₂	1,698.56	133.60	0.0012	0.0038	0.9416
2C1	Iron and steel production	CO ₂	5,582.62	4,945.17	0.0012	0.0037	0.9454
2C6	Zinc production	CO ₂	1,358.83	-	0.0011	0.0034	0.9488
1A3c	Railways: liquid fuels	CO ₂	1,455.18	1,925.85	0.0011	0.0033	0.9520
1A4	Other sectors: solid fuels	CH ₄	1,477.49	208.43	0.0010	0.0030	0.9550
5B	Biological treatment of solid waste	CH ₄	5.48	715.72	0.0008	0.0026	0.9576
1A3a	Domestic aviation: liquid fuels	CO ₂	1,859.57	1,961.73	0.0008	0.0024	0.9599
1A3d	Domestic Navigation: liquid fuels	CO ₂	2,168.02	2,165.99	0.0008	0.0023	0.9623
5C	Incineration and open burning of waste	CO ₂	1,304.38	264.30	0.0008	0.0023	0.9646
5B	Biological treatment of solid waste	N ₂ O	4.90	608.32	0.0007	0.0022	0.9668
1A3b	Road transportation: liquid fuels	CH ₄	758.31	55.62	0.0006	0.0017	0.9685
1A4	Other sectors: liquid fuels	CO ₂	18,559.20	12,541.14	0.0005	0.0016	0.9701
5D	Wastewater treatment and discharge	CH ₄	4,169.64	3,359.33	0.0005	0.0016	0.9717
3B1	Manure management from Cattle	CH ₄	3,072.25	2,549.91	0.0005	0.0014	0.9731

IPCC Code	IPCC Category	Greenhouse Gas	Base year emissions (Gg CO ₂ e)	2013 emissions (Gg CO ₂ e)	Trend Assessment	Contribution to Trend (%)	Cumulative Total
1A2	Manufacturing industries and construction: other fuels	CO ₂	1.07	386.12	0.0005	0.0014	0.9745
2F2	Foam blowing agents	HFCs, PFCs, SF ₆ and NF ₃	184.55	495.71	0.0004	0.0013	0.9758
3G	Liming	CO ₂	1,579.17	774.69	0.0004	0.0012	0.9770
2A4	Other process uses of carbonates	CO ₂	646.76	771.83	0.0004	0.0011	0.9781
1A3e	Other transportation: liquid fuels	CO ₂	224.74	458.95	0.0004	0.0011	0.9792
1A4	Other sectors: peat	CO ₂	475.59	46.96	0.0003	0.0010	0.9802
5D	Wastewater treatment and discharge	N ₂ O	1,111.13	1,063.03	0.0003	0.0010	0.9813
3A4	Enteric fermentation from Other livestock	CH ₄	295.24	491.61	0.0003	0.0010	0.9823
2F3	Fire protection	HFCs, PFCs, SF ₆ and NF ₃	1.42	269.86	0.0003	0.0010	0.9833
1B2	Oil and gas extraction	CO ₂	5,777.92	3,789.72	0.0003	0.0009	0.9842
2C3	Aluminium production	CO ₂	450.32	68.14	0.0003	0.0009	0.9851
3A2	Enteric fermentation from Sheep	CH ₄	5,727.52	4,243.16	0.0003	0.0008	0.9859
2C3	Aluminium production	HFCs, PFCs, SF ₆ and NF ₃	333.43	6.95	0.0003	0.0008	0.9867
2A2	Lime production	CO ₂	1,462.05	1,239.15	0.0003	0.0008	0.9875
2G1	Electrical equipment	HFCs, PFCs, SF ₆ and NF ₃	747.79	318.54	0.0002	0.0007	0.9882
1A4	Other sectors: biomass	CH ₄	75.42	238.17	0.0002	0.0007	0.9889
2D	Non-energy products from fuels and solvent use	CO ₂	1,232.62	1,032.92	0.0002	0.0006	0.9895
1A1	Energy industries: solid fuels	N ₂ O	1,553.80	935.10	0.0002	0.0005	0.9900
1A4	Other sectors: solid fuels	N ₂ O	277.04	43.36	0.0002	0.0005	0.9906
3F	Field burning of agricultural residues	CH ₄	205.48	-	0.0002	0.0005	0.9911

IPCC Code	IPCC Category	Greenhouse Gas	Base year emissions (Gg CO ₂ e)	2013 emissions (Gg CO ₂ e)	Trend Assessment	Contribution to Trend (%)	Cumulative Total
3B3	Manure management from Swine	CH ₄	1,091.66	633.80	0.0002	0.0005	0.9916
2B8	Petrochemical and carbon black production	CO ₂	4,036.08	2,948.24	0.0001	0.0004	0.9920
2B7	Soda ash production	CO ₂	231.55	285.27	0.0001	0.0004	0.9925
3B1	Manure management from Cattle	N ₂ O	1,107.19	897.98	0.0001	0.0004	0.9929
2C4	Magnesium production	HFCs, PFCs, SF ₆ and NF ₃	387.17	148.80	0.0001	0.0004	0.9933
1A3b	Road transportation: liquid fuels	N ₂ O	1,142.41	909.33	0.0001	0.0004	0.9937
2A3	Glass production	CO ₂	408.14	389.18	0.0001	0.0004	0.9941
3B4	Manure management from Other livestock	N ₂ O	1,053.80	837.18	0.0001	0.0004	0.9945
1A2	Manufacturing industries and construction: solid fuels	N ₂ O	387.27	171.54	0.0001	0.0004	0.9948
1A2	Manufacturing industries and construction: liquid fuels	N ₂ O	910.70	538.92	0.0001	0.0004	0.9952
5C	Incineration and open burning of waste	CH ₄	137.22	9.95	0.0001	0.0003	0.9955
1A1	Energy industries: gaseous fuels	N ₂ O	198.68	221.69	0.0001	0.0003	0.9958
1A4	Other sectors: gaseous fuels	CH ₄	157.20	191.56	0.0001	0.0003	0.9961
2G2	SF ₆ and PFCs from other product use	HFCs, PFCs, SF ₆ and NF ₃	278.58	271.13	0.0001	0.0003	0.9963
1A1	Energy industries: biomass	N ₂ O	0.25	66.71	0.0001	0.0002	0.9966
2B10	Other Chemical Industry	CH ₄	185.65	66.42	0.0001	0.0002	0.9968
3B4	Manure management from Other livestock	CH ₄	86.40	122.88	0.0001	0.0002	0.9970
1A4	Other sectors: liquid fuels	N ₂ O	595.99	475.43	0.0001	0.0002	0.9972
2B6	Titanium dioxide production	CO ₂	104.63	123.43	0.0001	0.0002	0.9974
3F	Field burning of agricultural residues	N ₂ O	63.50	-	0.0001	0.0002	0.9976

IPCC Code	IPCC Category	Greenhouse Gas	Base year emissions (Gg CO ₂ e)	2013 emissions (Gg CO ₂ e)	Trend Assessment	Contribution to Trend (%)	Cumulative Total
1A1	Energy industries: other fuels	N ₂ O	5.96	48.24	0.0001	0.0002	0.9977
1A1	Energy industries: biomass	CH ₄	0.47	42.77	0.0000	0.0002	0.9979
1B1	Coal mining and handling liquid fuels	CO ₂	-	41.68	0.0000	0.0001	0.9980
1A3e	Other transportation: liquid fuels	N ₂ O	27.86	56.82	0.0000	0.0001	0.9982
1A3c	Railways: solid fuels	CO ₂	-	32.60	0.0000	0.0001	0.9983
1A2	Manufacturing industries and construction: biomass	N ₂ O	10.44	39.02	0.0000	0.0001	0.9984
1A1	Energy industries: gaseous fuels	CH ₄	92.19	95.30	0.0000	0.0001	0.9985
1A4	Other sectors: biomass	N ₂ O	11.20	36.24	0.0000	0.0001	0.9986
1A1	Energy industries: other fuels	CH ₄	13.57	37.57	0.0000	0.0001	0.9987
3H	Urea application to land	CO ₂	385.46	245.22	0.0000	0.0001	0.9988
5C	Incineration and open burning of waste	N ₂ O	29.85	44.28	0.0000	0.0001	0.9989
2F5	Solvents	HFCs, PFCs, SF ₆ and NF ₃	-	21.93	0.0000	0.0001	0.9990
1A4	Other sectors: peat	CH ₄	33.65	3.32	0.0000	0.0001	0.9991
1A2	Manufacturing industries and construction: biomass	CH ₄	6.57	24.61	0.0000	0.0001	0.9991
1A4	Other sectors: gaseous fuels	N ₂ O	37.48	45.67	0.0000	0.0001	0.9992
2B1	Ammonia production	CO ₂	2,004.50	1,383.38	0.0000	0.0001	0.9993
2B8	Petrochemical and carbon black production	CH ₄	27.79	37.45	0.0000	0.0001	0.9993
2A4	Other process uses of carbonates	CH ₄	31.27	4.52	0.0000	0.0001	0.9994
1A1	Energy industries: liquid fuels	N ₂ O	156.12	93.43	0.0000	0.0001	0.9994
3B3	Manure management from Swine	N ₂ O	196.57	121.88	0.0000	0.0001	0.9995
3A3	Enteric fermentation from Swine	CH ₄	283.32	183.08	0.0000	0.0001	0.9996

IPCC Code	IPCC Category	Greenhouse Gas	Base year emissions (Gg CO ₂ e)	2013 emissions (Gg CO ₂ e)	Trend Assessment	Contribution to Trend (%)	Cumulative Total
1A1	Energy industries: liquid fuels	CH ₄	43.71	17.70	0.0000	0.0000	0.9996
1A5	Other: liquid fuels	N ₂ O	47.24	20.30	0.0000	0.0000	0.9996
3B2	Manure management from Sheep	CH ₄	222.48	164.91	0.0000	0.0000	0.9997
1A2	Manufacturing industries and construction: solid fuels	CH ₄	21.01	5.94	0.0000	0.0000	0.9997
2E1	Integrated circuit or semiconductor	HFCs, PFCs, SF ₆ and NF ₃	9.56	14.83	0.0000	0.0000	0.9997
1A2	Manufacturing industries and construction: other fuels	N ₂ O	0.02	6.55	0.0000	0.0000	0.9998
1A4	Other sectors: liquid fuels	CH ₄	56.73	33.31	0.0000	0.0000	0.9998
1A3a	Domestic aviation: liquid fuels	N ₂ O	17.60	18.56	0.0000	0.0000	0.9998
2F6	Other product uses as substitutes for ODS	HFCs, PFCs, SF ₆ and NF ₃	165.54	110.00	0.0000	0.0000	0.9998
2C1	Iron and steel production	CH ₄	36.89	20.49	0.0000	0.0000	0.9999
1A3d	Domestic Navigation: liquid fuels	N ₂ O	16.26	16.22	0.0000	0.0000	0.9999
1A2	Manufacturing industries and construction: other fuels	CH ₄	0.01	4.12	0.0000	0.0000	0.9999
1A1	Energy industries: solid fuels	CH ₄	41.90	25.28	0.0000	0.0000	0.9999
1A2	Manufacturing industries and construction: liquid fuels	CH ₄	43.69	26.69	0.0000	0.0000	0.9999
1B1	Coal mining and handling biomass	CH ₄	0.10	3.82	0.0000	0.0000	0.9999
1A3d	Domestic Navigation: liquid fuels	CH ₄	2.22	4.90	0.0000	0.0000	0.9999
1B2	Oil and gas extraction	N ₂ O	40.75	30.79	0.0000	0.0000	0.9999
3B2	Manure management from Sheep	N ₂ O	45.91	34.24	0.0000	0.0000	1.0000

IPCC Code	IPCC Category	Greenhouse Gas	Base year emissions (Gg CO ₂ e)	2013 emissions (Gg CO ₂ e)	Trend Assessment	Contribution to Trend (%)	Cumulative Total
1A3a	Domestic aviation: liquid fuels	CH ₄	4.17	0.86	0.0000	0.0000	1.0000
1A3c	Railways: liquid fuels	N ₂ O	3.27	4.32	0.0000	0.0000	1.0000
2C1	Iron and steel production	N ₂ O	62.46	45.23	0.0000	0.0000	1.0000
1A2	Manufacturing industries and construction: gaseous fuels	N ₂ O	14.99	11.96	0.0000	0.0000	1.0000
1A2	Manufacturing industries and construction: gaseous fuels	CH ₄	12.57	10.04	0.0000	0.0000	1.0000
1A4	Other sectors: peat	N ₂ O	1.87	0.18	0.0000	0.0000	1.0000
1A5	Other: liquid fuels	CH ₄	3.73	1.59	0.0000	0.0000	1.0000
1A3c	Railways: solid fuels	CH ₄	-	0.86	0.0000	0.0000	1.0000
1A3c	Railways: liquid fuels	CH ₄	2.46	2.12	0.0000	0.0000	1.0000
1A3e	Other transportation: liquid fuels	CH ₄	0.29	0.48	0.0000	0.0000	1.0000
2B8	Petrochemical and carbon black production	N ₂ O	1.79	1.53	0.0000	0.0000	1.0000
2D	Non-energy products from fuels and solvent use	N ₂ O	1.64	0.91	0.0000	0.0000	1.0000
2G3	N ₂ O from product uses	N ₂ O	0.44	0.49	0.0000	0.0000	1.0000
2D	Non-energy products from fuels and solvent use	CH ₄	0.69	0.38	0.0000	0.0000	1.0000
1A3c	Railways: solid fuels	N ₂ O	-	0.08	0.0000	0.0000	1.0000
2B1	Ammonia production	N ₂ O	0.31	0.27	0.0000	0.0000	1.0000
2B1	Ammonia production	CH ₄	0.26	0.23	0.0000	0.0000	1.0000
1B1	Coal mining and handling solid fuels	N ₂ O	0.09	0.04	0.0000	0.0000	1.0000
1B1	Coal mining and handling solid fuels	CH ₄	0.08	0.03	0.0000	0.0000	1.0000
Total			811,160.90	575,695.98	0.3258	1.0000	

A 1.4 QUANTITATIVE APPROACH 2 KCA FOLLOWING IPCC 2006 GUIDELINES

Following the 2006 IPCC Guidelines, the UK has also completed an Approach 2 KCA for both level and trend, which takes into account uncertainties, using the Approach 1 method for

uncertainty estimates. This analysis has been performed using the data shown in **Table A 2.3.1** to **Table A 2.3.4** using the same categorisation and the same estimates of uncertainty.

The results of the **level assessment** (based on Approach 2) with and without LULUCF for the base year and the latest reported year are shown in **Table A 1.4.1** to **Table A 1.4.4**. The key source categories are highlighted by the shaded cells in the table. The source categories (i.e. rows of the table) were sorted in descending order of magnitude based on the results of the “Level Parameter”, and then the cumulative total was included in the final column of the table. The key source categories are those whose contributions add up to 90% of the sum of the level parameter in the final column after this sorting process, which accounts for 90% of the uncertainty in level.

The results of the **trend assessment** (based on Approach 2) with and without LULUCF for the base year to the latest reported year to the latest reported year, are shown in **Table A 1.4.5** and **Table A 1.4.6**. The key source categories are highlighted by the shaded cells in the table. The trend parameter was calculated using the absolute value of the result; an absolute function is used since Land Use, Land Use Change and Forestry contains negative sources (sinks) and the absolute function is necessary to produce positive uncertainty contributions for these sinks. The source categories (i.e. rows of the table) were sorted in descending order of magnitude based on the results of the trend parameter, and then the cumulative total was included in the final column of the table. The key source categories are those whose contributions add up to 90% of the sum of the level parameter in the final column after this sorting process, which accounts for 90% of the uncertainty in trend.

Any methodological improvements to the uncertainty analysis are discussed in **Annex 2**.

Table A 1.4.1 Approach 2 Level Assessment for Base year (including LULUCF) with Key Categories Shaded in Grey

IPCC Code	IPCC Category	Greenhouse Gas	Base year emissions (Gg CO ₂ e)	Absolute value of Base year emissions (Gg CO ₂ e)	Level Assessment	Cumul ative Total
3D	3D Agricultural Soils	N ₂ O	23153.96	23153.96	0.3261	0.3261
5A	5A Solid Waste Disposal	CH ₄	62749.72	62749.72	0.1651	0.4912
2B3	2B3 Adipic Acid Production	N ₂ O	19934.61	19934.61	0.1084	0.5996
4B	4B Cropland	CO ₂	15130.79	15130.79	0.0370	0.6366
3B	3B Manure Management	N ₂ O	2403.46	2403.46	0.0332	0.6698
1A	1A (Stationary) Oil	CO ₂	93760.90	93760.90	0.0310	0.7008
3A	3A Enteric Fermentation	CH ₄	28227.49	28227.49	0.0307	0.7315
1A	1A Coal	CO ₂	244317.59	244317.59	0.0284	0.7599
4A	4A Forest Land	CO ₂	-16016.07	16016.07	0.0261	0.7861
1B1	1B1 Coal Mining	CH ₄	21788.17	21788.17	0.0238	0.8099
2B2	2B2 Nitric Acid Production	N ₂ O	3860.26	3860.26	0.0211	0.8310
4E	4E Settlements	CO ₂	6919.10	6919.10	0.0188	0.8498

IPCC Code	IPCC Category	Greenhouse Gas	Base year emissions (Gg CO ₂ e)	Absolute value of Base year emissions (Gg CO ₂ e)	Level Assessment	Cumulative Total
5D	5D Wastewater Handling	N ₂ O	1111.13	1111.13	0.0150	0.8648
1A1 & 1A2 & 1A4 & 1A5	1A1 & 1A2 & 1A4 & 1A5 Other Combustion	N ₂ O	4209.03	4209.03	0.0139	0.8787
1A	1A Natural Gas	CO ₂	107734.56	107734.56	0.0128	0.8915
1B2	1B2 Natural Gas Transmission	CH ₄	10168.29	10168.29	0.0112	0.9027
2B	2B Chemical industry	HFCs	17680.04	17680.04	0.0096	0.9123
1A3b	1A3b Gasoline/ LPG	CO ₂	75557.88	75557.88	0.0091	0.9214
4C	4C Grassland	CO ₂	-3677.20	3677.20	0.0090	0.9304
3B	3B Manure Management	CH ₄	4472.78	4472.78	0.0073	0.9377
2B	2B Chemical industries	CO ₂	6376.75	6376.75	0.0066	0.9443
5D	5D Wastewater Handling	CH ₄	4169.64	4169.64	0.0061	0.9504
1A1 & 1A2 & 1A4 & 1A5	1A1 & 1A2 & 1A4 & 1A5 Other Combustion	CH ₄	2079.93	2079.93	0.0042	0.9546
1A3b	1A3b DERV	CO ₂	33012.63	33012.63	0.0040	0.9586
1A3b	1A3b Gasoline/ LPG	N ₂ O	862.61	862.61	0.0035	0.9620
1A3b	1A3b Gasoline/ LPG	CH ₄	630.55	630.55	0.0025	0.9646
1B2	1B2 Oil & Natural Gas	CO ₂	5777.92	5777.92	0.0022	0.9668
2D	2D Non Energy Products from Fuels and Solvent Use	CO ₂	1232.62	1232.62	0.0022	0.9690
1A3d	1A3d Marine fuel	CO ₂	2168.02	2168.02	0.0021	0.9711
2C	2C Metal Industries	CO ₂	7391.77	7391.77	0.0021	0.9732
1B2	1B2 Offshore Oil& Gas	CH ₄	2164.64	2164.64	0.0020	0.9752
1A3a	1A3a Aviation Fuel	CO ₂	1859.57	1859.57	0.0020	0.9772
1A3b	1A3b DERV	N ₂ O	279.80	279.80	0.0020	0.9792
4B	4B Cropland	N ₂ O	649.15	649.15	0.0019	0.9812
5C	5C Waste Incineration	CO ₂	1304.38	1304.38	0.0018	0.9830

IPCC Code	IPCC Category	Greenhouse Gas	Base year emissions (Gg CO ₂ e)	Absolute value of Base year emissions (Gg CO ₂ e)	Level Assessment	Cumulative Total
3G	3G Liming	CO ₂	1579.17	1579.17	0.0018	0.9847
1A3	1A3 Other diesel	CO ₂	1679.92	1679.92	0.0014	0.9861
4D	4D Wetland	CO ₂	481.73	481.73	0.0013	0.9874
2A1	2A1 Cement Production	CO ₂	7295.26	7295.26	0.0013	0.9887
2F	2F Product Uses as Substitutes for ODS	HFCs	1870.17	1870.17	0.0012	0.9899
3H	3H Urea application to agriculture	CO ₂	385.46	385.46	0.0010	0.9909
1A3b	1A3b DERV	CH ₄	127.77	127.77	0.0009	0.9918
1A4	1A4 Peat	CO ₂	475.59	475.59	0.0008	0.9926
3F	3F Field Burning	N ₂ O	63.50	63.50	0.0008	0.9934
1B1	1B1 Solid Fuel Transformation	CO ₂	1698.56	1698.56	0.0006	0.9941
3F	3F Field Burning	CH ₄	205.48	205.48	0.0006	0.9947
4E	4E Settlements	N ₂ O	369.67	369.67	0.0004	0.9951
2C	2C Iron & Steel	N ₂ O	62.46	62.46	0.0004	0.9955
2A2	2A2 Lime Production	CO ₂	1462.05	1462.05	0.0004	0.9959
2G	2G Other Product Manufacture and Use	PFCs	149.17	149.17	0.0004	0.9963
5C	5C Waste Incineration	CH ₄	137.22	137.22	0.0004	0.9966
5C	5C Waste Incineration	N ₂ O	29.85	29.85	0.0004	0.9970
2C	2C Metal Industries	PFCs	333.43	333.43	0.0004	0.9974
2G	2G Other Product Manufacture and Use	SF ₆	877.20	877.20	0.0003	0.9977
1B2	1B2 Oil & Natural Gas	N ₂ O	40.75	40.75	0.0002	0.9979
2B	2B Chemical Industry	CH ₄	213.70	213.70	0.0002	0.9981
1A3	1A3 Other diesel	N ₂ O	31.12	31.12	0.0002	0.9984
2A	2A Minerals industry	CH ₄	31.27	31.27	0.0002	0.9985
1A	1A Other (waste)	CO ₂	228.07	228.07	0.0002	0.9987
2C	2C Metal Industries	SF ₆	387.17	387.17	0.0001	0.9988
2A4	2A4 Other process uses of carbonates	CO ₂	646.76	646.76	0.0001	0.9990
1A4	1A4 Petroleum Coke	CO ₂	81.64	81.64	0.0001	0.9991
2A3	2A3 Glass production	CO ₂	408.14	408.14	0.0001	0.9992

IPCC Code	IPCC Category	Greenhouse Gas	Base year emissions (Gg CO ₂ e)	Absolute value of Base year emissions (Gg CO ₂ e)	Level Assessment	Cumulative Total
4A	4A Forest land	N ₂ O	48.26	48.26	0.0001	0.9993
1A3a	1A3a Aviation Fuel	N ₂ O	17.60	17.60	0.0001	0.9994
1A3d	1A3d Marine fuel	N ₂ O	16.26	16.26	0.0001	0.9995
4G	4G Other Activities	CO ₂	40.72	40.72	0.0001	0.9996
2C	2C Iron & Steel Production	CH ₄	36.89	36.89	0.0001	0.9997
2B	2B Chemical industry	PFCs	113.90	113.90	0.0001	0.9997
4C	4C Grassland	CH ₄	11.29	11.29	0.0000	0.9998
5B	5B Biological treatment of solid waste	CH ₄	5.48	5.48	0.0000	0.9998
5B	5B Biological treatment of solid waste	N ₂ O	4.90	4.90	0.0000	0.9998
4C	4C Grassland	N ₂ O	12.10	12.10	0.0000	0.9999
2E	2E Electronics Industry	HFCs	8.73	8.73	0.0000	0.9999
4D	4D Grassland	N ₂ O	3.83	3.83	0.0000	0.9999
1A3	1A3 Other diesel	CH ₄	2.75	2.75	0.0000	0.9999
1A3d	1A3d Marine fuel	CH ₄	2.22	2.22	0.0000	0.9999
1A3a	1A3a Aviation Fuel	CH ₄	4.17	4.17	0.0000	1.0000
4A	4A Forest Land	CH ₄	3.72	3.72	0.0000	1.0000
2D	2D Non-energy Products from Fuels and Solvent Use	N ₂ O	1.64	1.64	0.0000	1.0000
2B8	2B8 Petrochemical and Carbon Black Production	N ₂ O	1.79	1.79	0.0000	1.0000
4E	4E Settlements	CH ₄	5.89	5.89	0.0000	1.0000
2G	2G Other Product Manufacture and Use	N ₂ O	0.44	0.44	0.0000	1.0000
2D	2D Non-energy Products from Fuels and Solvent Use	CH ₄	0.69	0.69	0.0000	1.0000
2E	2E Electronics Industry	NF ₃	0.83	0.83	0.0000	1.0000
2B1	2B1 Ammonia Production	N ₂ O	0.31	0.31	0.0000	1.0000
2F	2F Product Uses as Substitutes for ODS	PFCs	0.44	0.44	0.0000	1.0000
1B1	1B1 Fugitive Emissions from Solid Fuels	N ₂ O	0.09	0.09	0.0000	1.0000
1B1	1B1 Solid Fuel Transformation	CH ₄	0.18	0.18	0.0000	1.0000

IPCC Code	IPCC Category	Greenhouse Gas	Base year emissions (Gg CO ₂ e)	Absolute value of Base year emissions (Gg CO ₂ e)	Level Assessment	Cumulative Total
4B	4B Cropland	CH ₄	0.11	0.11	0.0000	1.0000
1A3c	1A3c Coal	CO ₂	0.00	0.00	0.0000	1.0000
4F	4F Other Land	CO ₂	0.00	0.00	0.0000	1.0000
1A3c	1A3c Coal	CH ₄	0.00	0.00	0.0000	1.0000
1A3c	1A3c Coal	N ₂ O	0.00	0.00	0.0000	1.0000
2C	2C Metal Industries	HFCs	0.00	0.00	0.0000	1.0000

Table A 1.4.2 Approach 2 Level Assessment for 2013 (including LULUCF) with Key Categories Shaded in Grey

IPCC Code	IPCC Category	Greenhouse Gas	2013 emissions (Gg CO ₂ e)	Absolute value of 2013 emissions (Gg CO ₂ e)	Level Assessment	Cumulative Total
3D	3D Agricultural Soils	N ₂ O	19513.90	19513.90	0.4450	0.4450
5A	5A Solid Waste Disposal	CH ₄	16681.83	16681.83	0.0711	0.5161
4B	4B Cropland	CO ₂	12150.39	12150.39	0.0482	0.5642
4A	4A Forest Land	CO ₂	-17297.63	17297.63	0.0457	0.6100
3B	3B Manure Management	N ₂ O	1891.28	1891.28	0.0423	0.6523
3A	3A Enteric Fermentation	CH ₄	23621.80	23621.80	0.0416	0.6938
1A	1A Natural Gas	CO ₂	154663.64	154663.64	0.0297	0.7236
4E	4E Settlements	CO ₂	5876.98	5876.98	0.0259	0.7495
1A	1A Coal	CO ₂	136994.51	136994.51	0.0258	0.7753
1A	1A (Stationary) Oil	CO ₂	44829.31	44829.31	0.0240	0.7993
4C	4C Grassland	CO ₂	-5897.14	5897.14	0.0234	0.8226
5D	5D Wastewater Handling	N ₂ O	1063.03	1063.03	0.0232	0.8459
2F	2F Product Uses as Substitutes for ODS	HFCs	16200.73	16200.73	0.0165	0.8624
1A1 & 1A2 & 1A4 & 1A5	1A1 & 1A2 & 1A4 & 1A5 Other Combustion	N ₂ O	2754.34	2754.34	0.0148	0.8771
1A3b	1A3b DERV	CO ₂	68350.74	68350.74	0.0135	0.8906

IPCC Code	IPCC Category	Greenhouse Gas	2013 emissions (Gg CO ₂ e)	Absolute value of 2013 emissions (Gg CO ₂ e)	Level Assessment	Cumulative Total
3B	3B Manure Management	CH ₄	3471.50	3471.50	0.0092	0.8998
1A3b	1A3b DERV	N ₂ O	758.86	758.86	0.0087	0.9084
5D	5D Wastewater Handling	CH ₄	3359.33	3359.33	0.0080	0.9164
2B	2B Chemical industries	CO ₂	4740.31	4740.31	0.0079	0.9243
1B2	1B2 Natural Gas Transmission	CH ₄	4396.86	4396.86	0.0078	0.9321
1A3b	1A3b Gasoline/ LPG	CO ₂	38697.84	38697.84	0.0076	0.9397
5B	5B Biological treatment of solid waste	CH ₄	715.72	715.72	0.0065	0.9462
5B	5B Biological treatment of solid waste	N ₂ O	608.32	608.32	0.0051	0.9513
4G	4G Other Activities	CO ₂	-1121.03	1121.03	0.0044	0.9558
1A3a	1A3a Aviation Fuel	CO ₂	1961.73	1961.73	0.0034	0.9592
1A3d	1A3d Marine fuel	CO ₂	2165.99	2165.99	0.0034	0.9625
1A3	1A3 Other diesel	CO ₂	2384.81	2384.81	0.0032	0.9657
1A1 & 1A2 & 1A4 & 1A5	1A1 & 1A2 & 1A4 & 1A5 Other Combustion	CH ₄	966.42	966.42	0.0031	0.9689
2D	2D Non Energy Products from Fuels and Solvent Use	CO ₂	1032.92	1032.92	0.0030	0.9719
1B1	1B1 Coal Mining	CH ₄	1682.41	1682.41	0.0030	0.9748
1A	1A Other (waste)	CO ₂	2426.36	2426.36	0.0027	0.9776
1B2	1B2 Oil & Natural Gas	CO ₂	3789.72	3789.72	0.0024	0.9800
2C	2C Metal Industries	CO ₂	5013.31	5013.31	0.0023	0.9822
1B2	1B2 Offshore Oil& Gas	CH ₄	1290.40	1290.40	0.0020	0.9842
4B	4B Cropland	N ₂ O	329.94	329.94	0.0016	0.9858
3G	3G Liming	CO ₂	774.69	774.69	0.0014	0.9872
4D	4D Wetland	CO ₂	300.49	300.49	0.0013	0.9885
2A1	2A1 Cement Production	CO ₂	4029.11	4029.11	0.0011	0.9897
3H	3H Urea application to agriculture	CO ₂	245.22	245.22	0.0011	0.9907
1A3b	1A3b Gasoline/ LPG	N ₂ O	150.48	150.48	0.0010	0.9917

IPCC Code	IPCC Category	Greenhouse Gas	2013 emissions (Gg CO ₂ e)	Absolute value of 2013 emissions (Gg CO ₂ e)	Level Assessment	Cumulative Total
5C	5C Waste Incineration	N ₂ O	44.28	44.28	0.0009	0.9926
1A3	1A3 Other diesel	N ₂ O	61.14	61.14	0.0007	0.9933
5C	5C Waste Incineration	CO ₂	264.30	264.30	0.0006	0.9939
4E	4E Settlements	N ₂ O	318.07	318.07	0.0006	0.9945
2G	2G Other Product Manufacture and Use	PFCs	134.30	134.30	0.0006	0.9950
2A2	2A2 Lime Production	CO ₂	1239.15	1239.15	0.0005	0.9956
2C	2C Iron & Steel	N ₂ O	45.23	45.23	0.0005	0.9961
2B2	2B2 Nitric Acid Production	N ₂ O	42.91	42.91	0.0004	0.9964
1A4	1A4 Petroleum Coke	CO ₂	160.19	160.19	0.0004	0.9968
1B2	1B2 Oil & Natural Gas	N ₂ O	30.79	30.79	0.0003	0.9971
1A3b	1A3b Gasoline/ LPG	CH ₄	38.64	38.64	0.0003	0.9973
2A4	2A4 Other process uses of carbonates	CO ₂	771.83	771.83	0.0002	0.9976
2G	2G Other Product Manufacture and Use	SF ₆	455.37	455.37	0.0002	0.9978
1A3b	1A3b DERV	CH ₄	16.98	16.98	0.0002	0.9980
2B	2B Chemical Industry	CH ₄	104.11	104.11	0.0002	0.9982
1A3a	1A3a Aviation Fuel	N ₂ O	18.56	18.56	0.0002	0.9984
4A	4A Forest land	N ₂ O	50.29	50.29	0.0002	0.9985
2A3	2A3 Glass production	CO ₂	389.18	389.18	0.0002	0.9987
1A3d	1A3d Marine fuel	N ₂ O	16.22	16.22	0.0002	0.9989
4C	4C Grassland	CH ₄	28.46	28.46	0.0001	0.9990
1A4	1A4 Peat	CO ₂	46.96	46.96	0.0001	0.9991
1B1	1B1 Solid Fuel Transformation	CO ₂	175.28	175.28	0.0001	0.9993
2B	2B Chemical industry	PFCs	112.13	112.13	0.0001	0.9994
2C	2C Metal Industries	SF ₆	146.18	146.18	0.0001	0.9994
2C	2C Iron & Steel Production	CH ₄	20.49	20.49	0.0001	0.9995
2E	2E Electronics Industry	HFCs	14.47	14.47	0.0001	0.9996
1A3c	1A3c Coal	CO ₂	32.60	32.60	0.0001	0.9996
1A3d	1A3d Marine fuel	CH ₄	4.90	4.90	0.0001	0.9997
5C	5C Waste Incineration	CH ₄	9.95	9.95	0.0000	0.9997

IPCC Code	IPCC Category	Greenhouse Gas	2013 emissions (Gg CO ₂ e)	Absolute value of 2013 emissions (Gg CO ₂ e)	Level Assessment	Cumulative Total
2B	2B Chemical industry	HFCs	46.11	46.11	0.0000	0.9998
2A	2A Minerals industry	CH ₄	4.52	4.52	0.0000	0.9998
1A3	1A3 Other diesel	CH ₄	2.60	2.60	0.0000	0.9999
4C	4C Grassland	N ₂ O	9.15	9.15	0.0000	0.9999
4A	4A Forest Land	CH ₄	5.04	5.04	0.0000	0.9999
1B1	1B1 Solid Fuel Transformation	CH ₄	3.85	3.85	0.0000	0.9999
2B8	2B8 Petrochemical and Carbon Black Production	N ₂ O	1.53	1.53	0.0000	0.9999
2C	2C Metal Industries	PFCs	6.95	6.95	0.0000	1.0000
2D	2D Non-energy Products from Fuels and Solvent Use	N ₂ O	0.91	0.91	0.0000	1.0000
1A3c	1A3c Coal	CH ₄	0.86	0.86	0.0000	1.0000
2G	2G Other Product Manufacture and Use	N ₂ O	0.49	0.49	0.0000	1.0000
1A3a	1A3a Aviation Fuel	CH ₄	0.86	0.86	0.0000	1.0000
4D	4D Grassland	N ₂ O	0.50	0.50	0.0000	1.0000
4E	4E Settlements	CH ₄	1.99	1.99	0.0000	1.0000
2C	2C Metal Industries	HFCs	2.61	2.61	0.0000	1.0000
2D	2D Non-energy Products from Fuels and Solvent Use	CH ₄	0.38	0.38	0.0000	1.0000
2E	2E Electronics Industry	NF ₃	0.36	0.36	0.0000	1.0000
2B1	2B1 Ammonia Production	N ₂ O	0.27	0.27	0.0000	1.0000
1A3c	1A3c Coal	N ₂ O	0.08	0.08	0.0000	1.0000
4B	4B Cropland	CH ₄	0.14	0.14	0.0000	1.0000
1B1	1B1 Fugitive Emissions from Solid Fuels	N ₂ O	0.04	0.04	0.0000	1.0000
4F	4F Other Land	CO ₂	0.02	0.02	0.0000	1.0000
3F	3F Field Burning	CH ₄	0.00	0.00	0.0000	1.0000
2B3	2B3 Adipic Acid Production	N ₂ O	0.00	0.00	0.0000	1.0000
3F	3F Field Burning	N ₂ O	0.00	0.00	0.0000	1.0000
2F	2F Product Uses as Substitutes for ODS	PFCs	0.00	0.00	0.0000	1.0000

Table A 1.4.3 Approach 2 Level Assessment for Base year (not including LULUCF) with Key Categories Shaded in Grey

IPCC Code	IPCC Category	Greenhouse Gas	Base year emissions (Gg CO ₂ e)	Absolute value of Base year emissions (Gg CO ₂ e)	Level Assessment	Cumulative Total
3D	3D Agricultural Soils	N ₂ O	23153.96	23153.96	0.3603	0.3603
5A	5A Solid Waste Disposal	CH ₄	62749.72	62749.72	0.1824	0.5427
2B3	2B3 Adipic Acid Production	N ₂ O	19934.61	19934.61	0.1198	0.6625
3B	3B Manure Management	N ₂ O	2403.46	2403.46	0.0367	0.6991
1A	1A (Stationary) Oil	CO ₂	93760.90	93760.90	0.0343	0.7334
3A	3A Enteric Fermentation	CH ₄	28227.49	28227.49	0.0339	0.7673
1A	1A Coal	CO ₂	244317.59	244317.59	0.0314	0.7987
1B1	1B1 Coal Mining	CH ₄	21788.17	21788.17	0.0263	0.8250
2B2	2B2 Nitric Acid Production	N ₂ O	3860.26	3860.26	0.0233	0.8483
5D	5D Wastewater Handling	N ₂ O	1111.13	1111.13	0.0166	0.8649
1A1 & 1A2 & 1A4 & 1A5	1A1 & 1A2 & 1A4 & 1A5 Other Combustion	N ₂ O	4209.03	4209.03	0.0154	0.8803
1A	1A Natural Gas	CO ₂	107734.56	107734.56	0.0141	0.8944
1B2	1B2 Natural Gas Transmission	CH ₄	10168.29	10168.29	0.0124	0.9068
2B	2B Chemical industry	HFCs	17680.04	17680.04	0.0106	0.9174
1A3b	1A3b Gasoline/ LPG	CO ₂	75557.88	75557.88	0.0101	0.9275
3B	3B Manure Management	CH ₄	4472.78	4472.78	0.0081	0.9356
2B	2B Chemical industries	CO ₂	6376.75	6376.75	0.0072	0.9428
5D	5D Wastewater Handling	CH ₄	4169.64	4169.64	0.0067	0.9495
1A1 & 1A2 & 1A4 & 1A5	1A1 & 1A2 & 1A4 & 1A5 Other Combustion	CH ₄	2079.93	2079.93	0.0046	0.9542
1A3b	1A3b DERV	CO ₂	33012.63	33012.63	0.0044	0.9586
1A3b	1A3b Gasoline/ LPG	N ₂ O	862.61	862.61	0.0038	0.9624
1A3b	1A3b Gasoline/ LPG	CH ₄	630.55	630.55	0.0028	0.9652

IPCC Code	IPCC Category	Greenhouse Gas	Base year emissions (Gg CO ₂ e)	Absolute value of Base year emissions (Gg CO ₂ e)	Level Assessment	Cumulative Total
1B2	1B2 Oil & Natural Gas	CO ₂	5777.92	5777.92	0.0025	0.9677
2D	2D Non Energy Products from Fuels and Solvent Use	CO ₂	1232.62	1232.62	0.0024	0.9702
1A3d	1A3d Marine fuel	CO ₂	2168.02	2168.02	0.0023	0.9725
2C	2C Metal Industries	CO ₂	7391.77	7391.77	0.0023	0.9747
1B2	1B2 Offshore Oil& Gas	CH ₄	2164.64	2164.64	0.0023	0.9770
1A3a	1A3a Aviation Fuel	CO ₂	1859.57	1859.57	0.0022	0.9792
1A3b	1A3b DERV	N ₂ O	279.80	279.80	0.0022	0.9814
5C	5C Waste Incineration	CO ₂	1304.38	1304.38	0.0020	0.9834
3G	3G Liming	CO ₂	1579.17	1579.17	0.0020	0.9854
1A3	1A3 Other diesel	CO ₂	1679.92	1679.92	0.0015	0.9869
2A1	2A1 Cement Production	CO ₂	7295.26	7295.26	0.0014	0.9883
2F	2F Product Uses as Substitutes for ODS	HFCs	1870.17	1870.17	0.0013	0.9896
3H	3H Urea application to agriculture	CO ₂	385.46	385.46	0.0012	0.9907
1A3b	1A3b DERV	CH ₄	127.77	127.77	0.0010	0.9917
1A4	1A4 Peat	CO ₂	475.59	475.59	0.0009	0.9926
3F	3F Field Burning	N ₂ O	63.50	63.50	0.0009	0.9935
1B1	1B1 Solid Fuel Transformation	CO ₂	1698.56	1698.56	0.0007	0.9942
3F	3F Field Burning	CH ₄	205.48	205.48	0.0007	0.9949
2C	2C Iron & Steel	N ₂ O	62.46	62.46	0.0004	0.9953
2A2	2A2 Lime Production	CO ₂	1462.05	1462.05	0.0004	0.9958
2G	2G Other Product Manufacture and Use	PFCs	149.17	149.17	0.0004	0.9962
5C	5C Waste Incineration	CH ₄	137.22	137.22	0.0004	0.9966
5C	5C Waste Incineration	N ₂ O	29.85	29.85	0.0004	0.9970
2C	2C Metal Industries	PFCs	333.43	333.43	0.0004	0.9974
2G	2G Other Product Manufacture and Use	SF ₆	877.20	877.20	0.0003	0.9977
1B2	1B2 Oil & Natural Gas	N ₂ O	40.75	40.75	0.0003	0.9980
2B	2B Chemical Industry	CH ₄	213.70	213.70	0.0003	0.9983
1A3	1A3 Other diesel	N ₂ O	31.12	31.12	0.0002	0.9985

IPCC Code	IPCC Category	Greenhouse Gas	Base year emissions (Gg CO ₂ e)	Absolute value of Base year emissions (Gg CO ₂ e)	Level Assessment	Cumulative Total
2A	2A Minerals industry	CH ₄	31.27	31.27	0.0002	0.9987
1A	1A Other (waste)	CO ₂	228.07	228.07	0.0002	0.9989
2C	2C Metal Industries	SF ₆	387.17	387.17	0.0002	0.9990
2A4	2A4 Other process uses of carbonates	CO ₂	646.76	646.76	0.0001	0.9992
1A4	1A4 Petroleum Coke	CO ₂	81.64	81.64	0.0001	0.9993
2A3	2A3 Glass production	CO ₂	408.14	408.14	0.0001	0.9994
1A3a	1A3a Aviation Fuel	N ₂ O	17.60	17.60	0.0001	0.9995
1A3d	1A3d Marine fuel	N ₂ O	16.26	16.26	0.0001	0.9997
2C	2C Iron & Steel Production	CH ₄	36.89	36.89	0.0001	0.9998
2B	2B Chemical industry	PFCs	113.90	113.90	0.0001	0.9998
5B	5B Biological treatment of solid waste	CH ₄	5.48	5.48	0.0000	0.9999
5B	5B Biological treatment of solid waste	N ₂ O	4.90	4.90	0.0000	0.9999
2E	2E Electronics Industry	HFCs	8.73	8.73	0.0000	0.9999
1A3	1A3 Other diesel	CH ₄	2.75	2.75	0.0000	0.9999
1A3d	1A3d Marine fuel	CH ₄	2.22	2.22	0.0000	1.0000
1A3a	1A3a Aviation Fuel	CH ₄	4.17	4.17	0.0000	1.0000
2D	2D Non-energy Products from Fuels and Solvent Use	N ₂ O	1.64	1.64	0.0000	1.0000
2B8	2B8 Petrochemical and Carbon Black Production	N ₂ O	1.79	1.79	0.0000	1.0000
2G	2G Other Product Manufacture and Use	N ₂ O	0.44	0.44	0.0000	1.0000
2D	2D Non-energy Products from Fuels and Solvent Use	CH ₄	0.69	0.69	0.0000	1.0000
2E	2E Electronics Industry	NF ₃	0.83	0.83	0.0000	1.0000
2B1	2B1 Ammonia Production	N ₂ O	0.31	0.31	0.0000	1.0000
2F	2F Product Uses as Substitutes for ODS	PFCs	0.44	0.44	0.0000	1.0000
1B1	1B1 Fugitive Emissions from Solid Fuels	N ₂ O	0.09	0.09	0.0000	1.0000
1B1	1B1 Solid Fuel Transformation	CH ₄	0.18	0.18	0.0000	1.0000

IPCC Code	IPCC Category	Greenhouse Gas	Base year emissions (Gg CO ₂ e)	Absolute value of Base year emissions (Gg CO ₂ e)	Level Assessment	Cumulative Total
1A3c	1A3c Coal	CO ₂	0.00	0.00	0.0000	1.0000
1A3c	1A3c Coal	CH ₄	0.00	0.00	0.0000	1.0000
1A3c	1A3c Coal	N ₂ O	0.00	0.00	0.0000	1.0000
2C	2C Metal Industries	HFCs	0.00	0.00	0.0000	1.0000

Table A 1.4.4 Approach 2 Level Assessment for 2013 (not including LULUCF) with Key Categories Shaded in Grey

IPCC Code	IPCC Category	Greenhouse Gas	2013 emissions (Gg CO ₂ e)	Absolute value of 2013 emissions (Gg CO ₂ e)	Level Assessment	Cumulative Total
3D	3D Agricultural Soils	N ₂ O	19513.90	19513.90	0.5244	0.5244
5A	5A Solid Waste Disposal	CH ₄	16681.83	16681.83	0.0837	0.6082
3B	3B Manure Management	N ₂ O	1891.28	1891.28	0.0498	0.6580
3A	3A Enteric Fermentation	CH ₄	23621.80	23621.80	0.0490	0.7070
1A	1A Natural Gas	CO ₂	154663.64	154663.64	0.0350	0.7421
1A	1A Coal	CO ₂	136994.51	136994.51	0.0304	0.7725
1A	1A (Stationary) Oil	CO ₂	44829.31	44829.31	0.0283	0.8008
5D	5D Wastewater Handling	N ₂ O	1063.03	1063.03	0.0274	0.8281
2F	2F Product Uses as Substitutes for ODS	HFCs	16200.73	16200.73	0.0194	0.8476
1A1 & 1A2 & 1A4 & 1A5	1A1 & 1A2 & 1A4 & 1A5 Other Combustion	N ₂ O	2754.34	2754.34	0.0174	0.8650
1A3b	1A3b DERV	CO ₂	68350.74	68350.74	0.0159	0.8808
3B	3B Manure Management	CH ₄	3471.50	3471.50	0.0108	0.8916
1A3b	1A3b DERV	N ₂ O	758.86	758.86	0.0102	0.9019
5D	5D Wastewater Handling	CH ₄	3359.33	3359.33	0.0094	0.9113
2B	2B Chemical industries	CO ₂	4740.31	4740.31	0.0093	0.9206
1B2	1B2 Natural Gas Transmission	CH ₄	4396.86	4396.86	0.0092	0.9298
1A3b	1A3b Gasoline/ LPG	CO ₂	38697.84	38697.84	0.0089	0.9387

IPCC Code	IPCC Category	Greenhouse Gas	2013 emissions (Gg CO ₂ e)	Absolute value of 2013 emissions (Gg CO ₂ e)	Level Assessment	Cumulative Total
5B	5B Biological treatment of solid waste	CH ₄	715.72	715.72	0.0077	0.9464
5B	5B Biological treatment of solid waste	N ₂ O	608.32	608.32	0.0060	0.9524
1A3a	1A3a Aviation Fuel	CO ₂	1961.73	1961.73	0.0040	0.9564
1A3d	1A3d Marine fuel	CO ₂	2165.99	2165.99	0.0040	0.9604
1A3	1A3 Other diesel	CO ₂	2384.81	2384.81	0.0037	0.9642
1A1 & 1A2 & 1A4 & 1A5	1A1 & 1A2 & 1A4 & 1A5 Other Combustion	CH ₄	966.42	966.42	0.0037	0.9679
2D	2D Non Energy Products from Fuels and Solvent Use	CO ₂	1032.92	1032.92	0.0035	0.9714
1B1	1B1 Coal Mining	CH ₄	1682.41	1682.41	0.0035	0.9749
1A	1A Other (waste)	CO ₂	2426.36	2426.36	0.0032	0.9781
1B2	1B2 Oil & Natural Gas	CO ₂	3789.72	3789.72	0.0028	0.9809
2C	2C Metal Industries	CO ₂	5013.31	5013.31	0.0027	0.9836
1B2	1B2 Offshore Oil& Gas	CH ₄	1290.40	1290.40	0.0023	0.9859
3G	3G Liming	CO ₂	774.69	774.69	0.0017	0.9876
2A1	2A1 Cement Production	CO ₂	4029.11	4029.11	0.0013	0.9889
3H	3H Urea application to agriculture	CO ₂	245.22	245.22	0.0013	0.9902
1A3b	1A3b Gasoline/ LPG	N ₂ O	150.48	150.48	0.0012	0.9914
5C	5C Waste Incineration	N ₂ O	44.28	44.28	0.0011	0.9924
1A3	1A3 Other diesel	N ₂ O	61.14	61.14	0.0008	0.9932
5C	5C Waste Incineration	CO ₂	264.30	264.30	0.0007	0.9939
2G	2G Other Product Manufacture and Use	PFCs	134.30	134.30	0.0007	0.9946
2A2	2A2 Lime Production	CO ₂	1239.15	1239.15	0.0006	0.9952
2C	2C Iron & Steel	N ₂ O	45.23	45.23	0.0006	0.9958
2B2	2B2 Nitric Acid Production	N ₂ O	42.91	42.91	0.0004	0.9962
1A4	1A4 Petroleum Coke	CO ₂	160.19	160.19	0.0004	0.9967
1B2	1B2 Oil & Natural Gas	N ₂ O	30.79	30.79	0.0003	0.9970

IPCC Code	IPCC Category	Greenhouse Gas	2013 emissions (Gg CO ₂ e)	Absolute value of 2013 emissions (Gg CO ₂ e)	Level Assessment	Cumulative Total
1A3b	1A3b Gasoline/ LPG	CH ₄	38.64	38.64	0.0003	0.9973
2A4	2A4 Other process uses of carbonates	CO ₂	771.83	771.83	0.0003	0.9976
2G	2G Other Product Manufacture and Use	SF ₆	455.37	455.37	0.0003	0.9979
1A3b	1A3b DERV	CH ₄	16.98	16.98	0.0002	0.9981
2B	2B Chemical Industry	CH ₄	104.11	104.11	0.0002	0.9983
1A3a	1A3a Aviation Fuel	N ₂ O	18.56	18.56	0.0002	0.9985
2A3	2A3 Glass production	CO ₂	389.18	389.18	0.0002	0.9987
1A3d	1A3d Marine fuel	N ₂ O	16.22	16.22	0.0002	0.9989
1A4	1A4 Peat	CO ₂	46.96	46.96	0.0002	0.9991
1B1	1B1 Solid Fuel Transformation	CO ₂	175.28	175.28	0.0001	0.9992
2B	2B Chemical industry	PFCs	112.13	112.13	0.0001	0.9993
2C	2C Metal Industries	SF ₆	146.18	146.18	0.0001	0.9994
2C	2C Iron & Steel Production	CH ₄	20.49	20.49	0.0001	0.9995
2E	2E Electronics Industry	HFCs	14.47	14.47	0.0001	0.9996
1A3c	1A3c Coal	CO ₂	32.60	32.60	0.0001	0.9997
1A3d	1A3d Marine fuel	CH ₄	4.90	4.90	0.0001	0.9997
5C	5C Waste Incineration	CH ₄	9.95	9.95	0.0001	0.9998
2B	2B Chemical industry	HFCs	46.11	46.11	0.0000	0.9998
2A	2A Minerals industry	CH ₄	4.52	4.52	0.0000	0.9999
1A3	1A3 Other diesel	CH ₄	2.60	2.60	0.0000	0.9999
1B1	1B1 Solid Fuel Transformation	CH ₄	3.85	3.85	0.0000	0.9999
2B8	2B8 Petrochemical and Carbon Black Production	N ₂ O	1.53	1.53	0.0000	0.9999
2C	2C Metal Industries	PFCs	6.95	6.95	0.0000	1.0000
2D	2D Non-energy Products from Fuels and Solvent Use	N ₂ O	0.91	0.91	0.0000	1.0000
1A3c	1A3c Coal	CH ₄	0.86	0.86	0.0000	1.0000
2G	2G Other Product Manufacture and Use	N ₂ O	0.49	0.49	0.0000	1.0000
1A3a	1A3a Aviation Fuel	CH ₄	0.86	0.86	0.0000	1.0000
2C	2C Metal Industries	HFCs	2.61	2.61	0.0000	1.0000

IPCC Code	IPCC Category	Greenhouse Gas	2013 emissions (Gg CO ₂ e)	Absolute value of 2013 emissions (Gg CO ₂ e)	Level Assessment	Cumulative Total
2D	2D Non-energy Products from Fuels and Solvent Use	CH ₄	0.38	0.38	0.0000	1.0000
2E	2E Electronics Industry	NF ₃	0.36	0.36	0.0000	1.0000
2B1	2B1 Ammonia Production	N ₂ O	0.27	0.27	0.0000	1.0000
1A3c	1A3c Coal	N ₂ O	0.08	0.08	0.0000	1.0000
1B1	1B1 Fugitive Emissions from Solid Fuels	N ₂ O	0.04	0.04	0.0000	1.0000
3F	3F Field Burning	CH ₄	0.00	0.00	0.0000	1.0000
2B3	2B3 Adipic Acid Production	N ₂ O	0.00	0.00	0.0000	1.0000
3F	3F Field Burning	N ₂ O	0.00	0.00	0.0000	1.0000
2F	2F Product Uses as Substitutes for ODS	PFCs	0.00	0.00	0.0000	1.0000

Table A 1.4.5 Approach 2 Assessment for Trend (including LULUCF) with Key Categories Shaded in Grey

IPCC Code	IPCC Category	Greenhouse Gas	Base year emissions (Gg CO ₂ e)	2013 emissions (Gg CO ₂ e)	Trend Assessment with Uncertainty	Contribution to Trend Uncertainty (%)	Cumulative Total
2B3	2B3 Adipic Acid Production	N ₂ O	19934.61	0.00	0.0163	23.2%	0.2316
5A	5A Solid Waste Disposal	CH ₄	62749.72	16681.83	0.0154	21.9%	0.4503
3D	3D Agricultural Soils	N ₂ O	23153.96	19513.90	0.0100	14.2%	0.5927
1B1	1B1 Coal Mining	CH ₄	21788.17	1682.41	0.0032	4.5%	0.6379
2B2	2B2 Nitric Acid Production	N ₂ O	3860.26	42.91	0.0031	4.4%	0.6823
1A	1A Natural Gas	CO ₂	107734.56	154663.64	0.0020	2.9%	0.7110
2F	2F Product Uses as Substitutes for ODS	HFCs	1870.17	16200.73	0.0020	2.9%	0.7396
1A	1A (Stationary) Oil	CO ₂	93760.90	44829.31	0.0015	2.1%	0.7605
2B	2B Chemical industry	HFCs	17680.04	46.11	0.0014	2.0%	0.7810
4A	4A Forest Land	CO ₂	-16016.07	-17297.63	0.0012	1.8%	0.7986
1A3b	1A3b DERV	CO ₂	33012.63	68350.74	0.0012	1.7%	0.8154

IPCC Code	IPCC Category	Greenhouse Gas	Base year emissions (Gg CO ₂ e)	2013 emissions (Gg CO ₂ e)	Trend Assessment with Uncertainty	Contribution to Trend Uncertainty (%)	Cumulative Total
3A	3A Enteric Fermentation	CH ₄	28227.49	23621.80	0.0009	1.3%	0.8282
5B	5B Biological treatment of solid waste	CH ₄	5.48	715.72	0.0009	1.2%	0.8405
1A3b	1A3b DERV	N ₂ O	279.80	758.86	0.0009	1.2%	0.8526
1A	1A Coal	CO ₂	244317.59	136994.51	0.0009	1.2%	0.8647
5D	5D Wastewater Handling	N ₂ O	1111.13	1063.03	0.0008	1.2%	0.8765
4B	4B Cropland	CO ₂	15130.79	12150.39	0.0008	1.2%	0.8881
5B	5B Biological treatment of solid waste	N ₂ O	4.90	608.32	0.0007	1.0%	0.8977
1B2	1B2 Natural Gas Transmission	CH ₄	10168.29	4396.86	0.0006	0.9%	0.9068
3B	3B Manure Management	N ₂ O	2403.46	1891.28	0.0006	0.9%	0.9156
4E	4E Settlements	CO ₂	6919.10	5876.98	0.0006	0.9%	0.9242
4G	4G Other Activities	CO ₂	40.72	-1121.03	0.0006	0.9%	0.9328
4C	4C Grassland	CO ₂	-3677.20	-5897.14	0.0006	0.8%	0.9411
1A3b	1A3b Gasoline/ LPG	N ₂ O	862.61	150.48	0.0004	0.6%	0.9467
1A3b	1A3b Gasoline/ LPG	CO ₂	75557.88	38697.84	0.0004	0.5%	0.9519
1A3b	1A3b Gasoline/ LPG	CH ₄	630.55	38.64	0.0003	0.5%	0.9569
1A	1A Other (waste)	CO ₂	228.07	2426.36	0.0003	0.5%	0.9617
1A3	1A3 Other diesel	CO ₂	1679.92	2384.81	0.0002	0.3%	0.9647
1A1 & 1A2 & 1A4 & 1A5	1A1 & 1A2 & 1A4 & 1A5 Other Combustion	CH ₄	2079.93	966.42	0.0002	0.3%	0.9677
5C	5C Waste Incineration	CO ₂	1304.38	264.30	0.0002	0.3%	0.9705
1A3a	1A3a Aviation Fuel	CO ₂	1859.57	1961.73	0.0002	0.2%	0.9726
5D	5D Wastewater Handling	CH ₄	4169.64	3359.33	0.0001	0.2%	0.9746
1A1 & 1A2 &	1A1 & 1A2 & 1A4 & 1A5 Other Combustion	N ₂ O	4209.03	2754.34	0.0001	0.2%	0.9765

IPCC Code	IPCC Category	Greenhouse Gas	Base year emissions (Gg CO ₂ e)	2013 emissions (Gg CO ₂ e)	Trend Assessment with Uncertainty	Contribution to Trend Uncertainty (%)	Cumulative Total
1A4 & 1A5							
1A3d	1A3d Marine fuel	CO ₂	2168.02	2165.99	0.0001	0.2%	0.9784
3F	3F Field Burning	N ₂ O	63.50	0.00	0.0001	0.2%	0.9802
3B	3B Manure Management	CH ₄	4472.78	3471.50	0.0001	0.2%	0.9819
1A3b	1A3b DERV	CH ₄	127.77	16.98	0.0001	0.2%	0.9834
1A4	1A4 Peat	CO ₂	475.59	46.96	0.0001	0.2%	0.9849
3F	3F Field Burning	CH ₄	205.48	0.00	0.0001	0.1%	0.9863
3G	3G Liming	CO ₂	1579.17	774.69	0.0001	0.1%	0.9874
1B1	1B1 Solid Fuel Transformation	CO ₂	1698.56	175.28	0.0001	0.1%	0.9885
4B	4B Cropland	N ₂ O	649.15	329.94	0.0001	0.1%	0.9897
2D	2D Non Energy Products from Fuels and Solvent Use	CO ₂	1232.62	1032.92	0.0001	0.1%	0.9906
5C	5C Waste Incineration	N ₂ O	29.85	44.28	0.0001	0.1%	0.9915
2B	2B Chemical industries	CO ₂	6376.75	4740.31	0.0001	0.1%	0.9924
1A3	1A3 Other diesel	N ₂ O	31.12	61.14	0.0001	0.1%	0.9932
2C	2C Metal Industries	PFCs	333.43	6.95	0.0001	0.1%	0.9940
5C	5C Waste Incineration	CH ₄	137.22	9.95	0.0001	0.1%	0.9947
1B2	1B2 Offshore Oil & Gas	CH ₄	2164.64	1290.40	0.0000	0.1%	0.9953
2A1	2A1 Cement Production	CO ₂	7295.26	4029.11	0.0000	0.1%	0.9959
1A4	1A4 Petroleum Coke	CO ₂	81.64	160.19	0.0000	0.0%	0.9963
4D	4D Wetland	CO ₂	481.73	300.49	0.0000	0.0%	0.9966
1B2	1B2 Oil & Natural Gas	CO ₂	5777.92	3789.72	0.0000	0.0%	0.9969
2A	2A Minerals industry	CH ₄	31.27	4.52	0.0000	0.0%	0.9972
2G	2G Other Product Manufacture and Use	PFCs	149.17	134.30	0.0000	0.0%	0.9975

IPCC Code	IPCC Category	Greenhouse Gas	Base year emissions (Gg CO ₂ e)	2013 emissions (Gg CO ₂ e)	Trend Assessment with Uncertainty	Contribution to Trend Uncertainty (%)	Cumulative Total
3H	3H Urea application to agriculture	CO ₂	385.46	245.22	0.0000	0.0%	0.9977
4E	4E Settlements	N ₂ O	369.67	318.07	0.0000	0.0%	0.9979
2A4	2A4 Other process uses of carbonates	CO ₂	646.76	771.83	0.0000	0.0%	0.9980
4C	4C Grassland	CH ₄	11.29	28.46	0.0000	0.0%	0.9982
2A2	2A2 Lime Production	CO ₂	1462.05	1239.15	0.0000	0.0%	0.9984
2G	2G Other Product Manufacture and Use	SF ₆	877.20	455.37	0.0000	0.0%	0.9986
2B	2B Chemical Industry	CH ₄	213.70	104.11	0.0000	0.0%	0.9987
2C	2C Metal Industries	SF ₆	387.17	146.18	0.0000	0.0%	0.9989
2C	2C Metal Industries	CO ₂	7391.77	5013.31	0.0000	0.0%	0.9990
1A3a	1A3a Aviation Fuel	N ₂ O	17.60	18.56	0.0000	0.0%	0.9991
1A3c	1A3c Coal	CO ₂	0.00	32.60	0.0000	0.0%	0.9992
4A	4A Forest land	N ₂ O	48.26	50.29	0.0000	0.0%	0.9993
1A3d	1A3d Marine fuel	N ₂ O	16.26	16.22	0.0000	0.0%	0.9994
2A3	2A3 Glass production	CO ₂	408.14	389.18	0.0000	0.0%	0.9995
1A3d	1A3d Marine fuel	CH ₄	2.22	4.90	0.0000	0.0%	0.9996
2E	2E Electronics Industry	HFCs	8.73	14.47	0.0000	0.0%	0.9997
2B	2B Chemical industry	PFCs	113.90	112.13	0.0000	0.0%	0.9997
2C	2C Iron & Steel Production	CH ₄	36.89	20.49	0.0000	0.0%	0.9998
1B2	1B2 Oil & Natural Gas	N ₂ O	40.75	30.79	0.0000	0.0%	0.9998
4D	4D Grassland	N ₂ O	3.83	0.50	0.0000	0.0%	0.9998
1B1	1B1 Solid Fuel Transformation	CH ₄	0.18	3.85	0.0000	0.0%	0.9999
2C	2C Iron & Steel	N ₂ O	62.46	45.23	0.0000	0.0%	0.9999
4A	4A Forest Land	CH ₄	3.72	5.04	0.0000	0.0%	0.9999
1A3a	1A3a Aviation Fuel	CH ₄	4.17	0.86	0.0000	0.0%	0.9999
1A3c	1A3c Coal	CH ₄	0.00	0.86	0.0000	0.0%	0.9999
1A3	1A3 Other diesel	CH ₄	2.75	2.60	0.0000	0.0%	1.0000

IPCC Code	IPCC Category	Greenhouse Gas	Base year emissions (Gg CO ₂ e)	2013 emissions (Gg CO ₂ e)	Trend Assessment with Uncertainty	Contribution to Trend Uncertainty (%)	Cumulative Total
4E	4E Settlements	CH ₄	5.89	1.99	0.0000	0.0%	1.0000
2C	2C Metal Industries	HFCs	0.00	2.61	0.0000	0.0%	1.0000
2B8	2B8 Petrochemical and Carbon Black Production	N ₂ O	1.79	1.53	0.0000	0.0%	1.0000
2D	2D Non-energy Products from Fuels and Solvent Use	N ₂ O	1.64	0.91	0.0000	0.0%	1.0000
2G	2G Other Product Manufacture and Use	N ₂ O	0.44	0.49	0.0000	0.0%	1.0000
4C	4C Grassland	N ₂ O	12.10	9.15	0.0000	0.0%	1.0000
2E	2E Electronics Industry	NF ₃	0.83	0.36	0.0000	0.0%	1.0000
1A3c	1A3c Coal	N ₂ O	0.00	0.08	0.0000	0.0%	1.0000
2F	2F Product Uses as Substitutes for ODS	PFCs	0.44	0.00	0.0000	0.0%	1.0000
2D	2D Non-energy Products from Fuels and Solvent Use	CH ₄	0.69	0.38	0.0000	0.0%	1.0000
4B	4B Cropland	CH ₄	0.11	0.14	0.0000	0.0%	1.0000
2B1	2B1 Ammonia Production	N ₂ O	0.31	0.27	0.0000	0.0%	1.0000
1B1	1B1 Fugitive Emissions from Solid Fuels	N ₂ O	0.09	0.04	0.0000	0.0%	1.0000
4F	4F Other Land	CO ₂	0.00	0.02	0.0000	0.0%	1.0000

Table A 1.4.6 Approach 2 Assessment for Trend (not including LULUCF) with Key Categories Shaded in Grey

IPCC Code	IPCC Category	Greenhouse Gas	Base year emissions (Gg CO ₂ e)	2013 emissions (Gg CO ₂ e)	Trend Assessment with Uncertainty	Contribution to Trend Uncertainty (%)	Cumulative Total
2B3	2B3 Adipic Acid Production	N ₂ O	19934.61	0.00	0.0163	24.6%	0.2456
5A	5A Solid Waste Disposal	CH ₄	62749.72	16681.83	0.0154	23.2%	0.4774
3D	3D Agricultural Soils	N ₂ O	23153.96	19513.90	0.0100	15.1%	0.6283

IPCC Code	IPCC Category	Greenhouse Gas	Base year emissions (Gg CO ₂ e)	2013 emissions (Gg CO ₂ e)	Trend Assessment with Uncertainty	Contribution to Trend Uncertainty (%)	Cumulative Total
1B1	1B1 Coal Mining	CH ₄	21788.17	1682.41	0.0032	4.8%	0.6763
2B2	2B2 Nitric Acid Production	N ₂ O	3860.26	42.91	0.0031	4.7%	0.7233
1A	1A Natural Gas	CO ₂	107734.56	154663.64	0.0020	3.0%	0.7538
2F	2F Product Uses as Substitutes for ODS	HFCs	1870.17	16200.73	0.0020	3.0%	0.7841
1A	1A (Stationary) Oil	CO ₂	93760.90	44829.31	0.0015	2.2%	0.8063
2B	2B Chemical industry	HFCs	17680.04	46.11	0.0014	2.2%	0.8280
1A3b	1A3b DERV	CO ₂	33012.63	68350.74	0.0012	1.8%	0.8458
3A	3A Enteric Fermentation	CH ₄	28227.49	23621.80	0.0009	1.4%	0.8594
5B	5B Biological treatment of solid waste	CH ₄	5.48	715.72	0.0009	1.3%	0.8724
1A3b	1A3b DERV	N ₂ O	279.80	758.86	0.0009	1.3%	0.8853
1A	1A Coal	CO ₂	244317.59	136994.51	0.0009	1.3%	0.8981
5D	5D Wastewater Handling	N ₂ O	1111.13	1063.03	0.0008	1.2%	0.9106
5B	5B Biological treatment of solid waste	N ₂ O	4.90	608.32	0.0007	1.0%	0.9207
1B2	1B2 Natural Gas Transmission	CH ₄	10168.29	4396.86	0.0006	1.0%	0.9304
3B	3B Manure Management	N ₂ O	2403.46	1891.28	0.0006	0.9%	0.9397
1A3b	1A3b Gasoline/ LPG	N ₂ O	862.61	150.48	0.0004	0.6%	0.9456
1A3b	1A3b Gasoline/ LPG	CO ₂	75557.88	38697.84	0.0004	0.6%	0.9512
1A3b	1A3b Gasoline/ LPG	CH ₄	630.55	38.64	0.0003	0.5%	0.9564
1A	1A Other (waste)	CO ₂	228.07	2426.36	0.0003	0.5%	0.9615
1A3	1A3 Other diesel	CO ₂	1679.92	2384.81	0.0002	0.3%	0.9647
1A1 & 1A2 & 1A4 & 1A5	1A1 & 1A2 & 1A4 & 1A5 Other Combustion	CH ₄	2079.93	966.42	0.0002	0.3%	0.9679
5C	5C Waste Incineration	CO ₂	1304.38	264.30	0.0002	0.3%	0.9708

IPCC Code	IPCC Category	Greenhouse Gas	Base year emissions (Gg CO ₂ e)	2013 emissions (Gg CO ₂ e)	Trend Assessment with Uncertainty	Contribution to Trend Uncertainty (%)	Cumulative Total
1A3a	1A3a Aviation Fuel	CO ₂	1859.57	1961.73	0.0002	0.2%	0.9731
5D	5D Wastewater Handling	CH ₄	4169.64	3359.33	0.0001	0.2%	0.9752
1A1 & 1A2 & 1A4 & 1A5	1A1 & 1A2 & 1A4 & 1A5 Other Combustion	N ₂ O	4209.03	2754.34	0.0001	0.2%	0.9773
1A3d	1A3d Marine fuel	CO ₂	2168.02	2165.99	0.0001	0.2%	0.9793
3F	3F Field Burning	N ₂ O	63.50	0.00	0.0001	0.2%	0.9811
3B	3B Manure Management	CH ₄	4472.78	3471.50	0.0001	0.2%	0.9829
1A3b	1A3b DERV	CH ₄	127.77	16.98	0.0001	0.2%	0.9845
1A4	1A4 Peat	CO ₂	475.59	46.96	0.0001	0.2%	0.9861
3F	3F Field Burning	CH ₄	205.48	0.00	0.0001	0.1%	0.9875
3G	3G Liming	CO ₂	1579.17	774.69	0.0001	0.1%	0.9888
1B1	1B1 Solid Fuel Transformation	CO ₂	1698.56	175.28	0.0001	0.1%	0.9900
2D	2D Non Energy Products from Fuels and Solvent Use	CO ₂	1232.62	1032.92	0.0001	0.1%	0.9910
5C	5C Waste Incineration	N ₂ O	29.85	44.28	0.0001	0.1%	0.9919
2B	2B Chemical industries	CO ₂	6376.75	4740.31	0.0001	0.1%	0.9928
1A3	1A3 Other diesel	N ₂ O	31.12	61.14	0.0001	0.1%	0.9937
2C	2C Metal Industries	PFCs	333.43	6.95	0.0001	0.1%	0.9945
5C	5C Waste Incineration	CH ₄	137.22	9.95	0.0001	0.1%	0.9953
1B2	1B2 Offshore Oil& Gas	CH ₄	2164.64	1290.40	0.0000	0.1%	0.9960
2A1	2A1 Cement Production	CO ₂	7295.26	4029.11	0.0000	0.1%	0.9966
1A4	1A4 Petroleum Coke	CO ₂	81.64	160.19	0.0000	0.0%	0.9970
1B2	1B2 Oil & Natural Gas	CO ₂	5777.92	3789.72	0.0000	0.0%	0.9973
2A	2A Minerals industry	CH ₄	31.27	4.52	0.0000	0.0%	0.9977

IPCC Code	IPCC Category	Greenhouse Gas	Base year emissions (Gg CO ₂ e)	2013 emissions (Gg CO ₂ e)	Trend Assessment with Uncertainty	Contribution to Trend Uncertainty (%)	Cumulative Total
2G	2G Other Product Manufacture and Use	PFCs	149.17	134.30	0.0000	0.0%	0.9979
3H	3H Urea application to agriculture	CO ₂	385.46	245.22	0.0000	0.0%	0.9981
2A4	2A4 Other process uses of carbonates	CO ₂	646.76	771.83	0.0000	0.0%	0.9983
2A2	2A2 Lime Production	CO ₂	1462.05	1239.15	0.0000	0.0%	0.9985
2G	2G Other Product Manufacture and Use	SF ₆	877.20	455.37	0.0000	0.0%	0.9987
2B	2B Chemical Industry	CH ₄	213.70	104.11	0.0000	0.0%	0.9988
2C	2C Metal Industries	SF ₆	387.17	146.18	0.0000	0.0%	0.9990
2C	2C Metal Industries	CO ₂	7391.77	5013.31	0.0000	0.0%	0.9991
1A3a	1A3a Aviation Fuel	N ₂ O	17.60	18.56	0.0000	0.0%	0.9993
1A3c	1A3c Coal	CO ₂	0.00	32.60	0.0000	0.0%	0.9994
1A3d	1A3d Marine fuel	N ₂ O	16.26	16.22	0.0000	0.0%	0.9995
2A3	2A3 Glass production	CO ₂	408.14	389.18	0.0000	0.0%	0.9996
1A3d	1A3d Marine fuel	CH ₄	2.22	4.90	0.0000	0.0%	0.9996
2E	2E Electronics Industry	HFCs	8.73	14.47	0.0000	0.0%	0.9997
2B	2B Chemical industry	PFCs	113.90	112.13	0.0000	0.0%	0.9998
2C	2C Iron & Steel Production	CH ₄	36.89	20.49	0.0000	0.0%	0.9998
1B2	1B2 Oil & Natural Gas	N ₂ O	40.75	30.79	0.0000	0.0%	0.9999
1B1	1B1 Solid Fuel Transformation	CH ₄	0.18	3.85	0.0000	0.0%	0.9999
2C	2C Iron & Steel	N ₂ O	62.46	45.23	0.0000	0.0%	0.9999
1A3a	1A3a Aviation Fuel	CH ₄	4.17	0.86	0.0000	0.0%	0.9999
1A3c	1A3c Coal	CH ₄	0.00	0.86	0.0000	0.0%	1.0000
1A3	1A3 Other diesel	CH ₄	2.75	2.60	0.0000	0.0%	1.0000
2C	2C Metal Industries	HFCs	0.00	2.61	0.0000	0.0%	1.0000
2B8	2B8 Petrochemical and Carbon Black Production	N ₂ O	1.79	1.53	0.0000	0.0%	1.0000

IPCC Code	IPCC Category	Greenhouse Gas	Base year emissions (Gg CO ₂ e)	2013 emissions (Gg CO ₂ e)	Trend Assessment with Uncertainty	Contribution to Trend Uncertainty (%)	Cumulative Total
2D	2D Non-energy Products from Fuels and Solvent Use	N ₂ O	1.64	0.91	0.0000	0.0%	1.0000
2G	2G Other Product Manufacture and Use	N ₂ O	0.44	0.49	0.0000	0.0%	1.0000
2E	2E Electronics Industry	NF ₃	0.83	0.36	0.0000	0.0%	1.0000
1A3c	1A3c Coal	N ₂ O	0.00	0.08	0.0000	0.0%	1.0000
2F	2F Product Uses as Substitutes for ODS	PFCs	0.44	0.00	0.0000	0.0%	1.0000
2D	2D Non-energy Products from Fuels and Solvent Use	CH ₄	0.69	0.38	0.0000	0.0%	1.0000
2B1	2B1 Ammonia Production	N ₂ O	0.31	0.27	0.0000	0.0%	1.0000
1B1	1B1 Fugitive Emissions from Solid Fuels	N ₂ O	0.09	0.04	0.0000	0.0%	1.0000

A 1.5 KEY CATEGORY ANALYSIS (KCA) RANKING SYSTEM

The Key Category Analysis (KCA) ranking system is an additional tool that the UK has developed to aid in the prioritisation of improvement work. The KCA ranking system works by allocating a score based on how high categories rank in the base year and most recent year level assessments and the trend assessment for the approach 1 KCA including LULUCF. For example if CO₂ from road transport liquid fuel use is the 4th highest by the base year level assessment, 3rd highest by the most recent year level assessment and has the 5th highest trend assessment then it's score would be 4+3+5=12. The categories are then ranked from lowest score to highest, with scores that are equal resolved by the most recent year level assessment.

The assessments excluding LULUCF are ignored for this exercise, as the LULUCF sectors would only be included in half of the assessments and would therefore give an unrepresentative weighting.

The results of this ranking are presented in **Table A 1.5.1**.

Table A 1.5.1 KCA Ranking

KCA rank	IPCC Code	IPCC Category	Greenhouse Gas
1	1A1	Energy industries: solid fuels	CO ₂
2	1A3b	Road transportation: liquid fuels	CO ₂
3	1A4	Other sectors: gaseous fuels	CO ₂
4	5A	Solid waste disposal	CH ₄

KCA rank	IPCC Code	IPCC Category	Greenhouse Gas
5	1A1	Energy industries: gaseous fuels	CO ₂
6	1A1	Energy industries: liquid fuels	CO ₂
7	1A2	Manufacturing industries and construction: solid fuels	CO ₂
8	1A2	Manufacturing industries and construction: gaseous fuels	CO ₂
9	3D	Agricultural soils	N ₂ O
10	1A2	Manufacturing industries and construction: liquid fuels	CO ₂
11	3A1	Enteric fermentation from Cattle	CH ₄
12	4A	Forest land	CO ₂
13	1A4	Other sectors: solid fuels	CO ₂
14	1B1	Coal mining and handling	CH ₄
15	4B	Cropland	CO ₂
16	1B2	Oil and gas extraction	CH ₄
17	1A4	Other sectors: liquid fuels	CO ₂
18	4E	Settlements	CO ₂
19	2A1	Cement production	CO ₂
20	2F1	Refrigeration and air conditioning	HFCs, PFCs, SF ₆ and NF ₃
21	4C	Grassland	CO ₂
22	2C1	Iron and steel production	CO ₂
23	1A5	Other: liquid fuels	CO ₂
24	5D	Wastewater treatment and discharge	CH ₄
25	1A3d	Domestic Navigation: liquid fuels	CO ₂
26	1B2	Oil and gas extraction	CO ₂
27	2B9	Fluorochemical production	HFCs, PFCs, SF ₆ and NF ₃
28	3A2	Enteric fermentation from Sheep	CH ₄
29	3B1	Manure management from Cattle	CH ₄
30	2F4	Aerosols	HFCs, PFCs, SF ₆ and NF ₃
31	1A3c	Railways: liquid fuels	CO ₂
32	2B8	Petrochemical and carbon black production	CO ₂
33	1A1	Energy industries: other fuels	CO ₂
34	1B1	Coal mining and handling solid fuels	CO ₂
35	2B2	Nitric acid production	N ₂ O
36	4G	Harvested wood products	CO ₂
29	1A3a	Domestic aviation: liquid fuels	CO ₂
33	3G	Liming	CO ₂
34	1A4	Other sectors: solid fuels	CH ₄
35	2A4	Other process uses of carbonates	CO ₂
36	3B4	Manure management from Other livestock	N ₂ O
33	2A2	Lime production	CO ₂
34	2D	Non-energy products from fuels and solvent use	CO ₂
35	1A3b	Road transportation: liquid fuels	N ₂ O
36	2G1	Electrical equipment	HFCs, PFCs, SF ₆ and NF ₃

A 1.6 APPROACH USED TO IDENTIFY KP-LULUCF KEY CATEGORIES

From the 2010 NIR onwards, the NIR contains a list of the Key Categories for Land Use, Land-Use Change and Forestry Activities under the Kyoto Protocol. The description below explains the Key Category analysis for Article 3.3 activities and any elected activities under Article 3.4.

Three categories are considered to be key: Article 3.3 Afforestation and Reforestation (CO₂), Article 3.3 Deforestation (CO₂) and Article 3.4 Forest Management (CO₂). These have been assessed according to the 2006 IPCC good practice guidance for KP (Chapter 2, Section 2.3.6). The numbers have been compared with **Table A 1.4.2** Key category analysis for the latest reported year (2013) based on level of emissions (including LULUCF).

Article 3.3 Afforestation and Reforestation (CO₂): The associated UNFCCC category 4A (-17 298 Gg CO₂) is a key category and the AR component (forest planted since 1990) is key on its own (i.e. its category contribution (-3 134 Gg CO₂) is greater than the smallest UNFCCC key category (1A3b 1A3b_DERV)). Removals from this category are also predicted to increase over time as a result of tree planting schemes partially focussed on climate change mitigation.

Article 3.3 Deforestation (CO₂): The associated UNFCCC categories (4B, 4C and 4E) are key categories (12 150, -5 897 and 5 877 Gg CO₂ respectively). The Deforestation category contribution (776 Gg CO₂) to these UNFCCC categories is also greater than the smallest UNFCCC key category (1A3b 1A3b_DERV). The data used in the calculation of deforestation emissions are the most uncertain of the data sources in the KP-LULUCF inventory and are a priority for improvement.

Article 3.4 Forest Management (CO₂): The associated UNFCCC category 4A is a key category (-17 298 Gg CO₂). The Forest Management category contribution (-14 348 Gg CO₂) is also greater than other categories in the UNFCCC key category analysis.

A 1.7 USING THE UNCERTAINTY ANALYSIS TO PLAN IMPROVEMENTS IN THE PREPARATION OF THE INVENTORY

The key category analysis is used to prioritise and plan improvements. The approach the UK takes to achieve this is described in **Chapter 1, Section 1.2.2.4. Table 1.7** to **Table 1.11** in **Chapter 1** show the key category summary tables.

A 1.8 TABLE NIR 3, AS CONTAINED IN THE ANNEX TO DECISION 6/CMP.3

Table A 1.8.1 below is Table NIR 3, containing a summary overview for Key Categories for Land Use, Land-Use Change and Forestry Activities under the Kyoto Protocol³.

Table A 1.8.1 Table NIR 3. Summary overview for Key Categories for Land Use, Land-Use Change and Forestry Activities under the Kyoto Protocol

KEY CATEGORIES OF EMISSIONS AND REMOVALS	GAS	CRITERIA USED FOR KEY CATEGORY IDENTIFICATION			COMMENTS ⁽³⁾
		Associated category in UNFCCC inventory ⁽¹⁾ is key (indicate which category)	Category contribution is greater than the smallest category considered key in the UNFCCC inventory ^{(1), (4)} (including LULUCF)	Other ⁽²⁾	
Specify key categories according to the national level of disaggregation used ⁽¹⁾					
Afforestation and Reforestation	CO ₂	Conversion to forest land	Yes	Associated UNFCCC category (4A) is key	The associated UNFCCC inventory category is a key category and the Afforestation and Reforestation category contribution is greater than the smallest UNFCCC key category.

³ Table NIR 3 can be found in FCCC/KP/CMP/2007/9/Add.2.

KEY CATEGORIES OF EMISSIONS AND REMOVALS	GAS	CRITERIA USED FOR KEY CATEGORY IDENTIFICATION			COMMENTS ⁽³⁾
		Associated category in UNFCCC inventory ⁽¹⁾ is key (indicate which category)	Category contribution is greater than the smallest category considered key in the UNFCCC inventory ^{(1), (4)} (including LULUCF)	Other ⁽²⁾	
Deforestation	CO ₂	Conversion to cropland, Conversion to grassland, Conversion to settlements	No	Associated UNFCCC categories (4B, 4C and 4E) are key	The associated UNFCCC categories (4B Cropland, 4C Grassland and 4E Settlements) are key categories and the Deforestation category contribution is greater than the smallest UNFCCC category.
Forest Management	CO ₂	Conversion to forest land	Yes	Associated UNFCCC category (4A) is key	The associated UNFCCC inventory category is a key category and the Forest Management category contribution is greater than the smallest UNFCCC key category.

(1) See section 5.4 of the IPCC good practice guidance for LULUCF

(2) This should include qualitative consideration as per Section 5.4.3 of the IPCC Good Practice Guidance for LULUCF or any other criteria

(3) Describe the criteria identifying the category as key

- (4) If the emissions or removals of the category exceed the emissions of the smallest category identified as key in the UNFCCC inventory (including LULUCF), Parties should indicate YES. If not, Parties should indicate NO

ANNEX 2: Assessment of Uncertainty

Uncertainty estimates are calculated using two methods: Approach 1 (error propagation) and Approach 2 (Monte Carlo simulation).

The uncertainty assessment in this NIR continues a number of improvements that were introduced in the 2007 submission, including presenting estimates of uncertainties according to IPCC sector in addition to presenting estimates by direct greenhouse gas. Estimated uncertainty presented in National Communication categories (which are consistent with the UK's Carbon Budgets sectors) are not reported here, since the categories are not consistent with the requirements of the UK's commitments under the UNFCCC and Kyoto Protocol.

The Monte Carlo method was reviewed and revised in the 2007 NIR, taking into account guidance from the 2006 Guidelines (IPCC, 2006), a summary of recommendations from the EUMM Workshop on Uncertainties held in Finland in 2005, and from an internal review of the uncertainty work. In the 2008 NIR, there was also a major review of the correlations used in the Monte Carlo simulation, which included discussions with the LULUCF sector experts.

A further review of the uncertainty parameters used within the industrial processes sector was carried out in 2010; the recommendations from this review were included in the 2011 submission of the NIR. The review followed recommendations from the UNFCCC ERT.

In 2013, a review was carried out of the uncertainty parameters ascribed to activity data, emission factors or emissions in the following sectors: energy (selection of subcategories), industry (selection of subcategories), agriculture (all subcategories), LULUCF (cropland, grassland & settlements) and waste (selection of categories). Changes were made to the uncertainty parameters in both the error propagation and Monte Carlo model, ensuring the parameters used were the same in both approaches where possible. Methodological changes were made to the error propagation method to optimise the use of the emission factor uncertainty data where categories are aggregated. Changes were made to the Monte Carlo model to integrate new UK specific uncertainty data about the probability distribution functions associated with emissions in some categories of the agriculture sector.

The overall method used to estimate uncertainties is described below, and the work to improve the accuracy of the uncertainty analysis continues. The key category analysis used data from the uncertainty analysis, and the results of the key category analysis are given in **Annex 1**.

A 2.1 ESTIMATION OF UNCERTAINTIES USING AN ERROR PROPAGATION APPROACH (APPROACH 1)

The IPCC 2006 Guidelines defines error propagation and Monte Carlo modelling approaches to estimating uncertainties in national greenhouse gas inventories. The results of the error propagation approach are shown in **Tables of uncertainty** estimates from the error propagation approach

Table A 2.1.1. Uncertainties are then estimated for these categories. The uncertainties used in the error propagation approach are not exactly the same as those used in the Monte Carlo Simulation since the error propagation source categorisation is less detailed and has a more simplistic approach to uncertainties. The approach 1 uncertainties is not concerned with

asymmetric uncertainties, or variations in uncertainty in the time series unlike the Monte Carlo approach which takes into account these factors.

A 2.1.1 Previous Improvements to the Error Propagation Model

The uncertainty analysis is kept under review and is refined to reflect improvements in the scientific understanding of uncertainties, and in methods. There have been a number of major improvements in the uncertainty analysis. Improvements made prior to the 2014NIR are not listed here, but can be found in the 2014NIR.

- **2014 NIR peer review (Abbott, 2014).** Uncertainty parameters ascribed to activity data, emission factors or emissions in the following sectors were updated: energy (selection of subcategories), industry (selection of subcategories), agriculture (all subcategories) and waste (categories relating to waste combustion). Several sources of data were used: DECC DUKES publication, EU ETS detailed returns, and expert elicitation.
- **2014 NIR peer review (Abbott, 2014).** Methodological changes were made to the error propagation method to optimise the use of the emission factor uncertainty data where categories are aggregated. In our uncertainty analysis, we used an aggregated emission factor uncertainty for each fuel type. The aggregated emission factor was selected as the most representative value from the component sectors. We have improved on this by calculating the aggregate emission factor uncertainties as:

$$U_f = \sqrt{\frac{\sum_i U_{fi}^2 E_i^2}{\sum_i E_i^2}}$$

where U_f are the emission factor uncertainties and E is the emissions from the component sectors, i .

A 2.1.2 Review of Changes Made to the Error Propagation Model since the last NIR

In response to ERT recommendations and significant changes to sectoral breakdown and new sources due to the changes introduced by the 2006 IPCC guidelines the UK has reviewed its approach towards the Approach 1 uncertainties. The main results of the review are as follows:

- F-gases are considered at the 2nd sector level instead of the 1st (i.e. 2B, 2C, 2E, 2F and 2G instead of just 2). This change is in response to ERT recommendations to treat sources of F-gases in more sectoral detail in the uncertainty model.
- A number of new sectors have been included, and the coverage of many sectors has changed. In light of this the UK reviewed most of the uncertainty values used and made estimates of uncertainty for previously unconsidered sectors. A number of expert judgements were requested, followed by a discussion involving the sector experts to determine the final values of uncertainties to use.

A 2.1.3 Key Categories

In the UK inventory, certain source categories are particularly significant in terms of their contribution to the overall uncertainty of the inventory. These key source categories have been identified so that the resources available for inventory preparation may be prioritised, and the

best possible estimates prepared for the most significant source categories. We have used the method described in Section 4.3.2 of the 2006 IPCC Guidelines Volume 1 General Guidance and Reporting (*Approach 2 to identify key categories*).

The results of this key category analysis can be found in **Annex 1**.

A 2.1.4 Tables of uncertainty estimates from the error propagation approach

Table A 2.1.1 Summary of error propagation uncertainty estimates including LULUCF, base year to the latest reported year

IPCC Category	Gas	Base year emissions (Gg CO ₂ e)	2013 emissions (Gg CO ₂ e)	Activity data uncertainty (%)	Emission factor uncertainty (%)	Combined uncertainty (%)	Contribution to variance by Category in 2013	Type A sensitivity	Type B sensitivity	Uncertainty in trend in national emissions introduced by emission factor / estimation parameter uncertainty	Uncertainty in trend in national emissions introduced by activity data uncertainty	Uncertainty introduced into the trend in total national emissions
1A_Coal	CO ₂	244,317.59	136,994.51	0.86%	1.96%	2.1%	0.0000	4.1565%	16.8062%	0.0813%	0.2055%	0.0005%
1A_Natural Gas	CO ₂	107,734.56	154,663.64	1.17%	1.84%	2.2%	0.0000	9.7117%	18.9738%	0.1788%	0.3148%	0.0013%
1A_(Stationary)_Oil	CO ₂	93,760.90	44,829.31	5.48%	2.65%	6.1%	0.0000	2.5471%	5.4996%	0.0675%	0.4260%	0.0019%
1A_Other (waste)	CO ₂	228.07	2,426.36	1.51%	12.69%	12.8%	0.0000	0.2781%	0.2977%	0.0353%	0.0064%	0.0000%
2D_Non Energy Products from Fuels and Solvent Use	CO ₂	1,232.62	1,032.92	6.36%	32.32%	32.9%	0.0000	0.0209%	0.1267%	0.0068%	0.0114%	0.0000%
1A3a_Aviation Fuel	CO ₂	1,859.57	1,961.73	19.53%	3.22%	19.8%	0.0000	0.0810%	0.2407%	0.0026%	0.0665%	0.0000%
1A3_Other diesel	CO ₂	1,679.92	2,384.81	15.00%	2.00%	15.1%	0.0000	0.1483%	0.2926%	0.0030%	0.0621%	0.0000%
1A3c_Coal	CO ₂	-	32.60	20.00%	6.00%	20.9%	0.0000	0.0040%	0.0040%	0.0002%	0.0011%	0.0000%
1A3b_DERV	CO ₂	33,012.63	68,350.74	1.00%	2.00%	2.2%	0.0000	5.5487%	8.3851%	0.1110%	0.1186%	0.0003%
1A3b_Gasoline/ LPG	CO ₂	75,557.88	38,697.84	0.99%	1.99%	2.2%	0.0000	1.7378%	4.7474%	0.0345%	0.0667%	0.0001%
1A3d_Marine fuel	CO ₂	2,168.02	2,165.99	17.57%	1.76%	17.7%	0.0000	0.0796%	0.2657%	0.0014%	0.0660%	0.0000%
1A4_Peat	CO ₂	475.59	46.96	30.00%	10.00%	31.6%	0.0000	0.0351%	0.0058%	0.0035%	0.0024%	0.0000%

IPCC Category	Gas	Base year emissions (Gg CO ₂ e)	2013 emissions (Gg CO ₂ e)	Activity data uncertainty (%)	Emission factor uncertainty (%)	Combined uncertainty (%)	Contribution to variance by Category in 2013	Type A sensitivity	Type B sensitivity	Uncertainty in trend in national emissions introduced by emission factor / estimation parameter uncertainty	Uncertainty in trend in national emissions introduced by activity data uncertainty	Uncertainty introduced into the trend in total national emissions
1A4_Petroleum Coke	CO ₂	81.64	160.19	20.00%	15.00%	25.0%	0.0000	0.0126%	0.0197%	0.0019%	0.0056%	0.0000%
1B1_Solid Fuel Transformation	CO ₂	1,698.56	175.28	4.81%	4.75%	6.8%	0.0000	0.1243%	0.0215%	0.0059%	0.0015%	0.0000%
1B2_Oil & Natural Gas	CO ₂	5,777.92	3,789.72	4.57%	5.50%	7.1%	0.0000	0.0311%	0.4649%	0.0017%	0.0301%	0.0000%
2A1_Cement Production	CO ₂	7,295.26	4,029.11	1.00%	3.00%	3.2%	0.0000	0.1320%	0.4943%	0.0040%	0.0070%	0.0000%
2A2_Lime Production	CO ₂	1,462.05	1,239.15	0.00%	5.00%	5.0%	0.0000	0.0265%	0.1520%	0.0013%	0.0000%	0.0000%
2A4_Other_process_uses_of_carbonates	CO ₂	646.76	771.83	2.00%	3.00%	3.6%	0.0000	0.0392%	0.0947%	0.0012%	0.0027%	0.0000%
2A3_Glass_production	CO ₂	408.14	389.18	0.00%	5.00%	5.0%	0.0000	0.0127%	0.0477%	0.0006%	0.0000%	0.0000%
2B_Chemical industries	CO ₂	6,376.75	4,740.31	18.63%	3.15%	18.9%	0.0000	0.0341%	0.5815%	0.0011%	0.1532%	0.0002%
2C_Metal_Industries	CO ₂	7,391.77	5,013.31	1.20%	5.01%	5.2%	0.0000	0.0196%	0.6150%	0.0010%	0.0105%	0.0000%
3G_Liming	CO ₂	1,579.17	774.69	0.00%	20.90%	20.9%	0.0000	0.0405%	0.0950%	0.0085%	0.0000%	0.0000%
3H_Urea_application_to_agriculture	CO ₂	385.46	245.22	0.00%	50.00%	50.0%	0.0000	0.0030%	0.0301%	0.0015%	0.0000%	0.0000%
4A Forest Land	CO ₂	-16,016.07	-17,297.63	1.00%	30.00%	30.0%	0.0001	0.7472%	2.1220%	0.2242%	0.0300%	0.0005%
4B_Cropland	CO ₂	15,130.79	12,150.39	1.00%	45.00%	45.0%	0.0001	0.1915%	1.4906%	0.0862%	0.0211%	0.0001%
4C_Grassland	CO ₂	-3,677.20	-5,897.14	1.00%	45.00%	45.0%	0.0000	0.4078%	0.7234%	0.1835%	0.0102%	0.0003%
4D Wetland	CO ₂	481.73	300.49	1.00%	50.00%	50.0%	0.0000	0.0045%	0.0369%	0.0022%	0.0005%	0.0000%

IPCC Category	Gas	Base year emissions (Gg CO ₂ e)	2013 emissions (Gg CO ₂ e)	Activity data uncertainty (%)	Emission factor uncertainty (%)	Combined uncertainty (%)	Contribution to variance by Category in 2013	Type A sensitivity	Type B sensitivity	Uncertainty in trend in national emissions introduced by emission factor / estimation parameter uncertainty	Uncertainty in trend in national emissions introduced by activity data uncertainty	Uncertainty introduced into the trend in total national emissions
4E Settlements	CO ₂	6,919.10	5,876.98	1.00%	50.00%	50.0%	0.0000	0.1269%	0.7210%	0.0635%	0.0102%	0.0000%
4F Other Land (NOT REPORTED)	CO ₂	-	0.02	0.00%	50.00%	50.0%	0.0000	0.0000%	0.0000%	0.0000%	0.0000%	0.0000%
4G Other Activities	CO ₂	40.72	-1,121.03	1.00%	45.00%	45.0%	0.0000	0.1410%	0.1375%	0.0635%	0.0019%	0.0000%
5C_Waste Incineration	CO ₂	1,304.38	264.30	15.07%	20.41%	25.4%	0.0000	0.0796%	0.0324%	0.0162%	0.0069%	0.0000%
1A1 & 1A2 & 1A4 & 1A5_Other Combustion	CH ₄	2,079.93	966.42	0.74%	36.88%	36.9%	0.0000	0.0600%	0.1186%	0.0221%	0.0012%	0.0000%
1A3a_Aviation Fuel	CH ₄	4.17	0.86	14.78%	58.00%	59.8%	0.0000	0.0003%	0.0001%	0.0001%	0.0000%	0.0000%
1A3_Other diesel	CH ₄	2.75	2.60	15.00%	130.00%	130.9%	0.0000	0.0001%	0.0003%	0.0001%	0.0001%	0.0000%
1A3c_Coal	CH ₄	-	0.86	20.00%	110.00%	111.8%	0.0000	0.0001%	0.0001%	0.0001%	0.0000%	0.0000%
1A3b_DERV	CH ₄	127.77	16.98	1.00%	130.00%	130.0%	0.0000	0.0089%	0.0021%	0.0116%	0.0000%	0.0000%
1A3b_Gasoline/ LPG	CH ₄	630.55	38.64	0.99%	74.11%	74.1%	0.0000	0.0494%	0.0047%	0.0366%	0.0001%	0.0000%
1A3d_Marine fuel	CH ₄	2.22	4.90	19.55%	127.11%	128.6%	0.0000	0.0004%	0.0006%	0.0005%	0.0002%	0.0000%
1B1_Solid Fuel Transformation	CH ₄	0.18	3.85	0.01%	49.60%	49.6%	0.0000	0.0005%	0.0005%	0.0002%	0.0000%	0.0000%
1B1_Coal Mining	CH ₄	21,788.17	1,682.41	2.00%	20.00%	20.1%	0.0000	1.6637%	0.2064%	0.3327%	0.0058%	0.0011%
1B2_Offshore Oil& Gas	CH ₄	2,164.64	1,290.40	4.20%	16.82%	17.3%	0.0000	0.0275%	0.1583%	0.0046%	0.0094%	0.0000%

IPCC Category	Gas	Base year emissions (Gg CO ₂ e)	2013 emissions (Gg CO ₂ e)	Activity data uncertainty (%)	Emission factor uncertainty (%)	Combined uncertainty (%)	Contribution to variance by Category in 2013	Type A sensitivity	Type B sensitivity	Uncertainty in trend in national emissions introduced by emission factor / estimation parameter uncertainty	Uncertainty in trend in national emissions introduced by activity data uncertainty	Uncertainty introduced into the trend in total national emissions
1B2_Natural Gas Transmission	CH ₄	10,168.29	4,396.86	3.00%	20.00%	20.2%	0.0000	0.3335%	0.5394%	0.0667%	0.0229%	0.0000%
2A_Minerals_industry	CH ₄	31.27	4.52	0.00%	100.00%	100.0%	0.0000	0.0021%	0.0006%	0.0021%	0.0000%	0.0000%
2B_Chemical Industry	CH ₄	213.70	104.11	0.00%	20.00%	20.0%	0.0000	0.0056%	0.0128%	0.0011%	0.0000%	0.0000%
2C_Iron & Steel Production	CH ₄	36.89	20.49	1.93%	48.38%	48.4%	0.0000	0.0007%	0.0025%	0.0003%	0.0001%	0.0000%
2D_Non-energy_Products_from_Fuels_and_Solvent_Use	CH ₄	0.69	0.38	50.00%	50.00%	70.7%	0.0000	0.0000%	0.0000%	0.0000%	0.0000%	0.0000%
3A_Enteric Fermentation	CH ₄	28,227.49	23,621.80	0.10%	20.00%	20.0%	0.0001	0.4743%	2.8979%	0.0949%	0.0041%	0.0001%
3B_Manure Management	CH ₄	4,472.78	3,471.50	0.10%	30.00%	30.0%	0.0000	0.0419%	0.4259%	0.0126%	0.0006%	0.0000%
3F_Field Burning	CH ₄	205.48	-	25.00%	50.00%	55.9%	-	0.0176%	0.0000%	0.0088%	0.0000%	0.0000%
4A_Forest_Land	CH ₄	3.72	5.04	1.00%	55.00%	55.0%	0.0000	0.0003%	0.0006%	0.0002%	0.0000%	0.0000%
4B_Cropland	CH ₄	0.11	0.14	1.00%	55.00%	55.0%	0.0000	0.0000%	0.0000%	0.0000%	0.0000%	0.0000%
4C_Grassland	CH ₄	11.29	28.46	1.00%	55.00%	55.0%	0.0000	0.0025%	0.0035%	0.0014%	0.0000%	0.0000%
4E_Settlements	CH ₄	5.89	1.99	1.00%	20.00%	20.0%	0.0000	0.0003%	0.0002%	0.0001%	0.0000%	0.0000%

IPCC Category	Gas	Base year emissions (Gg CO ₂ e)	2013 emissions (Gg CO ₂ e)	Activity data uncertainty (%)	Emission factor uncertainty (%)	Combined uncertainty (%)	Contribution to variance by Category in 2013	Type A sensitivity	Type B sensitivity	Uncertainty in trend in national emissions introduced by emission factor / estimation parameter uncertainty	Uncertainty in trend in national emissions introduced by activity data uncertainty	Uncertainty introduced into the trend in total national emissions
5A_Solid Waste Disposal	CH ₄	62,749.72	16,681.83	15.00%	46.00%	48.4%	0.0002	3.3381%	2.0465%	1.5355%	0.4341%	0.0255%
5D_Wastewater Handling	CH ₄	4,169.64	3,359.33	10.00%	25.00%	26.9%	0.0000	0.0541%	0.4121%	0.0135%	0.0583%	0.0000%
5C_Waste Incineration	CH ₄	137.22	9.95	5.00%	50.00%	50.2%	0.0000	0.0106%	0.0012%	0.0053%	0.0001%	0.0000%
5B_Biological_treatment_of_solid_waste	CH ₄	5.48	715.72	30.00%	99.50%	103.9%	0.0000	0.0873%	0.0878%	0.0869%	0.0373%	0.0001%
1A1 & 1A2 & 1A4 & 1A5_Other Combustion	N ₂ O	4,209.03	2,754.34	0.61%	60.91%	60.9%	0.0000	0.0235%	0.3379%	0.0143%	0.0029%	0.0000%
1A3a_Aviation Fuel	N ₂ O	17.60	18.56	19.52%	107.37%	109.1%	0.0000	0.0008%	0.0023%	0.0008%	0.0006%	0.0000%
1A3_Other diesel	N ₂ O	31.12	61.14	15.00%	130.00%	130.9%	0.0000	0.0048%	0.0075%	0.0063%	0.0016%	0.0000%
1A3c_Coal	N ₂ O	-	0.08	20.00%	110.00%	111.8%	0.0000	0.0000%	0.0000%	0.0000%	0.0000%	0.0000%
1A3b_DERV	N ₂ O	279.80	758.86	1.00%	130.00%	130.0%	0.0000	0.0691%	0.0931%	0.0898%	0.0013%	0.0001%
1A3b_Gasoline/ LPG	N ₂ O	862.61	150.48	0.99%	74.14%	74.2%	0.0000	0.0556%	0.0185%	0.0412%	0.0003%	0.0000%
1A3d_Marine fuel	N ₂ O	16.26	16.22	17.60%	114.38%	115.7%	0.0000	0.0006%	0.0020%	0.0007%	0.0005%	0.0000%
1B1_Fugitive_Emissions_from_Solid_Fuels	N ₂ O	0.09	0.04	1.00%	118.00%	118.0%	0.0000	0.0000%	0.0000%	0.0000%	0.0000%	0.0000%
1B2_Oil & Natural Gas	N ₂ O	40.75	30.79	4.81%	105.83%	105.9%	0.0000	0.0003%	0.0038%	0.0003%	0.0003%	0.0000%

IPCC Category	Gas	Base year emissions (Gg CO ₂ e)	2013 emissions (Gg CO ₂ e)	Activity data uncertainty (%)	Emission factor uncertainty (%)	Combined uncertainty (%)	Contribution to variance by Category in 2013	Type A sensitivity	Type B sensitivity	Uncertainty in trend in national emissions introduced by emission factor / estimation parameter uncertainty	Uncertainty in trend in national emissions introduced by activity data uncertainty	Uncertainty introduced into the trend in total national emissions
2B3_Adipic Acid Production	N ₂ O	19,934.61	-	2.00%	100.00%	100.0%	-	1.7110%	0.0000%	1.7110%	0.0000%	0.0293%
2B2_Nitric Acid Production	N ₂ O	3,860.26	42.91	10.00%	100.00%	100.5%	0.0000	0.3261%	0.0053%	0.3261%	0.0007%	0.0011%
2B8_Petrochemical_and_Carbon_Black_Production	N ₂ O	1.79	1.53	10.00%	100.00%	100.5%	0.0000	0.0000%	0.0002%	0.0000%	0.0000%	0.0000%
2C_Iron & Steel	N ₂ O	62.46	45.23	1.00%	118.00%	118.0%	0.0000	0.0002%	0.0055%	0.0002%	0.0001%	0.0000%
2D_Non-energy_Products_from_Fuels_and_Solvent_Use	N ₂ O	1.64	0.91	50.00%	100.00%	111.8%	0.0000	0.0000%	0.0001%	0.0000%	0.0001%	0.0000%
2G_Other_Product_Manufacture_and_Use	N ₂ O	0.44	0.49	100.00%	100.00%	141.4%	0.0000	0.0000%	0.0001%	0.0000%	0.0001%	0.0000%
3B_Manure Management	N ₂ O	2,403.46	1,891.28	1.00%	254.00%	254.0%	0.0001	0.0257%	0.2320%	0.0652%	0.0033%	0.0000%
3D_Agricultural Soils	N ₂ O	23,153.96	19,513.90	1.00%	259.00%	259.0%	0.0078	0.4060%	2.3939%	1.0515%	0.0339%	0.0111%
3F_Field Burning	N ₂ O	63.50	-	25.00%	230.00%	231.4%	-	0.0055%	0.0000%	0.0125%	0.0000%	0.0000%
4A_Forest_land	N ₂ O	48.26	50.29	1.00%	40.00%	40.0%	0.0000	0.0020%	0.0062%	0.0008%	0.0001%	0.0000%
4D_Grassland	N ₂ O	3.83	0.50	1.00%	100.00%	100.0%	0.0000	0.0003%	0.0001%	0.0003%	0.0000%	0.0000%
4E_Settlements	N ₂ O	369.67	318.07	1.00%	20.00%	20.0%	0.0000	0.0073%	0.0390%	0.0015%	0.0006%	0.0000%

IPCC Category	Gas	Base year emissions (Gg CO ₂ e)	2013 emissions (Gg CO ₂ e)	Activity data uncertainty (%)	Emission factor uncertainty (%)	Combined uncertainty (%)	Contribution to variance by Category in 2013	Type A sensitivity	Type B sensitivity	Uncertainty in trend in national emissions introduced by emission factor / estimation parameter uncertainty	Uncertainty in trend in national emissions introduced by activity data uncertainty	Uncertainty introduced into the trend in total national emissions
5C_Waste Incineration	N ₂ O	29.85	44.28	7.00%	230.00%	230.1%	0.0000	0.0029%	0.0054%	0.0066%	0.0005%	0.0000%
5B_Biological_treatment_of_solid_waste	N ₂ O	4.90	608.32	30.00%	90.00%	94.9%	0.0000	0.0742%	0.0746%	0.0668%	0.0317%	0.0001%
5D_Wastewater Handling	N ₂ O	1,111.13	1,063.03	10.00%	248.00%	248.2%	0.0000	0.0350%	0.1304%	0.0868%	0.0184%	0.0001%
2C_Metal_Industries	SF ₆	387.17	146.18	5.00%	5.00%	7.1%	0.0000	0.0153%	0.0179%	0.0008%	0.0013%	0.0000%
2G_Other_Product_Manufacture_and_Use	SF ₆	877.20	455.37	0.00%	5.92%	5.9%	0.0000	0.0194%	0.0559%	0.0012%	0.0000%	0.0000%
2B_Chemical_industry	HFCs	17,680.04	46.11	0.00%	10.00%	10.0%	0.0000	1.5119%	0.0057%	0.1512%	0.0000%	0.0002%
2C_Metal_Industries	HFCs	-	2.61	5.00%	10.00%	11.2%	0.0000	0.0003%	0.0003%	0.0000%	0.0000%	0.0000%
2E_Electronics_Industry	HFCs	8.73	14.47	0.00%	47.15%	47.1%	0.0000	0.0010%	0.0018%	0.0005%	0.0000%	0.0000%
2F_Product_Uses_as_Substitutes_for_ODS	HFCs	1,870.17	16,200.73	8.12%	8.22%	11.6%	0.0000	1.8269%	1.9875%	0.1502%	0.2282%	0.0007%
2B_Chemical_industry	PFCs	113.90	112.13	0.00%	10.00%	10.0%	0.0000	0.0040%	0.0138%	0.0004%	0.0000%	0.0000%
2C_Metal_Industries	PFCs	333.43	6.95	0.00%	20.00%	20.0%	0.0000	0.0278%	0.0009%	0.0056%	0.0000%	0.0000%
2F_Product_Uses_as_Substitutes_for_ODS	PFCs	0.44	-	0.00%	25.00%	25.0%	-	0.0000%	0.0000%	0.0000%	0.0000%	0.0000%

IPCC Category	Gas	Base year emissions (Gg CO ₂ e)	2013 emissions (Gg CO ₂ e)	Activity data uncertainty (%)	Emission factor uncertainty (%)	Combined uncertainty (%)	Contribution to variance by Category in 2013	Type A sensitivity	Type B sensitivity	Uncertainty in trend in national emissions introduced by emission factor / estimation parameter uncertainty	Uncertainty in trend in national emissions introduced by activity data uncertainty	Uncertainty introduced into the trend in total national emissions	
2G_Other_Product_Manufacture_and_Use	PFCs	149.17	134.30	0.00%	47.15%	47.1%	0.0000	0.0037%	0.0165%	0.0017%	0.0000%	0.0000%	
2E_Electronics_Industry	NF ₃	0.83	0.36	0.00%	47.15%	47.1%	0.0000	0.0000%	0.0000%	0.0000%	0.0000%	0.0000%	
2B1_Ammonia_Production	N ₂ O	0.31	0.27	2.00%	50.00%	50.0%	0.0000	0.0000%	0.0000%	0.0000%	0.0000%	0.0000%	
4B_Cropland	N ₂ O	649.15	329.94	1.00%	55.00%	55.0%	0.0000	0.0153%	0.0405%	0.0084%	0.0006%	0.0000%	
4C_Grassland	N ₂ O	12.10	9.15	1.00%	35.00%	35.0%	0.0000	0.0001%	0.0011%	0.0000%	0.0000%	0.0000%	
Total		815,143.98	570,451.65				0.0086					0.0749%	
							Percentage uncertainty in total inventory:	9.3%				Trend uncertainty:	2.7%

A 2.2 ESTIMATION OF UNCERTAINTY BY SIMULATION (APPROACH 2)

A 2.2.1 Overview of the Method

Quantitative estimates of the uncertainties in the emissions were calculated using a Monte Carlo simulation. This corresponds to the IPCC Approach 2 method, discussed in the 2006 Guidelines (IPCC, 2006). The background to the implementation of the Monte Carlo simulation is described in detail by Eggleston *et al* (1998), with the estimates reported here revised to reflect changes in the latest inventory and improvements made in the model. This section gives a brief summary of the methodology, assumptions and results of the simulation.

The computational procedure is detailed below.

- A probability distribution function (PDF) was allocated to each unique emission factor and piece of activity data. The PDFs were mostly normal or log-normal, with one log-logistic PDF. The parameters of the PDFs were set by analysing the available data on emission factors and activity data, and by expert judgement.
- A calculation was set up to estimate the total emissions of each gas for the years 1990 and the latest reported year.
- Using the software tool @RISK™, each PDF was sampled at least 20,000 times, such that the emission calculations performed produced a converged output distribution.
- The distribution of errors in the parameter values was calculated from the difference between 2.5 and 97.5 percentile values in the distribution, as a percentage of the distribution mean.
- The uncertainties used for the fuel activity data were estimated from the statistical difference between the total supply and demand for each fuel. Data on the statistical difference between supply and demand for individual sectors are not available. This means that the quoted uncertainties in

- **Table A 2.3.1** refer to the total fuel consumption rather than the consumption by a particular sector, e.g. coal consumed in the residential sector. Hence, to avoid underestimating uncertainties, it was necessary to correlate the uncertainties used for the same fuel in different sectors.
- The uncertainty in the trend between 1990 and the latest reported year, according to gas, was also estimated.

A 2.2.2 Summary of Recent Improvements to the Monte Carlo Model

In 2013, changes were made to the Monte Carlo model to integrate new UK specific uncertainty data to activity data, emission factors or emissions. The PDFs ascribed to each category in the model were also reviewed, and the PDFs associated with emissions in some categories of the agriculture sector were changed.

A 2.2.3 Review of changes made to the Monte Carlo model since the last NIR

Four main changes have been introduced into the model for the 2014 submission:

- Uncertainty parameters ascribed to activity data, emission factors or emissions in the following sectors were updated in the model: energy (selection of subcategories), industry (selection of subcategories), agriculture (all subcategories), LULUCF (cropland, grassland & settlements) and waste (selection of categories). Several sources of data were used: DECC DUKES publication, EU ETS detailed returns, and expert elicitation.
- The PDF for category 4D agricultural soils was changed, using information based on recent UK research.
- Uncertainties introduced on trends, 1990 and the latest inventory totals are calculated from the difference between 2.5 and 97.5 percentiles as a percentage of the trend, 1990 or latest inventory year annual emission distribution mean. This is in contrast to previous versions of these data in which uncertainties were calculated from the standard deviation of the results, and is considered an improvement on the previous approach particularly in relation to non-normally distributed uncertainties, for which the standard deviation is less convergent than the 95 percentile confidence limits. The impact of this change is most notable for IPCC sectors 4D and 6B2, for which uncertainties follow log-normal and log-logistic distribution functions respectively. The measure of uncertainty expressed for 2012 total emissions is reduced by this change from 95% to 82% for 'Agriculture - N₂O' (IPCC sectors 4B12-14 & 4D), and from 260% to 189% for 6B2. Similarly, the uncertainty expressed for N₂O and Agriculture total emissions is reduced when derived via this approach, from 82% to 69% and 53% to 44% respectively.
- For LULUCF, the calculation of the 'Range of likely % change' in these results is based on the difference between 1990 and 2012 emissions in each iteration as a percentage of the average 1990 emission across all iterations. The previous approach, which calculated the percentage change from 1990 to the latest year in each iteration, led to

non-convergence of results for the trend in LULUCF emissions, where statistical characteristics were dominated by iterations with 1990 emissions close to zero resulting in disproportionately high percentage changes.

A 2.2.4 Methodological details of the Monte Carlo model

A 2.2.4.1 Uncertainty Distributions

A 2.2.4.1.1 Distributions

Nearly all of the distributions of emissions from sources in the inventory are modelled using normal or log normal distributions, with one log-logistic PDF and one custom PDF. The log-logistic function was provided by Rothamsted Research for N₂O emissions from agricultural soils, and is based on the data presented in Milne *et al.* (2014).

A 2.2.4.1.2 Custom distributions

Emissions from landfill have been modelled using a custom distribution. Aitchison *et al.* (cited in Eggelston *et al.*, 1998) estimated the uncertainty for landfill emissions using Monte Carlo analysis and found it to be skewed. The distribution histogram was used to generate an empirical distribution of emissions. For this study we examined the distribution and fitted a log normal distribution to Aitchison's data. The emissions are scaled according to the mean estimate of landfill emissions for each year.

A 2.2.4.2 Correlations

The Monte Carlo model contains a number of correlations. Omitting these correlations will not affect uncertainties on emission totals in 1990 or the latest inventory year, but would lead to the uncertainties on trends being overestimated, by negating the dependence on base year emissions in the methodology for estimating subsequent emissions. These correlations were not included in the very early versions of the Monte Carlo model used in the UK NIR, and were introduced over the years to improve the accuracy of the predicted uncertainties. The trend uncertainty in the Monte Carlo model is particularly sensitive to some correlations, for example, the correlation across years in emissions of N₂O from agricultural soils. Other correlations have a less marked influence.

The type and implementation of the correlations has been examined as part of a review (Abbott *et al.*, 2007). The sensitivity analysis that we have completed on the Monte Carlo model suggests that the uncertainties are not sensitive to the correlations between emission factors for fuel used, and for LULUCF sources.

A 2.2.4.2.1 Across years

In running this simulation it was necessary to make assumptions about the degree of correlation between sources in 1990 and the latest reported year. If source emission factors are correlated this will have the effect of reducing the trend uncertainty.

The model has been designed to aggregate activities and emission factors where possible, and the correlations included are listed at the start of the sections presenting uncertainties according to gas.

The trend estimated by the Monte Carlo model is particularly sensitive to N₂O emissions from agricultural soils. Correlations are also included for N₂O emissions from sewage sludge,

calculated from a lognormal distribution. The LULUCF correlations are discussed below. Other correlations are listed at the start of the sections presenting uncertainties according to gas.

All correlations in the LULUCF sector were reviewed (see 2008 NIR for full details) and the revised and assumptions have been implemented in the current Monte Carlo model. This review found that the emission sources and carbon sinks in this sector were not correlated with each other, but were correlated across inventory years.

A 2.2.4.2.2 *Between Sources in the same year*

Where we have estimated the uncertainty on the activity data based on statistical difference produced by DECC and reported in DUKES, it has been necessary to correlate the fuel use for all sources using the same fuel.

A 2.2.4.2.3 *Simulation Method*

Following recommendations in the 2006 IPCC Guidelines, the model uses a true Monte Carlo sampling method.

A 2.2.4.2.4 *Treatment of categories where emissions are zero*

The original Monte Carlo model contained a number of sources where the emissions were zero, but uncertainties were still allocated to the activity data and emission factors. These zero emissions existed for several reasons:

- Emissions occurred in 1990 but were absent in later years;
 - The activity had been banned (for example, burning of agricultural straw residues);
 - Emissions had been transferred to another sector (for example MSW emissions from waste from IPCC category 6C to 1A1a.); and
- Because data had been included in the analysis for completeness where either the emission factor or the activity data were zero thus leading to a zero emission.

The estimated uncertainties were unaffected when the 'zero emissions' were removed from the model.

A 2.2.4.2.5 *Aggregation of categories*

For the new Monte Carlo model, the detailed data from the GHG inventory was aggregated where appropriate in order to minimise the number of sources used in the calculation. Emissions were aggregated where possible for fuels (any emission arising from combustion), by activity data type e.g. coal, petrol, natural gas, and by emission factor. In doing so, the data are also being correlated as any uncertainty in the emission factor is then applied once, to all appropriate emissions, and the same is true of the activity data. Minimising the number of calculations performed in the Monte Carlo simulation ensures that the overall uncertainty is more accurately estimated by the model.

A 2.2.4.3 *F-gas uncertainties*

Estimated emissions and projections of F-gases were reviewed and updated (AEA, 2008). This work also included an update to the uncertainty analysis, which has been taken into account in the overall uncertainty analysis for the greenhouse gas inventory. Uncertainties from the Refrigeration and Air Conditioning (RAC) sector were taken from ICF (2011).

A 2.2.5 Quality Control Checks on the Monte Carlo Model Output

A number of quality control checks are completed as part of the uncertainty analysis.

a) Checks against totals of the national emissions

To ensure the emissions in the Monte Carlo model closely agree with the reported totals in the NIR, the emissions in the model were checked against the national totals both before and after the simulation was run. The central estimates from the model are expected to be similar to the reported emissions totals, but are not expected to match exactly.

b) Inter-comparison between the output of the error propagation and Monte Carlo models

A formal check to compare the output of the error propagation and Monte Carlo model is completed. The results of this comparison are discussed in **Section A 2.6**.

c) Calculation of uncertainty on the total

The uncertainty on the 1990 and the 2011 emissions was calculated using two different methods;

- i) Using $\frac{2s.d}{\mu}$
- ii) Using $\frac{(97.5 \text{ percentile} - 2.5 \text{ percentile})}{2 \times \mu}$

The first method uses the standard deviation calculated by @RISK and the mean to give an overall uncertainty, while the second method averages out the implied standard deviation(s) given by the percentiles quoted. When a distribution is completely normally distributed, the two methods will give approximately the same results. However, when a distribution is skewed the first method converges to a much lower precision, since the variance is dominated by outliers. The overall uncertainty quoted in **Table A 2.4.1** is calculated using the second method so that uncertainties in sectors that show a skewed distribution (such as agricultural soils and N₂O) are better represented.

Calculating the uncertainty using both of these methods allows us to check that the Monte Carlo analysis is behaving in the way we would expect, and that convergence of the distributions is being achieved. Comparing the results using both calculations showed that the uncertainties were almost the same for gases where the distributions used were predominantly normal, but higher for N₂O and the GWP weighted total, as expected.

A 2.3 UNCERTAINTIES ACCORDING TO GAS

The following sections present the uncertainties in emissions, and the trend in emissions according to gas.

A 2.3.1 Carbon Dioxide Emission Uncertainties

A 2.3.1.1 General Considerations

The uncertainty in the trend between 1990 and 2013 is given in **Section A 2.4.2**. In running this simulation it was necessary to make assumptions about the degree of correlation between

sources in 1990 and 2013. If source emission factors are correlated this will have the effect of reducing the trend uncertainty.

A 2.3.1.2 Uncertainty Parameters

Two tables are provided in this section – a table of uncertainties in the activity data and emission factors for the major fuels used to estimate emissions of carbon dioxide, and a table of the same parameters for “non-fuels”. These non-fuels relate to emissions from a range of sources, including the following:

- The release of carbon from the breakdown of pesticides and detergents.
- Use of natural gas for the production of ammonia.

In some cases the individual uncertainties for the activity data and the emission factor are unknown, but the uncertainty on the total emission is known. In these cases, the uncertainties are listed in the column marked “uncertainty in emission”.

Table A 2.3.1 Uncertainties in the activity data and emission factors for fuels used in the carbon dioxide (CO₂) inventory

Category	Fuel	1990		2013		Notes
		Activity uncertainty (%)	Emission factor uncertainty (%)	Activity uncertainty (%)	Emission factor uncertainty (%)	
1A1	Blast Furnace Gas	1.5%	10.0%	1.0%	10.0%	
1A1	Coke Oven Coke	7.8%	10.0%	1.0%	10.0%	
1A1	Coke Oven Gas	1.5%	10.0%	1.0%	10.0%	
1A1	Colliery Methane	5.0%	5.0%	5.0%	5.0%	
1A1	Gas/Diesel Oil	1.8%	2.1%	1.8%	2.1%	
1A1	Liquefied Petroleum Gas	25.7%	2.1%	2.5%	2.1%	
1A1	Motor Gasoline	7.8%	2.1%	2.5%	2.1%	
1A1	Municipal Solid Waste	1.0%	15.0%	1.0%	15.0%	
1A1	Naphtha	10.0%	5.0%	10.0%	5.0%	
1A1	Natural Gas	2.8%	2.0%	1.0%	2.0%	
1A1	Orimulsion	5.0%	5.0%	5.0%	5.0%	
1A1	Other Bituminous Coal	1.5%	2.0%	1.0%	2.0%	
1A1	Other Kerosene	7.8%	5.0%	1.3%	5.0%	
1A1	Other Oil: Other	11.9%	5.0%	10.0%	5.0%	
1A1	Petroleum Coke	7.8%	10.0%	5.0%	10.0%	
1A1	Refinery Gas	50.0%	20.0%	25.0%	15.0%	
1A1	Residual Fuel Oil	5.5%	2.6%	1.3%	2.6%	

Category	Fuel	1990		2013		Notes
		Activity uncertainty (%)	Emission factor uncertainty (%)	Activity uncertainty (%)	Emission factor uncertainty (%)	
1A1	Scrap Tyres	15.0%	10.0%	15.0%	10.0%	
1A2	Blast Furnace Gas	1.5%	10.0%	1.0%	10.0%	
1A2	Coke Oven Coke	3.0%	10.0%	1.0%	10.0%	
1A2	Coke Oven Gas	3.0%	10.0%	1.0%	10.0%	
1A2	Colliery Methane	5.0%	5.0%	5.0%	5.0%	
1A2	Gas/Diesel Oil	20.0%	2.0%	20.0%	2.0%	
1A2	Liquefied Petroleum Gas	25.7%	2.1%	2.5%	2.1%	
1A2	Motor Gasoline	20.0%	2.1%	20.0%	2.1%	
1A2	Municipal Solid Waste	25.0%	15.0%	5.0%	15.0%	
1A2	Natural Gas	2.8%	3.0%	1.0%	3.0%	
1A2	non-fuel combustion	50.0%	100.0%	50.0%	100.0%	
1A2	Other Bituminous Coal	5.0%	10.0%	5.0%	10.0%	
1A2	Other Kerosene	6.0%	2.0%	6.0%	2.0%	
1A2	Other Oil: Other	5.0%	50.0%	5.0%	3.0%	
1A2	Patent Fuel	10.0%	3.0%	10.0%	3.0%	
1A2	Petroleum Coke	25.0%	15.0%	20.0%	15.0%	
1A2	Refinery Gas	50.0%	15.0%	50.0%	15.0%	
1A2	Residual Fuel Oil	5.5%	2.1%	1.5%	2.1%	

Category	Fuel	1990		2013		Notes
		Activity uncertainty (%)	Emission factor uncertainty (%)	Activity uncertainty (%)	Emission factor uncertainty (%)	
1A2	Scrap Tyres	15.0%	10.0%	15.0%	10.0%	
1A3	Aviation Gasoline	20.0%	3.3%	20.0%	3.3%	
1A3	Jet Gasoline	20.0%	3.3%	20.0%	3.3%	
1A3	Other Bituminous Coal	20.0%	6.0%	20.0%	6.0%	
1A3	Other Gas/Diesel Oil	15.0%	2.0%	15.0%	2.0%	
1A3b	Gas/Diesel Oil	1.8%	2.0%	1.0%	2.0%	
1A3b	Liquefied Petroleum Gas	5.0%	2.0%	5.0%	2.0%	
1A3b	Motor Gasoline	1.0%	2.0%	1.0%	2.0%	
1A3d	Gas/Diesel Oil	20.0%	2.0%	20.0%	2.0%	
1A3d	Residual Fuel Oil	20.0%	2.0%	20.0%	2.0%	
1A4	Anthracite	1.5%	6.0%	1.0%	6.0%	
1A4	Coke Oven Coke	3.0%	10.0%	1.0%	10.0%	
1A4	Gas Works Gas	5.0%	5.0%	5.0%	5.0%	
1A4	Gas/Diesel Oil	30.0%	2.0%	30.0%	2.0%	
1A4	Liquefied Petroleum Gas	25.7%	2.1%	2.5%	2.1%	
1A4	Motor Gasoline	50.0%	2.0%	50.0%	2.0%	
1A4	Natural Gas	2.8%	3.0%	2.0%	3.0%	
1A4	Other Bituminous Coal	3.0%	10.0%	3.0%	10.0%	

Category	Fuel	1990		2013		Notes
		Activity uncertainty (%)	Emission factor uncertainty (%)	Activity uncertainty (%)	Emission factor uncertainty (%)	
1A4	Other Kerosene	3.0%	2.0%	3.0%	2.0%	
1A4	Patent Fuel	3.3%	3.0%	2.0%	3.0%	
1A4	Peat	30.0%	10.0%	30.0%	10.0%	
1A4	Petroleum Coke	20.0%	15.0%	20.0%	15.0%	
1A4	Residual Fuel Oil	5.5%	2.1%	3.0%	2.1%	
1A5	Gas/Diesel Oil	6.3%	2.1%	6.3%	2.1%	
1A5	Jet Gasoline	10.0%	3.0%	10.0%	3.0%	
1B1	Coke Oven Gas	1.5%	10.0%	1.0%	10.0%	
1B1	petroleum coke	20.0%	10.0%	20.0%	10.0%	
1B1	Other Bituminous Coal	1.5%	6.0%	1.5%	6.0%	
1B2a	non-fuel combustion	5.0%	6.0%	5.0%	6.0%	
1B2b	non-fuel combustion		14.1%		14.1%	<i>Approach 2 based on estimates of uncertainty in components of gas leakage estimates given in Williams, 1993, Table 4.7</i>
1B2c	non-fuel combustion	5.0%	6.0%	5.0%	6.0%	
2A1	non-fuel combustion	1.0%	3.0%	1.0%	3.0%	
2A2	non-fuel combustion	10.0%	5.0%	0.0%	5.0%	
2A3	non-fuel combustion	0.0%	5.0%	0.0%	5.0%	

Category	Fuel	1990		2013		Notes
		Activity uncertainty (%)	Emission factor uncertainty (%)	Activity uncertainty (%)	Emission factor uncertainty (%)	
2A4	non-fuel combustion	2.0%	3.0%	2.0%	3.0%	
2B	Coke	1.0%	10.0%	1.0%	10.0%	
2B	coke oven coke	0.0%	20.0%	0.0%	20.0%	
2B	Natural Gas	2.8%	1.3%	1.8%	1.3%	
2B	non-fuel combustion	2.0%	5.0%	2.0%	5.0%	
2B	OPG	0.0%	5.0%	0.0%	5.0%	
2B	petroleum coke	1.0%	10.0%	1.0%	10.0%	
2B	refinery gas	30.0%	5.0%	30.0%	5.0%	
2C	Blast Furnace Gas	2.0%	10.0%	2.0%	10.0%	
2C	Coke	2.0%	10.0%	2.0%	10.0%	
2C	coke oven coke	2.0%	5.0%	2.0%	5.0%	
2C	non-fuel combustion	2.0%	10.0%	2.0%	10.0%	
2C	Petroleum Coke	10.0%	7.5%	10.0%	7.5%	
2D	Lubricants	2.0%	50.0%	2.0%	50.0%	
2D	non-fuel combustion	25.0%	2.0%	25.0%	2.0%	
2D	Petroleum Coke	20.0%	30.0%	20.0%	30.0%	
2D	Petroleum Waxes	10.0%	50.0%	10.0%	50.0%	
3G	non-fuel combustion	0.0%	20.9%	0.0%	20.9%	

Category	Fuel	1990		2013		Notes
		Activity uncertainty (%)	Emission factor uncertainty (%)	Activity uncertainty (%)	Emission factor uncertainty (%)	
3H	non-fuel combustion	0.0%	50.0%	0.0%	50.0%	
4A	non-fuel combustion	1.0%	30.0%	1.0%	30.0%	
4B	non-fuel combustion	1.0%	45.0%	1.0%	45.0%	
4C	non-fuel combustion	1.0%	45.0%	1.0%	45.0%	
4D	non-fuel combustion	1.0%	50.0%	1.0%	50.0%	
4G	non-fuel combustion	1.0%	45.0%	1.0%	45.0%	
5C	Chemical waste	10.0%	30.0%	10.0%	30.0%	
5C	Clinical waste	5.0%	20.0%	5.0%	20.0%	
5C	Municipal Solid Waste	1.0%	15.0%	1.0%	15.0%	
5C	non-fuel combustion	300.0%	40.0%	300.0%	40.0%	

Notes

1. Uncertainties expressed as $0.5 \cdot R/E$ where R is the difference between 2.5 and 97.5 percentiles and E is the mean.

Not used = Fuel not used

(r) Revised in comparison to previous NIR
(Because of the large number of changes due to the implementation of the 2006 GLs, this notation was not used in the 2015 NIR)

Table A 2.3.2 Estimated uncertainties in the activity data and emission factors used in the methane (CH₄) inventory

Category	Fuel	1990		2013		Notes
		Activity uncertainty (%)	Emission factor uncertainty (%)	Activity uncertainty (%)	Emission factor uncertainty (%)	
1A1	All fuels	1.5%	50.0%	1.0%	50.0%	
1A2	All fuels	1.5%	50.0%	1.0%	50.0%	
1A3	Aviation Gasoline	20.0%	78.5%	20.0%	78.5%	
1A3	Jet Gasoline	20.0%	78.5%	20.0%	78.5%	
1A3	Other Bituminous Coal	20.0%	110.0%	20.0%	110.0%	
1A3	Other Gas/Diesel Oil	15.0%	130.0%	15.0%	130.0%	
1A3b	Gas/Diesel Oil	1.8%	130.0%	1.0%	130.0%	
1A3b	Liquefied Petroleum Gas	5.0%	130.0%	5.0%	130.0%	
1A3b	Motor Gasoline	1.0%	75.0%	1.0%	75.0%	
1A3d	Gas/Diesel Oil	20.0%	130.0%	20.0%	130.0%	
1A3d	Residual Fuel Oil	20.0%	130.0%	20.0%	130.0%	
1A4	All fuels	1.5%	50.0%	1.0%	50.0%	

Category	Fuel	1990		2013		Notes
		Activity uncertainty (%)	Emission factor uncertainty (%)	Activity uncertainty (%)	Emission factor uncertainty (%)	
1A5	All fuels	7.1%	65.6%	7.1%	65.6%	
1B1	Coke Oven Gas	1.5%	50.0%	1.0%	50.0%	
1B1	non-fuel combustion	2.0%	20.0%	2.0%	20.0%	
1B1	Wood		50.0%		50.0%	<i>All uncertainty ascribed to emission factor.</i>
1B2a	non-fuel combustion	5.0%	20.0%	5.0%	20.0%	
1B2b	non-fuel combustion		14.1%		14.1%	<i>Approach 2 based on estimates of uncertainty in components of gas leakage estimates given in Williams, 1993, Table 4.7. All uncertainty ascribed to emission factor.</i>
1B2c	non-fuel combustion	5.0%	20.0%	5.0%	20.0%	
2A4	All fuels		100.0%		100.0%	<i>All uncertainty ascribed to emission factor.</i>
2B	All fuels		20.0%		20.0%	<i>All uncertainty ascribed to emission factor.</i>
2C	Blast Furnace Gas	2.0%	50.0%	2.0%	50.0%	
2C	coke oven coke	2.0%	50.0%	2.0%	50.0%	
2C	non-fuel combustion	1.0%	50.0%	1.0%	50.0%	
2D	All fuels	50.0%	50.0%	50.0%	50.0%	
3A	non-fuel combustion	1.0%	20.0%	0.1%	20.0%	

Category	Fuel	1990		2013		Notes
		Activity uncertainty (%)	Emission factor uncertainty (%)	Activity uncertainty (%)	Emission factor uncertainty (%)	
3B	non-fuel combustion	1.0%	30.0%	0.1%	30.0%	
3F	non-fuel combustion	25.0%	50.0%	25.0%	50.0%	
4A	non-fuel combustion	1.0%	55.0%	1.0%	55.0%	
4B	non-fuel combustion	1.0%	55.0%	1.0%	55.0%	
4C	non-fuel combustion	1.0%	55.0%	1.0%	55.0%	
4E	non-fuel combustion	1.0%	20.0%	1.0%	20.0%	
5A	non-fuel combustion	15.0%	70.0%	15.0%	70.0%	<i>Approach 2 implemented as generalised log-logistic function</i>
5B	All fuels	30.0%	99.5%	30.0%	99.5%	
5C	Municipal Solid Waste	1.0%	75.0%	1.0%	75.0%	
5C	non-fuel combustion	5.0%	50.0%	5.0%	50.0%	
5D	non-fuel combustion	10.0%	25.0%	10.0%	25.0%	

Notes

1. Uncertainties expressed as $0.5 \cdot R/E$ where R is the difference between 2.5 and 97.5 percentiles and E is the mean.

Not used = Fuel not used

(r) Revised in comparison to previous NIR

(Because of the large number of changes due to the implementation of the 2006 GLs, this notation was not used in the 2015 NIR)

Table A 2.3.3 Estimated uncertainties in the activity data and emission factors used in the nitrous oxide (N₂O) inventory

Category	Fuel	1990		2013		Notes
		Activity uncertainty (%)	Emission factor uncertainty (%)	Activity uncertainty (%)	Emission factor uncertainty (%)	
1A1	All fuels	1.5%	100.0%	1.0%	100.0%	
1A2	All fuels	1.5%	100.0%	1.0%	100.0%	
1A3	Aviation Gasoline	20.0%	110.0%	20.0%	110.0%	
1A3	Jet Gasoline	20.0%	110.0%	20.0%	110.0%	
1A3	Other Bituminous Coal	20.0%	110.0%	20.0%	110.0%	
1A3	Other Gas/Diesel Oil	15.0%	130.0%	15.0%	130.0%	
1A3b	Gas/Diesel Oil	1.8%	130.0%	1.0%	130.0%	
1A3b	Liquefied Petroleum Gas	5.0%	130.0%	5.0%	130.0%	
1A3b	Motor Gasoline	1.0%	75.0%	1.0%	75.0%	
1A3d	Gas/Diesel Oil	20.0%	130.0%	20.0%	130.0%	
1A3d	Residual Fuel Oil	20.0%	130.0%	20.0%	130.0%	
1A4	All fuels	1.5%	100.0%	1.0%	100.0%	

Category	Fuel	1990		2013		Notes
		Activity uncertainty (%)	Emission factor uncertainty (%)	Activity uncertainty (%)	Emission factor uncertainty (%)	
1A5	All fuels	7.1%	85.1%	7.1%	85.1%	
1B1	All fuels	1.5%	118.0%	1.0%	118.0%	
1B2a	All fuels	5.0%	110.0%	5.0%	110.0%	
1B2b	All fuels	5.0%	110.0%	5.0%	110.0%	
1B2c	All fuels	5.0%	110.0%	5.0%	110.0%	
2B1	All fuels	2.0%	50.0%	2.0%	50.0%	
2B2	All fuels	10.0%	100.0%	10.0%	100.0%	
2B3	All fuels	2.0%	100.0%	2.0%	100.0%	
2B8	All fuels	10.0%	100.0%	10.0%	100.0%	
2C	All fuels	1.5%	118.0%	1.0%	118.0%	
2D	All fuels	50.0%	100.0%	50.0%	100.0%	
2G	All fuels	100.0%	100.0%	100.0%	100.0%	
3B	All fuels	1.0%	254.0%	1.0%	254.0%	
3D	All fuels	1.0%	259.0%	1.0%	259.0%	<i>Approach 2 implemented as generalised log-logistic function</i>
3F	All fuels	25.0%	230.0%	25.0%	230.0%	
4A	non-fuel combustion	1.0%	40.0%	1.0%	40.0%	
4B	non-fuel combustion	1.0%	55.0%	1.0%	55.0%	
4C	non-fuel combustion	1.0%	35.0%	1.0%	35.0%	

Category	Fuel	1990		2013		Notes
		Activity uncertainty (%)	Emission factor uncertainty (%)	Activity uncertainty (%)	Emission factor uncertainty (%)	
4D	non-fuel combustion	1.0%	100.0%	1.0%	100.0%	
4E	non-fuel combustion	1.0%	20.0%	1.0%	20.0%	
5B	All fuels	30.0%	90.0%	30.0%	90.0%	
5C	All fuels	7.0%	230.0%	7.0%	230.0%	
5D	All fuels	10.0%	248.0%	10.0%	248.0%	<i>Approach 2 implemented as generalised log-logistic function</i>

Notes

1. Uncertainties expressed as $0.5 \cdot R/E$ where R is the difference between 2.5 and 97.5 percentiles and E is the mean.

Not used = Fuel not used

(r) Revised in comparison to previous NIR

(Because of the large number of changes due to the implementation of the 2006 GLs, this notation was not used in the 2015 NIR)

Table A 2.3.4 Estimated uncertainties in the activity data and emission factors used in the F-gas inventory

Gas	Category	1990		2013		Notes
		Activity uncertainty (%)	Emission factor uncertainty (%)	Activity uncertainty (%)	Emission factor uncertainty (%)	
SF ₆	2C4	5.0%	5.0%	5.0%	5.0%	
SF ₆	2G1		5.0%		5.0%	<i>All uncertainty ascribed to emission factor.</i>
SF ₆	2G2a		17.5%		17.5%	<i>All uncertainty ascribed to emission factor.</i>
SF ₆	2G2b		40.0%		40.0%	<i>Approach 2 implemented as generalised log-logistic function. All uncertainty ascribed to emission factor.</i>
SF ₆	2G2e		47.1%		47.1%	<i>Approach 2 implemented as generalised log-logistic function. All uncertainty ascribed to emission factor.</i>
HFCs	2B9		10.0%		10.0%	<i>All uncertainty ascribed to emission factor.</i>
HFCs	2C4	5.0%	10.0%	5.0%	10.0%	
HFCs	2E1		47.1%		47.1%	<i>Approach 2 implemented as generalised log-logistic function. All uncertainty ascribed to emission factor.</i>
HFCs	2F1	10.0%	10.0%	10.0%	10.0%	
HFCs	2F2		25.0%		25.0%	<i>All uncertainty ascribed to emission factor.</i>
HFCs	2F3		25.0%		25.0%	<i>All uncertainty ascribed to emission factor.</i>
HFCs	2F4a	5.0%	10.0%	5.0%	10.0%	

Gas	Category	1990		2013		Notes
		Activity uncertainty (%)	Emission factor uncertainty (%)	Activity uncertainty (%)	Emission factor uncertainty (%)	
HFCs	2F4b		10.0%		10.0%	<i>All uncertainty ascribed to emission factor.</i>
HFCs	2F5		25.5%		25.5%	<i>Approach 2 implemented as generalised log-logistic function. All uncertainty ascribed to emission factor.</i>
HFCs	2F6		53.0%		53.0%	<i>All uncertainty ascribed to emission factor.</i>
NF ₃	2E1		47.1%		47.1%	<i>Approach 2 implemented as generalised log-logistic function. All uncertainty ascribed to emission factor.</i>
PFCs	2B9		10.0%		10.0%	<i>All uncertainty ascribed to emission factor.</i>
PFCs	2C3		20.0%		20.0%	<i>All uncertainty ascribed to emission factor.</i>
PFCs	2F1	10.0%	10.0%	10.0%	10.0%	
PFCs	2F3		25.0%		25.0%	<i>All uncertainty ascribed to emission factor.</i>
PFCs	2G2e		47.1%		47.1%	<i>Approach 2 implemented as generalised log-logistic function. All uncertainty ascribed to emission factor.</i>

A 2.3.1.3 Uncertainty in the Emissions

The overall uncertainty and central estimate from the Monte Carlo model for 2013 is given in **Section A 2.4.1**.

A 2.3.1.4 Uncertainty in the Trend

The uncertainty in the trend between 1990 and 2013 is given in **Section A 2.4.2**. In running this simulation it was necessary to make assumptions about the degree of correlation between sources in 1990 and 2013. If source emission factors are correlated this will have the effect of reducing the trend uncertainty. The assumptions were as follows:

- Activity data are uncorrelated;
- Emission factors of some similar fuels are correlated;
- Land Use Change and forestry emissions are correlated (e.g. 1990 5A CO₂ with 2013 5A CO₂);
- Offshore emissions are not correlated since they are based on separate studies using emission factors appropriate for the time;
- Emission factors covered by the Carbon Factors Review (Baggott et al, 2004) are not correlated; and
- Process emissions from blast furnaces, coke ovens and ammonia plant were not correlated.

A 2.3.2 Methane Emission Uncertainties**A 2.3.2.1 General Considerations**

The uncertainty in the trend between 1990 and 2013 is given in **Section A 2.4.2**. In running this simulation it was necessary to make assumptions about the degree of correlation between sources in 1990 and 2013. If source emission factors are correlated this will have the effect of reducing the trend uncertainty.

A 2.3.2.2 Uncertainty in the Emissions

The overall uncertainty and central estimate from the Monte Carlo model for 2013 is given in **Section A 2.4.1**.

A 2.3.2.3 Uncertainty in the Trend

The uncertainty in the trend between 1990 and 2013 is given in **Section A 2.4.2**. In running this simulation it was necessary to make assumptions about the degree of correlation between sources in 1990 and 2013. If source emission factors are correlated this will have the effect of reducing uncertainty in the emissions trend. The assumptions were:

Activity data are uncorrelated between years, but activity data for major fuels were correlated in the same year in a similar manner to that described above for carbon;

- Landfill emissions were partly correlated across years in the simulation. It is likely that the emission factors used in the model will be correlated, and also the historical estimates of waste arisings will be correlated since they are estimated by extrapolation from the year of the study. However, the reduction in emissions is due to flaring and

utilisation systems installed since 1990 and this is unlikely to be correlated. As a simple estimate it was assumed that the degree of correlation should reflect the reduction. Emissions have reduced by 57% hence the degree of correlation was 43%;

- Offshore emissions are not correlated across years since they are based on separate studies using emission factors that reflected the processes in use at the time;
- Gas leakage emissions were partially correlated across years. As a simple estimate it was assumed that the degree of correlation should reflect the reduction in emissions. Emissions have reduced by 53% hence the degree of correlation was 47%; and
- Emissions from deep mines were not correlated across years as they were based on different studies, and a different selection of mines. Open cast and coal storage and transport were correlated since they are based on default emission factors.

A 2.3.3 Nitrous Oxide Emission Uncertainties

A 2.3.3.1 General Considerations

The analysis of the uncertainties in the nitrous oxide emissions is particularly difficult because emissions sources are diverse, and few data are available to form an assessment of the uncertainties in each source. Emission factor data for the combustion sources are scarce and for some fuels are not available. The parameter uncertainties are shown in **Table A 2.3.3**. The uncertainty assumed for agricultural soils (IPCC category 4D) uses a log-logistic distribution defined such that the 97.5 percentile is greater than the 2.5 percentile by a factor of around 4.2-4.3. These parameterised functions have been defined and provided by Rothamsted Research (from research based on Milne *et al.*, 2014) as the best possible fit to the expected distribution of uncertainties in 1990 and 2010 emissions, and are normalised in the Approach 2 methodology such that the resultant mean is consistent with the current inventory emissions in 1990 and the most recent year.

A 2.3.3.2 Uncertainty in the Emissions

The overall uncertainty and central estimate from the Monte Carlo model for 2013 is given in **Section A 2.4.1**.

A 2.3.3.3 Uncertainty in the Trend

The uncertainty in the trend between 1990 and 2013 is given in **Section A 2.4.2**. In running this simulation it was necessary to make assumptions about the degree of correlation between sources across years. If sources are correlated this will have the effect of reducing the emissions. The assumptions were as follows:

- Activity data are uncorrelated between years, but similar fuels are correlated in the same year;
- Emissions from agricultural soils are correlated in the base and inventory year;
- The emission factor used for sewage treatment was assumed to be correlated between years, though the protein consumption data used as activity data were assumed not to be correlated;
- Nitric acid production emission factors were assumed not to be correlated, since the mix of operating plant is very different in 2013 compared with 1990 – only 2 of the

original 8 units are still operating in the latest inventory year, all of which now have differing levels of abatement fitted.

A 2.3.4 Halocarbons, NF₃ and SF₆

A 2.3.4.1 Uncertainty Parameters

The uncertainties in the emissions of HFCs, PFCs, NF₃ and SF₆ are based on the recent study to update emissions and projections of F-gases (Mathis, P. et al. 2014), for all sources. For these sources, the uncertainty parameters were taken from this report and information from the HFC/HCFC plant operator.

A 2.3.4.2 Uncertainty in the Emissions

The uncertainty in the trend between 1990 and 2013 is given in **Section A 2.4.2**.

A 2.3.4.3 Uncertainty in the Trend

The uncertainty in the trend between 1990 and 2013 is given in **Section A 2.4.2**.

A 2.4 UNCERTAINTIES IN GWP WEIGHTED EMISSIONS

A 2.4.1 Uncertainty in the emissions

The uncertainty in the combined GWP weighted emission is given in **Table A 2.4.1**, along with uncertainties for each of the 7 categorised GHGs. This is calculated as half of the 95% confidence range, i.e. the limits between which there is a 95% probability that the actual value of emissions falls.

A 2.4.2 Uncertainty in the Trend

The uncertainty estimates for all gases are summarised in **Table A 2.4.1** under 'Range of likely % change'. This indicates the range between which there is a 95% probability that the actual trend in inventory emissions falls.

The source which makes the major contribution to the overall uncertainty is 3D Agricultural Soils.

Table A 2.4.1 Summary of Monte Carlo Uncertainty Estimates

IPCC Source Category	Gas	1990 Emissions	2013 Emissions	Range of uncertainty in 1990 emissions		Uncertainty introduced on national total in 1990	Range of uncertainty in 2013 emissions		Uncertainty introduced on national total in 2013	% change in emissions between 2013 and 1990	Range of likely % change between 2013 and 1990	
				2.5 percentile	97.5 percentile		2.5 percentile	97.5 percentile			2.5 percentile	97.5 percentile
		Gg CO ₂ e	Gg CO ₂ e	Gg CO ₂ e	Gg CO ₂ e	%	Gg CO ₂ e	Gg CO ₂ e	%	%	%	%
Total	CO ₂ (net)	599,340	469,206	586,146	612,614	2%	459,154	479,338	2%	-22%	-24%	-19%
	CH ₄	137,231	56,385	109,014	176,641	25%	46,867	68,078	19%	-58%	-70%	-45%
	N ₂ O	57,151	27,648	39,062	89,116	44%	17,512	49,742	58%	-49%	-75%	-3%
	HFC	14,539	16,262	12,386	16,703	15%	14,858	17,657	9%	13%	-5%	34%
	PFC	1,652	253	1,571	1,734	5%	197	320	24%	-85%	-88%	-81%
	SF ₆	1,279	602	1,147	1,413	10%	539	664	10%	-53%	-59%	-45%
	NF ₃	0.4	0.4	0.2	0.6	45%	0.2	0.5	46%	-13%	-55%	76%
	All	811,192	570,356	772,997	862,066	5%	551,887	596,105	4%	-30%	-34%	-25%

Uncertainty calculated as 0.5*R/E where R is the difference between 2.5 and 97.5 percentiles and E is the mean calculated in the simulation.

Emissions of CO₂ are net emissions (i.e. sum of emissions and removals).

Important - Emissions in this table are taken from the Monte Carlo model output. The central estimates, according to gas, for 1990 and the latest inventory year are very similar but not identical to the emission estimates in the inventory. The Executive Summary of this NIR and the accompanying CRF tables present the agreed national GHG emissions and removals reported to the UNFCCC.

A 2.5 SECTORAL UNCERTAINTIES

A 2.5.1 Overview of the Method

Sectoral uncertainties were calculated from the same base data used for the “by gas” analysis. The emissions and uncertainties per sector are presented in **Table A 2.5.1**. We recommend that the estimates in the table are taken only as indicative, since the uncertainties for fuels are based on the overall statistical difference for the fuel total and it is assumed in this analysis that additional proportional uncertainty due to disaggregation is distributed evenly at sector level. The estimates are presented in IPCC categories, which is consistent with the reporting format used within this submission to the UNFCCC.

A 2.5.2 Review of Changes made to the Monte Carlo Model since the last NIR

No changes that are specific to the sectoral uncertainty analysis have been made. The changes made to the uncertainty parameters used to estimate the uncertainties by gas are all reflected within the sectoral analysis.

Table A 2.5.1 Sectoral Uncertainty Estimates

IPCC Source Category	Gas	1990 Emissions	2013 Emissions	Uncertainty in 2013 emissions as % of emissions in category		Uncertainty Introduced on national total in 2013	% change in emissions between 1990 and 2013	Range of likely % change between 1990 and 2013	
				2.5 percentile	97.5 percentile			2.5 percentile	97.5 percentile
				1A1a	GWP weighted total			205,895	149,087
1A1b	GWP weighted total	17,879	14,723	12,690	16,930	14.4%	-18%	-37%	11%
1A1c	GWP weighted total	14,136	15,245	14,892	15,621	2.4%	8%	2%	14%
1A2a	GWP weighted total	21,592	14,674	13,382	15,953	8.8%	-32%	-40%	-24%
1A2b	GWP weighted total	4,811	935	844	1,031	10.0%	-81%	-84%	-75%
1A2c	GWP weighted total	12,230	5,242	4,918	5,577	6.3%	-57%	-61%	-53%
1A2d	GWP weighted total	4,662	1,979	1,870	2,090	5.6%	-58%	-62%	-52%
1A2e	GWP weighted total	7,693	4,507	4,281	4,734	5.0%	-41%	-46%	-36%
1A2f	GWP weighted total	7,171	2,361	2,108	2,626	11.0%	-67%	-74%	-58%
1A2g	GWP weighted total	39,490	27,560	26,108	29,059	5.4%	-30%	-36%	-23%
1A3a	GWP weighted total	1,882	1,981	1,591	2,368	19.6%	5%	-20%	40%
1A3b	GWP weighted total	110,472	108,017	106,220	109,841	1.7%	-2%	-5%	0%
1A3c	GWP weighted total	1,462	1,964	1,612	2,309	17.7%	34%	4%	73%
1A3d	GWP weighted total	2,188	2,189	1,810	2,574	17.4%	0%	-21%	25%
1A3e	GWP weighted total	253	516	422	619	19.1%	104%	56%	166%
1A4a	GWP weighted total	25,658	24,321	23,512	25,144	3.4%	-5%	-11%	1%
1A4b	GWP weighted total	80,559	75,390	72,332	78,479	4.1%	-6%	-12%	-1%
1A4c	GWP weighted total	5,801	4,519	3,148	5,907	30.5%	-22%	-53%	41%
1A5b	GWP weighted total	5,336	2,306	2,132	2,485	7.7%	-57%	-61%	-52%

IPCC Source Category	Gas	1990 Emissions	2013 Emissions	Uncertainty in 2013 emissions as % of emissions in category		Uncertainty Introduced on national total in 2013	% change in emissions between 1990 and 2013	Range of likely % change between 1990 and 2013	
				2.5 percentile	97.5 percentile			2.5 percentile	97.5 percentile
				1B1	GWP weighted total			23,473	1,862
1B2	GWP weighted total	18,150	9,520	6,464	12,560	32.0%	-48%	-63%	-32%
2A1	GWP weighted total	7,295	4,029	3,901	4,158	3.2%	-45%	-47%	-42%
2A2	GWP weighted total	1,462	1,239	1,176	1,301	5.0%	-15%	-25%	-4%
2A3	GWP weighted total	408	389	370	409	5.0%	-5%	-11%	2%
2A4	GWP weighted total	678	776	749	804	3.6%	15%	8%	21%
2B1	GWP weighted total	2,191	1,450	1,416	1,485	2.4%	-34%	-36%	-31%
2B2	GWP weighted total	3,870	43	20	81	71.3%	-99%	-100%	-97%
2B3	GWP weighted total	19,947	0	0	0	#DIV/0!	-100%	-100%	-100%
2B6	GWP weighted total	105	123	111	136	10.0%	18%	-5%	50%
2B7	GWP weighted total	232	285	270	301	5.5%	23%	11%	36%
2B8	GWP weighted total	4,063	2,988	2,097	3,882	29.9%	-26%	-52%	10%
2B9	GWP weighted total	14,391	158	146	170	7.5%	-99%	-99%	-99%
2C	GWP weighted total	9,432	5,234	4,976	5,499	5.0%	-45%	-48%	-41%
2D	GWP weighted total	1,235	1,034	789	1,348	27.0%	-16%	-47%	33%
2E	GWP weighted total	5	15	9	22	46.1%	213%	65%	494%
2F	GWP weighted total	161	16,199	14,795	17,597	8.6%	9953%	6484%	20419%
2G	GWP weighted total	974	591	508	677	14.3%	-39%	-50%	-26%
3A	GWP weighted total	28,238	23,608	18,886	28,335	20.0%	-16%	-37%	11%
3B	GWP weighted total	6,883	5,358	3,713	7,807	38.2%	-22%	-53%	31%
3D	GWP weighted total	23,102	19,458	10,009	41,271	80.3%	-16%	-70%	132%

IPCC Source Category	Gas	1990 Emissions	2013 Emissions	Uncertainty in 2013 emissions as % of emissions in category		Uncertainty Introduced on national total in 2013	% change in emissions between 1990 and 2013	Range of likely % change between 1990 and 2013	
				2.5 percentile	97.5 percentile			2.5 percentile	97.5 percentile
				3F	GWP weighted total			269	0
3G	GWP weighted total	1,580	774	615	937	20.8%	-51%	-64%	-34%
3H	GWP weighted total	385	245	160	360	40.8%	-36%	-64%	12%
4A	GWP weighted total	-15,978	-17,265	-22,494	-12,168	-29.9%	8%	-36%	53%
4B	GWP weighted total	15,768	12,518	7,122	17,929	43.2%	-21%	-76%	35%
4C	GWP weighted total	-3,653	-5,875	-8,517	-3,211	-45.2%	61%	-25%	146%
4D	GWP weighted total	485	301	198	444	40.8%	-38%	-90%	7%
4E	GWP weighted total	7,324	6,199	4,171	8,905	38.2%	-15%	-68%	34%
4F	GWP weighted total	0	0	0	0	40.6%	#DIV/0!	n/a	n/a
4G	GWP weighted total	41	-1,122	-1,628	-615	-45.1%	-2854%	-4097%	-1611%
5A	GWP weighted total	62,757	16,642	9,495	27,183	53.1%	-73%	-87%	-44%
5B	GWP weighted total	10	1,328	815	2,081	47.7%	12679%	6304%	26238%
5C	GWP weighted total	1,472	320	-264	925	185.5%	-78%	-296%	148%
5D	GWP weighted total	5,273	4,413	2,857	8,596	65.0%	-16%	-60%	82%
Grand Total	GWP weighted total	811,193	570,357	551,887	596,106	3.9%	-30%	-34%	-25%

Note: Although the range of likely trend for some sectors are large (e.g. ~14,000% for 2F), these are comparable to the trend from 1990 to 2013 (~-10,000%). Such instances arise where 1990 emissions are orders of magnitude lower than 2013 emissions.

Important - Emissions in this table are taken from the Monte Carlo model output. The central estimates, according to gas, for 1990 and the latest inventory year are very similar but not identical to the emission estimates in the inventory. The Executive Summary of this NIR and the accompanying CRF tables present the agreed national GHG emissions and removals reported to the UNFCCC.

A 2.6 COMPARISON OF UNCERTAINTIES FROM THE ERROR PROPAGATION AND MONTE CARLO ANALYSES

Comparing the results of the error propagation approach, and the Monte Carlo estimation of uncertainty by simulation, is a useful quality control check on the behaviour of the Monte Carlo model.

The reason that the error propagation approach is used as a reference is because the mathematical approach to the error propagation approach has been defined and checked by the IPCC, and is clearly set out in the IPCC 2000 Good Practice Guidance and the 2006 Guidelines. The UK has implemented the IPCC error propagation approach as set out in this guidance. The implementation of an uncertainty estimation by simulation cannot be prescriptive, and will depend on the Monte Carlo software a country chooses to use, how the country constructs its model, and the correlations included within that model. Therefore, there is a greater likelihood of errors being introduced in the model used to estimate uncertainty by simulation.

If all the distributions in the Monte Carlo model were normal, and the assumed correlations were identical, the estimated errors on the trend from the Monte Carlo model should be identical to those estimated by the error propagation approach. The Error Propagation approach assumes 100% correlation between EFs in the base and inventory year, and no correlation between sources, however in reality the nature and degree of correlation varies by source, and many distributions are not normal but heavily skewed, particularly those with very high uncertainty. These differences interact in various ways, but would be expected broadly to result in higher trend uncertainty, and lower uncertainty on the 2013 total.

Table A 2.6.1 shows differences in the trend uncertainty between the error propagation and Monte Carlo approaches. These differences mostly arise from the fact that the error propagation approach only uses normal distributions, cannot account for different uncertainty parameters between the 1990 and the latest inventory year, cannot account for correlations within years, and automatically assumes a correlation between the emission factor uncertainty in 1990 and 2013.

The central estimates of emissions generated by the Monte Carlo model in 1990, and those in the latest inventory year, are very close. Mathematically we would not expect the central estimates from the two methods to be identical.

Table A 2.6.1 Comparison of the central estimates and trends in emissions from the error propagation (Approach 1) and Monte Carlo (Approach 2) uncertainty analyses

Method of uncertainty estimation	Central estimate (Gg CO ₂ equivalent) ^b		Uncertainty on trend, 95% CI (1990 to 2013)
	1990	2013	
Error propagation	815,144	570,452	2.7
Monte Carlo	811,222	570,565	3.3 ^a

Notes

CI Confidence Interval

- ^a Calculated as half the difference between 2.5 and 97.5 percentiles , assuming a normal distribution is equal to ± 1.96 standard deviations on the central estimate.
- ^b Net emissions, including emissions and removals from LULUCF

ANNEX 3: Other Detailed Methodological Descriptions for Individual Source or Sink Categories, Including for KP-LULUCF Activities.

This Annex contains background information about methods used to estimate emissions in the UK GHG inventory. This information has not been incorporated in the main body of the report because of the level of detail, and because the methods used to estimate emissions cut across sectors.

This Annex provides background information on the fuels used in the UK GHG inventory, mapping between IPCC and NAEI source categories and detailed description of methods used to estimate GHG emissions, and emission factors used in those methods – presented in **Section A 3.3** onwards.

A 3.1 ENERGY

Methods for calculating emissions within the energy sector are detailed in the method statements set out in **Chapter 3**. This Annex details the emission factors used and their source, and elaborates on references commonly used within the Energy sector. The national energy balance (and how it is used) is described in **Annex 4**.

A 3.1.1 Emission factors

Table A 3.1.1 Emission factors used for the 2015 submission (sector 1A)

Category	Source Name	Activity Name	Fuel (CRF)	Gas	Unit	1990	2013	Max	Min	EF Method Reference
1A1ai	Miscellaneous industrial/commercial combustion	Landfill gas	Biomass	CH ₄	kg/TJ	5.1	5.1	5.1	5.1	IPCC 2006 Guidelines
1A1ai	Miscellaneous industrial/commercial combustion	Landfill gas	Biomass	N ₂ O	kg/TJ	0.1	0.1	0.1	0.1	IPCC 2006 Guidelines
1A1ai	Miscellaneous industrial/commercial combustion	MSW	Other	CO ₂	t/TJ	33.7	37.4	41.0	22.8	Biodegradability of municipal solid waste (WR1003) - resource futures, 2012
1A1ai	Miscellaneous industrial/commercial combustion	MSW	Other	CH ₄	kg/TJ	222.7	222.7	222.7	220.4	IPCC 2006 Guidelines
1A1ai	Miscellaneous industrial/commercial combustion	MSW	Other	N ₂ O	kg/TJ	3.0	3.0	3.0	2.9	IPCC 2006 Guidelines
1A1ai	Power stations	Burning oil	Liquid	CO ₂	t/TJ	71.8	71.7	71.8	71.7	Carbon Factors Review 2004
1A1ai	Power stations	Burning oil	Liquid	CH ₄	kg/TJ	3.0	3.0	3.0	3.0	IPCC 2006 Guidelines
1A1ai	Power stations	Burning oil	Liquid	N ₂ O	kg/TJ	0.6	0.6	0.6	0.6	IPCC 2006 Guidelines
1A1ai	Power stations	Coal	Solid	CO ₂	t/TJ	94.2	89.8	94.6	89.8	EU ETS data from EA, SEPA, NIEA (2014)
1A1ai	Power stations	Coal	Solid	CH ₄	kg/TJ	0.8	0.8	0.8	0.8	Brain et al, 1994
1A1ai	Power stations	Coal	Solid	N ₂ O	kg/TJ	2.7	2.5	2.7	2.5	Based on Fynes & Sage, 1994

Category	Source Name	Activity Name	Fuel (CRF)	Gas	Unit	1990	2013	Max	Min	EF Method Reference
1A1ai	Power stations	Coke	Solid	CO ₂	t/TJ	114.3	111.3	114.3	101.8	Ricardo-AEA estimate based on carbon balance for integrated steelwork processes (2014)
1A1ai	Power stations	Coke	Solid	CH ₄	kg/TJ	0.7	0.7	0.7	0.7	Brain et al, 1994
1A1ai	Power stations	Coke	Solid	N ₂ O	kg/TJ	2.7	2.5	2.7	2.5	Based on Fynes & Sage, 1994
1A1ai	Power stations	Fuel oil	Liquid	CO ₂	t/TJ	77.5	77.7	78.4	76.3	EU ETS data from EA, SEPA, NIEA (2014)
1A1ai	Power stations	Fuel oil	Liquid	CH ₄	kg/TJ	3.0	3.0	3.0	3.0	IPCC 2006 Guidelines
1A1ai	Power stations	Fuel oil	Liquid	N ₂ O	kg/TJ	0.6	0.6	0.6	0.6	IPCC 2006 Guidelines
1A1ai	Power stations	Gas oil	Liquid	CO ₂	t/TJ	73.9	74.1	74.2	73.5	Carbon Factors Review 2004
1A1ai	Power stations	Gas oil	Liquid	CH ₄	kg/TJ	3.0	3.0	3.0	3.0	IPCC 2006 Guidelines
1A1ai	Power stations	Gas oil	Liquid	N ₂ O	kg/TJ	0.6	0.6	0.6	0.6	IPCC 2006 Guidelines
1A1ai	Power stations	Landfill gas	Biomass	CH ₄	kg/TJ	1.0	1.0	1.0	1.0	IPCC 2006 Guidelines
1A1ai	Power stations	Landfill gas	Biomass	N ₂ O	kg/TJ	0.1	0.1	0.1	0.1	IPCC 2006 Guidelines
1A1ai	Power stations	Liquid bio-fuels	Biomass	CH ₄	kg/TJ	0.0	30.0	30.0	0.0	IPCC 2006 Guidelines
1A1ai	Power stations	Liquid bio-fuels	Biomass	N ₂ O	kg/TJ	0.0	4.0	4.0	0.0	IPCC 2006 Guidelines
1A1ai	Power stations	LPG	Liquid	CO ₂	t/TJ	64.7	64.7	64.7	64.7	Carbon Factors Review 2004
1A1ai	Power stations	LPG	Liquid	CH ₄	kg/TJ	1.0	1.0	1.0	1.0	IPCC 2006 Guidelines
1A1ai	Power stations	LPG	Liquid	N ₂ O	kg/TJ	0.1	0.1	0.1	0.1	IPCC 2006 Guidelines
1A1ai	Power stations	MSW	Other	CO ₂	t/TJ	33.7	37.4	41.0	22.8	Biodegradability of municipal solid waste (WR1003) - resource futures, 2012
1A1ai	Power stations	MSW	Other	CH ₄	kg/TJ	22.3	22.3	22.3	22.0	IPCC 2006 Guidelines

Other Detailed Methodological Descriptions **A3**

Category	Source Name	Activity Name	Fuel (CRF)	Gas	Unit	1990	2013	Max	Min	EF Method Reference
1A1ai	Power stations	MSW	Other	N ₂ O	kg/TJ	3.0	3.0	3.0	2.9	IPCC 2006 Guidelines
1A1ai	Power stations	Natural gas	Gaseous	CO ₂	t/TJ	56.0	56.5	57.2	56.0	EU ETS data from EA, SEPA, NIEA (2014)
1A1ai	Power stations	Natural gas	Gaseous	CH ₄	kg/TJ	1.0	1.0	1.0	1.0	IPCC 2006 Guidelines
1A1ai	Power stations	Natural gas	Gaseous	N ₂ O	kg/TJ	0.1	0.1	0.1	0.1	IPCC 2006 Guidelines
1A1ai	Power stations	OPG	Liquid	CO ₂	t/TJ	56.3	56.2	59.8	54.9	EU ETS data from EA, SEPA, NIEA (2014)
1A1ai	Power stations	OPG	Liquid	CH ₄	kg/TJ	1.0	1.0	1.0	1.0	IPCC 2006 Guidelines
1A1ai	Power stations	OPG	Liquid	N ₂ O	kg/TJ	0.1	0.1	0.1	0.1	IPCC 2006 Guidelines
1A1ai	Power stations	Orimulsion	Liquid	CO ₂	t/TJ	77.6	77.6	77.6	77.6	Manufacturer's specification, BITOR (1995)
1A1ai	Power stations	Orimulsion	Liquid	CH ₄	kg/TJ	4.3	4.3	4.3	4.3	Use 2006 IPCC default for fuel oil
1A1ai	Power stations	Orimulsion	Liquid	N ₂ O	kg/TJ	0.9	0.9	0.9	0.9	Use 2006 IPCC default for fuel oil
1A1ai	Power stations	Petroleum coke	Liquid	CO ₂	t/TJ	82.7	111.2	111.2	82.7	EU ETS data from EA, SEPA, NIEA (2014)
1A1ai	Power stations	Petroleum coke	Liquid	CH ₄	kg/TJ	3.2	3.8	3.8	3.2	IPCC 2006 Guidelines
1A1ai	Power stations	Petroleum coke	Liquid	N ₂ O	kg/TJ	2.0	2.6	2.6	2.0	Based on Fynes & Sage, 1994
1A1ai	Power stations	Poultry litter	Biomass	CH ₄	kg/TJ	0.0	35.6	37.4	0.0	IPCC 2006 Guidelines
1A1ai	Power stations	Poultry litter	Biomass	N ₂ O	kg/TJ	0.0	4.8	5.0	0.0	IPCC 2006 Guidelines
1A1ai	Power stations	Scrap tyres	Other	CO ₂	t/TJ	C	C	C	C	BCA/Lafarge Cement (2006,2007,2008, 2009, 2010,2011,2012) Mineral Products Association (2013, 2014)
1A1ai	Power stations	Scrap tyres	Other	CH ₄	kg/TJ	31.0	30.0	31.0	30.0	IPCC 2006 Guidelines
1A1ai	Power stations	Scrap tyres	Other	N ₂ O	kg/TJ	4.1	4.0	4.1	4.0	IPCC 2006 Guidelines
1A1ai	Power stations	Sewage gas	Biomass	CH ₄	kg/TJ	1.0	1.0	1.0	1.0	IPCC 2006 Guidelines

Category	Source Name	Activity Name	Fuel (CRF)	Gas	Unit	1990	2013	Max	Min	EF Method Reference
1A1ai	Power stations	Sewage gas	Biomass	N ₂ O	kg/TJ	0.1	0.1	0.1	0.1	IPCC 2006 Guidelines
1A1ai	Power stations	Sour gas	Gaseous	CO ₂	t/TJ	74.0	74.6	74.6	70.4	EU ETS data from EA, SEPA, NIEA (2014)
1A1ai	Power stations	Sour gas	Gaseous	CH ₄	kg/TJ	1.0	1.0	1.0	1.0	IPCC 2006 Guidelines
1A1ai	Power stations	Sour gas	Gaseous	N ₂ O	kg/TJ	0.1	0.1	0.1	0.1	IPCC 2006 Guidelines
1A1ai	Power stations	Straw	Biomass	CH ₄	kg/TJ	0.0	30.0	30.0	0.0	IPCC 2006 Guidelines
1A1ai	Power stations	Straw	Biomass	N ₂ O	kg/TJ	0.0	4.0	4.0	0.0	IPCC 2006 Guidelines
1A1ai	Power stations	Waste oils	Liquid	CO ₂	t/TJ	73.5	73.7	73.8	73.0	Based on analysis of eight samples of waste oil - see Passant, 2004
1A1ai	Power stations	Waste oils	Liquid	CH ₄	kg/TJ	3.0	3.0	3.0	3.0	IPCC 2006 Guidelines
1A1ai	Power stations	Waste oils	Liquid	N ₂ O	kg/TJ	3.8	3.8	3.8	3.8	IPCC 2006 Guidelines
1A1ai	Power stations	Wood	Biomass	CH ₄	kg/TJ	0.0	21.4	21.4	0.0	IPCC 2006 Guidelines
1A1ai	Power stations	Wood	Biomass	N ₂ O	kg/TJ	0.0	2.9	2.9	0.0	IPCC 2006 Guidelines
1A1ai	Public sector combustion	Sewage gas	Biomass	CH ₄	kg/TJ	5.1	5.1	5.1	5.1	IPCC 2006 Guidelines
1A1ai	Public sector combustion	Sewage gas	Biomass	N ₂ O	kg/TJ	0.1	0.1	0.1	0.1	IPCC 2006 Guidelines
1A1b	Refineries - combustion	Burning oil	Liquid	CO ₂	t/TJ	71.8	71.7	71.8	71.7	Carbon Factors Review 2004
1A1b	Refineries - combustion	Burning oil	Liquid	CH ₄	kg/TJ	3.0	3.0	3.0	3.0	IPCC 2006 Guidelines
1A1b	Refineries - combustion	Burning oil	Liquid	N ₂ O	kg/TJ	0.6	0.6	0.6	0.6	IPCC 2006 Guidelines
1A1b	Refineries - combustion	Fuel oil	Liquid	CO ₂	t/TJ	77.5	77.9	79.1	77.2	EU ETS data from EA, SEPA, NIEA (2014)
1A1b	Refineries - combustion	Fuel oil	Liquid	CH ₄	kg/TJ	3.0	3.0	3.0	3.0	IPCC 2006 Guidelines
1A1b	Refineries - combustion	Fuel oil	Liquid	N ₂ O	kg/TJ	0.6	0.6	0.6	0.6	IPCC 2006 Guidelines

Category	Source Name	Activity Name	Fuel (CRF)	Gas	Unit	1990	2013	Max	Min	EF Method Reference
1A1b	Refineries - combustion	Gas oil	Liquid	CO ₂	t/TJ	73.9	74.1	74.2	73.5	Carbon Factors Review 2004
1A1b	Refineries - combustion	Gas oil	Liquid	CH ₄	kg/TJ	3.0	3.0	3.0	3.0	IPCC 2006 Guidelines
1A1b	Refineries - combustion	Gas oil	Liquid	N ₂ O	kg/TJ	0.6	0.6	0.6	0.6	IPCC 2006 Guidelines
1A1b	Refineries - combustion	LPG	Liquid	CO ₂	t/TJ	64.7	64.7	64.7	64.7	Carbon Factors Review 2004
1A1b	Refineries - combustion	LPG	Liquid	CH ₄	kg/TJ	1.0	1.0	1.0	1.0	IPCC 2006 Guidelines
1A1b	Refineries - combustion	LPG	Liquid	N ₂ O	kg/TJ	0.1	0.1	0.1	0.1	IPCC 2006 Guidelines
1A1b	Refineries - combustion	Naphtha	Liquid	CO ₂	t/TJ	68.8	69.0	69.8	68.8	Carbon Factors Review 2004
1A1b	Refineries - combustion	Naphtha	Liquid	CH ₄	kg/TJ	3.0	3.0	3.0	3.0	IPCC 2006 Guidelines
1A1b	Refineries - combustion	Naphtha	Liquid	N ₂ O	kg/TJ	0.6	0.6	0.6	0.6	IPCC 2006 Guidelines
1A1b	Refineries - combustion	Natural gas	Gaseous	CO ₂	t/TJ	56.0	56.4	57.2	55.7	EU ETS data from EA, SEPA, NIEA (2014)
1A1b	Refineries - combustion	Natural gas	Gaseous	CH ₄	kg/TJ	1.0	1.0	1.0	1.0	IPCC 2006 Guidelines
1A1b	Refineries - combustion	Natural gas	Gaseous	N ₂ O	kg/TJ	0.1	0.1	0.1	0.1	IPCC 2006 Guidelines
1A1b	Refineries - combustion	OPG	Liquid	CO ₂	t/TJ	56.3	56.2	59.8	54.9	EU ETS data from EA, SEPA, NIEA (2014)
1A1b	Refineries - combustion	OPG	Liquid	CH ₄	kg/TJ	1.0	1.0	1.0	1.0	IPCC 2006 Guidelines
1A1b	Refineries - combustion	OPG	Liquid	N ₂ O	kg/TJ	0.1	0.1	0.1	0.1	IPCC 2006 Guidelines
1A1b	Refineries - combustion	Petrol	Liquid	CO ₂	t/TJ	70.2	70.0	70.2	70.0	Carbon Factors Review 2004
1A1b	Refineries - combustion	Petrol	Liquid	CH ₄	kg/TJ	3.0	3.0	3.0	3.0	IPCC 2006 Guidelines
1A1b	Refineries - combustion	Petrol	Liquid	N ₂ O	kg/TJ	0.6	0.6	0.6	0.6	IPCC 2006 Guidelines
1A1b	Refineries - combustion	Petroleum coke	Liquid	CO ₂	t/TJ	90.9	100.4	100.4	90.9	Carbon Factors Review 2004

Category	Source Name	Activity Name	Fuel (CRF)	Gas	Unit	1990	2013	Max	Min	EF Method Reference
1A1b	Refineries - combustion	Petroleum coke	Liquid	CH ₄	kg/TJ	3.2	3.2	3.5	3.2	IPCC 2006 Guidelines
1A1b	Refineries - combustion	Petroleum coke	Liquid	N ₂ O	kg/TJ	2.0	2.2	2.2	2.0	Based on Fynes & Sage, 1994
1A1b	Refineries - combustion	Refinery miscellaneous	Liquid	CO ₂	t/TJ	71.5	71.6	72.3	70.5	UKPIA (1989)
1A1b	Refineries - combustion	Refinery miscellaneous	Liquid	CH ₄	kg/TJ	3.3	3.3	3.3	3.3	IPCC 2006 Guidelines
1A1b	Refineries - combustion	Refinery miscellaneous	Liquid	N ₂ O	kg/TJ	0.7	0.7	0.7	0.7	IPCC 2006 Guidelines
1A1ci	Coke production	Blast furnace gas	Solid	CO ₂	t/TJ	280.5	278.3	311.8	249.4	Ricardo-AEA estimate based on carbon balance for integrated steelwork processes (2014)
1A1ci	Coke production	Blast furnace gas	Solid	CH ₄	kg/TJ	1.0	1.0	1.0	1.0	IPCC 2006 Guidelines
1A1ci	Coke production	Blast furnace gas	Solid	N ₂ O	kg/TJ	0.1	0.1	0.1	0.1	IPCC 2006 Guidelines
1A1ci	Coke production	Coke oven gas	Solid	CO ₂	t/TJ	44.5	42.9	48.8	42.5	Ricardo-AEA estimate based on carbon balance for integrated steelwork processes (2014)
1A1ci	Coke production	Coke oven gas	Solid	CH ₄	kg/TJ	1.0	1.0	1.0	1.0	IPCC 2006 Guidelines
1A1ci	Coke production	Coke oven gas	Solid	N ₂ O	kg/TJ	0.1	0.1	0.1	0.1	IPCC 2006 Guidelines
1A1ci	Coke production	Colliery methane	Gaseous	CO ₂	t/TJ	69.4	72.0	72.0	67.2	EU ETS data from EA, SEPA, NIEA (2014)
1A1ci	Coke production	Colliery methane	Gaseous	CH ₄	kg/TJ	1.0	1.0	1.0	1.0	IPCC 2006 Guidelines

Other Detailed Methodological Descriptions **A3**

Category	Source Name	Activity Name	Fuel (CRF)	Gas	Unit	1990	2013	Max	Min	EF Method Reference
1A1ci	Coke production	Colliery methane	Gaseous	N ₂ O	kg/TJ	0.1	0.1	0.1	0.1	IPCC 2006 Guidelines
1A1ci	Coke production	Natural gas	Gaseous	CO ₂	t/TJ	56.0	56.9	57.2	56.0	Natural gas carbon emission factor is derived from Gas Network Operator data on gas composition within each Local Distribution Zone, prepared for use by operators within the EU ETS. (DECC, 2014)
1A1ci	Coke production	Natural gas	Gaseous	CH ₄	kg/TJ	1.0	1.0	1.0	1.0	IPCC 2006 Guidelines
1A1ci	Coke production	Natural gas	Gaseous	N ₂ O	kg/TJ	0.1	0.1	0.1	0.1	IPCC 2006 Guidelines
1A1ci	Solid smokeless fuel production	Coke	Solid	CO ₂	t/TJ	114.3	111.3	114.3	101.8	Ricardo-AEA estimate based on carbon balance for integrated steelwork processes (2014)
1A1ci	Solid smokeless fuel production	Coke	Solid	CH ₄	kg/TJ	0.4	0.4	0.4	0.4	Brain et al, 1994
1A1ci	Solid smokeless fuel production	Coke	Solid	N ₂ O	kg/TJ	8.6	8.1	8.6	8.1	Based on Fynes & Sage, 1994
1A1ci	Solid smokeless fuel production	Natural gas	Gaseous	CO ₂	t/TJ	56.0	56.9	57.2	56.0	Natural gas carbon emission factor is derived from Gas Network Operator data on gas composition within each Local Distribution Zone, prepared for use by operators within the EU ETS. (DECC, 2014)
1A1ci	Solid smokeless fuel production	Natural gas	Gaseous	CH ₄	kg/TJ	1.0	1.0	1.0	1.0	IPCC 2006 Guidelines
1A1ci	Solid smokeless fuel production	Natural gas	Gaseous	N ₂ O	kg/TJ	0.1	0.1	0.1	0.1	IPCC 2006 Guidelines
1A1cii	Upstream Gas Production - fuel combustion	Gas oil	Liquid	CO ₂	t/TJ	73.9	74.1	74.2	73.5	Carbon Factors Review 2004

Category	Source Name	Activity Name	Fuel (CRF)	Gas	Unit	1990	2013	Max	Min	EF Method Reference
1A1cii	Upstream Gas Production - fuel combustion	Gas oil	Liquid	CH ₄	kg/TJ	3.4	5.1	6.6	2.1	EEMS 2013 atmospheric emissions data (DECC Offshore Inspectorate, 2014)
1A1cii	Upstream Gas Production - fuel combustion	Gas oil	Liquid	N ₂ O	kg/TJ	5.1	4.9	5.2	3.8	EEMS 2013 atmospheric emissions data (DECC Offshore Inspectorate, 2014)
1A1cii	Upstream Gas Production - fuel combustion	Natural gas	Gaseous	CO ₂	t/TJ	65.0	64.3	67.4	59.7	EEMS 2013 atmospheric emissions data (DECC Offshore Inspectorate, 2014)
1A1cii	Upstream Gas Production - fuel combustion	Natural gas	Gaseous	CH ₄	kg/TJ	40.4	20.8	56.5	20.8	EEMS 2013 atmospheric emissions data (DECC Offshore Inspectorate, 2014)
1A1cii	Upstream Gas Production - fuel combustion	Natural gas	Gaseous	N ₂ O	kg/TJ	5.9	4.9	7.9	4.7	EEMS 2013 atmospheric emissions data (DECC Offshore Inspectorate, 2014)
1A1cii	Upstream oil and gas production - combustion at gas separation plant	LPG	Liquid	CO ₂	t/TJ	64.7	64.7	64.7	64.7	Carbon Factors Review 2004
1A1cii	Upstream oil and gas production - combustion at gas separation plant	LPG	Liquid	CH ₄	kg/TJ	24.0	20.9	25.2	18.5	EEMS 2013 atmospheric emissions data (DECC Offshore Inspectorate, 2014)
1A1cii	Upstream oil and gas production - combustion at gas separation plant	LPG	Liquid	N ₂ O	kg/TJ	4.8	4.3	5.0	4.0	EEMS 2013 atmospheric emissions data (DECC Offshore Inspectorate, 2014)
1A1cii	Upstream oil and gas production - combustion at gas separation plant	OPG	Liquid	CO ₂	t/TJ	67.0	67.0	67.0	67.0	IPCC 2006 Guidelines
1A1cii	Upstream oil and gas production - combustion at gas separation plant	OPG	Liquid	CH ₄	kg/TJ	24.0	20.9	25.2	18.5	EEMS 2013 atmospheric emissions data (DECC Offshore Inspectorate, 2014)

Other Detailed Methodological Descriptions **A3**

Category	Source Name	Activity Name	Fuel (CRF)	Gas	Unit	1990	2013	Max	Min	EF Method Reference
1A1cii	Upstream oil and gas production - combustion at gas separation plant	OPG	Liquid	N ₂ O	kg/TJ	4.8	4.3	5.0	4.0	EEMS 2013 atmospheric emissions data (DECC Offshore Inspectorate, 2014)
1A1cii	Upstream Oil Production - fuel combustion	Gas oil	Liquid	CO ₂	t/TJ	73.9	74.1	74.2	73.5	Carbon Factors Review 2004
1A1cii	Upstream Oil Production - fuel combustion	Gas oil	Liquid	CH ₄	kg/TJ	2.3	2.4	3.7	1.8	EEMS 2013 atmospheric emissions data (DECC Offshore Inspectorate, 2014)
1A1cii	Upstream Oil Production - fuel combustion	Gas oil	Liquid	N ₂ O	kg/TJ	8.4	4.5	8.4	4.5	EEMS 2013 atmospheric emissions data (DECC Offshore Inspectorate, 2014)
1A1cii	Upstream Oil Production - fuel combustion	Natural gas	Gaseous	CO ₂	t/TJ	64.6	63.1	66.3	60.9	EEMS 2013 atmospheric emissions data (DECC Offshore Inspectorate, 2014)
1A1cii	Upstream Oil Production - fuel combustion	Natural gas	Gaseous	CH ₄	kg/TJ	24.5	21.3	25.7	19.0	EEMS 2013 atmospheric emissions data (DECC Offshore Inspectorate, 2014)
1A1cii	Upstream Oil Production - fuel combustion	Natural gas	Gaseous	N ₂ O	kg/TJ	4.9	4.3	5.1	4.0	EEMS 2013 atmospheric emissions data (DECC Offshore Inspectorate, 2014)
1A1ciii	Collieries - combustion	Coal	Solid	CO ₂	t/TJ	91.4	91.4	91.4	91.4	Based on Fynes & Sage, 1994
1A1ciii	Collieries - combustion	Coal	Solid	CH ₄	kg/TJ	0.4	0.4	0.4	0.4	Brain et al, 1994
1A1ciii	Collieries - combustion	Coal	Solid	N ₂ O	kg/TJ	5.5	5.4	6.0	5.2	Based on Fynes & Sage, 1994
1A1ciii	Collieries - combustion	Coke oven gas	Solid	CO ₂	t/TJ	44.5	42.9	48.8	42.5	Ricardo-AEA estimate based on carbon balance for integrated steelwork processes (2014)
1A1ciii	Collieries - combustion	Coke oven gas	Solid	CH ₄	kg/TJ	1.0	1.0	1.0	1.0	IPCC 2006 Guidelines
1A1ciii	Collieries - combustion	Coke oven gas	Solid	N ₂ O	kg/TJ	0.1	0.1	0.1	0.1	IPCC 2006 Guidelines

Other Detailed Methodological Descriptions **A3**

Category	Source Name	Activity Name	Fuel (CRF)	Gas	Unit	1990	2013	Max	Min	EF Method Reference
1A1ciii	Collieries - combustion	Colliery methane	Gaseous	CO ₂	t/TJ	69.4	72.0	72.0	67.2	EU ETS data from EA, SEPA, NIEA (2014)
1A1ciii	Collieries - combustion	Colliery methane	Gaseous	CH ₄	kg/TJ	1.0	1.0	1.0	1.0	IPCC 2006 Guidelines
1A1ciii	Collieries - combustion	Colliery methane	Gaseous	N ₂ O	kg/TJ	0.1	0.1	0.1	0.1	IPCC 2006 Guidelines
1A1ciii	Collieries - combustion	Natural gas	Gaseous	CO ₂	t/TJ	56.0	56.9	57.2	56.0	Natural gas carbon emission factor is derived from Gas Network Operator data on gas composition within each Local Distribution Zone, prepared for use by operators within the EU ETS. (DECC, 2014)
1A1ciii	Collieries - combustion	Natural gas	Gaseous	CH ₄	kg/TJ	1.0	1.0	1.0	1.0	IPCC 2006 Guidelines
1A1ciii	Collieries - combustion	Natural gas	Gaseous	N ₂ O	kg/TJ	0.1	0.1	0.1	0.1	IPCC 2006 Guidelines
1A1ciii	Nuclear fuel production	Natural gas	Gaseous	CO ₂	t/TJ	56.0	56.9	57.2	56.0	Natural gas carbon emission factor is derived from Gas Network Operator data on gas composition within each Local Distribution Zone, prepared for use by operators within the EU ETS. (DECC, 2014)
1A1ciii	Nuclear fuel production	Natural gas	Gaseous	CH ₄	kg/TJ	1.0	1.0	1.0	1.0	IPCC 2006 Guidelines
1A1ciii	Nuclear fuel production	Natural gas	Gaseous	N ₂ O	kg/TJ	0.1	0.1	0.1	0.1	IPCC 2006 Guidelines
1A1ciii	Gas production	Colliery methane	Gaseous	CO ₂	t/TJ	69.4	72.0	72.0	67.2	EU ETS data from EA, SEPA, NIEA (2014)
1A1ciii	Gas production	Colliery methane	Gaseous	CH ₄	kg/TJ	1.0	1.0	1.0	1.0	IPCC 2006 Guidelines
1A1ciii	Gas production	Colliery methane	Gaseous	N ₂ O	kg/TJ	0.1	0.1	0.1	0.1	IPCC 2006 Guidelines

Other Detailed Methodological Descriptions **A3**

Category	Source Name	Activity Name	Fuel (CRF)	Gas	Unit	1990	2013	Max	Min	EF Method Reference
1A1ciii	Gas production	LPG	Liquid	CO ₂	t/TJ	64.7	64.7	64.7	64.7	Carbon Factors Review 2004
1A1ciii	Gas production	LPG	Liquid	CH ₄	kg/TJ	1.0	1.0	1.0	1.0	IPCC 2006 Guidelines
1A1ciii	Gas production	LPG	Liquid	N ₂ O	kg/TJ	0.1	0.1	0.1	0.1	IPCC 2006 Guidelines
1A1ciii	Gas production	Natural gas	Gaseous	CO ₂	t/TJ	56.0	56.9	57.2	56.0	Natural gas carbon emission factor is derived from Gas Network Operator data on gas composition within each Local Distribution Zone, prepared for use by operators within the EU ETS. (DECC, 2014)
1A1ciii	Gas production	Natural gas	Gaseous	CH ₄	kg/TJ	1.0	1.0	1.0	1.0	IPCC 2006 Guidelines
1A1ciii	Gas production	Natural gas	Gaseous	N ₂ O	kg/TJ	0.1	0.1	0.1	0.1	IPCC 2006 Guidelines
1A1ciii	Gas production	OPG	Liquid	CO ₂	t/TJ	56.3	56.2	59.8	54.9	EU ETS data from EA, SEPA, NIEA (2014)
1A1ciii	Gas production	OPG	Liquid	CH ₄	kg/TJ	1.0	1.0	1.0	1.0	IPCC 2006 Guidelines
1A1ciii	Gas production	OPG	Liquid	N ₂ O	kg/TJ	0.1	0.1	0.1	0.1	IPCC 2006 Guidelines
1A1ciii	Gas production	Town gas	Solid	CO ₂	t/TJ	61.7	61.7	61.7	61.7	Ricardo-AEA estimate based on carbon balance for town gas manufacture (2014)
1A1ciii	Gas production	Town gas	Solid	CH ₄	kg/TJ	1.0	1.0	1.0	1.0	IPCC 2006 Guidelines
1A1ciii	Gas production	Town gas	Solid	N ₂ O	kg/TJ	0.1	0.1	0.1	0.1	IPCC 2006 Guidelines
1A2a	Blast furnaces	Blast furnace gas	Solid	CO ₂	t/TJ	280.5	278.3	311.8	249.4	Ricardo-AEA estimate based on carbon balance for integrated steelwork processes (2014)
1A2a	Blast furnaces	Blast furnace gas	Solid	CH ₄	kg/TJ	1.0	1.0	1.0	1.0	IPCC 2006 Guidelines
1A2a	Blast furnaces	Blast furnace gas	Solid	N ₂ O	kg/TJ	0.1	0.1	0.1	0.1	IPCC 2006 Guidelines

Other Detailed Methodological Descriptions **A3**

Category	Source Name	Activity Name	Fuel (CRF)	Gas	Unit	1990	2013	Max	Min	EF Method Reference
1A2a	Blast furnaces	Coke oven gas	Solid	CO ₂	t/TJ	44.5	42.9	48.8	42.5	Ricardo-AEA estimate based on carbon balance for integrated steelwork processes (2014)
1A2a	Blast furnaces	Coke oven gas	Solid	CH ₄	kg/TJ	1.0	1.0	1.0	1.0	IPCC 2006 Guidelines
1A2a	Blast furnaces	Coke oven gas	Solid	N ₂ O	kg/TJ	0.1	0.1	0.1	0.1	IPCC 2006 Guidelines
1A2a	Blast furnaces	Gas oil	Liquid	CO ₂	t/TJ	73.9	74.1	74.2	73.5	Carbon Factors Review 2004
1A2a	Blast furnaces	Gas oil	Liquid	CH ₄	kg/TJ	2.4	2.4	2.4	2.4	IPCC 2006 Guidelines
1A2a	Blast furnaces	Gas oil	Liquid	N ₂ O	kg/TJ	0.6	0.6	0.6	0.6	IPCC 2006 Guidelines
1A2a	Blast furnaces	LPG	Liquid	CO ₂	t/TJ	64.7	64.7	64.7	64.7	Carbon Factors Review 2004
1A2a	Blast furnaces	LPG	Liquid	CH ₄	kg/TJ	1.0	1.0	1.0	1.0	IPCC 2006 Guidelines
1A2a	Blast furnaces	LPG	Liquid	N ₂ O	kg/TJ	0.1	0.1	0.1	0.1	IPCC 2006 Guidelines
1A2a	Blast furnaces	Natural gas	Gaseous	CO ₂	t/TJ	56.0	56.9	57.2	56.0	Natural gas carbon emission factor is derived from Gas Network Operator data on gas composition within each Local Distribution Zone, prepared for use by operators within the EU ETS. (DECC, 2014)
1A2a	Blast furnaces	Natural gas	Gaseous	CH ₄	kg/TJ	1.0	1.0	1.0	1.0	IPCC 2006 Guidelines
1A2a	Blast furnaces	Natural gas	Gaseous	N ₂ O	kg/TJ	0.1	0.1	0.1	0.1	IPCC 2006 Guidelines
1A2a	Iron and steel - combustion plant	Blast furnace gas	Solid	CH ₄	kg/TJ	1.0	1.0	1.0	1.0	IPCC 2006 Guidelines
1A2a	Iron and steel - combustion plant	Coal	Solid	CO ₂	t/TJ	90.4	90.4	90.4	90.4	Based on Fynes & Sage, 1994
1A2a	Iron and steel - combustion plant	Coal	Solid	CH ₄	kg/TJ	0.4	0.4	0.4	0.4	Brain et al, 1994

Other Detailed Methodological Descriptions **A3**

Category	Source Name	Activity Name	Fuel (CRF)	Gas	Unit	1990	2013	Max	Min	EF Method Reference
1A2a	Iron and steel - combustion plant	Coal	Solid	N ₂ O	kg/TJ	8.6	8.2	8.7	7.9	Based on Fynes & Sage, 1994
1A2a	Iron and steel - combustion plant	Coke	Solid	CO ₂	t/TJ	114.3	111.3	114.3	101.8	Ricardo-AEA estimate based on carbon balance for integrated steelwork processes (2014)
1A2a	Iron and steel - combustion plant	Coke	Solid	CH ₄	kg/TJ	0.4	0.4	0.4	0.4	Brain et al, 1994
1A2a	Iron and steel - combustion plant	Coke	Solid	N ₂ O	kg/TJ	8.6	8.1	8.6	8.1	Based on Fynes & Sage, 1994
1A2a	Iron and steel - combustion plant	Coke oven gas	Solid	CO ₂	t/TJ	44.5	42.9	48.8	42.5	Ricardo-AEA estimate based on carbon balance for integrated steelwork processes (2014)
1A2a	Iron and steel - combustion plant	Coke oven gas	Solid	CH ₄	kg/TJ	1.0	1.0	1.0	1.0	IPCC 2006 Guidelines
1A2a	Iron and steel - combustion plant	Coke oven gas	Solid	N ₂ O	kg/TJ	0.1	0.1	0.1	0.1	IPCC 2006 Guidelines
1A2a	Iron and steel - combustion plant	Fuel oil	Liquid	CO ₂	t/TJ	77.5	78.4	78.4	77.3	Carbon Factors Review 2004
1A2a	Iron and steel - combustion plant	Fuel oil	Liquid	CH ₄	kg/TJ	3.0	3.0	3.0	3.0	IPCC 2006 Guidelines
1A2a	Iron and steel - combustion plant	Fuel oil	Liquid	N ₂ O	kg/TJ	0.6	0.6	0.6	0.6	IPCC 2006 Guidelines
1A2a	Iron and steel - combustion plant	Gas oil	Liquid	CO ₂	t/TJ	73.9	74.1	74.2	73.5	Carbon Factors Review 2004
1A2a	Iron and steel - combustion plant	Gas oil	Liquid	CH ₄	kg/TJ	3.0	3.0	3.0	3.0	IPCC 2006 Guidelines

Other Detailed Methodological Descriptions **A3**

Category	Source Name	Activity Name	Fuel (CRF)	Gas	Unit	1990	2013	Max	Min	EF Method Reference
1A2a	Iron and steel - combustion plant	Gas oil	Liquid	N ₂ O	kg/TJ	0.6	0.6	0.6	0.6	IPCC 2006 Guidelines
1A2a	Iron and steel - combustion plant	LPG	Liquid	CO ₂	t/TJ	64.7	64.7	64.7	64.7	Carbon Factors Review 2004
1A2a	Iron and steel - combustion plant	LPG	Liquid	CH ₄	kg/TJ	1.0	1.0	1.0	1.0	IPCC 2006 Guidelines
1A2a	Iron and steel - combustion plant	LPG	Liquid	N ₂ O	kg/TJ	0.1	0.1	0.1	0.1	IPCC 2006 Guidelines
1A2a	Iron and steel - combustion plant	Natural gas	Gaseous	CO ₂	t/TJ	56.0	56.9	57.2	56.0	Natural gas carbon emission factor is derived from Gas Network Operator data on gas composition within each Local Distribution Zone, prepared for use by operators within the EU ETS. (DECC, 2014)
1A2a	Iron and steel - combustion plant	Natural gas	Gaseous	CH ₄	kg/TJ	1.0	1.0	1.0	1.0	IPCC 2006 Guidelines
1A2a	Iron and steel - combustion plant	Natural gas	Gaseous	N ₂ O	kg/TJ	0.1	0.1	0.1	0.1	IPCC 2006 Guidelines
1A2a	Iron and steel - combustion plant	Town gas	Solid	CO ₂	t/TJ	61.7	61.7	61.7	61.7	Ricardo-AEA estimate based on carbon balance for town gas manufacture (2014)
1A2a	Iron and steel - combustion plant	Town gas	Solid	CH ₄	kg/TJ	1.0	1.0	1.0	1.0	IPCC 2006 Guidelines
1A2a	Iron and steel - combustion plant	Town gas	Solid	N ₂ O	kg/TJ	0.1	0.1	0.1	0.1	IPCC 2006 Guidelines
1A2b	Autogeneration - exported to grid	Coal	Solid	CO ₂	t/TJ	94.0	85.6	94.0	83.2	Based on Fynes & Sage, 1994

Other Detailed Methodological Descriptions **A3**

Category	Source Name	Activity Name	Fuel (CRF)	Gas	Unit	1990	2013	Max	Min	EF Method Reference
1A2b	Autogeneration - exported to grid	Coal	Solid	CH ₄	kg/TJ	0.8	0.8	0.8	0.7	Brain et al, 1994
1A2b	Autogeneration - exported to grid	Coal	Solid	N ₂ O	kg/TJ	2.5	2.6	2.6	2.4	Based on Fynes & Sage, 1994
1A2b	Autogenerators	Coal	Solid	CO ₂	t/TJ	94.0	85.6	94.0	83.2	Based on Fynes & Sage, 1994
1A2b	Autogenerators	Coal	Solid	CH ₄	kg/TJ	0.8	0.8	0.8	0.7	Brain et al, 1994
1A2b	Autogenerators	Coal	Solid	N ₂ O	kg/TJ	2.5	2.6	2.6	2.4	Based on Fynes & Sage, 1994
1A2b	Non-Ferrous Metal (combustion)	Coal	Solid	CO ₂	t/TJ	94.0	94.0	94.0	94.0	Based on Fynes & Sage, 1994
1A2b	Non-Ferrous Metal (combustion)	Coal	Solid	CH ₄	kg/TJ	0.4	0.4	0.4	0.4	Brain et al, 1994
1A2b	Non-Ferrous Metal (combustion)	Coal	Solid	N ₂ O	kg/TJ	8.1	8.4	8.5	7.8	Based on Fynes & Sage, 1994
1A2b	Non-Ferrous Metal (combustion)	Coke	Solid	CO ₂	t/TJ	114.3	111.3	114.3	101.8	Ricardo-AEA estimate based on carbon balance for integrated steelwork processes (2014)
1A2b	Non-Ferrous Metal (combustion)	Coke	Solid	CH ₄	kg/TJ	0.4	0.4	0.4	0.4	Brain et al, 1994
1A2b	Non-Ferrous Metal (combustion)	Coke	Solid	N ₂ O	kg/TJ	8.6	8.1	8.6	8.1	Based on Fynes & Sage, 1994
1A2b	Non-Ferrous Metal (combustion)	Fuel oil	Liquid	CO ₂	t/TJ	77.5	78.4	78.4	77.3	Carbon Factors Review 2004
1A2b	Non-Ferrous Metal (combustion)	Fuel oil	Liquid	CH ₄	kg/TJ	3.0	3.0	3.0	3.0	IPCC 2006 Guidelines

Other Detailed Methodological Descriptions **A3**

Category	Source Name	Activity Name	Fuel (CRF)	Gas	Unit	1990	2013	Max	Min	EF Method Reference
1A2b	Non-Ferrous Metal (combustion)	Fuel oil	Liquid	N ₂ O	kg/TJ	0.6	0.6	0.6	0.6	IPCC 2006 Guidelines
1A2b	Non-Ferrous Metal (combustion)	Gas oil	Liquid	CO ₂	t/TJ	73.9	74.1	74.2	73.5	Carbon Factors Review 2004
1A2b	Non-Ferrous Metal (combustion)	Gas oil	Liquid	CH ₄	kg/TJ	3.0	3.0	3.0	3.0	IPCC 2006 Guidelines
1A2b	Non-Ferrous Metal (combustion)	Gas oil	Liquid	N ₂ O	kg/TJ	0.6	0.6	0.6	0.6	IPCC 2006 Guidelines
1A2b	Non-Ferrous Metal (combustion)	Natural gas	Gaseous	CO ₂	t/TJ	56.0	56.9	57.2	56.0	Natural gas carbon emission factor is derived from Gas Network Operator data on gas composition within each Local Distribution Zone, prepared for use by operators within the EU ETS. (DECC, 2014)
1A2b	Non-Ferrous Metal (combustion)	Natural gas	Gaseous	CH ₄	kg/TJ	1.0	1.0	1.0	1.0	IPCC 2006 Guidelines
1A2b	Non-Ferrous Metal (combustion)	Natural gas	Gaseous	N ₂ O	kg/TJ	0.1	0.1	0.1	0.1	IPCC 2006 Guidelines
1A2c	Chemicals (combustion)	Coal	Solid	CO ₂	t/TJ	94.0	94.0	94.0	94.0	Based on Fynes & Sage, 1994
1A2c	Chemicals (combustion)	Coal	Solid	CH ₄	kg/TJ	0.4	0.4	0.4	0.4	Brain et al, 1994
1A2c	Chemicals (combustion)	Coal	Solid	N ₂ O	kg/TJ	8.1	8.4	8.5	7.8	Based on Fynes & Sage, 1994
1A2c	Chemicals (combustion)	Fuel oil	Liquid	CO ₂	t/TJ	77.5	78.4	78.4	77.3	Carbon Factors Review 2004
1A2c	Chemicals (combustion)	Fuel oil	Liquid	CH ₄	kg/TJ	3.0	3.0	3.0	3.0	IPCC 2006 Guidelines
1A2c	Chemicals (combustion)	Fuel oil	Liquid	N ₂ O	kg/TJ	0.6	0.6	0.6	0.6	IPCC 2006 Guidelines
1A2c	Chemicals (combustion)	Gas oil	Liquid	CO ₂	t/TJ	73.9	74.1	74.2	73.5	Carbon Factors Review 2004

Category	Source Name	Activity Name	Fuel (CRF)	Gas	Unit	1990	2013	Max	Min	EF Method Reference
1A2c	Chemicals (combustion)	Gas oil	Liquid	CH ₄	kg/TJ	3.0	3.0	3.0	3.0	IPCC 2006 Guidelines
1A2c	Chemicals (combustion)	Gas oil	Liquid	N ₂ O	kg/TJ	0.6	0.6	0.6	0.6	IPCC 2006 Guidelines
1A2c	Chemicals (combustion)	Natural gas	Gaseous	CO ₂	t/TJ	56.0	56.9	57.2	56.0	Natural gas carbon emission factor is derived from Gas Network Operator data on gas composition within each Local Distribution Zone, prepared for use by operators within the EU ETS. (DECC, 2014)
1A2c	Chemicals (combustion)	Natural gas	Gaseous	CH ₄	kg/TJ	1.0	1.0	1.0	1.0	IPCC 2006 Guidelines
1A2c	Chemicals (combustion)	Natural gas	Gaseous	N ₂ O	kg/TJ	0.1	0.1	0.1	0.1	IPCC 2006 Guidelines
1A2d	Pulp, Paper and Print (combustion)	Coal	Solid	CO ₂	t/TJ	94.0	94.0	94.0	94.0	Based on Fynes & Sage, 1994
1A2d	Pulp, Paper and Print (combustion)	Coal	Solid	CH ₄	kg/TJ	0.4	0.4	0.4	0.4	Brain et al, 1994
1A2d	Pulp, Paper and Print (combustion)	Coal	Solid	N ₂ O	kg/TJ	8.1	8.4	8.5	7.8	Based on Fynes & Sage, 1994
1A2d	Pulp, Paper and Print (combustion)	Fuel oil	Liquid	CO ₂	t/TJ	77.5	78.4	78.4	77.3	Carbon Factors Review 2004
1A2d	Pulp, Paper and Print (combustion)	Fuel oil	Liquid	CH ₄	kg/TJ	3.0	3.0	3.0	3.0	IPCC 2006 Guidelines
1A2d	Pulp, Paper and Print (combustion)	Fuel oil	Liquid	N ₂ O	kg/TJ	0.6	0.6	0.6	0.6	IPCC 2006 Guidelines
1A2d	Pulp, Paper and Print (combustion)	Gas oil	Liquid	CO ₂	t/TJ	73.9	74.1	74.2	73.5	Carbon Factors Review 2004
1A2d	Pulp, Paper and Print (combustion)	Gas oil	Liquid	CH ₄	kg/TJ	3.0	3.0	3.0	3.0	IPCC 2006 Guidelines

Category	Source Name	Activity Name	Fuel (CRF)	Gas	Unit	1990	2013	Max	Min	EF Method Reference
1A2d	Pulp, Paper and Print (combustion)	Gas oil	Liquid	N ₂ O	kg/TJ	0.6	0.6	0.6	0.6	IPCC 2006 Guidelines
1A2d	Pulp, Paper and Print (combustion)	Natural gas	Gaseous	CO ₂	t/TJ	56.0	56.9	57.2	56.0	Natural gas carbon emission factor is derived from Gas Network Operator data on gas composition within each Local Distribution Zone, prepared for use by operators within the EU ETS. (DECC, 2014)
1A2d	Pulp, Paper and Print (combustion)	Natural gas	Gaseous	CH ₄	kg/TJ	1.0	1.0	1.0	1.0	IPCC 2006 Guidelines
1A2d	Pulp, Paper and Print (combustion)	Natural gas	Gaseous	N ₂ O	kg/TJ	0.1	0.1	0.1	0.1	IPCC 2006 Guidelines
1A2e	Food & drink, tobacco (combustion)	Coal	Solid	CO ₂	t/TJ	94.0	94.0	94.0	94.0	Based on Fynes & Sage, 1994
1A2e	Food & drink, tobacco (combustion)	Coal	Solid	CH ₄	kg/TJ	0.4	0.4	0.4	0.4	Brain et al, 1994
1A2e	Food & drink, tobacco (combustion)	Coal	Solid	N ₂ O	kg/TJ	8.1	8.4	8.5	7.8	Based on Fynes & Sage, 1994
1A2e	Food & drink, tobacco (combustion)	Fuel oil	Liquid	CO ₂	t/TJ	77.5	78.4	78.4	77.3	Carbon Factors Review 2004
1A2e	Food & drink, tobacco (combustion)	Fuel oil	Liquid	CH ₄	kg/TJ	3.0	3.0	3.0	3.0	IPCC 2006 Guidelines
1A2e	Food & drink, tobacco (combustion)	Fuel oil	Liquid	N ₂ O	kg/TJ	0.6	0.6	0.6	0.6	IPCC 2006 Guidelines
1A2e	Food & drink, tobacco (combustion)	Gas oil	Liquid	CO ₂	t/TJ	73.9	74.1	74.2	73.5	Carbon Factors Review 2004

Other Detailed Methodological Descriptions **A3**

Category	Source Name	Activity Name	Fuel (CRF)	Gas	Unit	1990	2013	Max	Min	EF Method Reference
1A2e	Food & drink, tobacco (combustion)	Gas oil	Liquid	CH ₄	kg/TJ	3.0	3.0	3.0	3.0	IPCC 2006 Guidelines
1A2e	Food & drink, tobacco (combustion)	Gas oil	Liquid	N ₂ O	kg/TJ	0.6	0.6	0.6	0.6	IPCC 2006 Guidelines
1A2e	Food & drink, tobacco (combustion)	Natural gas	Gaseous	CO ₂	t/TJ	56.0	56.9	57.2	56.0	Natural gas carbon emission factor is derived from Gas Network Operator data on gas composition within each Local Distribution Zone, prepared for use by operators within the EU ETS. (DECC, 2014)
1A2e	Food & drink, tobacco (combustion)	Natural gas	Gaseous	CH ₄	kg/TJ	1.0	1.0	1.0	1.0	IPCC 2006 Guidelines
1A2e	Food & drink, tobacco (combustion)	Natural gas	Gaseous	N ₂ O	kg/TJ	0.1	0.1	0.1	0.1	IPCC 2006 Guidelines
1A2f	Cement production - combustion	Coal	Solid	CO ₂	t/TJ	C	C	C	C	BCA/Lafarge Cement (2006,2007,2008, 2009, 2010,2011,2012) Mineral Products Association (2013, 2014)
1A2f	Cement production - combustion	Coal	Solid	CH ₄	kg/TJ	11.4	11.8	11.9	11.0	Brain et al, 1994
1A2f	Cement production - combustion	Coal	Solid	N ₂ O	kg/TJ	4.1	4.3	4.3	4.0	Based on Fynes & Sage, 1994
1A2f	Cement production - combustion	Fuel oil	Liquid	CO ₂	t/TJ	77.5	78.4	78.4	77.3	Carbon Factors Review 2004
1A2f	Cement production - combustion	Fuel oil	Liquid	CH ₄	kg/TJ	3.0	3.0	3.0	3.0	IPCC 2006 Guidelines
1A2f	Cement production - combustion	Fuel oil	Liquid	N ₂ O	kg/TJ	0.6	0.6	0.6	0.6	IPCC 2006 Guidelines

Other Detailed Methodological Descriptions **A3**

Category	Source Name	Activity Name	Fuel (CRF)	Gas	Unit	1990	2013	Max	Min	EF Method Reference
1A2f	Cement production - combustion	Gas oil	Liquid	CO ₂	t/TJ	C	C	C	C	BCA/Lafarge Cement (2006,2007,2008, 2009, 2010,2011,2012) Mineral Products Association (2013, 2014)
1A2f	Cement production - combustion	Gas oil	Liquid	CH ₄	kg/TJ	3.0	3.0	3.0	3.0	IPCC 2006 Guidelines
1A2f	Cement production - combustion	Gas oil	Liquid	N ₂ O	kg/TJ	0.6	0.6	0.6	0.6	IPCC 2006 Guidelines
1A2f	Cement production - combustion	Natural gas	Gaseous	CO ₂	t/TJ	56.0	56.9	57.2	56.0	Natural gas carbon emission factor is derived from Gas Network Operator data on gas composition within each Local Distribution Zone, prepared for use by operators within the EU ETS. (DECC, 2014)
1A2f	Cement production - combustion	Natural gas	Gaseous	CH ₄	kg/TJ	1.0	1.0	1.0	1.0	IPCC 2006 Guidelines
1A2f	Cement production - combustion	Natural gas	Gaseous	N ₂ O	kg/TJ	0.1	0.1	0.1	0.1	IPCC 2006 Guidelines
1A2f	Cement production - combustion	Petroleum coke	Liquid	CO ₂	t/TJ	C	C	C	C	BCA/Lafarge Cement (2006,2007,2008, 2009, 2010,2011,2012) Mineral Products Association (2013, 2014)
1A2f	Cement production - combustion	Petroleum coke	Liquid	CH ₄	kg/TJ	3.2	3.2	3.5	3.2	IPCC 2006 Guidelines
1A2f	Cement production - combustion	Petroleum coke	Liquid	N ₂ O	kg/TJ	3.8	4.2	4.2	3.8	Based on Fynes & Sage, 1994
1A2f	Cement production - combustion	Scrap tyres	Other	CO ₂	t/TJ	C	C	C	C	BCA/Lafarge Cement (2006,2007,2008, 2009, 2010,2011,2012) Mineral Products Association (2013, 2014)
1A2f	Cement production - combustion	Scrap tyres	Other	CH ₄	kg/TJ	31.0	30.0	31.0	30.0	IPCC 2006 Guidelines

Other Detailed Methodological Descriptions **A3**

Category	Source Name	Activity Name	Fuel (CRF)	Gas	Unit	1990	2013	Max	Min	EF Method Reference
1A2f	Cement production - combustion	Scrap tyres	Other	N ₂ O	kg/TJ	4.1	4.0	4.1	4.0	IPCC 2006 Guidelines
1A2f	Cement production - combustion	Waste oils	Liquid	CO ₂	t/TJ	C	C	C	C	Based on data supplied by the Mineral Products Association, 2012 - 2014
1A2f	Cement production - combustion	Waste oils	Liquid	CH ₄	kg/TJ	3.0	3.0	3.0	3.0	Use 2006 IPCC default for gas oil
1A2f	Cement production - combustion	Waste oils	Liquid	N ₂ O	kg/TJ	3.8	3.8	3.8	3.8	IPCC 2006 Guidelines
1A2f	Cement production - combustion	Waste solvent	Liquid	CO ₂	t/TJ	C	C	C	C	Based on data supplied by the Mineral Products Association, 2012 - 2014
1A2f	Cement production - combustion	Waste solvent	Liquid	CH ₄	kg/TJ	2.8	2.8	2.8	2.8	IPCC 2006 Guidelines
1A2f	Cement production - combustion	Waste solvent	Liquid	N ₂ O	kg/TJ	0.6	0.6	0.6	0.6	IPCC 2006 Guidelines
1A2f	Lime production - non decarbonising	Coal	Solid	CO ₂	t/TJ	94.0	99.2	104.3	88.4	EU ETS data from EA, SEPA, NIEA (2014)
1A2f	Lime production - non decarbonising	Coal	Solid	CH ₄	kg/TJ	0.4	0.4	0.4	0.4	Brain et al, 1994
1A2f	Lime production - non decarbonising	Coal	Solid	N ₂ O	kg/TJ	8.1	8.4	8.5	7.8	Based on Fynes & Sage, 1994
1A2f	Lime production - non decarbonising	Coke	Solid	CO ₂	t/TJ	114.3	111.3	114.3	101.8	Ricardo-AEA estimate based on carbon balance for integrated steelwork processes (2014)
1A2f	Lime production - non decarbonising	Coke	Solid	CH ₄	kg/TJ	0.4	0.4	0.4	0.4	Brain et al, 1994

Other Detailed Methodological Descriptions **A3**

Category	Source Name	Activity Name	Fuel (CRF)	Gas	Unit	1990	2013	Max	Min	EF Method Reference
1A2f	Lime production - non decarbonising	Coke	Solid	N ₂ O	kg/TJ	8.6	8.1	8.6	8.1	Based on Fynes & Sage, 1994
1A2f	Lime production - non decarbonising	Natural gas	Gaseous	CO ₂	t/TJ	56.0	56.9	57.2	56.0	Natural gas carbon emission factor is derived from Gas Network Operator data on gas composition within each Local Distribution Zone, prepared for use by operators within the EU ETS. (DECC, 2014)
1A2f	Lime production - non decarbonising	Natural gas	Gaseous	CH ₄	kg/TJ	1.0	1.0	1.0	1.0	IPCC 2006 Guidelines
1A2f	Lime production - non decarbonising	Natural gas	Gaseous	N ₂ O	kg/TJ	0.1	0.1	0.1	0.1	IPCC 2006 Guidelines
1A2gvii	Industrial off-road mobile machinery	DERV	Liquid	CO ₂	t/TJ	73.5	73.0	73.5	72.8	Carbon Factors Review 2004
1A2gvii	Industrial off-road mobile machinery	DERV	Liquid	CH ₄	kg/TJ	3.8	2.8	3.8	2.8	Developed based on T3 approach considering emission limits on total hydrocarbons
1A2gvii	Industrial off-road mobile machinery	DERV	Liquid	N ₂ O	kg/TJ	31.4	28.9	31.4	28.8	EMEP/EEA Guidebook, 2006
1A2gvii	Industrial off-road mobile machinery	Gas oil	Liquid	CO ₂	t/TJ	73.9	74.1	74.2	73.5	Carbon Factors Review 2004
1A2gvii	Industrial off-road mobile machinery	Gas oil	Liquid	CH ₄	kg/TJ	3.8	2.8	3.8	2.8	Developed based on T3 approach considering emission limits on total hydrocarbons
1A2gvii	Industrial off-road mobile machinery	Gas oil	Liquid	N ₂ O	kg/TJ	31.4	29.1	31.4	29.1	EMEP/EEA Guidebook, 2006
1A2gvii	Industrial off-road mobile machinery	Petrol	Liquid	CO ₂	t/TJ	70.2	70.0	70.2	70.0	Carbon Factors Review 2004

Other Detailed Methodological Descriptions **A3**

Category	Source Name	Activity Name	Fuel (CRF)	Gas	Unit	1990	2013	Max	Min	EF Method Reference
1A2gvii	Industrial off-road mobile machinery	Petrol	Liquid	CH ₄	kg/TJ	84.2	83.9	84.2	83.9	Developed based on T3 approach considering emission limits on total hydrocarbons
1A2gvii	Industrial off-road mobile machinery	Petrol	Liquid	N ₂ O	kg/TJ	1.2	1.1	1.2	1.1	EMEP/EEA Guidebook, 2006
1A2gviii	Autogeneration - exported to grid	Natural gas	Gaseous	CO ₂	t/TJ	56.0	56.9	57.2	56.0	Natural gas carbon emission factor is derived from Gas Network Operator data on gas composition within each Local Distribution Zone, prepared for use by operators within the EU ETS. (DECC, 2014)
1A2gviii	Autogeneration - exported to grid	Natural gas	Gaseous	CH ₄	kg/TJ	1.0	1.0	1.0	1.0	IPCC 2006 Guidelines
1A2gviii	Autogeneration - exported to grid	Natural gas	Gaseous	N ₂ O	kg/TJ	0.1	0.1	0.1	0.1	IPCC 2006 Guidelines
1A2gviii	Autogenerators	Biogas	Biomass	CH ₄	kg/TJ	1.0	1.0	1.0	1.0	IPCC 2006 Guidelines
1A2gviii	Autogenerators	Biogas	Biomass	N ₂ O	kg/TJ	0.1	0.1	0.1	0.1	IPCC 2006 Guidelines
1A2gviii	Autogenerators	Natural gas	Gaseous	CO ₂	t/TJ	56.0	56.9	57.2	56.0	Natural gas carbon emission factor is derived from Gas Network Operator data on gas composition within each Local Distribution Zone, prepared for use by operators within the EU ETS. (DECC, 2014)
1A2gviii	Autogenerators	Natural gas	Gaseous	CH ₄	kg/TJ	1.0	1.0	1.0	1.0	IPCC 2006 Guidelines
1A2gviii	Autogenerators	Natural gas	Gaseous	N ₂ O	kg/TJ	0.1	0.1	0.1	0.1	IPCC 2006 Guidelines
1A2gviii	Other industrial combustion	Biomass	Biomass	CH ₄	kg/TJ	26.8	26.8	26.8	26.8	IPCC 2006 Guidelines
1A2gviii	Other industrial combustion	Biomass	Biomass	N ₂ O	kg/TJ	3.6	3.6	3.6	3.6	IPCC 2006 Guidelines
1A2gviii	Other industrial combustion	Burning oil	Liquid	CO ₂	t/TJ	71.8	71.7	71.8	71.7	Carbon Factors Review 2004

Other Detailed Methodological Descriptions **A3**

Category	Source Name	Activity Name	Fuel (CRF)	Gas	Unit	1990	2013	Max	Min	EF Method Reference
1A2gviii	Other industrial combustion	Burning oil	Liquid	CH ₄	kg/TJ	3.0	3.0	3.0	3.0	IPCC 2006 Guidelines
1A2gviii	Other industrial combustion	Burning oil	Liquid	N ₂ O	kg/TJ	0.6	0.6	0.6	0.6	IPCC 2006 Guidelines
1A2gviii	Other industrial combustion	Coal	Solid	CO ₂	t/TJ	94.0	94.0	94.0	94.0	Based on Fynes & Sage, 1994
1A2gviii	Other industrial combustion	Coal	Solid	CH ₄	kg/TJ	0.4	0.4	0.4	0.4	Brain et al, 1994
1A2gviii	Other industrial combustion	Coal	Solid	N ₂ O	kg/TJ	8.1	8.4	8.5	7.8	Based on Fynes & Sage, 1994
1A2gviii	Other industrial combustion	Coke	Solid	CO ₂	t/TJ	114.3	111.3	114.3	101.8	Ricardo-AEA estimate based on carbon balance for integrated steelwork processes (2014)
1A2gviii	Other industrial combustion	Coke	Solid	CH ₄	kg/TJ	0.4	0.4	0.4	0.4	Brain et al, 1994
1A2gviii	Other industrial combustion	Coke	Solid	N ₂ O	kg/TJ	8.6	8.1	8.6	8.1	Based on Fynes & Sage, 1994
1A2gviii	Other industrial combustion	Coke oven gas	Solid	CO ₂	t/TJ	44.5	42.9	48.8	42.5	Ricardo-AEA estimate based on carbon balance for integrated steelwork processes (2014)
1A2gviii	Other industrial combustion	Coke oven gas	Solid	CH ₄	kg/TJ	1.0	1.0	1.0	1.0	IPCC 2006 Guidelines
1A2gviii	Other industrial combustion	Coke oven gas	Solid	N ₂ O	kg/TJ	0.1	0.1	0.1	0.1	IPCC 2006 Guidelines
1A2gviii	Other industrial combustion	Colliery methane	Gaseous	CO ₂	t/TJ	69.4	72.0	72.0	67.2	EU ETS data from EA, SEPA, NIEA (2014)
1A2gviii	Other industrial combustion	Colliery methane	Gaseous	CH ₄	kg/TJ	1.0	1.0	1.0	1.0	IPCC 2006 Guidelines
1A2gviii	Other industrial combustion	Colliery methane	Gaseous	N ₂ O	kg/TJ	0.1	0.1	0.1	0.1	IPCC 2006 Guidelines
1A2gviii	Other industrial combustion	Fuel oil	Liquid	CO ₂	t/TJ	77.5	78.4	78.4	77.3	Carbon Factors Review 2004
1A2gviii	Other industrial combustion	Fuel oil	Liquid	CH ₄	kg/TJ	3.0	3.0	3.0	3.0	IPCC 2006 Guidelines

Other Detailed Methodological Descriptions **A3**

Category	Source Name	Activity Name	Fuel (CRF)	Gas	Unit	1990	2013	Max	Min	EF Method Reference
1A2gviii	Other industrial combustion	Fuel oil	Liquid	N ₂ O	kg/TJ	0.6	0.6	0.6	0.6	IPCC 2006 Guidelines
1A2gviii	Other industrial combustion	Gas oil	Liquid	CO ₂	t/TJ	73.9	74.1	74.2	73.5	Carbon Factors Review 2004
1A2gviii	Other industrial combustion	Gas oil	Liquid	CH ₄	kg/TJ	3.0	3.0	3.0	3.0	IPCC 2006 Guidelines
1A2gviii	Other industrial combustion	Gas oil	Liquid	N ₂ O	kg/TJ	0.6	0.6	0.6	0.6	IPCC 2006 Guidelines
1A2gviii	Other industrial combustion	LPG	Liquid	CO ₂	t/TJ	64.7	64.7	64.7	64.7	UKPIA (1989)
1A2gviii	Other industrial combustion	LPG	Liquid	CH ₄	kg/TJ	1.0	1.0	1.0	1.0	IPCC 2006 Guidelines
1A2gviii	Other industrial combustion	LPG	Liquid	N ₂ O	kg/TJ	0.1	0.1	0.1	0.1	IPCC 2006 Guidelines
1A2gviii	Other industrial combustion	Natural gas	Gaseous	CO ₂	t/TJ	56.0	56.9	57.2	56.0	Natural gas carbon emission factor is derived from Gas Network Operator data on gas composition within each Local Distribution Zone, prepared for use by operators within the EU ETS. (DECC, 2014)
1A2gviii	Other industrial combustion	Natural gas	Gaseous	CH ₄	kg/TJ	1.0	1.0	1.0	1.0	IPCC 2006 Guidelines
1A2gviii	Other industrial combustion	Natural gas	Gaseous	N ₂ O	kg/TJ	0.1	0.1	0.1	0.1	IPCC 2006 Guidelines
1A2gviii	Other industrial combustion	OPG	Liquid	CO ₂	t/TJ	56.3	56.2	59.8	54.9	EU ETS data from EA, SEPA, NIEA (2014)
1A2gviii	Other industrial combustion	OPG	Liquid	CH ₄	kg/TJ	1.0	1.0	1.0	1.0	IPCC 2006 Guidelines
1A2gviii	Other industrial combustion	OPG	Liquid	N ₂ O	kg/TJ	0.1	0.1	0.1	0.1	IPCC 2006 Guidelines
1A2gviii	Other industrial combustion	Petroleum coke	Liquid	CO ₂	t/TJ	79.6	87.6	89.7	79.6	EU ETS data from EA, SEPA, NIEA (2014)
1A2gviii	Other industrial combustion	Petroleum coke	Liquid	CH ₄	kg/TJ	3.2	3.2	3.5	3.2	IPCC 2006 Guidelines
1A2gviii	Other industrial combustion	Petroleum coke	Liquid	N ₂ O	kg/TJ	2.3	2.6	2.6	2.3	Based on Fynes & Sage, 1994
1A2gviii	Other industrial combustion	SSF	Solid	CO ₂	t/TJ	110.5	102.3	110.5	93.6	British Coal (1989)

Other Detailed Methodological Descriptions **A3**

Category	Source Name	Activity Name	Fuel (CRF)	Gas	Unit	1990	2013	Max	Min	EF Method Reference
1A2gviii	Other industrial combustion	SSF	Solid	CH ₄	kg/TJ	0.4	0.4	0.4	0.4	Brain et al, 1994
1A2gviii	Other industrial combustion	SSF	Solid	N ₂ O	kg/TJ	9.0	8.4	9.0	7.7	Based on Fynes & Sage, 1994
1A2gviii	Other industrial combustion	Town gas	Solid	CO ₂	t/TJ	61.7	61.7	61.7	61.7	Ricardo-AEA estimate based on carbon balance for town gas manufacture (2014)
1A2gviii	Other industrial combustion	Town gas	Solid	CH ₄	kg/TJ	1.0	1.0	1.0	1.0	IPCC 2006 Guidelines
1A2gviii	Other industrial combustion	Town gas	Solid	N ₂ O	kg/TJ	0.1	0.1	0.1	0.1	IPCC 2006 Guidelines
1A2gviii	Other industrial combustion	Waste solvent	Liquid	CO ₂	t/TJ	38.4	35.8	38.4	35.2	EU ETS data from EA, SEPA, NIEA (2014)
1A2gviii	Other industrial combustion	Waste solvent	Liquid	CH ₄	kg/TJ	2.8	2.8	2.8	2.8	IPCC 2006 Guidelines
1A2gviii	Other industrial combustion	Waste solvent	Liquid	N ₂ O	kg/TJ	0.6	0.6	0.6	0.6	IPCC 2006 Guidelines
1A2gviii	Other industrial combustion	Wood	Biomass	CH ₄	kg/TJ	27.9	27.9	27.9	27.9	IPCC 2006 Guidelines
1A2gviii	Other industrial combustion	Wood	Biomass	N ₂ O	kg/TJ	3.7	3.7	3.7	3.7	IPCC 2006 Guidelines
1A3a	Aircraft - domestic cruise	Aviation spirit	Liquid	CO ₂	t/TJ	69.6	69.5	69.6	69.3	Carbon Factors Review 2004
1A3a	Aircraft - domestic cruise	Aviation spirit	Liquid	N ₂ O	kg/TJ	2.2	2.2	2.2	2.2	Emissions from 1990 to the latest inventory year derived using IPCC Tier 3 methodology. Activity data provided by the CAA. Total consumptions of ATF and AS provided by DECC, via DUKES.
1A3a	Aircraft - domestic cruise	Aviation turbine fuel	Liquid	CO ₂	t/TJ	71.8	71.7	71.8	71.7	Carbon Factors Review 2004
1A3a	Aircraft - domestic cruise	Aviation turbine fuel	Liquid	N ₂ O	kg/TJ	2.3	2.3	2.3	2.3	Emissions from 1990 to the latest inventory year derived using IPCC Tier 3 methodology. Activity data provided by the CAA. Total consumptions of ATF and AS provided by DECC, via DUKES.

Other Detailed Methodological Descriptions **A3**

Category	Source Name	Activity Name	Fuel (CRF)	Gas	Unit	1990	2013	Max	Min	EF Method Reference
1A3a	Aircraft - domestic take-off and landing	Aviation spirit	Liquid	CO ₂	t/TJ	69.6	69.5	69.6	69.3	Carbon Factors Review 2004
1A3a	Aircraft - domestic take-off and landing	Aviation spirit	Liquid	CH ₄	kg/TJ	43.7	18.3	43.7	16.6	Emissions from 1990 to the latest inventory year derived using IPCC Tier 3 methodology. Activity data provided by the CAA. Total consumptions of ATF and AS provided by DECC, via DUKES.
1A3a	Aircraft - domestic take-off and landing	Aviation spirit	Liquid	N ₂ O	kg/TJ	2.2	2.2	2.2	2.2	Emissions from 1990 to the latest inventory year derived using IPCC Tier 3 methodology. Activity data provided by the CAA. Total consumptions of ATF and AS provided by DECC, via DUKES.
1A3a	Aircraft - domestic take-off and landing	Aviation turbine fuel	Liquid	CO ₂	t/TJ	71.8	71.7	71.8	71.7	Carbon Factors Review 2004
1A3a	Aircraft - domestic take-off and landing	Aviation turbine fuel	Liquid	CH ₄	kg/TJ	19.4	3.5	19.4	3.3	Emissions from 1990 to the latest inventory year derived using IPCC Tier 3 methodology. Activity data provided by the CAA. Total consumptions of ATF and AS provided by DECC, via DUKES.
1A3a	Aircraft - domestic take-off and landing	Aviation turbine fuel	Liquid	N ₂ O	kg/TJ	2.3	2.3	2.3	2.3	Emissions from 1990 to the latest inventory year derived using IPCC Tier 3 methodology. Activity data provided by the CAA. Total consumptions of ATF and AS provided by DECC, via DUKES.
1A3a	Aircraft between UK and CDs - Cruise	Aviation spirit	Liquid	CO ₂	t/TJ	69.6	69.5	69.6	69.3	Carbon Factors Review 2004
1A3a	Aircraft between UK and CDs - Cruise	Aviation spirit	Liquid	N ₂ O	kg/TJ	2.2	2.2	2.2	2.2	Emissions from 1990 to the latest inventory year derived using IPCC Tier 3 methodology. Activity data provided by the CAA. Total consumptions of ATF and AS provided by DECC, via DUKES.

Other Detailed Methodological Descriptions **A3**

Category	Source Name	Activity Name	Fuel (CRF)	Gas	Unit	1990	2013	Max	Min	EF Method Reference
1A3a	Aircraft between UK and CDs - Cruise	Aviation turbine fuel	Liquid	CO ₂	t/TJ	71.8	71.7	71.8	71.7	Carbon Factors Review 2004
1A3a	Aircraft between UK and CDs - Cruise	Aviation turbine fuel	Liquid	N ₂ O	kg/TJ	2.3	2.3	2.3	2.3	Emissions from 1990 to the latest inventory year derived using IPCC Tier 3 methodology. Activity data provided by the CAA. Total consumptions of ATF and AS provided by DECC, via DUKES.
1A3a	Aircraft between UK and CDs - TOL	Aviation spirit	Liquid	CO ₂	t/TJ	69.6	69.5	69.6	69.3	Carbon Factors Review 2004
1A3a	Aircraft between UK and CDs - TOL	Aviation spirit	Liquid	CH ₄	kg/TJ	43.2	26.8	47.9	18.2	Emissions from 1990 to the latest inventory year derived using IPCC Tier 3 methodology. Activity data provided by the CAA. Total consumptions of ATF and AS provided by DECC, via DUKES.
1A3a	Aircraft between UK and CDs - TOL	Aviation spirit	Liquid	N ₂ O	kg/TJ	2.2	2.2	2.2	2.2	Emissions from 1990 to the latest inventory year derived using IPCC Tier 3 methodology. Activity data provided by the CAA. Total consumptions of ATF and AS provided by DECC, via DUKES.
1A3a	Aircraft between UK and CDs - TOL	Aviation turbine fuel	Liquid	CO ₂	t/TJ	71.8	71.7	71.8	71.7	Carbon Factors Review 2004
1A3a	Aircraft between UK and CDs - TOL	Aviation turbine fuel	Liquid	CH ₄	kg/TJ	42.9	6.7	65.4	6.2	Emissions from 1990 to the latest inventory year derived using IPCC Tier 3 methodology. Activity data provided by the CAA. Total consumptions of ATF and AS provided by DECC, via DUKES.
1A3a	Aircraft between UK and CDs - TOL	Aviation turbine fuel	Liquid	N ₂ O	kg/TJ	2.3	2.3	2.3	2.3	Emissions from 1990 to the latest inventory year derived using IPCC Tier 3 methodology. Activity data provided by the CAA. Total consumptions of ATF and AS provided by DECC, via DUKES.

Other Detailed Methodological Descriptions **A3**

Category	Source Name	Activity Name	Fuel (CRF)	Gas	Unit	1990	2013	Max	Min	EF Method Reference
1A3a	Aircraft between UK and Gibraltar - Cruise	Aviation spirit	Liquid	CO ₂	t/TJ	69.6	69.5	69.6	69.3	Carbon Factors Review 2004
1A3a	Aircraft between UK and Gibraltar - Cruise	Aviation spirit	Liquid	N ₂ O	kg/TJ	0.0	0.0	2.2	2.2	Emissions from 1990 to the latest inventory year derived using IPCC Tier 3 methodology. Activity data provided by the CAA. Total consumptions of ATF and AS provided by DECC, via DUKES.
1A3a	Aircraft between UK and Gibraltar - Cruise	Aviation turbine fuel	Liquid	CO ₂	t/TJ	71.8	71.7	71.8	71.7	Carbon Factors Review 2004
1A3a	Aircraft between UK and Gibraltar - Cruise	Aviation turbine fuel	Liquid	N ₂ O	kg/TJ	2.3	2.3	2.3	2.3	Emissions from 1990 to the latest inventory year derived using IPCC Tier 3 methodology. Activity data provided by the CAA. Total consumptions of ATF and AS provided by DECC, via DUKES.
1A3a	Aircraft between UK and Gibraltar - TOL	Aviation spirit	Liquid	CO ₂	t/TJ	69.6	69.5	69.6	69.3	Carbon Factors Review 2004
1A3a	Aircraft between UK and Gibraltar - TOL	Aviation spirit	Liquid	CH ₄	kg/TJ	0.0	0.0	23.7	23.7	Emissions from 1990 to the latest inventory year derived using IPCC Tier 3 methodology. Activity data provided by the CAA. Total consumptions of ATF and AS provided by DECC, via DUKES.
1A3a	Aircraft between UK and Gibraltar - TOL	Aviation spirit	Liquid	N ₂ O	kg/TJ	0.0	0.0	2.2	2.2	Emissions from 1990 to the latest inventory year derived using IPCC Tier 3 methodology. Activity data provided by the CAA. Total consumptions of ATF and AS provided by DECC, via DUKES.
1A3a	Aircraft between UK and Gibraltar - TOL	Aviation turbine fuel	Liquid	CO ₂	t/TJ	71.8	71.7	71.8	71.7	Carbon Factors Review 2004

Other Detailed Methodological Descriptions **A3**

Category	Source Name	Activity Name	Fuel (CRF)	Gas	Unit	1990	2013	Max	Min	EF Method Reference
1A3a	Aircraft between UK and Gibraltar - TOL	Aviation turbine fuel	Liquid	CH ₄	kg/TJ	2.6	2.8	4.8	1.5	Emissions from 1990 to the latest inventory year derived using IPCC Tier 3 methodology. Activity data provided by the CAA. Total consumptions of ATF and AS provided by DECC, via DUKES.
1A3a	Aircraft between UK and Gibraltar - TOL	Aviation turbine fuel	Liquid	N ₂ O	kg/TJ	2.3	2.3	2.3	2.3	Emissions from 1990 to the latest inventory year derived using IPCC Tier 3 methodology. Activity data provided by the CAA. Total consumptions of ATF and AS provided by DECC, via DUKES.
1A3a	Aircraft between UK and other OTs (excl Gib.) - Cruise	Aviation spirit	Liquid	CO ₂	t/TJ	69.6	69.5	69.6	69.3	Carbon Factors Review 2004
1A3a	Aircraft between UK and other OTs (excl Gib.) - Cruise	Aviation turbine fuel	Liquid	CO ₂	t/TJ	71.8	71.7	71.8	71.7	Carbon Factors Review 2004
1A3a	Aircraft between UK and other OTs (excl Gib.) - Cruise	Aviation turbine fuel	Liquid	N ₂ O	kg/TJ	2.3	2.3	2.3	2.3	Emissions from 1990 to the latest inventory year derived using IPCC Tier 3 methodology. Activity data provided by the CAA. Total consumptions of ATF and AS provided by DECC, via DUKES.
1A3a	Aircraft between UK and other OTs (excl Gib.) - TOL	Aviation spirit	Liquid	CO ₂	t/TJ	69.6	69.5	69.6	69.3	Carbon Factors Review 2004
1A3a	Aircraft between UK and other OTs (excl Gib.) - TOL	Aviation turbine fuel	Liquid	CO ₂	t/TJ	71.8	71.7	71.8	71.7	Carbon Factors Review 2004
1A3a	Aircraft between UK and other OTs (excl Gib.) - TOL	Aviation turbine fuel	Liquid	CH ₄	kg/TJ	17.2	0.7	26.1	0.7	Emissions from 1990 to the latest inventory year derived using IPCC Tier 3 methodology. Activity data provided by the CAA. Total consumptions of ATF and AS provided by DECC, via DUKES.

Other Detailed Methodological Descriptions **A3**

Category	Source Name	Activity Name	Fuel (CRF)	Gas	Unit	1990	2013	Max	Min	EF Method Reference
1A3a	Aircraft between UK and other OTs (excl Gib.) - TOL	Aviation turbine fuel	Liquid	N ₂ O	kg/TJ	2.3	2.3	2.3	2.3	Emissions from 1990 to the latest inventory year derived using IPCC Tier 3 methodology. Activity data provided by the CAA. Total consumptions of ATF and AS provided by DECC, via DUKES.
1A3bi	Road transport - cars - motorway driving	DERV	Liquid	CO ₂	t/TJ	73.5	73.0	73.5	72.8	Carbon Factors Review 2004
1A3bi	Road transport - cars - motorway driving	DERV	Liquid	CH ₄	kg/TJ	4.0	0.4	4.0	0.4	COPERT 4 v10, DfT/TRL emission factors (Boulter et al, 2009) and EMEP/EEA Emission Inventory Guidebook (2009)
1A3bi	Road transport - cars - motorway driving	DERV	Liquid	N ₂ O	kg/TJ	0.0	2.2	2.2	0.0	EMEP/EEA Emissions Inventory Guidebook (2009)
1A3bi	Road transport - cars - motorway driving	Petrol	Liquid	CO ₂	t/TJ	70.2	70.0	70.2	70.0	Carbon Factors Review 2004
1A3bi	Road transport - cars - motorway driving	Petrol	Liquid	CH ₄	kg/TJ	17.5	1.8	17.5	1.8	COPERT 4 v10, DfT/TRL emission factors (Boulter et al, 2009) and EMEP/EEA Emission Inventory Guidebook (2009)
1A3bi	Road transport - cars - motorway driving	Petrol	Liquid	N ₂ O	kg/TJ	2.2	0.2	2.7	0.2	EMEP/EEA Emissions Inventory Guidebook (2009)
1A3bi	Road transport - cars - rural driving	DERV	Liquid	CO ₂	t/TJ	73.5	73.0	73.5	72.8	Carbon Factors Review 2004
1A3bi	Road transport - cars - rural driving	DERV	Liquid	CH ₄	kg/TJ	4.3	0.5	4.3	0.5	COPERT 4 v10, DfT/TRL emission factors (Boulter et al, 2009) and EMEP/EEA Emission Inventory Guidebook (2009)
1A3bi	Road transport - cars - rural driving	DERV	Liquid	N ₂ O	kg/TJ	0.0	2.3	2.3	0.0	EMEP/EEA Emissions Inventory Guidebook (2009)
1A3bi	Road transport - cars - rural driving	Petrol	Liquid	CO ₂	t/TJ	70.2	70.0	70.2	70.0	Carbon Factors Review 2004

Other Detailed Methodological Descriptions **A3**

Category	Source Name	Activity Name	Fuel (CRF)	Gas	Unit	1990	2013	Max	Min	EF Method Reference
1A3bi	Road transport - cars - rural driving	Petrol	Liquid	CH ₄	kg/TJ	12.8	1.5	12.8	1.5	COPERT 4 v10, DfT/TRL emission factors (Boulter et al, 2009) and EMEP/EEA Emission Inventory Guidebook (2009)
1A3bi	Road transport - cars - rural driving	Petrol	Liquid	N ₂ O	kg/TJ	2.3	0.3	4.2	0.3	EMEP/EEA Emissions Inventory Guidebook (2009)
1A3bi	Road transport - cars - urban driving	DERV	Liquid	CO ₂	t/TJ	73.5	73.0	73.5	72.8	Carbon Factors Review 2004
1A3bi	Road transport - cars - urban driving	DERV	Liquid	CH ₄	kg/TJ	5.9	0.8	5.9	0.8	COPERT 4 v10, DfT/TRL emission factors (Boulter et al, 2009) and EMEP/EEA Emission Inventory Guidebook (2009)
1A3bi	Road transport - cars - urban driving	DERV	Liquid	N ₂ O	kg/TJ	0.0	2.8	2.8	0.0	EMEP/EEA Emissions Inventory Guidebook (2009)
1A3bi	Road transport - cars - urban driving	Petrol	Liquid	CO ₂	t/TJ	70.2	70.0	70.2	70.0	Carbon Factors Review 2004
1A3bi	Road transport - cars - urban driving	Petrol	Liquid	CH ₄	kg/TJ	33.3	2.4	33.3	2.4	COPERT 4 v10, DfT/TRL emission factors (Boulter et al, 2009) and EMEP/EEA Emission Inventory Guidebook (2009)
1A3bi	Road transport - cars - urban driving	Petrol	Liquid	N ₂ O	kg/TJ	2.2	0.7	3.6	0.7	EMEP/EEA Emissions Inventory Guidebook (2009)
1A3bi	Road transport - cars non catalyst - motorway driving	Petrol	Liquid	CO ₂	t/TJ	70.2	70.0	70.2	70.0	Carbon Factors Review 2004
1A3bi	Road transport - cars non catalyst - rural driving	Petrol	Liquid	CO ₂	t/TJ	70.2	70.0	70.2	70.0	Carbon Factors Review 2004
1A3bi	Road transport - cars non catalyst - urban driving	Petrol	Liquid	CO ₂	t/TJ	70.2	70.0	70.2	70.0	Carbon Factors Review 2004
1A3bii	Road transport - LGVs - motorway driving	DERV	Liquid	CO ₂	t/TJ	73.5	73.0	73.5	72.8	Carbon Factors Review 2004

Category	Source Name	Activity Name	Fuel (CRF)	Gas	Unit	1990	2013	Max	Min	EF Method Reference
1A3bii	Road transport - LGVs - motorway driving	DERV	Liquid	CH ₄	kg/TJ	4.4	0.2	4.4	0.2	COPERT 4 v10, DfT/TRL emission factors (Boulter et al, 2009) and EMEP/EEA Emission Inventory Guidebook (2009)
1A3bii	Road transport - LGVs - motorway driving	DERV	Liquid	N ₂ O	kg/TJ	0.0	1.1	1.1	0.0	EMEP/EEA Emissions Inventory Guidebook (2009)
1A3bii	Road transport - LGVs - motorway driving	Petrol	Liquid	CO ₂	t/TJ	70.2	70.0	70.2	70.0	Carbon Factors Review 2004
1A3bii	Road transport - LGVs - motorway driving	Petrol	Liquid	CH ₄	kg/TJ	12.6	3.0	12.6	3.0	COPERT 4 v10, DfT/TRL emission factors (Boulter et al, 2009) and EMEP/EEA Emission Inventory Guidebook (2009)
1A3bii	Road transport - LGVs - motorway driving	Petrol	Liquid	N ₂ O	kg/TJ	1.6	0.8	1.9	0.8	EMEP/EEA Emissions Inventory Guidebook (2009)
1A3bii	Road transport - LGVs - rural driving	DERV	Liquid	CO ₂	t/TJ	73.5	73.0	73.5	72.8	Carbon Factors Review 2004
1A3bii	Road transport - LGVs - rural driving	DERV	Liquid	CH ₄	kg/TJ	3.1	0.2	3.1	0.2	COPERT 4 v10, DfT/TRL emission factors (Boulter et al, 2009) and EMEP/EEA Emission Inventory Guidebook (2009)
1A3bii	Road transport - LGVs - rural driving	DERV	Liquid	N ₂ O	kg/TJ	0.0	1.4	1.4	0.0	EMEP/EEA Emissions Inventory Guidebook (2009)
1A3bii	Road transport - LGVs - rural driving	Petrol	Liquid	CO ₂	t/TJ	70.2	70.0	70.2	70.0	Carbon Factors Review 2004
1A3bii	Road transport - LGVs - rural driving	Petrol	Liquid	CH ₄	kg/TJ	10.9	2.5	10.9	2.5	COPERT 4 v10, DfT/TRL emission factors (Boulter et al, 2009) and EMEP/EEA Emission Inventory Guidebook (2009)
1A3bii	Road transport - LGVs - rural driving	Petrol	Liquid	N ₂ O	kg/TJ	2.0	1.1	3.0	1.1	EMEP/EEA Emissions Inventory Guidebook (2009)
1A3bii	Road transport - LGVs - urban driving	DERV	Liquid	CO ₂	t/TJ	73.5	73.0	73.5	72.8	Carbon Factors Review 2004

Category	Source Name	Activity Name	Fuel (CRF)	Gas	Unit	1990	2013	Max	Min	EF Method Reference
1A3bii	Road transport - LGVs - urban driving	DERV	Liquid	CH ₄	kg/TJ	6.3	0.8	6.3	0.8	COPERT 4 v10, DfT/TRL emission factors (Boulter et al, 2009) and EMEP/EEA Emission Inventory Guidebook (2009)
1A3bii	Road transport - LGVs - urban driving	DERV	Liquid	N ₂ O	kg/TJ	0.0	2.1	2.1	0.0	EMEP/EEA Emissions Inventory Guidebook (2009)
1A3bii	Road transport - LGVs - urban driving	Petrol	Liquid	CO ₂	t/TJ	70.2	70.0	70.2	70.0	Carbon Factors Review 2004
1A3bii	Road transport - LGVs - urban driving	Petrol	Liquid	CH ₄	kg/TJ	32.3	5.5	32.3	5.5	COPERT 4 v10, DfT/TRL emission factors (Boulter et al, 2009) and EMEP/EEA Emission Inventory Guidebook (2009)
1A3bii	Road transport - LGVs - urban driving	Petrol	Liquid	N ₂ O	kg/TJ	2.1	1.5	3.3	1.5	EMEP/EEA Emissions Inventory Guidebook (2009)
1A3bii	Road transport - LGVs non catalyst - motorway driving	Petrol	Liquid	CO ₂	t/TJ	70.2	70.0	70.2	70.0	Carbon Factors Review 2004
1A3bii	Road transport - LGVs non catalyst - rural driving	Petrol	Liquid	CO ₂	t/TJ	70.2	70.0	70.2	70.0	Carbon Factors Review 2004
1A3bii	Road transport - LGVs non catalyst - urban driving	Petrol	Liquid	CO ₂	t/TJ	70.2	70.0	70.2	70.0	Carbon Factors Review 2004
1A3biii	Road transport - buses and coaches - motorway driving	DERV	Liquid	CO ₂	t/TJ	73.5	73.0	73.5	72.8	Carbon Factors Review 2004
1A3biii	Road transport - buses and coaches - motorway driving	DERV	Liquid	CH ₄	kg/TJ	18.6	1.5	18.6	1.5	COPERT 4 v10, DfT/TRL emission factors (Boulter et al, 2009) and EMEP/EEA Emission Inventory Guidebook (2009)
1A3biii	Road transport - buses and coaches - motorway driving	DERV	Liquid	N ₂ O	kg/TJ	3.6	1.9	3.6	0.5	EMEP/EEA Emissions Inventory Guidebook (2009)
1A3biii	Road transport - buses and coaches - rural driving	DERV	Liquid	CO ₂	t/TJ	73.5	73.0	73.5	72.8	Carbon Factors Review 2004

Category	Source Name	Activity Name	Fuel (CRF)	Gas	Unit	1990	2013	Max	Min	EF Method Reference
1A3biii	Road transport - buses and coaches - rural driving	DERV	Liquid	CH ₄	kg/TJ	24.1	2.1	24.1	2.1	COPERT 4 v10, DfT/TRL emission factors (Boulter et al, 2009) and EMEP/EEA Emission Inventory Guidebook (2009)
1A3biii	Road transport - buses and coaches - rural driving	DERV	Liquid	N ₂ O	kg/TJ	4.1	2.5	4.1	0.9	EMEP/EEA Emissions Inventory Guidebook (2009)
1A3biii	Road transport - buses and coaches - urban driving	DERV	Liquid	CO ₂	t/TJ	73.5	73.0	73.5	72.8	Carbon Factors Review 2004
1A3biii	Road transport - buses and coaches - urban driving	DERV	Liquid	CH ₄	kg/TJ	32.9	2.2	32.9	2.2	COPERT 4 v10, DfT/TRL emission factors (Boulter et al, 2009) and EMEP/EEA Emission Inventory Guidebook (2009)
1A3biii	Road transport - buses and coaches - urban driving	DERV	Liquid	N ₂ O	kg/TJ	2.6	1.5	2.6	0.6	EMEP/EEA Emissions Inventory Guidebook (2009)
1A3biii	Road transport - HGV articulated - motorway driving	DERV	Liquid	CO ₂	t/TJ	73.5	73.0	73.5	72.8	Carbon Factors Review 2004
1A3biii	Road transport - HGV articulated - motorway driving	DERV	Liquid	CH ₄	kg/TJ	10.3	0.8	10.3	0.8	COPERT 4 v10, DfT/TRL emission factors (Boulter et al, 2009) and EMEP/EEA Emission Inventory Guidebook (2009)
1A3biii	Road transport - HGV articulated - motorway driving	DERV	Liquid	N ₂ O	kg/TJ	2.0	3.0	3.0	0.7	EMEP/EEA Emissions Inventory Guidebook (2009)
1A3biii	Road transport - HGV articulated - rural driving	DERV	Liquid	CO ₂	t/TJ	73.5	73.0	73.5	72.8	Carbon Factors Review 2004
1A3biii	Road transport - HGV articulated - rural driving	DERV	Liquid	CH ₄	kg/TJ	12.0	0.9	12.0	0.9	COPERT 4 v10, DfT/TRL emission factors (Boulter et al, 2009) and EMEP/EEA Emission Inventory Guidebook (2009)
1A3biii	Road transport - HGV articulated - rural driving	DERV	Liquid	N ₂ O	kg/TJ	2.1	3.7	3.7	0.9	EMEP/EEA Emissions Inventory Guidebook (2009)

Other Detailed Methodological Descriptions **A3**

Category	Source Name	Activity Name	Fuel (CRF)	Gas	Unit	1990	2013	Max	Min	EF Method Reference
1A3biii	Road transport - HGV articulated - urban driving	DERV	Liquid	CO ₂	t/TJ	73.5	73.0	73.5	72.8	Carbon Factors Review 2004
1A3biii	Road transport - HGV articulated - urban driving	DERV	Liquid	CH ₄	kg/TJ	20.2	0.9	20.2	0.9	COPERT 4 v10, DfT/TRL emission factors (Boulter et al, 2009) and EMEP/EEA Emission Inventory Guidebook (2009)
1A3biii	Road transport - HGV articulated - urban driving	DERV	Liquid	N ₂ O	kg/TJ	1.6	2.2	2.2	0.8	EMEP/EEA Emissions Inventory Guidebook (2009)
1A3biii	Road transport - HGV rigid - motorway driving	DERV	Liquid	CO ₂	t/TJ	73.5	73.0	73.5	72.8	Carbon Factors Review 2004
1A3biii	Road transport - HGV rigid - motorway driving	DERV	Liquid	CH ₄	kg/TJ	4.4	0.5	4.4	0.5	COPERT 4 v10, DfT/TRL emission factors (Boulter et al, 2009) and EMEP/EEA Emission Inventory Guidebook (2009)
1A3biii	Road transport - HGV rigid - motorway driving	DERV	Liquid	N ₂ O	kg/TJ	3.0	1.8	3.0	0.5	EMEP/EEA Emissions Inventory Guidebook (2009)
1A3biii	Road transport - HGV rigid - rural driving	DERV	Liquid	CO ₂	t/TJ	73.5	73.0	73.5	72.8	Carbon Factors Review 2004
1A3biii	Road transport - HGV rigid - rural driving	DERV	Liquid	CH ₄	kg/TJ	5.3	0.7	5.3	0.7	COPERT 4 v10, DfT/TRL emission factors (Boulter et al, 2009) and EMEP/EEA Emission Inventory Guidebook (2009)
1A3biii	Road transport - HGV rigid - rural driving	DERV	Liquid	N ₂ O	kg/TJ	3.2	2.3	3.2	0.7	EMEP/EEA Emissions Inventory Guidebook (2009)
1A3biii	Road transport - HGV rigid - urban driving	DERV	Liquid	CO ₂	t/TJ	73.5	73.0	73.5	72.8	Carbon Factors Review 2004
1A3biii	Road transport - HGV rigid - urban driving	DERV	Liquid	CH ₄	kg/TJ	15.8	1.3	15.8	1.3	COPERT 4 v10, DfT/TRL emission factors (Boulter et al, 2009) and EMEP/EEA Emission Inventory Guidebook (2009)
1A3biii	Road transport - HGV rigid - urban driving	DERV	Liquid	N ₂ O	kg/TJ	2.6	1.4	2.6	0.6	EMEP/EEA Emissions Inventory Guidebook (2009)

Other Detailed Methodological Descriptions **A3**

Category	Source Name	Activity Name	Fuel (CRF)	Gas	Unit	1990	2013	Max	Min	EF Method Reference
1A3biv	Road transport - mopeds (<50cc 2st) - urban driving	Petrol	Liquid	CO ₂	t/TJ	70.2	70.0	70.2	70.0	Carbon Factors Review 2004
1A3biv	Road transport - mopeds (<50cc 2st) - urban driving	Petrol	Liquid	CH ₄	kg/TJ	196.2	64.5	196.2	64.5	COPERT 4 v10, DfT/TRL emission factors (Boulter et al, 2009) and EMEP/EEA Emission Inventory Guidebook (2009)
1A3biv	Road transport - mopeds (<50cc 2st) - urban driving	Petrol	Liquid	N ₂ O	kg/TJ	0.9	1.9	1.9	0.9	EMEP/EEA Emissions Inventory Guidebook (2009)
1A3biv	Road transport - motorcycle (>50cc 2st) - rural driving	Petrol	Liquid	CO ₂	t/TJ	70.2	70.0	70.2	70.0	Carbon Factors Review 2004
1A3biv	Road transport - motorcycle (>50cc 2st) - urban driving	Petrol	Liquid	CO ₂	t/TJ	70.2	70.0	70.2	70.0	Carbon Factors Review 2004
1A3biv	Road transport - motorcycle (>50cc 2st) - urban driving	Petrol	Liquid	CH ₄	kg/TJ	107.0	25.8	107.1	25.8	COPERT 4 v10, DfT/TRL emission factors (Boulter et al, 2009) and EMEP/EEA Emission Inventory Guidebook (2009)
1A3biv	Road transport - motorcycle (>50cc 2st) - urban driving	Petrol	Liquid	N ₂ O	kg/TJ	1.4	1.6	1.6	1.4	EMEP/EEA Emissions Inventory Guidebook (2009)
1A3biv	Road transport - motorcycle (>50cc 4st) - motorway driving	Petrol	Liquid	CO ₂	t/TJ	70.2	70.0	70.2	70.0	Carbon Factors Review 2004
1A3biv	Road transport - motorcycle (>50cc 4st) - motorway driving	Petrol	Liquid	CH ₄	kg/TJ	87.1	42.9	87.1	42.9	COPERT 4 v10, DfT/TRL emission factors (Boulter et al, 2009) and EMEP/EEA Emission Inventory Guidebook (2009)
1A3biv	Road transport - motorcycle (>50cc 4st) - motorway driving	Petrol	Liquid	N ₂ O	kg/TJ	0.9	1.0	1.0	0.9	EMEP/EEA Emissions Inventory Guidebook (2009)
1A3biv	Road transport - motorcycle (>50cc 4st) - rural driving	Petrol	Liquid	CO ₂	t/TJ	70.2	70.0	70.2	70.0	Carbon Factors Review 2004

Category	Source Name	Activity Name	Fuel (CRF)	Gas	Unit	1990	2013	Max	Min	EF Method Reference
1A3bv	Road transport - motorcycle (>50cc 4st) - rural driving	Petrol	Liquid	CH ₄	kg/TJ	139.4	60.0	139.4	60.0	COPERT 4 v10, DfT/TRL emission factors (Boulter et al, 2009) and EMEP/EEA Emission Inventory Guidebook (2009)
1A3bv	Road transport - motorcycle (>50cc 4st) - rural driving	Petrol	Liquid	N ₂ O	kg/TJ	1.4	1.4	1.4	1.3	EMEP/EEA Emissions Inventory Guidebook (2009)
1A3bv	Road transport - motorcycle (>50cc 4st) - urban driving	Petrol	Liquid	CO ₂	t/TJ	70.2	70.0	70.2	70.0	Carbon Factors Review 2004
1A3bv	Road transport - motorcycle (>50cc 4st) - urban driving	Petrol	Liquid	CH ₄	kg/TJ	129.5	69.3	129.5	69.3	COPERT 4 v10, DfT/TRL emission factors (Boulter et al, 2009) and EMEP/EEA Emission Inventory Guidebook (2009)
1A3bv	Road transport - motorcycle (>50cc 4st) - urban driving	Petrol	Liquid	N ₂ O	kg/TJ	1.3	1.3	1.3	1.2	EMEP/EEA Emissions Inventory Guidebook (2009)
1A3bv	Road transport - all vehicles LPG use	LPG	Liquid	CO ₂	t/TJ	64.7	64.7	64.7	64.7	Carbon Factors Review 2004
1A3bv	Road transport - all vehicles LPG use	LPG	Liquid	CH ₄	kg/TJ	0.0	4.4	17.2	4.4	DfT/TRL emission factors (Boulter et al, 2009)
1A3bv	Road transport - all vehicles LPG use	LPG	Liquid	N ₂ O	kg/TJ	0.0	1.4	3.5	1.4	DfT/TRL emission factors (Boulter et al, 2009)
1A3c	Rail - coal	Coal	Solid	CO ₂	t/TJ	95.0	95.0	95.0	95.0	Based on Fynes & Sage, 1994
1A3c	Rail - coal	Coal	Solid	CH ₄	kg/TJ	95.7	99.9	104.0	86.6	Emission factors based on AP42 figures for hand-fed stokers
1A3c	Rail - coal	Coal	Solid	N ₂ O	kg/TJ	0.8	0.8	0.8	0.7	Emission factors based on AP42 figures for hand-fed stokers
1A3c	Railways - freight	Gas oil	Liquid	CO ₂	t/TJ	73.9	74.1	74.2	73.5	Carbon Factors Review 2004
1A3c	Railways - freight	Gas oil	Liquid	CH ₄	kg/TJ	5.5	5.5	6.2	2.8	LRC emission factors & Pers comm, John Hawkins, First Great Western Railway
1A3c	Railways - freight	Gas oil	Liquid	N ₂ O	kg/TJ	0.6	0.6	0.6	0.6	N ₂ O EF taken from the 2009 EMEP / EEA Guidebook

Other Detailed Methodological Descriptions **A3**

Category	Source Name	Activity Name	Fuel (CRF)	Gas	Unit	1990	2013	Max	Min	EF Method Reference
1A3c	Railways - intercity	Gas oil	Liquid	CO ₂	t/TJ	73.9	74.1	74.2	73.5	Carbon Factors Review 2004
1A3c	Railways - intercity	Gas oil	Liquid	CH ₄	kg/TJ	3.7	2.9	6.4	2.5	Inter-city rail Efs derived from DfT Rail Emissions Model for train classes included in REM. Other Efs are as used previously, which are inferred from LRC emission factors
1A3c	Railways - intercity	Gas oil	Liquid	N ₂ O	kg/TJ	0.6	0.6	0.6	0.6	N ₂ O EF taken from the 2009 EMEP / EEA Guidebook
1A3c	Railways - regional	Gas oil	Liquid	CO ₂	t/TJ	73.9	74.1	74.2	73.5	Carbon Factors Review 2004
1A3c	Railways - regional	Gas oil	Liquid	CH ₄	kg/TJ	5.9	1.7	5.9	1.7	Regional passenger rail Efs derived from DfT Rail Emissions Model
1A3c	Railways - regional	Gas oil	Liquid	N ₂ O	kg/TJ	0.6	0.6	0.6	0.6	N ₂ O EF taken from the 2009 EMEP / EEA Guidebook
1A3d	Inland goods-carrying vessels	DERV	Liquid	CO ₂	t/TJ	73.5	73.0	73.5	72.8	Carbon Factors Review 2004
1A3d	Inland goods-carrying vessels	Gas oil	Liquid	CO ₂	t/TJ	73.9	74.1	74.2	73.5	Carbon Factors Review 2004
1A3d	Inland goods-carrying vessels	Gas oil	Liquid	CH ₄	kg/TJ	1.2	1.2	1.2	1.2	Greenhouse Gas Emissions from Inland Waterways and Recreational Craft in the UK, Task 25 of the 2010 DA / UK GHG Inventory Improvement Programme
1A3d	Inland goods-carrying vessels	Gas oil	Liquid	N ₂ O	kg/TJ	1.9	1.9	1.9	1.8	Greenhouse Gas Emissions from Inland Waterways and Recreational Craft in the UK, Task 25 of the 2010 DA / UK GHG Inventory Improvement Programme
1A3d	Inland goods-carrying vessels	Petrol	Liquid	CO ₂	t/TJ	70.2	70.0	70.2	70.0	Carbon Factors Review 2004
1A3d	Motorboats / workboats (e.g. canal boats, dredgers, service boats, tourist boats, river boats)	DERV	Liquid	CO ₂	t/TJ	73.5	73.0	73.5	72.8	Carbon Factors Review 2004

Category	Source Name	Activity Name	Fuel (CRF)	Gas	Unit	1990	2013	Max	Min	EF Method Reference
1A3d	Motorboats / workboats (e.g. canal boats, dredgers, service boats, tourist boats, river boats)	DERV	Liquid	CH ₄	kg/TJ	1.2	1.2	1.2	1.2	Greenhouse Gas Emissions from Inland Waterways and Recreational Craft in the UK, Task 25 of the 2010 DA / UK GHG Inventory Improvement Programme
1A3d	Motorboats / workboats (e.g. canal boats, dredgers, service boats, tourist boats, river boats)	DERV	Liquid	N ₂ O	kg/TJ	1.9	1.8	1.9	1.8	Greenhouse Gas Emissions from Inland Waterways and Recreational Craft in the UK, Task 25 of the 2010 DA / UK GHG Inventory Improvement Programme
1A3d	Motorboats / workboats (e.g. canal boats, dredgers, service boats, tourist boats, river boats)	Gas oil	Liquid	CO ₂	t/TJ	73.9	74.1	74.2	73.5	Carbon Factors Review 2004
1A3d	Motorboats / workboats (e.g. canal boats, dredgers, service boats, tourist boats, river boats)	Gas oil	Liquid	CH ₄	kg/TJ	1.2	1.2	1.2	1.2	Greenhouse Gas Emissions from Inland Waterways and Recreational Craft in the UK, Task 25 of the 2010 DA / UK GHG Inventory Improvement Programme
1A3d	Motorboats / workboats (e.g. canal boats, dredgers, service boats, tourist boats, river boats)	Gas oil	Liquid	N ₂ O	kg/TJ	1.9	1.9	1.9	1.8	Greenhouse Gas Emissions from Inland Waterways and Recreational Craft in the UK, Task 25 of the 2010 DA / UK GHG Inventory Improvement Programme
1A3d	Motorboats / workboats (e.g. canal boats, dredgers, service boats, tourist boats, river boats)	Petrol	Liquid	CO ₂	t/TJ	70.2	70.0	70.2	70.0	Carbon Factors Review 2004
1A3d	Motorboats / workboats (e.g. canal boats, dredgers, service boats, tourist boats, river boats)	Petrol	Liquid	CH ₄	kg/TJ	38.1	38.0	38.1	38.0	Greenhouse Gas Emissions from Inland Waterways and Recreational Craft in the UK, Task 25 of the 2010 DA / UK GHG Inventory Improvement Programme

Other Detailed Methodological Descriptions **A3**

Category	Source Name	Activity Name	Fuel (CRF)	Gas	Unit	1990	2013	Max	Min	EF Method Reference
1A3d	Motorboats / workboats (e.g. canal boats, dredgers, service boats, tourist boats, river boats)	Petrol	Liquid	N ₂ O	kg/TJ	1.8	1.8	1.8	1.8	Greenhouse Gas Emissions from Inland Waterways and Recreational Craft in the UK, Task 25 of the 2010 DA / UK GHG Inventory Improvement Programme
1A3d	Personal watercraft e.g. jet ski	DERV	Liquid	CO ₂	t/TJ	73.5	73.0	73.5	72.8	Carbon Factors Review 2004
1A3d	Personal watercraft e.g. jet ski	Gas oil	Liquid	CO ₂	t/TJ	73.9	74.1	74.2	73.5	Carbon Factors Review 2004
1A3d	Personal watercraft e.g. jet ski	Petrol	Liquid	CO ₂	t/TJ	70.2	70.0	70.2	70.0	Carbon Factors Review 2004
1A3d	Personal watercraft e.g. jet ski	Petrol	Liquid	CH ₄	kg/TJ	38.1	38.0	38.1	38.0	Greenhouse Gas Emissions from Inland Waterways and Recreational Craft in the UK, Task 25 of the 2010 DA / UK GHG Inventory Improvement Programme
1A3d	Personal watercraft e.g. jet ski	Petrol	Liquid	N ₂ O	kg/TJ	1.8	1.8	1.8	1.8	Greenhouse Gas Emissions from Inland Waterways and Recreational Craft in the UK, Task 25 of the 2010 DA / UK GHG Inventory Improvement Programme
1A3d	Sailing boats with auxiliary engines	DERV	Liquid	CO ₂	t/TJ	73.5	73.0	73.5	72.8	Carbon Factors Review 2004
1A3d	Sailing boats with auxiliary engines	DERV	Liquid	CH ₄	kg/TJ	1.2	1.2	1.2	1.2	Greenhouse Gas Emissions from Inland Waterways and Recreational Craft in the UK, Task 25 of the 2010 DA / UK GHG Inventory Improvement Programme
1A3d	Sailing boats with auxiliary engines	DERV	Liquid	N ₂ O	kg/TJ	1.9	1.8	1.9	1.8	Greenhouse Gas Emissions from Inland Waterways and Recreational Craft in the UK, Task 25 of the 2010 DA / UK GHG Inventory Improvement Programme
1A3d	Sailing boats with auxiliary engines	Gas oil	Liquid	CO ₂	t/TJ	73.9	74.1	74.2	73.5	Carbon Factors Review 2004

Other Detailed Methodological Descriptions **A3**

Category	Source Name	Activity Name	Fuel (CRF)	Gas	Unit	1990	2013	Max	Min	EF Method Reference
1A3d	Sailing boats with auxiliary engines	Petrol	Liquid	CO ₂	t/TJ	70.2	70.0	70.2	70.0	Carbon Factors Review 2004
1A3d	Shipping - coastal	Fuel oil	Liquid	CO ₂	t/TJ	77.5	78.4	78.4	77.3	Carbon Factors Review 2004
1A3d	Shipping - coastal	Fuel oil	Liquid	CH ₄	kg/TJ	1.2	1.2	1.2	1.2	EMEP/EEA Guidebook, 2006
1A3d	Shipping - coastal	Fuel oil	Liquid	N ₂ O	kg/TJ	1.9	1.9	2.0	1.9	EMEP/EEA Guidebook, 2006
1A3d	Shipping - coastal	Gas oil	Liquid	CO ₂	t/TJ	73.9	74.1	74.2	73.5	Carbon Factors Review 2004
1A3d	Shipping - coastal	Gas oil	Liquid	CH ₄	kg/TJ	1.2	1.2	1.2	1.2	EMEP/EEA Guidebook, 2006
1A3d	Shipping - coastal	Gas oil	Liquid	N ₂ O	kg/TJ	1.9	1.9	1.9	1.8	EMEP/EEA Guidebook, 2006
1A3d	Shipping between UK and Gibraltar	Fuel oil	Liquid	CO ₂	t/TJ	77.5	78.4	78.4	77.3	Carbon Factors Review 2004
1A3d	Shipping between UK and Gibraltar	Fuel oil	Liquid	CH ₄	kg/TJ	1.2	1.2	1.2	1.2	EMEP/EEA Guidebook, 2006
1A3d	Shipping between UK and Gibraltar	Fuel oil	Liquid	N ₂ O	kg/TJ	1.9	1.9	2.0	1.9	EMEP/EEA Guidebook, 2006
1A3d	Shipping between UK and OTs (excl. Gib)	Fuel oil	Liquid	CO ₂	t/TJ	77.5	78.4	78.4	77.3	Carbon Factors Review 2004
1A3d	Shipping between UK and OTs (excl. Gib)	Fuel oil	Liquid	CH ₄	kg/TJ	1.2	1.2	1.2	1.2	EMEP/EEA Guidebook, 2006
1A3d	Shipping between UK and OTs (excl. Gib)	Fuel oil	Liquid	N ₂ O	kg/TJ	1.9	1.9	2.0	1.9	EMEP/EEA Guidebook, 2006
1A3eii	Aircraft - support vehicles	Gas oil	All	CO ₂	t/TJ	73.9	74.1	74.2	73.5	Carbon Factors Review 2004
1A3eii	Aircraft - support vehicles	Gas oil	All	CH ₄	kg/TJ	3.9	3.1	3.9	3.1	Developed based on T3 approach considering emission limits on total hydrocarbons

Category	Source Name	Activity Name	Fuel (CRF)	Gas	Unit	1990	2013	Max	Min	EF Method Reference
1A3eii	Aircraft - support vehicles	Gas oil	All	N ₂ O	kg/TJ	30.7	30.8	30.8	30.5	EMEP/EEA Guidebook, 2006
1A4ai	Miscellaneous industrial/commercial combustion	Coal	Solid	CO ₂	t/TJ	95.0	95.0	95.0	95.0	Based on Fynes & Sage, 1994
1A4ai	Miscellaneous industrial/commercial combustion	Coal	Solid	CH ₄	kg/TJ	0.4	0.4	0.5	0.4	Brain et al, 1994
1A4ai	Miscellaneous industrial/commercial combustion	Coal	Solid	N ₂ O	kg/TJ	5.6	5.9	6.1	5.1	Based on Fynes & Sage, 1994
1A4ai	Miscellaneous industrial/commercial combustion	Coke	Solid	CO ₂	t/TJ	114.3	111.3	114.3	101.8	Ricardo-AEA estimate based on carbon balance for integrated steelwork processes (2014)
1A4ai	Miscellaneous industrial/commercial combustion	Coke	Solid	CH ₄	kg/TJ	0.4	0.4	0.4	0.4	Brain et al, 1994
1A4ai	Miscellaneous industrial/commercial combustion	Coke	Solid	N ₂ O	kg/TJ	5.6	5.3	5.6	5.3	Based on Fynes & Sage, 1994
1A4ai	Miscellaneous industrial/commercial combustion	Fuel oil	Liquid	CO ₂	t/TJ	77.5	78.4	78.4	77.3	Carbon Factors Review 2004
1A4ai	Miscellaneous industrial/commercial combustion	Fuel oil	Liquid	CH ₄	kg/TJ	9.9	9.9	10.0	9.8	IPCC 2006 Guidelines

Category	Source Name	Activity Name	Fuel (CRF)	Gas	Unit	1990	2013	Max	Min	EF Method Reference
1A4ai	Miscellaneous industrial/commercial combustion	Fuel oil	Liquid	N ₂ O	kg/TJ	0.6	0.6	0.6	0.6	IPCC 2006 Guidelines
1A4ai	Miscellaneous industrial/commercial combustion	Gas oil	Liquid	CO ₂	t/TJ	73.9	74.1	74.2	73.5	Carbon Factors Review 2004
1A4ai	Miscellaneous industrial/commercial combustion	Gas oil	Liquid	CH ₄	kg/TJ	9.9	9.9	9.9	9.8	IPCC 2006 Guidelines
1A4ai	Miscellaneous industrial/commercial combustion	Gas oil	Liquid	N ₂ O	kg/TJ	0.6	0.6	0.6	0.6	IPCC 2006 Guidelines
1A4ai	Miscellaneous industrial/commercial combustion	Natural gas	Gaseous	CO ₂	t/TJ	56.0	56.9	57.2	56.0	Natural gas carbon emission factor is derived from Gas Network Operator data on gas composition within each Local Distribution Zone, prepared for use by operators within the EU ETS. (DECC, 2014)
1A4ai	Miscellaneous industrial/commercial combustion	Natural gas	Gaseous	CH ₄	kg/TJ	5.0	5.0	5.0	5.0	IPCC 2006 Guidelines
1A4ai	Miscellaneous industrial/commercial combustion	Natural gas	Gaseous	N ₂ O	kg/TJ	0.1	0.1	0.1	0.1	IPCC 2006 Guidelines
1A4ai	Miscellaneous industrial/commercial combustion	SSF	Solid	CO ₂	t/TJ	110.5	102.3	110.5	93.6	British Coal (1989)

Other Detailed Methodological Descriptions **A3**

Category	Source Name	Activity Name	Fuel (CRF)	Gas	Unit	1990	2013	Max	Min	EF Method Reference
1A4ai	Miscellaneous industrial/commercial combustion	SSF	Solid	CH ₄	kg/TJ	0.4	0.4	0.4	0.4	Brain et al, 1994
1A4ai	Miscellaneous industrial/commercial combustion	SSF	Solid	N ₂ O	kg/TJ	5.9	5.5	5.9	5.0	Based on Fynes & Sage, 1994
1A4ai	Miscellaneous industrial/commercial combustion	Town gas	Solid	CO ₂	t/TJ	61.7	61.7	61.7	61.7	Ricardo-AEA estimate based on carbon balance for town gas manufacture (2014)
1A4ai	Miscellaneous industrial/commercial combustion	Town gas	Solid	CH ₄	kg/TJ	5.0	5.0	5.0	5.0	IPCC 2006 Guidelines
1A4ai	Public sector combustion	Burning oil	Liquid	CO ₂	t/TJ	71.8	71.7	71.8	71.7	Carbon Factors Review 2004
1A4ai	Public sector combustion	Burning oil	Liquid	CH ₄	kg/TJ	10.0	10.0	10.0	10.0	IPCC 2006 Guidelines
1A4ai	Public sector combustion	Burning oil	Liquid	N ₂ O	kg/TJ	0.6	0.6	0.6	0.6	IPCC 2006 Guidelines
1A4ai	Public sector combustion	Coal	Solid	CO ₂	t/TJ	95.0	95.0	95.0	95.0	Based on Fynes & Sage, 1994
1A4ai	Public sector combustion	Coal	Solid	CH ₄	kg/TJ	0.4	0.4	0.5	0.4	Brain et al., 1994
1A4ai	Public sector combustion	Coal	Solid	N ₂ O	kg/TJ	5.6	5.9	6.1	5.1	Based on Fynes & Sage, 1994
1A4ai	Public sector combustion	Coke	Solid	CO ₂	t/TJ	114.3	111.3	114.3	101.8	Ricardo-AEA estimate based on carbon balance for integrated steelwork processes (2014)
1A4ai	Public sector combustion	Coke	Solid	CH ₄	kg/TJ	0.4	0.4	0.4	0.4	Brain et al, 1994
1A4ai	Public sector combustion	Coke	Solid	N ₂ O	kg/TJ	5.6	5.3	5.6	5.3	Based on Fynes & Sage, 1994
1A4ai	Public sector combustion	Fuel oil	Liquid	CO ₂	t/TJ	77.5	78.4	78.4	77.3	Carbon Factors Review 2004

Other Detailed Methodological Descriptions **A3**

Category	Source Name	Activity Name	Fuel (CRF)	Gas	Unit	1990	2013	Max	Min	EF Method Reference
1A4ai	Public sector combustion	Fuel oil	Liquid	CH ₄	kg/TJ	9.9	9.9	10.0	9.8	IPCC 2006 Guidelines
1A4ai	Public sector combustion	Fuel oil	Liquid	N ₂ O	kg/TJ	0.6	0.6	0.6	0.6	IPCC 2006 Guidelines
1A4ai	Public sector combustion	Gas oil	Liquid	CO ₂	t/TJ	73.9	74.1	74.2	73.5	Carbon Factors Review 2004
1A4ai	Public sector combustion	Gas oil	Liquid	CH ₄	kg/TJ	9.9	9.9	9.9	9.8	IPCC 2006 Guidelines
1A4ai	Public sector combustion	Gas oil	Liquid	N ₂ O	kg/TJ	0.6	0.6	0.6	0.6	IPCC 2006 Guidelines
1A4ai	Public sector combustion	Natural gas	Gaseous	CO ₂	t/TJ	56.0	56.9	57.2	56.0	Natural gas carbon emission factor is derived from Gas Network Operator data on gas composition within each Local Distribution Zone, prepared for use by operators within the EU ETS. (DECC, 2014)
1A4ai	Public sector combustion	Natural gas	Gaseous	CH ₄	kg/TJ	5.0	5.0	5.0	5.0	IPCC 2006 Guidelines
1A4ai	Public sector combustion	Natural gas	Gaseous	N ₂ O	kg/TJ	0.1	0.1	0.1	0.1	IPCC 2006 Guidelines
1A4ai	Public sector combustion	Town gas	Solid	CO ₂	t/TJ	61.7	61.7	61.7	61.7	Ricardo-AEA estimate based on carbon balance for town gas manufacture (2014)
1A4ai	Public sector combustion	Town gas	Solid	CH ₄	kg/TJ	5.0	5.0	5.0	5.0	IPCC 2006 Guidelines
1A4ai	Public sector combustion	Town gas	Solid	N ₂ O	kg/TJ	0.1	0.1	0.1	0.1	IPCC 2006 Guidelines
1A4ai	Railways - stationary combustion	Burning oil	Liquid	CO ₂	t/TJ	71.8	71.7	71.8	71.7	Carbon Factors Review 2004
1A4ai	Railways - stationary combustion	Burning oil	Liquid	CH ₄	kg/TJ	10.0	10.0	10.0	10.0	IPCC 2006 Guidelines
1A4ai	Railways - stationary combustion	Burning oil	Liquid	N ₂ O	kg/TJ	0.6	0.6	0.6	0.6	IPCC 2006 Guidelines
1A4ai	Railways - stationary combustion	Coal	Solid	CO ₂	t/TJ	95.0	95.0	95.0	95.0	Based on Fynes & Sage, 1994

Category	Source Name	Activity Name	Fuel (CRF)	Gas	Unit	1990	2013	Max	Min	EF Method Reference
1A4ai	Railways - stationary combustion	Coal	Solid	CH ₄	kg/TJ	0.4	0.4	0.5	0.4	Brain et al, 1994
1A4ai	Railways - stationary combustion	Coal	Solid	N ₂ O	kg/TJ	5.6	5.9	6.1	5.1	Based on Fynes & Sage, 1994
1A4ai	Railways - stationary combustion	Coke	Solid	CO ₂	t/TJ	114.3	111.3	114.3	101.8	Ricardo-AEA estimate based on carbon balance for integrated steelwork processes (2014)
1A4ai	Railways - stationary combustion	Coke	Solid	CH ₄	kg/TJ	0.4	0.4	0.4	0.4	Brain et al, 1994
1A4ai	Railways - stationary combustion	Coke	Solid	N ₂ O	kg/TJ	5.6	5.3	5.6	5.3	Based on Fynes & Sage, 1994
1A4ai	Railways - stationary combustion	Fuel oil	Liquid	CO ₂	t/TJ	77.5	78.4	78.4	77.3	Carbon Factors Review 2004
1A4ai	Railways - stationary combustion	Fuel oil	Liquid	CH ₄	kg/TJ	9.9	9.9	10.0	9.8	IPCC 2006 Guidelines
1A4ai	Railways - stationary combustion	Fuel oil	Liquid	N ₂ O	kg/TJ	0.6	0.6	0.6	0.6	IPCC 2006 Guidelines
1A4ai	Railways - stationary combustion	Natural gas	Gaseous	CO ₂	t/TJ	56.0	56.9	57.2	56.0	Natural gas carbon emission factor is derived from Gas Network Operator data on gas composition within each Local Distribution Zone, prepared for use by operators within the EU ETS. (DECC, 2014)
1A4ai	Railways - stationary combustion	Natural gas	Gaseous	CH ₄	kg/TJ	5.0	5.0	5.0	5.0	IPCC 2006 Guidelines
1A4ai	Railways - stationary combustion	Natural gas	Gaseous	N ₂ O	kg/TJ	0.1	0.1	0.1	0.1	IPCC 2006 Guidelines
1A4bi	Domestic combustion	Anthracite	Solid	CO ₂	t/TJ	98.7	98.7	98.7	98.7	Based on Fynes & Sage, 1994

Category	Source Name	Activity Name	Fuel (CRF)	Gas	Unit	1990	2013	Max	Min	EF Method Reference
1A4bi	Domestic combustion	Anthracite	Solid	CH ₄	kg/TJ	62.7	61.3	62.8	60.6	Brain et al, 1994
1A4bi	Domestic combustion	Anthracite	Solid	N ₂ O	kg/TJ	4.5	4.4	4.5	4.3	Based on Fynes & Sage, 1994
1A4bi	Domestic combustion	Burning oil	Liquid	CO ₂	t/TJ	71.8	71.7	71.8	71.7	Carbon Factors Review 2004
1A4bi	Domestic combustion	Burning oil	Liquid	CH ₄	kg/TJ	9.7	9.7	9.7	9.7	IPCC 2006 Guidelines
1A4bi	Domestic combustion	Burning oil	Liquid	N ₂ O	kg/TJ	0.6	0.6	0.6	0.6	IPCC 2006 Guidelines
1A4bi	Domestic combustion	Charcoal	Biomass	CH ₄	kg/TJ	200.0	200.0	200.0	200.0	IPCC 2006 Guidelines
1A4bi	Domestic combustion	Charcoal	Biomass	N ₂ O	kg/TJ	1.0	1.0	1.0	1.0	IPCC 2006 Guidelines
1A4bi	Domestic combustion	Coal	Solid	CO ₂	t/TJ	92.0	92.0	92.0	92.0	Based on Fynes & Sage, 1994
1A4bi	Domestic combustion	Coal	Solid	CH ₄	kg/TJ	547.2	548.0	556.4	531.4	Brain et al, 1994
1A4bi	Domestic combustion	Coal	Solid	N ₂ O	kg/TJ	4.2	4.2	4.3	4.1	Based on Fynes & Sage, 1994
1A4bi	Domestic combustion	Coke	Solid	CO ₂	t/TJ	114.3	111.3	114.3	101.8	Ricardo-AEA estimate based on carbon balance for integrated steelwork processes (2014)
1A4bi	Domestic combustion	Coke	Solid	CH ₄	kg/TJ	217.3	204.9	217.3	204.9	Brain et al, 1994
1A4bi	Domestic combustion	Coke	Solid	N ₂ O	kg/TJ	4.4	4.1	4.4	4.1	Based on Fynes & Sage, 1994
1A4bi	Domestic combustion	Fuel oil	Liquid	CO ₂	t/TJ	77.5	78.4	78.4	77.3	Carbon Factors Review 2004
1A4bi	Domestic combustion	Fuel oil	Liquid	CH ₄	kg/TJ	9.9	9.9	10.0	9.8	IPCC 2006 Guidelines
1A4bi	Domestic combustion	Fuel oil	Liquid	N ₂ O	kg/TJ	0.6	0.6	0.6	0.6	IPCC 2006 Guidelines
1A4bi	Domestic combustion	Gas oil	Liquid	CO ₂	t/TJ	73.9	74.1	74.2	73.5	Carbon Factors Review 2004
1A4bi	Domestic combustion	Gas oil	Liquid	CH ₄	kg/TJ	9.9	9.9	9.9	9.8	IPCC 2006 Guidelines
1A4bi	Domestic combustion	Gas oil	Liquid	N ₂ O	kg/TJ	0.6	0.6	0.6	0.6	IPCC 2006 Guidelines

Other Detailed Methodological Descriptions **A3**

Category	Source Name	Activity Name	Fuel (CRF)	Gas	Unit	1990	2013	Max	Min	EF Method Reference
1A4bi	Domestic combustion	LPG	Liquid	CO ₂	t/TJ	64.7	64.7	64.7	64.7	Carbon Factors Review 2004
1A4bi	Domestic combustion	LPG	Liquid	CH ₄	kg/TJ	5.1	5.1	5.1	5.1	IPCC 2006 Guidelines
1A4bi	Domestic combustion	LPG	Liquid	N ₂ O	kg/TJ	0.1	0.1	0.1	0.1	IPCC 2006 Guidelines
1A4bi	Domestic combustion	Natural gas	Gaseous	CO ₂	t/TJ	56.0	56.9	57.2	56.0	Natural gas carbon emission factor is derived from Gas Network Operator data on gas composition within each Local Distribution Zone, prepared for use by operators within the EU ETS. (DECC, 2014)
1A4bi	Domestic combustion	Natural gas	Gaseous	CH ₄	kg/TJ	5.0	5.0	5.0	5.0	IPCC 2006 Guidelines
1A4bi	Domestic combustion	Natural gas	Gaseous	N ₂ O	kg/TJ	0.1	0.1	0.1	0.1	IPCC 2006 Guidelines
1A4bi	Domestic combustion	Peat	Peat	CO ₂	t/TJ	106.0	106.0	106.0	106.0	IPCC 2006 Guidelines
1A4bi	Domestic combustion	Peat	Peat	CH ₄	kg/TJ	300.0	300.0	300.0	300.0	Use emission factor for domestic wood combustion
1A4bi	Domestic combustion	Peat	Peat	N ₂ O	kg/TJ	1.4	1.4	1.4	1.4	Use emission factor for domestic wood combustion
1A4bi	Domestic combustion	Petroleum coke	Liquid	CO ₂	t/TJ	87.0	96.1	96.1	87.0	Carbon Factors Review 2004
1A4bi	Domestic combustion	Petroleum coke	Liquid	CH ₄	kg/TJ	9.1	10.1	10.1	9.1	IPCC 2006 Guidelines
1A4bi	Domestic combustion	Petroleum coke	Liquid	N ₂ O	kg/TJ	0.5	0.6	0.6	0.5	IPCC 2006 Guidelines
1A4bi	Domestic combustion	SSF	Solid	CO ₂	t/TJ	110.5	102.3	110.5	93.6	British Coal (1989)
1A4bi	Domestic combustion	SSF	Solid	CH ₄	kg/TJ	221.2	204.9	221.2	187.5	Brain et al, 1994
1A4bi	Domestic combustion	SSF	Solid	N ₂ O	kg/TJ	4.6	4.3	4.6	3.9	Based on Fynes & Sage, 1994
1A4bi	Domestic combustion	Town gas	Solid	CO ₂	t/TJ	61.7	61.7	61.7	61.7	Ricardo-AEA estimate based on carbon balance for town gas manufacture (2014)
1A4bi	Domestic combustion	Town gas	Solid	CH ₄	kg/TJ	5.0	5.0	5.0	5.0	IPCC 2006 Guidelines

Category	Source Name	Activity Name	Fuel (CRF)	Gas	Unit	1990	2013	Max	Min	EF Method Reference
1A4bi	Domestic combustion	Town gas	Solid	N ₂ O	kg/TJ	0.1	0.1	0.1	0.1	IPCC 2006 Guidelines
1A4bi	Domestic combustion	Wood	Biomass	CH ₄	kg/TJ	279.4	260.7	279.4	260.7	IPCC 2006 Guidelines
1A4bi	Domestic combustion	Wood	Biomass	N ₂ O	kg/TJ	3.7	3.5	3.7	3.5	IPCC 2006 Guidelines
1A4bii	House and garden machinery	DERV	Liquid	CO ₂	t/TJ	73.5	73.0	73.5	72.8	Carbon Factors Review 2004
1A4bii	House and garden machinery	DERV	Liquid	CH ₄	kg/TJ	3.7	3.7	3.7	3.6	Developed based on T3 approach considering emission limits on total hydrocarbons
1A4bii	House and garden machinery	DERV	Liquid	N ₂ O	kg/TJ	30.3	30.1	30.3	30.0	EMEP/EEA Guidebook, 2006
1A4bii	House and garden machinery	Petrol	Liquid	CO ₂	t/TJ	70.2	70.0	70.2	70.0	Carbon Factors Review 2004
1A4bii	House and garden machinery	Petrol	Liquid	CH ₄	kg/TJ	71.0	13.4	71.0	13.4	Developed based on T3 approach considering emission limits on total hydrocarbons
1A4bii	House and garden machinery	Petrol	Liquid	N ₂ O	kg/TJ	0.7	0.7	0.7	0.7	EMEP/EEA Guidebook, 2006
1A4ci	Agriculture - stationary combustion	Coal	Solid	CO ₂	t/TJ	90.4	90.4	90.4	90.4	Based on Fynes & Sage, 1994
1A4ci	Agriculture - stationary combustion	Coal	Solid	CH ₄	kg/TJ	0.4	0.4	0.4	0.4	Brain et al, 1994
1A4ci	Agriculture - stationary combustion	Coal	Solid	N ₂ O	kg/TJ	5.3	5.2	5.5	5.2	Based on Fynes & Sage, 1994
1A4ci	Agriculture - stationary combustion	Coke	Solid	CO ₂	t/TJ	114.3	111.3	114.3	101.8	Ricardo-AEA estimate based on carbon balance for integrated steelwork processes (2014)

Category	Source Name	Activity Name	Fuel (CRF)	Gas	Unit	1990	2013	Max	Min	EF Method Reference
1A4ci	Agriculture - stationary combustion	Coke	Solid	CH ₄	kg/TJ	0.4	0.4	0.4	0.4	Brain et al, 1994
1A4ci	Agriculture - stationary combustion	Coke	Solid	N ₂ O	kg/TJ	5.6	5.3	5.6	5.3	Based on Fynes & Sage, 1994
1A4ci	Agriculture - stationary combustion	Fuel oil	Liquid	CO ₂	t/TJ	77.5	78.4	78.4	77.3	Carbon Factors Review 2004
1A4ci	Agriculture - stationary combustion	Fuel oil	Liquid	CH ₄	kg/TJ	9.9	9.9	10.0	9.8	IPCC 2006 Guidelines
1A4ci	Agriculture - stationary combustion	Fuel oil	Liquid	N ₂ O	kg/TJ	0.6	0.6	0.6	0.6	IPCC 2006 Guidelines
1A4ci	Agriculture - stationary combustion	Gas oil	Liquid	CO ₂	t/TJ	73.9	74.1	74.2	73.5	Carbon Factors Review 2004
1A4ci	Agriculture - stationary combustion	Gas oil	Liquid	CH ₄	kg/TJ	9.9	9.9	9.9	9.8	IPCC 2006 Guidelines
1A4ci	Agriculture - stationary combustion	Gas oil	Liquid	N ₂ O	kg/TJ	0.6	0.6	0.6	0.6	IPCC 2006 Guidelines
1A4ci	Agriculture - stationary combustion	Natural gas	Gaseous	CO ₂	t/TJ	56.0	56.9	57.2	56.0	Natural gas carbon emission factor is derived from Gas Network Operator data on gas composition within each Local Distribution Zone, prepared for use by operators within the EU ETS. (DECC, 2014)
1A4ci	Agriculture - stationary combustion	Natural gas	Gaseous	CH ₄	kg/TJ	5.0	5.0	5.0	5.0	IPCC 2006 Guidelines
1A4ci	Agriculture - stationary combustion	Natural gas	Gaseous	N ₂ O	kg/TJ	0.1	0.1	0.1	0.1	IPCC 2006 Guidelines

Category	Source Name	Activity Name	Fuel (CRF)	Gas	Unit	1990	2013	Max	Min	EF Method Reference
1A4ci	Agriculture - stationary combustion	Straw	Biomass	CH ₄	kg/TJ	267.8	267.8	267.8	267.8	IPCC 2006 Guidelines
1A4ci	Agriculture - stationary combustion	Straw	Biomass	N ₂ O	kg/TJ	3.6	3.6	3.6	3.6	IPCC 2006 Guidelines
1A4ci	Agriculture - stationary combustion	Vaporising oil	Liquid	CO ₂	t/TJ	73.0	73.2	73.3	72.5	Assume as burning oil
1A4ci	Agriculture - stationary combustion	Vaporising oil	Liquid	CH ₄	kg/TJ	9.9	9.9	9.9	9.8	Use 2006 IPCC default for gas oil
1A4ci	Agriculture - stationary combustion	Vaporising oil	Liquid	N ₂ O	kg/TJ	0.6	0.6	0.6	0.6	IPCC 2006 Guidelines
1A4ci	Miscellaneous industrial/commercial combustion	Burning oil	Liquid	CO ₂	t/TJ	71.8	71.7	71.8	71.7	Carbon Factors Review 2004
1A4ci	Miscellaneous industrial/commercial combustion	Burning oil	Liquid	CH ₄	kg/TJ	10.0	10.0	10.0	10.0	IPCC 2006 Guidelines
1A4ci	Miscellaneous industrial/commercial combustion	Burning oil	Liquid	N ₂ O	kg/TJ	0.6	0.6	0.6	0.6	IPCC 2006 Guidelines
1A4cii	Agriculture - mobile machinery	Gas oil	Liquid	CO ₂	t/TJ	73.9	74.1	74.2	73.5	Carbon Factors Review 2004
1A4cii	Agriculture - mobile machinery	Gas oil	Liquid	CH ₄	kg/TJ	3.8	3.7	3.8	3.7	Developed based on T3 approach considering emission limits on total hydrocarbons
1A4cii	Agriculture - mobile machinery	Gas oil	Liquid	N ₂ O	kg/TJ	30.6	30.6	30.7	30.4	EMEP/EEA Guidebook, 2006

Other Detailed Methodological Descriptions **A3**

Category	Source Name	Activity Name	Fuel (CRF)	Gas	Unit	1990	2013	Max	Min	EF Method Reference
1A4cii	Agriculture - mobile machinery	Petrol	Liquid	CO ₂	t/TJ	70.2	70.0	70.2	70.0	Carbon Factors Review 2004
1A4cii	Agriculture - mobile machinery	Petrol	Liquid	CH ₄	kg/TJ	48.7	48.6	48.7	48.6	Developed based on T3 approach considering emission limits on total hydrocarbons
1A4cii	Agriculture - mobile machinery	Petrol	Liquid	N ₂ O	kg/TJ	0.3	0.3	0.3	0.3	EMEP/EEA Guidebook, 2006
1A4ciii	Fishing vessels	Fuel oil	Liquid	CO ₂	t/TJ	77.5	78.4	78.4	77.3	Carbon Factors Review 2004
1A4ciii	Fishing vessels	Fuel oil	Liquid	CH ₄	kg/TJ	1.2	1.2	1.2	1.2	EMEP/EEA Guidebook, 2006
1A4ciii	Fishing vessels	Fuel oil	Liquid	N ₂ O	kg/TJ	1.9	1.9	2.0	1.9	EMEP/EEA Guidebook, 2006
1A4ciii	Fishing vessels	Gas oil	Liquid	CO ₂	t/TJ	73.9	74.1	74.2	73.5	Carbon Factors Review 2004
1A4ciii	Fishing vessels	Gas oil	Liquid	CH ₄	kg/TJ	1.2	1.2	1.2	1.2	EMEP/EEA Guidebook, 2006
1A4ciii	Fishing vessels	Gas oil	Liquid	N ₂ O	kg/TJ	1.9	1.9	1.9	1.8	EMEP/EEA Guidebook, 2006
1A5b	Aircraft - military	Aviation spirit	Liquid	CO ₂	t/TJ	69.6	69.5	69.6	69.3	Carbon Factors Review 2004
1A5b	Aircraft - military	Aviation spirit	Liquid	CH ₄	kg/TJ	2.3	2.3	2.3	2.3	Emissions from 1990 to the latest inventory year derived using IPCC Tier 3 methodology. Activity data provided by the CAA. Total consumptions of ATF and AS provided by DECC, via DUKES.
1A5b	Aircraft - military	Aviation spirit	Liquid	N ₂ O	kg/TJ	2.2	2.2	2.2	2.2	Emissions from 1990 to the latest inventory year derived using IPCC Tier 3 methodology. Activity data provided by the CAA. Total consumptions of ATF and AS provided by DECC, via DUKES.
1A5b	Aircraft - military	Aviation turbine fuel	Liquid	CO ₂	t/TJ	71.8	71.7	71.8	71.7	Carbon Factors Review 2004

Category	Source Name	Activity Name	Fuel (CRF)	Gas	Unit	1990	2013	Max	Min	EF Method Reference
1A5b	Aircraft - military	Aviation turbine fuel	Liquid	CH ₄	kg/TJ	2.4	2.3	2.4	2.3	Emissions from 1990 to the latest inventory year derived using IPCC Tier 3 methodology. Activity data provided by the CAA. Total consumptions of ATF and AS provided by DECC, via DUKES.
1A5b	Aircraft - military	Aviation turbine fuel	Liquid	N ₂ O	kg/TJ	2.3	2.3	2.3	2.3	Emissions from 1990 to the latest inventory year derived using IPCC Tier 3 methodology. Activity data provided by the CAA. Total consumptions of ATF and AS provided by DECC, via DUKES.
1A5b	Shipping - naval	Gas oil	Liquid	CO ₂	t/TJ	73.9	74.1	74.2	73.5	Carbon Factors Review 2004
1A5b	Shipping - naval	Gas oil	Liquid	CH ₄	kg/TJ	1.2	1.2	1.2	1.2	EMEP/EEA Guidebook, 2006
1A5b	Shipping - naval	Gas oil	Liquid	N ₂ O	kg/TJ	1.9	1.9	1.9	1.8	EMEP/EEA Guidebook, 2006

Table A 3.1.2 Emission factors used for the 2015 submission (sector 1B)

Category	Source Name	Activity Name	Gas	Unit	1990	2013	Max	Min	EF Method Reference
1B1a1i	Deep-mined coal	Coal produced	CH ₄	kilotonne/ Mt material produced	10.0	10.1	14.0	9.0	Methane emission factor kt CH ₄ /Mt coal produced calculated based on data from UK Coal Mining Ltd
1B1a1ii	Coal storage and transport	Deep mined coal production	CH ₄	kilotonne/ Mt material produced	1.2	1.2	1.2	1.2	Emission factors kt CH ₄ /Mt deep mined coal. From CH ₄ _XX.xls: Bennett et al (1995) ETSU N/01/00006/REP
1B1a1iii	Closed Coal Mines	Non-fuel combustion	CH ₄	Modelled					Emission estimates from abandoned coal mines, WSP (2011)
1B1a2i	Open-cast coal	Coal produced	CH ₄	tonnes/ Mt material produced	339.8	339.8	339.8	339.8	kt CH ₄ / Mt open cast coal production. Williams , A,(1993), Methane Emissions, Watt Committee Report No 28

Other Detailed Methodological Descriptions **A3**

Category	Source Name	Activity Name	Gas	Unit	1990	2013	Max	Min	EF Method Reference
1B1b	Charcoal production	Charcoal produced	CH ₄	kilotonne/ Megatonne	30.0	30.0	30.0	30.0	IPCC, 1996, Revised Guidelines
1B1b	Coke production	Coke produced	CH ₄	tonnes/ Megatonne	80.2	80.2	80.2	80.2	EIPPCB, 2000
1B1b	Iron and steel - flaring	Coke oven gas	CO ₂	kilotonne/ Mth fuel consumed	4.2	4.1	4.6	4.0	Ricardo-AEA estimate based on carbon balance for integrated steelwork processes (2014)
1B1b	Iron and steel - flaring	Coke oven gas	CH ₄	tonnes/ Mth fuel consumed	0.1	0.1	0.1	0.1	IPCC 2006
1B1b	Iron and steel - flaring	Coke oven gas	N ₂ O	tonnes/ Mth fuel consumed	9.4E-03	9.4E-03	9.4E-03	9.4E-03	IPCC 2006
1B1b	Solid smokeless fuel production	Coal	CO ₂	kilotonne/ Mt fuel consumed	1012.8	375.6	1012.8	269.1	Ricardo-AEA estimate based on carbon balance for solid smokeless fuel manufacture
1B1b	Solid smokeless fuel production	Petroleum coke	CO ₂	kilotonne/ Mt fuel consumed	1012.8	375.6	1012.8	269.1	Factor calculated from difference between Ricardo-AEA estimates of carbon input to and carbon output from SSF manufacturing. From 2009 this includes new data (from DUKES) on petroleum coke inputs to SSF manufacture. (2014)
1B1b	Solid smokeless fuel production	SSF produced	CH ₄	tonnes/ Megatonne	80.2	80.2	80.2	80.2	EIPPCB, 2000
1B2a1	Upstream Oil Production - Offshore Well Testing	Exploration drilling :amount of gas flared	CO ₂	tonnes/ tonne	3.2	3.2	3.2	3.2	EEMS 2013 (DECC Offshore Inspectorate)
1B2a1	Upstream Oil Production - Offshore Well Testing	Exploration drilling :amount of gas flared	CH ₄	tonnes/ tonne	1.6E-02	2.5E-02	3.7E-02	1.1E-02	EEMS 2013 (DECC Offshore Inspectorate)

Other Detailed Methodological Descriptions **A3**

Category	Source Name	Activity Name	Gas	Unit	1990	2013	Max	Min	EF Method Reference
1B2a1	Upstream Oil Production - Offshore Well Testing	Exploration drilling :amount of gas flared	N ₂ O	tonnes/ tonne	2.9E-05	7.9E-05	8.5E-05	2.2E-05	EEMS 2013 (DECC Offshore Inspectorate)
1B2a2	Petroleum processes	Oil production	CH ₄	tonnes/ Megatonne	21.7	149.5	149.5	13.7	Based on data reported in the Pollution Inventory (Environment Agency, 2013)
1B2a2	Upstream Oil Production - process emissions	Non-fuel combustion	CO ₂	Operator reported emissions					EEMS 2013 (DECC Offshore Inspectorate)
1B2a2	Upstream Oil Production - process emissions	Non-fuel combustion	CH ₄	Operator reported emissions					EEMS 2013 (DECC Offshore Inspectorate)
1B2a3	Upstream Oil Production - Offshore Oil Loading	Crude oil	CH ₄	tonnes/ t	4.3E-05	5.5E-05	1.2E-04	4.3E-05	EEMS 2013 (DECC Offshore Inspectorate)
1B2a3	Upstream Oil Production - Onshore Oil Loading	Crude oil	CH ₄	tonnes/ t	1.3E-05	8.0E-06	1.3E-05	7.2E-06	EEMS 2013 (DECC Offshore Inspectorate)
1B2a4	Upstream Oil Production - Oil terminal storage	Non-fuel combustion	CH ₄	Operator reported emissions					EEMS 2013 (DECC Offshore Inspectorate)
1B2b1	Upstream Gas Production - Offshore Well Testing	Exploration drilling :amount of gas flared	CO ₂	tonnes/ t	2.7	2.8	2.9	2.7	EEMS 2013 (DECC Offshore Inspectorate)
1B2b1	Upstream Gas Production - Offshore Well Testing	Exploration drilling :amount of gas flared	CH ₄	tonnes/ t	1.6E-02	4.5E-02	4.5E-02	1.1E-02	EEMS 2013 (DECC Offshore Inspectorate)
1B2b1	Upstream Gas Production - Offshore Well Testing	Exploration drilling :amount of gas flared	N ₂ O	tonnes/ t	2.9E-05	8.1E-05	8.5E-05	2.2E-05	EEMS 2013 (DECC Offshore Inspectorate)

Other Detailed Methodological Descriptions **A3**

Category	Source Name	Activity Name	Gas	Unit	1990	2013	Max	Min	EF Method Reference
1B2b3	Upstream Gas Production - process emissions	Non-fuel combustion	CO ₂	Operator reported emissions					EEMS 2013 (DECC Offshore Inspectorate)
1B2b3	Upstream Gas Production - process emissions	Non-fuel combustion	CH ₄	Operator reported emissions					EEMS 2013 (DECC Offshore Inspectorate)
1B2b4	Upstream Gas Production - Gas terminal storage	Non-fuel combustion	CH ₄	Operator reported emissions					EEMS 2013 (DECC Offshore Inspectorate)
1B2b5	Gas leakage	Natural Gas (leakage at point of use)	CO ₂	tonnes/ Mth fuel consumed	9.2E-03	8.0E-03	9.5E-03	6.2E-03	Gas leakage calculated based on boiler numbers, supplied by DECC (Dunbabin, pers. comm.)
1B2b5	Gas leakage	Natural Gas (leakage at point of use)	CH ₄	tonnes/ Mth fuel consumed	0.2	0.2	0.2	0.2	Gas leakage calculated based on boiler numbers, supplied by DECC (Dunbabin, pers. comm.)
1B2b5	Gas leakage	Natural gas supply	CO ₂	tonnes/ kt Ngas leaked	39.2	34.6	39.2	26.0	NMVOC, CH ₄ and CO ₂ Efs from annual gas composition data from gas distribution networks (2014)
1B2b5	Gas leakage	Natural gas supply	CH ₄	tonnes/ kt Ngas leaked	843.4	798.9	843.4	763.3	C. Rose, British Gas (1993) & S. Pearson British Gas(1994)
1B2c1i	Upstream Oil Production - venting	Non-fuel combustion	CO ₂	Operator reported emissions					EEMS 2013 (DECC Offshore Inspectorate)
1B2c1i	Upstream Oil Production - venting	Non-fuel combustion	CH ₄	Operator reported emissions					EEMS 2013 (DECC Offshore Inspectorate)

Other Detailed Methodological Descriptions **A3**

Category	Source Name	Activity Name	Gas	Unit	1990	2013	Max	Min	EF Method Reference
1B2c1ii	Upstream Gas Production - venting	Non-fuel combustion	CO ₂	Operator reported emissions					EEMS 2013 (DECC Offshore Inspectorate)
1B2c1ii	Upstream Gas Production - venting	Non-fuel combustion	CH ₄	Operator reported emissions					EEMS 2013 (DECC Offshore Inspectorate)
1B2c2i	Upstream Oil Production - flaring	Non-fuel combustion	CO ₂	tonnes/ tonne	1.4	2.6	2.7	1.4	EEMS 2013 (DECC Offshore Inspectorate)
1B2c2i	Upstream Oil Production - flaring	Non-fuel combustion	CH ₄	tonnes/ tonne	9.1E-03	9.7E-03	1.1E-02	6.7E-03	EEMS 2013 (DECC Offshore Inspectorate)
1B2c2i	Upstream Oil Production - flaring	Non-fuel combustion	N ₂ O	tonnes/ tonne	4.5E-05	7.4E-05	1.0E-04	4.5E-05	EEMS 2013 (DECC Offshore Inspectorate)
1B2c2ii	Upstream Gas Production - flaring	Non-fuel combustion	CO ₂	tonnes/ tonne	1.4	2.3	2.7	1.4	EEMS 2013 (DECC Offshore Inspectorate)
1B2c2ii	Upstream Gas Production - flaring	Non-fuel combustion	CH ₄	tonnes/ tonne	9.1E-03	7.0E-03	1.8E-02	5.6E-03	EEMS 2013 (DECC Offshore Inspectorate)
1B2c2ii	Upstream Gas Production - flaring	Non-fuel combustion	N ₂ O	tonnes/ tonne	4.5E-05	7.4E-05	8.6E-05	4.5E-05	EEMS 2013 (DECC Offshore Inspectorate)

A 3.1.2 Commonly used references

This section describes data sources that are used across multiple emission sources within the energy sector, and how they are used.

A 3.1.2.1 Baggott et al., 2004 – Carbon factors review

A review of the carbon factors used in the UK GHG inventory was carried out in 2004. The report detailing this study is available from:

[http://uk-air.defra.gov.uk/assets/documents/reports/cat07/0611061401-417_Review_of Carbon Emission Factors 2004NIR Issue1 v1.3.2.pdf](http://uk-air.defra.gov.uk/assets/documents/reports/cat07/0611061401-417_Review_of_Carbon_Emission_Factors_2004NIR_Issue1_v1.3.2.pdf)

This aimed to validate existing emission factors and seek new data for country specific emission factors for the UK. At the time of publication this reference provided new emission factors for:

- coal from power stations
- fuels used in the cement industry
- a number of petroleum based fuels
- natural gas
- coke oven and blast furnace gas.

Since then following updates are made to the following emission factors based on new information:

1. Coal emission factors are adjusted based on the annual variations in the GCV of the fuels using methods developed as part of the 2004 analysis (Baggott et al., 2004).

$$EF_y = EF_{ref} / GCV_{ref} * GCV_y$$

Where

EF_y is the emission factor in year y

EF_{ref} is the EF in the reference year (the year for which data are available)

GCV_{ref} is the GCV in the reference year

GCV_y is the GCV in year y

2. Since the advent of EU ETS in 2005, a number of sources of emissions from coal which had previously been reliant on Baggott et al., 2004 have now been replaced with data from the ETS, where the data set was considered suitable (high proportion of source included, and high proportion of T3 plant specific data). In addition, in 2014 the use of oxidation factors from this report was reviewed, and where suitable background evidence to support the factors used were not available, the IPCC default (of 1, IPCC 2006) has been used.
3. Emission factors for petroleum based fuels (where ETS data are not available) are still largely based on Baggott et al., 2004. These were reviewed in 2014 and compared with the defaults in the 2006 IPCC Guidelines and found to be largely within the range

of the 2006 Guidelines. No new data for the UK has been identified and the emission factors from Baggott et al., 2004 are considered to continue to be relevant country specific emission factors.

4. Emission factors for natural gas are updated annually based on analyses from the gas network operators (Personal Communications from network operators, 2014).
5. Emission factors for coke oven gas and blast furnace gas are estimated based on a carbon balance approach (as described in Chapter 3, section XX).
6. The Mineral Products Association provide data for fuels used in the cement industry annually on a confidential basis, and these are validated with EU ETS data (Personal Communication, MPA, 2014).

A 3.1.2.2 Entec (2010) UK Ship Emissions Inventory

A method has been developed for estimating emissions and fuel consumption from shipping in UK waters and ports by Entec (now AMEC) (Entec, 2010) under contract to Defra. The method uses a bottom-up procedure based on detailed shipping movement data for different vessel types, fuels and journeys. The approach represents a Tier 3 method for estimating emissions from domestic water-borne navigation in the IPCC Guidelines for national inventories.

The detailed study included the following methods, data sources and assumptions.

1. It covered a detailed study of movements, types, abatement technology, fuels used in only one year, 2007. A gridded emissions inventory from ship movements within waters surrounding the UK including the North Sea, English Channel, Irish Sea and North East Atlantic was developed based on ship movement data. The study area was 200 nautical miles from the UK coastline and fuel consumption and emissions were resolved to a 5x5km grid and included emissions from vessels cruising at sea and manoeuvring and at berth in port.
2. The study used proxy data to back cast movements and fuel consumption to 1990 and forward cast to 2009 (of all time series of the inventory at the time). The variables that were considered were:
 - Trends in vessel movements over time affected by changes in the number of vessels and their size;
 - Trends in fuel type in use over time reflecting the era before the introduction of SECAs which would have permitted higher sulphur content fuel to be used;
 - A time-series of tonnes fish landed in the UK provided in UK Sea Fisheries Statistics by the Marine Management Organisation was used for the domestic fishing vessels category (MMO, 2013).

The key consideration was the trend in vessel movements over time. For this, DfT's annual published Maritime Statistics were used as proxies for activity rate changes which were taken to be indicators of fuel consumed. A range of time-series trends back to 1990 from the DfT statistics are available and appropriate data were assigned to different vessel categories, differentiating between international and domestic movements. Details are given in the Entec (2010) report, but in brief:

- All ports traffic data based on tonnes cargo for domestic and international movements was assigned as an indicator for the bulk carrier, general cargo and tanker vessel categories. Trends were available from full the full time series 1990 to the latest inventory year;
 - All ports main unitised statistics reported as number of units for domestic and international movements was assigned as an indicator for the container ship and Ro-Ro cargo vessel categories. Trends were available for the full time series;
 - International and domestic sea passenger movements reported as number of passengers was assigned to the passenger vessel category.
3. Individual vessel movements and characteristics data provided by Lloyd's Marine Intelligence Unit (LMIU) for the year 2007 supplemented by Automatic Identification System (AIS) data transmitted by vessels to shore with information about a ship's position and course were integrated. The LMIU movement data included vessel type and speed. The vessel types were grouped into eight vessel categories, which differentiated between engine size and vessel operation for vessel types, giving an indication of the likely fuel used.
 4. The LMIU database does not capture all vessel movements. The LMIU data alone underestimated port arrivals for each port most likely from missing vessels <300 GT with multiple callings each day. Movement and fuel consumption estimates are scaled up based on an inter-comparison with the DfT's marine statistics⁴ (DfT, 2008c), which also include data from MDS-Transmodal which includes small vessels and those with multiple callings to the same port each day, such as cross-channel passenger ferries and fishing vessels.
 5. Missing from both the DfT and LMIU data sets are movements by tugs, dredgers, research vessels and other vessels employed within the limit of the port or estuary as well as small pleasure craft, which are included in the GHGI as part of the estimates for inland waterways.
 6. Entec only covered emissions from fishing activities occurring within the UK waters study area extending 200 nautical miles from the UK coast.
 7. Fuel consumption and emissions were calculated for each of these vessel categories for different operations. Vessel speeds were combined with distance travelled to determine the time spent at sea by each vessel. Entec undertook a detailed analysis of port callings where a significant proportion of emissions occur. The analysis considered time-in-mode for manoeuvring, hoteling in ports and loading and unloading operations.
 8. The Entec study developed a series of assumptions on the fuel type, engine size and S content of fuels what have been included in the estimates

⁴ The DfT port statistics comprise annual totals by port and ship type in Maritime Statistics and refer to movement of all sea-going vessels >100 Gross Tonnage (GT) involved in the movement of goods or passengers derived from LMIU and a number of other datasets.

Table A 3.1.3 below shows how the Entec figures are used, and how this is reconciled with total fuel use for shipping from DUKES.

Table A 3.1.3 Approaches of estimating activity and emission factors

	Source	IPCC	Activity data			Emission factors	
			Source	Base year	Time-series		
DUKES total marine fuel consumption (A)	Domestic (B)	Domestic coastal	1A3dii	Entec (2010) based on detailed vessel movement data (LMIU and AIS)	2007	DfT port movement data to scale from 2007 to other years	Entec (2010), EMEP/EEA Guidebook, UKPIA (2013)
		Fishing in UK sea territories	1A4ciii	Entec (2010) based on detailed vessel movement data	2007	MMO fish landing statistics to scale from 2007 to other years	Entec (2010), EMEP/EEA Guidebook, UKPIA (2013)
		Fishing in non-UK sea territories	1A4ciii	MMO data on fish landings by sea territory from 1994-2012 and estimates of fish landed per trip			Entec (2010), EMEP/EEA Guidebook, UKPIA (2013)
		Naval	1A5b	MoD data on fuel consumption by naval vessels			Assumed same as international shipping vessels using gas oil
		Shipping between UK and OTs	1A3dii	DfT Maritime Statistics and OT port authorities: number of sailings between UK and OT	2000-2012	Trends for years before 2000 based on trends in fuel consumption derived by Entec for international shipping and trends in DfT data on number of cruise passengers	Assumed same as international shipping vessels using fuel oil
		Inland waterways	1A3dii	Based on estimates of vessel population and usage estimates using data from various sources.	2008	Statistics on expenditure on recreation (ONS), tourism (Visit England), port freight traffic (DfT), inland waterways goods lifted (DfT) used to scale from 2008	EMEP/EEA Guidebook, UKPIA (2013)

		Source	IPCC	Activity data			Emission factors
				Source	Base year	Time-series	
	International (C)		1A3di	Fuel consumption difference between DUKES total marine fuel consumption and domestic navigation calculated above $(C=A-B)$			Implied emission factor for international shipping from Entec (2010)

A 3.1.2.3 The Pollution Inventory and other regulators' inventories.

The Pollution Inventory (PI) has, since 1998, provided emission data for the six Kyoto gases and other air pollutant for installations regulated by the Environment Agency (EA) in England and Natural Resources Wales (NRW) in Wales. This is part of the UK's process for managing regulated emissions from industry processes under IPPC permitting system. The PI does contain some earlier data as well, with carbon dioxide emissions at some sites reported from 1994 onwards. The Scottish Pollutant Release Inventory (SPRI) covers processes regulated by the Scottish Environment Protection Agency (SEPA), and contains data from 2002 and 2004 onwards. The Northern Ireland Pollution Inventory (NIPI) covers processes regulated by the Department of the Environment (Northern Ireland) and includes data for 1999 onwards.

These three sets of data are subject to some very significant limitations:

- emissions of each pollutant are reported for each permitted installation as a whole, so emissions data for carbon dioxide, for example, can cover emissions from fuel use as well as from an industrial process. No information is given on what the source of emissions is, so a judgement has to be made about the scope of reporting.
- permitting arrangements have changed over time, so the reporting of data is not on a consistent basis across the time-series. In general, the tendency has been to reduce the number of permits, so that whereas in the early 1990s there might have been separate permits at an industrial installation covering the boiler plant and the chemical processes, from the late 1990s onwards the tendency would be to issue a single permit to cover both. Therefore, the problems with the scope of emissions data mentioned in the first bullet point are most severe for the second half of the GHGI time series.
- Since 1998, process operators need only report emissions of each pollutant if those emissions exceed a reporting threshold. So, if the emissions from an installation are less than 10,000 tonnes in the case of CO₂, or 10 tonnes in the case of methane, data are not available. Reporting thresholds are irrelevant for many of the sectors of interest to this study, since emissions would be many times higher than the thresholds, but the reporting thresholds do mean that it is necessary to consider whether the data available in the PI (and in the SPRI & NIPI for later years) will be complete,

Despite these limitations, these data are still a useful source of information for the UK GHG inventory. A considerable amount of effort is put into manually interpreting the individual returns and allocating these to appropriate categories for use in the inventory estimates by the Inventory Agency.

A 3.1.2.4 The EEMS Reporting System

Emissions from upstream oil and gas production facilities, including onshore terminals, are estimated based on operator reporting via EEMS, regulated by the DECC Offshore Inspectorate and developed in conjunction with the trade association Oil & Gas UK (formerly the UK Offshore Operators' Association, UKOOA). The EEMS data provides a detailed inventory of point source emissions estimates, based on operator returns for the years 1995-2012. However, the EEMS data for 1995 to 1997 are not complete, frequently exhibiting duplicate entries with identical submissions by operators across years, Since the 1995 – 1997

data are not considered reliable, the EEMS dataset is only used directly to inform national inventory estimates from 1998 onwards for the following sources:

- Gas flaring;
- Own gas combustion;
- Well testing; and
- Oil loading (onshore and offshore)

[Activity data are not routinely collected via EEMS for sources including: fugitive releases, direct process activities, oil storage or gas venting. The emissions from these sources are reported as annual estimates by operators and used directly within the inventory.]

These EEMS-derived activity data enable greater analysis of the oil & gas emissions and related emission factors at the installation level, providing a high degree of data transparency and improving the level of detail for performing quality checks by source, by site, by year. For those sources, this has led to an improvement in data transparency and easier query of Implied Emission Factor trends. However, the EEMS activity data are only available back to 1998, and hence the activity data back to 1990 are extrapolated using the oil and gas production time-series that were collected at that time for the purposes of energy data reporting.

A 3.1.3 Feedstocks and NEU of fuels

The estimation methods are described within individual sections of the NIR, but are summarised here. The general approach adopted in the UK GHG inventory is to assume that emissions from all non-energy uses of fuels are zero (i.e. the carbon is assumed to be sequestered as products), except for cases where emission sources can be identified and emission estimates included in the inventory.

The UK Inventory Agency conducts periodic studies into the fate of fuels reported as non-energy use, in order to assess the levels of stored carbon and carbon emitted for different fuels over the time series. These detailed studies are supplemented through annual data gathering and consultation with stakeholders to maintain an accurate representation of the emitted and stored carbon in the inventory.

The assumptions and estimates for individual sources are based on a review conducted in 2013-14 (Ricardo-AEA, 2014b) which included research into UK-specific activities and data sources as well as a review of the National Inventory Reports (NIRs) of other countries.

The sections below outline the emission sources from feedstock and NEU of fuels that are included in the UK GHGI, the source data and estimation methods and a summary of the time series for each of the fuel types where there is a stored carbon component in the UK energy balance. The estimates are all presented in CRF Tables 1.Ab and 1.Ad.

Table A 3.1.4 Summary of Emission Sources for UK Fuels Allocated as Non Energy Use in UK Energy Statistics

Fuel	IPCC	Source Category
<i>Liquid Fossil</i>		
Naphtha, Liquid Petroleum Gases (LPG), Refinery Fuel Gas (RFG) / Other Petroleum Gases (OPG), gas oil and Ethane	1A1a	Scrap tyre combustion in power stations (1994 to 2000 only). Fossil carbon in MSW combustion in energy from waste plant. <i>Emissions of carbon from chemical feedstock via combustion of products such as synthetic rubbers and plastics.</i>
	1A1b	Other petroleum gas use in refineries (2004 to 2012 only). <i>Re-allocated from non-energy use as EU ETS and trade association data indicates that DUKES data on OPG combustion are an under-report.</i>
	1A2c	Other petroleum gas use in petrochemical facility combustion. <i>Re-allocated from non-energy use as EU ETS and operator data indicates that DUKES data on OPG combustion are an under-report. These emissions were reported under 1A2f in the 2013 submission, but have now been re-allocated to 1A2c as these are entirely emissions from chemical and petrochemical production facilities.</i>
	1A2f	Containing fossil carbon, in cement kilns. Scrap tyre combustion in cement kilns.
	1A2g	Carbon in energy recovery from waste solvent and mixed general waste Industrial combustion of waste solvents. <i>Emissions of carbon from chemical feedstock via combustion of products such as synthetic rubbers and solvents.</i>
	2B5	Energy recovery from process gases in the chemical industry. Release of carbon from breakdown of chemical products such as soaps, detergents and pesticides after use. <i>Emissions of carbon from chemical feedstock via breakdown of products.</i>
	5C	Fossil carbon in chemical waste incineration. Fossil carbon in MSW incineration. Fossil carbon in clinical waste incineration. <i>Emissions of carbon from chemical feedstock via combustion of products such as synthetic rubbers and plastics.</i>

Fuel	IPCC	Source Category
Lubricants	1A1a	Waste oil combustion in power stations.
	1A2f	Waste oil combustion in cement kilns.
	1A2g	Waste oil combustion in unclassified industry (including road-stone coating plant)
	2D1	Lubricant combustion in aircraft, industrial, road vehicle, marine shipping and agricultural engines.
	5C	Incineration of waste oil.
Bitumen	n/a	<i>No known UK applications that lead to GHG emissions.</i>
Petroleum coke	1A2f, 1A4b, 2C1, 2C3	<p>Based on reported energy use data by specific industries within datasets such as EU ETS and also from direct dialogue with industry representatives, the Inventory Agency re-allocates a small proportion of the reported “NEU” allocation from DUKES, and reports emissions within the UK GHG inventory. This re-allocation generates emissions for the mineral processing sector (1A2f) and for petcoke use in the domestic sector (1A4b).</p> <p>There are also non-combustion, emissive uses of petcoke in the UK through the use of petcoke-derived anodes in the metal processing industries. Emissions from these uses of petcoke are reported in 2C1 (electrode use in electric arc furnaces) and 2C3 (anode use in aluminium manufacture).</p> <p><i>Note that DUKES already includes allocations of petcoke use as a fuel in combustion in power stations (1A1a) and refineries (1A1b), which are included in the UK GHG inventory.</i></p>
Other Oil	2B5	Carbon released from use of petroleum waxes. Uses of petroleum waxes includes candles, with carbon emitted during use.
<i>Solid Fossil</i>		
Coking coal (coal oils and tars)	n/a	<i>Unknown quantities of coal tar pitch are used in the manufacture of anodes for industrial processes. In the UK inventory the emissions from the use of these anodes are allocated only against petroleum coke (also used in anode production). This is a small mis-allocation of emissions between the two fuels since the carbon emitted is likely to arise from both petroleum coke and the coal tar pitch, but it is due to lack of detailed data, and does not affect the accuracy of UK inventory emissions.</i>
<i>Gaseous Fossil</i>		
Natural Gas	2B1	Ammonia production leading to either direct release of CO ₂ or associated chemical production (of methanol) with subsequent release of carbon originating in the natural gas feedstock.

A 3.1.3.1 Naphtha, Ethane, Gas Oil, Refinery/Other Fuel Gas (RFG/OPG) Propane and Butane (LPG)

Ethane, LPG (given separately as propane & butane in the energy statistics), gas oil (“middle distillate oil” when used as a chemical feedstock), refinery / other fuel gas (RFG/OPG) and naphtha are all consumed in very significant quantities for non-energy uses, primarily as feedstock in chemical manufacturing. In the UK, several major petrochemical production facilities are supplied with Natural Gas Liquid (NGL) feedstock directly from upstream production pipelines, and then utilise NGL fractions such as ethane, propane and butane in their manufacturing processes. In addition, several integrated refinery / petrochemical

complexes in the UK use a proportion of the refinery fuel gas as a feedstock in petrochemical production.

The NEU allocations presented in DUKES reflect the reported disposals of these commodities as feedstocks to chemical and petrochemical companies. There are several sources of GHG emissions from this stock of “NEU” feedstock carbon, although a high proportion of carbon is stored into products and not emitted.

One large emission source known to occur in the UK is the use of carbon-containing process off-gases as a fuel within the chemical facilities. Whilst the exact source of the carbon cannot be traced directly to a specific feedstock commodity within the UK sectoral approach, the available information from EU ETS and from consultation with operators enables the Inventory Agency to derive estimates of the GHG emissions across the time series from this emission source.

During the 2013 research study into NEU of fuels, the estimates of emissions and activity from this source were revised, leading to a number of small recalculations compared to the 2013 submission. Estimates for the ethylene manufacturing sites in the UK were improved through new information provided by the plant operators, whilst analysis of EU ETS data for a number of other chemical sites identified small additional emission sources that could be attributed to the combustion of process off-gases and residues derived from the chemical feedstock. As a result, the UK inventory emissions in 1A2c now include estimates of emissions from use of process off-gases and residues at 5 ethylene manufacturing sites and 4 other chemical manufacturing sites in the UK. The derivation of a time series of emission estimates from these sources is based as far as possible on reported data by plant operators within trading scheme data and other regulatory reporting mechanisms. For the early part of the time series, data on changes in plant capacity over time is used to derive the best estimates of activity and emissions by extrapolation back from later emission estimates, whilst for later years the completeness and transparency of operator reporting is greater. Therefore, whilst the uncertainty for the emission estimates in the early part of the time series is significantly greater than for those in recent years. The Inventory Agency has made best use of the available data to derive the time series estimates of emissions from “NEU” activity. Consultation with a sector trade association has also confirmed that there are no other sector estimates of this activity, or of production data across the time series, that could be used to further improve the time series (Personal communication: Chemical Industries Association, 2014).

Other emissions included within the UK GHG inventory (2B5) and derived from the NEU allocations as chemical feedstock include emission sources from the use or destruction of chemical products, e.g. when waste chemical products are incinerated. Although emissions from incineration and combustion of wastes are estimated, we cannot relate the carbon in these wastes back to individual feedstock, so it is not possible to generate reliable UK estimates of the proportion of carbon that is ultimately emitted from each individual fuel. Some butane is used as a propellant in aerosols and is emitted as VOC. The UK inventory contains estimates of these VOC emissions, combined with emissions of solvents used in aerosols.

Emissions can also occur from breakdown or combustion of products from the chemical industry that contain “stored carbon” from feedstock that in DUKES is allocated to non-energy use. Sources of emissions include burning of waste products and final products (e.g. flaring and use of wastes as fuels, or burning of candles, firelighters and other products etc.) or

degradation of products after disposal resulting in CO₂ emissions (including breakdown of consumer products such as detergents etc.).

Other emission sources from the use or degradation of chemical products that contain carbon from ethane, naphtha, gas oil, RFG/OPG and LPG feedstocks are:

- Carbon emitted during energy recovery - chemical industry;
- Carbon in products - soaps, shampoos, detergents etc.; and
- Carbon in products – pesticides.

A full time series of emissions is included in the inventory, and details of the methodology for these sectors are given in Passant, Watterson & Jackson, 2007. Emissions are reported under 2B5.

[Emissions in 2B5 from petroleum waxes are accounted for in the UK inventory under the fuel category “Other Oils” in CRF table 1Ad (see below).]

Emissions from products that contain fossil carbon that are oxidised in incinerators (MSW, chemical, clinical) are included in the GHG inventory and reported under 6C; where the use of such wastes as fuel includes power generation, the emissions are allocated in 1A1a. In the UK there have been short periods where scrap tyres have been used as a power station fuel. Tyres contain a mixture of natural and synthetic rubbers, however, and the emission estimates for combustion of scrap tyres in the UK inventory takes into account that only some of the carbon emitted is derived from fossil fuels.

In the reporting of the **Reference Approach** within the 2013 submission, the Inventory Agency had derived UK-specific carbon storage fractions using the available data on emissions (from the emission sources outlined above and reported in the Sectoral Approach) and the aggregated NEU allocations for ethane, LPG and naphtha. However, the 2013 research study engaged two UNFCCC Lead Reviewers to review the UK inventory compilation and reporting approach. The method used to calculate and report the Reference Approach was critically reviewed, compared against the approach taken by other Parties and the method overhauled for the 2014 submission. The UK inventory Reference Approach now follows the recommended approach from the 1996 IPCC Guidelines to apply default carbon storage fractions. In recent UNFCCC expert reviews, the Inventory Agency has received a range of advice on the most correct interpretation of the 1996 IPCC Guidelines. The two UNFCCC Lead Reviewers considered the available data and emission sources in the UK and determined that the best approach was to apply default carbon fractions to aid comparability against other reporting parties.

The UK energy balance for **gas oil** indicates a relatively high allocation of gas oil use in non-energy uses (130kt in 2012, which is around 2.9% of total UK supply) and these data are reflected in the UK GHG inventory, and the carbon stored is reported in CRF table 1A(d). Consultation with DECC energy statisticians in 2013 has confirmed that gas oil is delivered from UK refineries to petrochemical production facilities, and is used a feedstock material, consistent with the NEU allocation in DUKES (DECC, 2013). Furthermore, consultation with the Health and Safety Executive, who regulate the UK explosives industry, has confirmed that no UK installations manufacture explosives using gas oil or fuel oil as a feedstock (HSE, 2013).

A 3.1.3.2 Lubricants

Lubricants are listed separately in the UK energy statistics and are used in vehicles and in machinery. The inventory includes estimates of emissions of carbon due to oxidation of lubricants during use, and also includes estimates of emissions from the combustion of waste lubricants and other oils used as fuel.

UK GHG inventory estimates of the quantities of lubricants burnt are based on data from Recycling Advisory Unit, 1999; BLF/UKPIA/CORA, 1994; Oakdene Hollins Ltd, 2001 & ERM, 2008, as well as recent research to access information regarding the UK market for waste oils and the impact of European Directives to consolidate industrial emission regulations such as the Waste Incineration Directive (Oil Recycling Association, 2010). Estimates of waste oil combustion are derived for the following source categories:

- 1A1a Power stations
- 1A2f Cement kilns
- 1A2f Other (unclassified) industry

The estimated emissions for other industry assume that waste oils are used by two sectors: road-stone coating plant and garages. Other sectors may use waste oils as a fuel or as a reductant, but research to date provides no compelling evidence that there is a gap in the UK inventory for waste oil use by industrial operators.

The emission trends from power station use of waste lubricants reflect the fact that the Waste Incineration Directive (WID) had a profound impact on the market for waste oil, used as a fuel. It is assumed that no waste oil was burnt in power stations for the years 2006-2008. In 2009 a Quality Protocol⁵ was introduced that allowed compliant fuel produced from waste oils to be burned as non-waste and this has encouraged a resumption in the consumption of waste oil-derived fuels from 2009 onwards.

Carbon dioxide emission estimates for the oxidation of lubricants within vehicle engines and machinery, and the use of waste oils for energy are all based on a single carbon emission factor derived from analysis of the elemental composition of a series of UK-sourced samples of waste oil (Passant, 2004). UK research into the fate (including combustion) of lubricating oils within engines has been conducted periodically, with one study providing an estimate of 16% of all lubricants burned in engines in 1989, and a 2005 study estimating the total to be 13% of all lubricant supply in 2004. The estimated percentage of total lubricant demand oxidised in engines for 1990-2003 is interpolated between the 1989 and 2004 values given in these studies, and the 2004 estimate is extrapolated forwards across all years to 2012.

This time series of the percentage of lubricant burnt in engines is then applied to the annual lubricant demand given in DUKES to give a time series of quantities burnt. The estimates of waste lubricants burnt are then further sub-divided into the different source categories (including road, rail, marine, off-road and air transport) using the more detailed source-specific analysis within the 2005 study.

⁵ <http://www.environment-agency.gov.uk/business/topics/waste/116133.aspx>

A 3.1.3.3 Bitumen

In the UK, bitumen is used only for applications where the carbon is stored. By far the most important of these is the use of bitumen in road dressings. The inventory does assume that a very small proportion of the carbon in the bitumen itself is emitted as VOC during road-stone coating but does not include any estimates of direct carbon emissions from uses of bitumen. Industry consultation in 2013 (UK Petroleum Industries Association, 2013; Refined Bitumen Association, 2013) has confirmed that there are no emissive applications of bitumen in the UK. Around 85% of bitumen is used in road paving, with the remaining proportion used almost entirely in the manufacture of weather-proofing materials.

A 3.1.3.4 Coal Oils and Tars

Coal-tars and benzoles are by-products of coke ovens. Within the UK energy balance published by DECC (DECC, 2013), these fuels are reported as used in combustion applications. However, consultation with the operators of coal ovens (Tata, 2013) and also the UK company that refines and processes coal tars and benzoles (Koppers UK, 2013) has confirmed that all of these materials are collected, refined and processed into a range of products that are not used as fuels. The carbon within coal tars and oils are entirely used within chemical processes. In some cases, the carbon is processed into anodes used in the ferrous and non-ferrous metals industries and then used (in the UK and overseas) within emissive applications. The UK inventory already includes estimates of emissions from UK consumption of carbon anodes within these industries, using methods based on UK metal production statistics.

Based on the evidence from process operators, the Inventory Agency deviates from the UK energy balance data and re-allocates all of the reported coal tars and oils to Non Energy Use, i.e. assuming that all carbon is stored and there are no GHG emissions from this source-activity.

In the reporting of the **Reference Approach** to the UK inventory, we assume the IPCC default assumption that 6% of coking coal is transformed into coal oils and tars, and then the carbon stored is recorded in the coking coal line of table 1A(b) and under “Coal oils and tars (from coking coal)” in table 1A(d).

Coal-tar pitch is used in the manufacture of electrodes, together with petroleum coke and a proportion of the carbon ultimately emitted, but details of input materials are scarce; emissions of carbon from these sources are included in the inventory attributed to petroleum coke. This may introduce a small mis-allocation of emissions between petroleum coke and coal oils and tars, but does not affect the UK inventory emissions total.

A 3.1.3.5 Natural Gas

Natural gas is used as a chemical feedstock for the manufacture of ammonia and methanol. Emissions either occur directly as a result of the manufacturing process or (in the case of methanol) are assumed to ultimately occur upon degradation of the chemical products. The emissions are reported under 2B1.

Most of the emissions from feedstock use of natural gas in ammonia production are at source, i.e. waste gases containing carbon are emitted directly from the ammonia plant. However, in keeping with IPCC guidance for the sector, the emissions also include carbon sequestered in methanol, which in the UK is produced as a by-product using the ammonia process CO₂ emissions. The methanol produced using carbon derived from natural gas feedstock will be

used for various applications, including in consumer products such as antifreeze and screen-wash, as well as a raw material for petrochemical manufacture. Further CO₂ is captured and sold for use elsewhere, for example, in carbonated drinks. This CO₂ is also assumed all to be emitted in the UK.

In the reporting of the **Reference Approach**, the reported carbon stored fraction has been revised to 100%, following feedback in the 2013 UNFCCC centralised review of the UK inventory. The ERT highlighted that as the comparison of the Reference Approach and Sectoral Approach is intended as a verification step for sources in 1A only, that the reporting of gas activity in 2B1 should not be considered in deriving the carbon storage fraction.

An overview of the time series of gas leak at point of use estimates in the UK, together with overall gas use by economic sector and appliance type is presented in **Table A 3.1.5** below.

Table A 3.1.5 Activity data and methane leakage estimates for Gas leakage at Point of Use, including cooking appliances, gas fires and boilers

Source / Appliance type	Units	1990	1995	2000	2005	2010	2012	2013
Annual Gas Use								
Domestic gas fires	ktoe	462	520	621	650	673	573	572
Domestic manual ignition hobs / cookers	ktoe	590	530	511	496	444	455	455
Domestic auto-ignition hobs / cookers	ktoe	211	190	183	177	159	163	163
Domestic auto-ignition space and water heating	ktoe	24572	26796	30491	31512	32223	28480	28433
Service sector catering (ovens and hobs)	ktoe	601	769	779	781	819	804	859
Other service sector appliances (boilers)	ktoe	6642	8502	9812	9283	9738	9530	10069
Methane Leakage								
Domestic cooking and gas fires	ktCH ₄	1.02	0.94	0.94	0.92	0.85	0.84	0.84
Domestic boilers and water heating	ktCH ₄	0.76	0.83	0.94	0.98	1.00	0.88	0.88
Service sector (all sources)	ktCH ₄	0.83	1.06	1.19	1.14	1.19	1.17	1.24

Source / Appliance type	Units	1990	1995	2000	2005	2010	2012	2013
Total	ktCH₄	2.61	2.83	3.07	3.03	3.04	2.89	2.96

A 3.1.3.6 Other Oil (industrial spirit, white spirit, petroleum wax, miscellaneous products)

White Spirit and Special Boiling Point (SBP) spirits are used exclusively for non-energy applications, and are listed in CRF Table 1.A(d) within the category ‘other oil’. They are used as solvents; SBP spirits are used for industrial applications where quick drying times are needed (e.g. adhesives and other coatings) while white spirit is used as a solvent for decorative paint, as a cleaning solvent and for other applications. Estimates of VOC emissions are included in the UK inventory but no estimates are made of direct emissions of carbon from these products, as they are regarded as “not occurring”.

The only emissions from this group of petroleum feedstock that are included in the UK GHG inventory are the releases of carbon from petroleum waxes which are reported under 2B5. These are accounted for in the UK inventory under the fuel category “Other Oils” in CRF Table 1Ad.

A 3.1.3.7 Petroleum Coke

The evidence from industrial reporting of fuel use and from periodic surveys of fuel producers that use petroleum coke to produce domestic fuels (including smokeless fuels) indicates that the allocation of petroleum coke to combustion activities in the UK energy balance is an underestimate across all years. Therefore, the Inventory Agency generates revised estimates for all combustion activities and effectively re-allocates some of the petroleum coke reported in DUKES as non-energy use to energy-related emission sources in the UK inventory.

Within the UK inventory, petroleum coke is included for the following energy and non-energy source categories:

- 1A1a: Power station use of petroleum coke, primarily within blends with coal at a small number of UK facilities;
- 1A1b: Refinery emissions from regeneration of catalysts;
- 1A2f: Cement industry use of petroleum coke as a fuel;
- 1A2g: Other industry use of petroleum coke as a fuel;
- 1A4b: Petroleum coke use within domestic fuels;
- 2C1: Carbon emissions from electrodes used in electric arc furnaces and ladle arc furnaces;
- 2C3: Carbon emissions from anode use in primary aluminium production.

The UK energy balance tables in DUKES contain data on the energy use in power stations (1A1a) and refineries (1A1b), although the former are only available for 2007 onwards, and both sets of data do not always agree with the available activity data from EU ETS. The remaining energy uses in other industrial combustion (1A2f) and the domestic sector (1A4b)

are not included in DUKES. The UK Inventory Agency therefore makes independent estimates of the consumption of petroleum coke in all of these sectors.

Petroleum coke is burnt in **cement kilns** (1A2f) and in a handful of **power stations** (1A1a). A few other **large industrial sites** (also 1A2f) have also used the fuel. Good estimates of the consumption of petroleum coke by these large sites are available from the operators themselves, from trade associations and from EU ETS data (from 2005 onwards).

Fuel grade petroleum coke is also used as a **domestic fuel** (both smokeless and non-smokeless types, reported in 1A4b). The Inventory Agency uses data supplied by the UK fuel supply industry to estimate petroleum coke consumption for domestic fuels over the period 1990 to 2012; these estimates are broadly consistent with fuel use data published in earlier editions of DUKES for a few years in the late 1990s.

Carbon deposits build up with time on catalysts used in **refinery** processes such as catalytic cracking. These deposits need to be burnt off to regenerate the surface area of the catalyst and ensure continued effectiveness of the catalyst; emissions from this process are reported within EU ETS since 2005, with the time series estimates provided by the trade association (UKPIA, 2013) and the catalyst regeneration is treated in the inventory as use of a fuel (since heat from the process is used) and are reported under 1A1b.

The remaining 'non-energy' consumption, following subtraction of these estimates from the DUKES UK supply total, is the best estimate for consumption of petroleum coke for non-energy uses, and these data are reported within the CRF table 1Ad.

Estimates of carbon released from electrodes and anodes during **metal processes** are estimated based on operator data and reported in 2C1 and 2C3. Petroleum coke content of these electrodes and anodes is estimated based on operator data and literature sources such as BREF notes.

The consumption estimates for industrial users of petcoke as a fuel or in anode use are associated with low uncertainty as they are primarily based on operator reported data within the EU ETS or other regulatory reporting mechanisms. Whilst it is conceivable that other sectors may also use petroleum coke as a fuel, there is no evidence from resources such as EU ETS and Climate Change Agreement reporting that this is the case in the UK. The estimates of petroleum coke used to generate fuels for the domestic sector are associated with higher uncertainty as they are based on periodic consultation with fuel suppliers to that market, and expert judgement of stakeholders.

As well as the total UK supply figure from UK energy statistics, DUKES has data on UK production, imports and exports of petroleum coke, which together provide more information on the nature of the UK consumption of petroleum coke. These data cover three distinct types of petroleum coke – catalyst coke, produced and consumed at refineries only (so no import/export or supply of fuel to other UK sectors), and then two products made in a refinery process known as coking: fuel grade (green) coke and anode-grade coke, with the former being used as a fuel, and the latter being a calcined version of the former, used in various non-energy processes. Consultation with the DECC energy statistics team and the only UK refinery with a coking process (DECC, 2013) has confirmed that the UK produces only anode-grade coke, and exports will also be anode-grade coke, whilst imports will be fuel grade coke for use as a cost-effective fuel source or raw material for production processes under NEU.

Carbon factors for petroleum coke use are derived from industry-specific data (including EU ETS fuel analysis) in the case of cement kilns (MPA, 2013), power stations and other industrial sites (EA, 2013; SEPA, 2013). The petroleum coke factor for refinery consumption is based on trade association analysis conducted as part of the 2004 Carbon Factors Review (UKPIA, 2004) while the factor for domestic consumption is based on compositional analysis of samples of petroleum coke sold as domestic fuels (Loader et al, 2008).

These factors do show quite a large variation from sector to sector: this is primarily a reflection of the different requirements of fuels for different sectors (higher quality, higher carbon for some, less so for others). The highest carbon factor is for ‘petroleum coke’ burnt in sector 1A1b, but this fuel is actually of a different nature from the fuel burnt as petroleum coke in sectors 1A1a, 1A2f and 1A4b. In the case of 1A1b, the fuel is a build-up of carbon on catalysts used in various refinery process units, while in the other three cases, the petroleum coke is a solid by-product of a totally different refinery process (coking) which has different characteristics.

The total emissions and carbon stored from petroleum coke for selected years across the time series are presented below. Note that the data on stored carbon are consistent with those reported as “100% stored” in the CRF:

Table A 3.1.6 Petroleum Coke Carbon Emissions and Carbon Stored across all UK Supply 1990-2012

	Units	1990	1995	2000	2005	2010	2011	2012
Petroleum coke emissions (Sum of: 1A1a, 1A1b, 1A2f, 1A4b, 2C1, 2C3)	kt Carbon	258	489	598	704	420	278	315
Petroleum coke – stored carbon	kt Carbon	350	344	58	43	288	170	220
Storage fraction <u>across all UK supply</u> of petroleum coke	%	58	41	9	6	41	38	41

A 3.1.3.8 Carbon Storage Fractions: Import-Export balance for Carbon-containing Materials

The analysis within the UK energy statistics or GHG inventory compilation system cannot accurately account for the variable (over time) import-export balance of carbon-containing materials in the UK economy. For example, where the Inventory Agency accounts for the carbon emissions from scrap tyres burned in cement kilns, power stations, incinerators and so on within the inventory estimates or from the degradation of plastics or released from detergents and other chemicals, there is no way of tracing the quantity that is derived from imported tyres/plastics/detergents.

The reported estimate of the fate of the reported NEU of fuels from the UK energy balance is based on an assumed “closed system”, whereby we account for all emissions from carbon-containing products and fuel types that are allocated as NEU as if they are derived from the fuel statistics in the UK energy balance. The source of the carbon emitted from feedstock and NEU of fuels will partly be carbon from imported materials, with UK feedstock carbon also exported and emitted elsewhere.

A 3.2 INDUSTRIAL PROCESSES (CRF SECTOR 2)**A 3.2.1 Potential Emissions of Halocarbons and SF₆**

Following current international reporting guidance, the UK now only reports actual emissions of F-gases. Potential emissions are neither estimated nor reported.

A 3.3 AGRICULTURE (CRF SECTOR 3)**A 3.3.1 Enteric Fermentation (3A)****Table A 3.3.1 Livestock Population Data for 2013 by Animal Type^a**

Animal type		Number
Cattle	Dairy cows	1,781,855
	Beef cows	1,611,008
	Dairy heifers	397,568
	Beef heifers	371,808
	Dairy replacements >1 year	530,901
	Beef all others >1 year	2,342,187
	Dairy calves <1 year	501,225
	Beef calves <1 year	2,306,990
Pigs	Sows	355,679
	Gilts	151,528
	Boars	14,915
	Fattening & other pigs 80 - >110 kg	737,331
	Fattening & other pigs 50-80 kg	1,029,924
	Other pigs 20-50 kg	1,224,582
	Pigs < 20 kg	1,365,012
Sheep	Breeding sheep	15,993,518
	Other sheep	482,027
	Lambs < 1 year	16,380,931
	Goats	97,643
	Deer	31,803
	Horses	1,023,615
Poultry	Growing pullets	8,474,482
	Laying fowls	27,526,541

Animal type		Number
	Breeding flock	11,213,186
	Table chicken	104,576,384
	Turkeys	4,125,060
	Total other poultry	6,882,964

^a Data derived as sum of totals for each Devolved Administration (i.e. England, Wales, Scotland and Northern Ireland), obtained from Devolved Administration statistical publications (June survey results: England: <https://www.gov.uk/government/statistical-data-sets/structure-of-the-agricultural-industry-in-england-and-the-uk-at-june> and Jennie Blackburn (DEFRA); Scotland: <http://www.scotland.gov.uk/Resource/0045/00455080.xls>; <http://www.scotland.gov.uk/Resource/0043/00436132.xls> and Graeme Kerr (The Scottish Government); Wales: <http://wales.gov.uk/docs/statistics/2013/131128-survey-agricultural-horticulture-june-2013-en1.xls> and John Bleasdale (Welsh Government); Northern Ireland: <http://www.dardni.gov.uk/index/statistics/crops-livestock-and-labour-numbers/crop-areas-and-production-1981-onwards.htm> and Conor McCormack (DARDNI).

Table A 3.3.2 Trends in Livestock Numbers ('000s) 1990-2013

Year	Dairy cows	Other cattle	Pigs	Sheep	Poultry
1990	2,848	9,344	7,548	44,469	127,952
1991	2,771	9,232	7,695	44,166	140,947
1992	2,683	9,242	7,707	44,540	137,613
1993	2,668	9,183	7,853	44,436	144,171
1994	2,716	9,238	7,892	43,813	140,447
1995	2,603	9,254	7,627	43,304	142,267
1996	2,587	9,452	7,590	42,086	148,936
1997	2,478	9,154	8,072	42,823	179,460
1998	2,439	9,080	8,146	44,471	165,087
1999	2,440	8,983	7,284	44,656	165,157
2000	2,336	8,799	6,482	42,264	169,773
2001	2,251	8,351	5,845	36,716	179,880
2002	2,227	8,118	5,588	35,834	168,996
2003	2,191	8,317	5,046	35,812	178,818
2004	2,129	8,459	5,159	35,817	181,759
2005	2,060	8,380	4,862	35,416	173,909
2006	2,054	8,269	4,933	34,722	173,081
2007	1,954	8,350	4,834	33,946	167,667
2008	1,909	8,198	4,714	33,131	166,200
2009	1,857	8,169	4,724	32,038	159,288

Year	Dairy cows	Other cattle	Pigs	Sheep	Poultry
2010	1,847	8,262	4,468	31,086	163,842
2011	1,814	8,119	4,441	31,634	162,551
2012	1,812	8,089	4,481	32,215	160,061
2013	1,782	8,062	4,879	32,856	162,799

^a Data derived as sum of totals for each Devolved Administration (i.e. England, Wales, Scotland and Northern Ireland), obtained from Devolved Administration statistical publications (June survey results: England: <https://www.gov.uk/government/statistical-data-sets/structure-of-the-agricultural-industry-in-england-and-the-uk-at-june> and Jennie Blackburn (DEFRA); Scotland: <http://www.scotland.gov.uk/Resource/0045/00455080.xls>; <http://www.scotland.gov.uk/Resource/0043/00436132.xls> and Graeme Kerr (The Scottish Government); Wales: <http://wales.gov.uk/docs/statistics/2013/131128-survey-agricultural-horticulture-june-2013-en1.xls> and John Bleasdale (Welsh Government); Northern Ireland: <http://www.dardni.gov.uk/index/statistics/crops-livestock-and-labour-numbers/crop-areas-and-production-1981-onwards.htm> and Conor McCormack (DARDNI).

Table A 3.3.3 Methane Emission Factors for Livestock Emissions for 2013

Animal type		Enteric methane ^a	Methane from manures ^{a, b}
		kg CH ₄ /head/year	kg CH ₄ /head/year
Cattle	Dairy cows ^{e, f}	126.3 ^b	17.1
	Beef cows ^f	94.2 ^b	10.8
	Dairy heifers	57	7.7
	Beef heifers	57	6.7
	Dairy replacements >1 year	57	7.7
	Beef all others >1 year	57	6.6
	Dairy calves <1 year	57	12.7
	Beef calves <1 year	57	9.5
Pigs	Sows	1.5	5.2
	Gilts	1.5	5.2
	Boars	1.5	5.2
	Fattening & other pigs 80 - >110 kg	1.5	5.5
	Fattening & other pigs 50-80 kg	1.5	5.5
	Other pigs 20-50 kg	1.5	5.5
	Pigs <20 kg	1.5	4.5
Sheep	Breeding sheep	8	0.31
	Other sheep	8 ^d	0.31 ^d
	Lambs < 1 year	2.2 ^{c f}	0.08 ^{c f}
Goats		5	0.31

Animal type		Enteric methane ^a kg CH ₄ /head/year	Methane from manures ^{a, b} kg CH ₄ /head/year
Horses		18	1.6
Deer		20.0	0.22
Poultry	Growing pullets	NE	0.028
	Laying fowls	NE	0.028
	Breeding flock	NE	0.029
	Table chicken	NE	0.013
	Turkeys	NE	0.091
	Total other poultry	NE	0.026

^a2006 IPCC guidelines; all manure EF's are tier 2 (with the exception of deer)

^bEmission factor for the year 2013 (with the exception of deer)

^cSneath et al. (1997)

^dFactor quoted assumes animal lives for a year; emission calculation assumes animal lives for 6 months

^e% time spent grazing 48% for dairy cows and 54% for beef cows (linked to NH₃ inventory)

^fFactor quoted assumes animal lives for 8.1 months

Table A 3.3.4 Dairy Cows Tier 2 Methane Emission Factors^a

Year	Average weight of cow (kg) ^b	Average milk yield per dairy cow (litres per annum) ^d	Average fat content (%)	Enteric emission factor (kg CH ₄ /head/y) ^c	Manure emission factor (kg CH ₄ /head/y)
1990	556	5151	4.01	100.9	14.1
1991	556	5133	4.04	100.9	14.1
1992	569	5237	4.06	102.9	14.4
1993	569	5259	4.07	103.2	14.4
1994	565	5300	4.05	103.1	14.3
1995	567	5398	4.05	104.2	14.5
1996	582	5545	4.08	106.8	14.8
1997	584	5790	4.07	109.0	15.1
1998	588	5775	4.07	109.2	15.1
1999	592	5964	4.03	110.9	15.3
2000	595	5979	4.03	111.3	15.3
2001	600	6346	4.01	114.8	15.7
2002	604	6493	3.97	116.1	15.9

Year	Average weight of cow (kg) ^b	Average milk yield per dairy cow (litres per annum) ^d	Average fat content (%)	Enteric emission factor (kg CH ₄ /head/y) ^c	Manure emission factor (kg CH ₄ /head/y)
2003	608	6621	3.96	117.4	16.1
2004	612	6763	4.00	119.3	16.3
2005	616	6986	4.02	121.7	16.6
2006	624	6977	4.04	122.4	16.7
2007	634	6913	4.06	122.7	16.7
2008	627	6943	4.06	123.1	16.7
2009	628	7031	3.99	123.5	16.7
2010	633	7273	3.95	125.1	16.9
2011	630	7528	4.06	127.6	17.3
2012	619	7442	4.07	125.3	17.0
2013	611	7535	4.02	126.3	17.1

^aIn 2013, 48% of animals graze on good quality pasture, rest confined

Gestation period 281 days

Digestible energy 74.52% (to 2 d.p.) (Bruce Cottrill, ADAS, pers. comm.)

Methane conversion rate 6.5%

Ash content of manure 8%

^bCattle Weights provided by Eileen Wall/Tracey Pritchard (SRUC).

^c2006 IPCC guidelines

^dMilk yield from AUK

Table A 3.3.5 Energy and digestibility coefficients of ruminant feeds (dairy cows) (MAFF, 1990)

Feed		GE (MJ/kg ODM)	DE (MJ/kg ODM)	DE/GE (%)	Proportion assumed in average annual diet.	GE (MJ/kg ODM) (weighted)	DE (MJ/kg ODM) (weighted)
Forages	Barley straw	18.4	8.2	44.6			
	Fresh grass (grazed) - all species	18.7	13.8	73.8	0.4	7.48	5.52
	Grass hay	18.4	10.7	58.2			
	Grass silage	19.0	13.6	71.6	0.5	9.5	6.8

Feed		GE (MJ/kg ODM)	DE (MJ/kg ODM)	DE/GE (%)	Proportion assumed in average annual diet.	GE (MJ/kg ODM) (weighted)	DE (MJ/kg ODM) (weighted)
	Grass silage (big bale)	18.9	12.7	67.2			
	Maize silage	18.2	11.7	64.3	0.1	1.82	1.17
	Totals				1.0	18.80	13.49
Concentrate energy feeds	Barley grain	18.4	15.8	85.9	0.38	6.992	6.004
	Citrus pulp	17.5	15.0	85.7			
	Fodder beet (fresh)	16.0	14.1	88.1			
	Maize gluten feed	19.2	15.8	82.3	0.05	0.96	0.79
	Oats	19.6	14.6	74.5			
	Rice bran (extracted)	16.7	NA				
	Sugar beet pulp (molassed)	17.1	15.0	87.7	0.05	0.855	0.75
	Wheat feed	19.1	14.3	74.9	0.1	1.91	1.43
	Wheat grain	18.3	16.4	89.6	0.08	1.464	1.312
Protein feeds	Brewers grains	20.9	NA				
	Cottonseed meal	20.4	13.9	68.1			
	Distillers grains (wheat)	21.7	15.8	72.8			
	Field beans						
	Field peas	18.5	NA				
	Rapeseed meal	19.7	15.2	77.2	0.2	3.94	3.04
	Soya bean meal	19.6	16.0	81.6	0.05	0.98	0.8
	Sunflower meal	19.5	12.8	65.6	0.04	0.78	0.512
Vitamins and minerals		0.0	0.0		0.05	0	0
	Totals				1.00	17.88	14.64

Table A 3.3.6 Calculation of digestibility (dairy cows)^a

1 Total annual energy requirement for 'average' UK dairy cow			
Assumed parameters - 7,000 litres, 600 kg live weight			
	ME _{lactation} (MJ)	36050	- Derived from 'Feed into Milk' (2004)
	ME _{pregnancy} (MJ)	2400	- See also Alderman and Cottrill (1993)
	ME _{maintenance} (MJ)	28760	
	Annual ME requirement (MJ)	67210	Sum of ME _{lactation} + ME _{pregnancy} + ME _{maintenance}
	Annual DE requirement (MJ)	82975	Assuming ME = 0.81 x DE (factor from Alderman, 1982)
2 Energy supplied from concentrate feed			
	Average annual milk yield (litres)	7000	
	Average concentrate use (kg FW/litre)	0.28	From Nix, 2009
	Annual concentrate use (kg FW)	1960	
	Annual concentrate use (kg DM)	1705.2	Assumes DM content of concentrate of 87% (MAFF, 1990)
	GE supplied by concentrates (MJ)	30491	Calculated from values given in MAFF, 1990 (see Table A 3.3.5)
	DE supplied by concentrates (MJ)	24961	Calculated from values given in MAFF, 1990 (see Table A 3.3.5)
3 Remaining energy supplied from forage			
	DE to be supplied by forage (MJ)	58015	
	Forage DM required (Kg)	4301	Proportion concentrate in diet: 28% ^a
	GE supplied by forage (MJ)	80851	Proportion forage in diet: 72% ^a
4 Diet digestibility			
	Total GE intake (MJ)	111341	
	Total DE intake (MJ)	82975	
	Digestibility (ME/GE), %	74.52341	

^aSee explanation of calculations in main chapter section 6.2.2.1

Feed into Milk: Agnew, R. E., Yan, T., France, J., Kebreab, E. and Thomas, C. (2004). Energy requirement and supply. In: ed. C. Thomas, Feed into milk: a new applied feeding system for dairy cows, pp.11-20. Nottingham University Press, Nottingham, UK.

Alderman, G and Cottrill, B (1993) Energy and Protein Requirements of Ruminants: An Advisory Manual prepared by the AFRC Technical Committee on Responses to Nutrients. CAB International.

Alderman G (1982) Comparison of rations calculated in different systems. In: Feed Evaluation and Protein Requirement Systems for Ruminants. Eds. Jarrige R and Alderman G. CEC Luxembourg, pp 238-296.

Nix 2009: Nix, J. (2009). Farm Management Pocketbook 2009.

MAFF (1990), UK Tables of nutritive value and chemical composition of feeding stuffs. Rowett Research Services Ltd, Greenburn Road, Bucksburn, Aberdeen, AB2 9SB, UK

Table A 3.3.7 Parameters used in the calculations of gross energy for dairy cows Tier 2 emission factors

Year	NE _m (Net energy for maintenance), MJ/d (eq. 10.3)	NE _a (Energy to obtain food), MJ/d (eq. 10.4)	NE _l (Net energy for lactation), MJ/d (eq. 10.8)	NE _{pregnancy} (Net energy for pregnancy) MJ/d (eq. 10.13)	REM (Ratio available energy for maintenance in a diet to digestible energy consumed) (eq. 10.14)	REG (Ratio available energy for growth in a diet to digestible energy consumed) (eq. 10.15)	GE (Gross energy intake), MJ/d (eq. 10.16)
1990	44.21	3.58	43.38	3.98	0.54	0.35	236.55
1991	44.17	3.58	43.40	3.98	0.54	0.35	236.49
1992	44.97	3.64	44.39	4.05	0.54	0.35	241.28
1993	44.99	3.65	44.64	4.05	0.54	0.35	241.95
1994	44.71	3.62	44.87	4.02	0.54	0.35	241.72
1995	44.88	3.64	45.70	4.04	0.54	0.35	244.27
1996	45.76	3.71	47.12	4.12	0.54	0.35	250.36
1997	45.84	3.72	49.14	4.13	0.54	0.35	255.63
1998	46.07	3.73	49.02	4.15	0.54	0.35	255.98
1999	46.30	3.75	50.36	4.17	0.54	0.35	259.99
2000	46.53	3.77	50.49	4.19	0.54	0.35	260.97
2001	46.81	3.79	53.41	4.21	0.54	0.35	269.07
2002	47.04	3.81	54.40	4.23	0.54	0.35	272.19
2003	47.27	3.83	55.40	4.25	0.54	0.35	275.33
2004	47.49	3.85	56.88	4.27	0.54	0.35	279.68
2005	47.72	3.87	58.91	4.29	0.54	0.35	285.39
2006	48.18	3.90	58.99	4.34	0.54	0.35	286.93
2007	48.79	3.95	58.60	4.39	0.54	0.35	287.73
2008	48.37	3.92	58.85	4.35	0.54	0.35	287.13

Year	NE _m (Net energy for maintenance), MJ/d (eq. 10.3)	NE _a (Energy to obtain food), MJ/d (eq. 10.4)	NE _l (Net energy for lactation), MJ/d (eq. 10.8)	NE _{pregnancy} (Net energy for pregnancy) MJ/d (eq. 10.13)	REM (Ratio available energy for maintenance in a diet to digestible energy consumed) (eq. 10.14)	REG (Ratio available energy for growth in a diet to digestible energy consumed) (eq. 10.15)	GE (Gross energy intake), MJ/d (eq. 10.16)
2009	48.40	3.92	59.06	4.36	0.54	0.35	287.73
2010	48.74	3.95	60.77	4.39	0.54	0.35	292.97
2011	48.54	3.93	63.81	4.37	0.54	0.35	299.96
2012	47.92	3.88	63.16	4.31	0.54	0.35	296.54
2013	47.45	3.85	63.54	4.27	0.54	0.35	296.12

Table A 3.3.8 Beef Cows Tier 2 Methane Emission Factors^a

Year	Average weight of cow (kg) ^a	Average milk yield per dairy cow (litres per annum) ^b	Average fat content (%)	Enteric emission factor (kg CH ₄ /head/y) ^c	Manure emission factor (kg CH ₄ /head/y)
1990	588	1906	3.6	89.0	10.3
1991	587	1906	3.6	89.0	10.3
1992	601	1906	3.6	90.2	10.4
1993	601	1906	3.6	90.3	10.4
1994	597	1906	3.6	89.8	10.4
1995	599	1906	3.6	90.1	10.4
1996	615	1906	3.6	91.5	10.6
1997	617	1906	3.6	91.6	10.6
1998	621	1906	3.6	92.0	10.6
1999	625	1906	3.6	92.3	10.7
2000	629	1906	3.6	92.7	10.7
2001	634	1906	3.6	93.1	10.7
2002	638	1906	3.6	93.5	10.7
2003	642	1906	3.6	93.9	10.8
2004	646	1906	3.6	94.2	10.8
2005	651	1906	3.6	94.6	10.9
2006	659	1906	3.6	95.3	11.0
2007	670	1906	3.6	96.3	11.1
2008	651	1906	3.6	94.7	10.9

Year	Average weight of cow (kg) ^a	Average milk yield per dairy cow (litres per annum) ^b	Average fat content (%)	Enteric emission factor (kg CH ₄ /head/y) ^c	Manure emission factor (kg CH ₄ /head/y)
2009	659	1906	3.6	95.3	11.0
2010	663	1906	3.6	95.8	11.0
2011	670	1906	3.6	96.2	11.1
2012	672	1906	3.6	96.3	11.1
2013	646	1906	3.6	94.2	10.8

^aA 2008-2012 time series was provided by Eileen Wall and Tracey Pritchard from SRUC. The 1990-2007 time-series was estimated by applying the ratio of the existing UK slaughter data to the dairy and beef live weights provided and the 2013 value was extrapolated.

^bMilk yield was from Energy and protein requirements of ruminants (1993).

Table A 3.3.9 Parameters used in the calculation of gross energy for beef cows Tier 2 emission factors

Year	NE _m (Net energy for maintenance), MJ/d (eq. 10.3)	NE _a (Energy for activity) (eq 10.4)	NE _{pregnancy} (Net energy for pregnancy) MJ/d (eq. 10.13)	REG (Ratio of net energy available for growth in a diet to digestible energy consumed) (eq. 10.15)	GE (Gross energy intake), MJ/d (eq. 10.16)
1990	46.1	4.2	4.1	0.31	208.6
1991	46.0	4.2	4.1	0.31	208.4
1992	46.9	4.3	4.2	0.31	211.4
1993	46.9	4.3	4.2	0.31	211.4
1994	46.6	4.3	4.2	0.31	210.4
1995	46.8	4.3	4.2	0.31	211.0
1996	47.7	4.4	4.3	0.31	214.3
1997	47.8	4.4	4.3	0.31	214.6
1998	48.0	4.4	4.3	0.31	215.4
1999	48.2	4.4	4.3	0.31	216.3
2000	48.5	4.5	4.4	0.31	217.1
2001	48.8	4.5	4.4	0.31	218.2
2002	49.0	4.5	4.4	0.31	219.0
2003	49.3	4.5	4.4	0.31	219.8
2004	49.5	4.6	4.5	0.31	220.7
2005	49.7	4.6	4.5	0.31	221.5
2006	50.2	4.6	4.5	0.31	223.2

Year	NE _m (Net energy for maintenance), MJ/d (eq. 10.3)	NE _a (Energy for activity (eq 10.4)	NE _{pregnancy} (Net energy for pregnancy) MJ/d (eq. 10.13)	REG (Ratio of net energy available for growth in a diet to digestible energy consumed) (eq. 10.15)	GE (Gross energy intake), MJ/d (eq. 10.16)
2007	50.8	4.7	4.6	0.31	225.5
2008	49.7	4.6	4.5	0.31	221.6
2009	50.2	4.6	4.5	0.31	223.2
2010	50.4	4.6	4.5	0.31	224.0
2011	50.8	4.7	4.6	0.31	225.4
2012	51.0	4.7	4.6	0.31	225.9
2013	49.4	4.6	4.4	0.31	220.5

NEI (Net energy for lactation), is 15.2 MJ/d (eq. 10.8)

REM (Ratio of net energy available in a diet for maintenance to digestible energy consumed), is 0.51 (eq. 10.14)

For beef cows digestible energy 65%, Ash content of manure 8.0%, Methane producing capacity of manure 0.17 m³/kg VS, % year spent grazing is 54%

The EF for other cattle is IPCC value of 57 kg CH₄ head⁻¹ yr⁻¹

Calculated following 2006 IPCC guidelines

Methane conversion rate is 6.5%

A 3.3.2 Manure Management (3B)

A 3.3.2.1 Methane emissions from animal manures

The emission factors for methane from manure management were given in **Table A 3.3.8**.

Table A 3.3.10 Methane conversion factors for Manure Management Systems in the UK

Manure Handling System	Methane Conversion Factor % ^a
Liquid - with natural crust cover ^b	10
Liquid - without natural crust cover ^b	17
Daily spread	0.1
Deep bedding	17
Pasture range and paddock	1
Poultry manure - with bedding	1.5
Poultry manure - without bedding	1.5

^a2006 IPCC guidelines

^bFraction of cattle and pig liquid MMS with a natural crust cover is 80% and 20%, respectively.

A 3.3.2.2 Nitrous Oxide emissions from Animal Waste Management Systems
Table A 3.3.11 Nitrogen Excretion Factors, kg N hd⁻¹ year⁻¹ for livestock in the UK (1990-2013)^a

Animal type		1990	1995	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013
Cattle	Dairy cows	97	100	106	110	112	113	115	117	117	117	117	118	121	123	123	124
	Beef cows	79	79	79	79	79	79	79	79	79	79	79	79	79	79	79	79
	Dairy heifers	67	67	67	67	67	67	67	67	67	67	67	67	67	67	67	67
	Beef heifers	56	56	56	56	56	56	56	56	56	56	56	56	56	56	56	56
	Dairy replacements >1 year	53	53	53	53	53	53	53	53	53	53	53	53	53	53	53	53
	Beef all others >1 year	53	53	53	53	53	53	53	53	53	53	53	53	53	53	53	53
	Dairy calves <1 year	38	38	38	38	38	38	38	38	38	38	38	38	38	38	38	38
	Beef calves <1 year	38	38	38	38	38	38	38	38	38	38	38	38	38	38	38	38
Pigs	Sows	23.6	22.5	21.4	21.2	20.9	20.7	20.5	20.1	19.7	19.3	18.9	18.5	18.1	18.1	18.1	18.1
	Gilts	15.5	15.5	15.5	15.5	15.5	15.5	15.5	15.5	15.5	15.5	15.5	15.5	15.5	15.5	15.5	15.5
	Boars	28.8	27.4	26.1	25.8	25.5	25.3	25.0	24.5	23.9	23.4	22.9	22.3	21.8	21.8	21.8	21.8
	Fattening & other pigs 80 - >110 kg	20.2	19.3	18.4	18.2	18.0	17.8	17.6	17.2	16.9	16.5	16.1	15.8	15.4	15.4	15.4	15.4
	Fattening & other pigs 50-80 kg	17.5	16.7	15.9	15.7	15.5	15.4	15.2	14.9	14.6	14.3	13.9	13.6	13.3	13.3	13.3	13.3
	Other pigs 20-50 kg	11.7	11.2	10.6	10.5	10.4	10.3	10.2	10.0	9.8	9.6	9.3	9.1	8.9	8.9	8.9	8.9
	Pigs <20 kg	4.6	4.4	4.2	4.1	4.1	4.0	4.0	3.9	3.8	3.7	3.6	3.5	3.4	3.4	3.4	3.4

Animal type		1990	1995	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013
Sheep	Breeding sheep	9.0	9.0	9.0	9.0	9.0	9.0	9.0	9.0	9.0	9.0	9.0	9.0	9.0	9.0	9.0	9.0
	Other sheep	9.0	9.0	9.0	9.0	9.0	9.0	9.0	9.0	9.0	9.0	9.0	9.0	9.0	9.0	9.0	9.0
	Lambs < 1 year	2.40	2.40	2.40	2.40	2.40	2.40	2.40	2.40	2.40	2.40	2.40	2.40	2.40	2.40	2.40	2.40
Goats		20.6	20.6	20.6	20.6	20.6	20.6	20.6	20.6	20.6	20.6	20.6	20.6	20.6	20.6	20.6	20.6
Deer		13	13	13	13	13	13	13	13	13	13	13	13	13	13	13	13
Horses		50	50	50	50	50	50	50	50	50	50	50	50	50	50	50	50
Poultry	Growing pullets	0.42	0.39	0.36	0.36	0.35	0.35	0.34	0.34	0.34	0.34	0.33	0.33	0.33	0.33	0.33	0.33
	Laying fowls	0.89	0.85	0.81	0.80	0.80	0.79	0.78	0.77	0.75	0.74	0.72	0.71	0.70	0.70	0.70	0.70
	Breeding flock	1.16	1.13	1.10	1.10	1.09	1.09	1.08	1.07	1.06	1.05	1.04	1.03	1.02	1.02	1.02	1.02
	Table chicken	0.64	0.59	0.55	0.54	0.53	0.52	0.51	0.49	0.47	0.46	0.44	0.42	0.40	0.40	0.40	0.40
	Turkeys	1.50	1.59	1.68	1.70	1.71	1.73	1.75	1.76	1.77	1.79	1.80	1.81	1.82	1.82	1.82	1.82
	Total other poultry	1.30	1.41	1.52	1.54	1.56	1.58	1.60	1.62	1.64	1.66	1.67	1.69	1.71	1.71	1.71	1.71

^aCottrill and Smith, ADAS

^bNex factors do not exclude $Frac_{LossMS}$

Table A 3.3.12 Distribution of Animal Waste Management Systems used for Different Animal types, 2013^a

Animal Type		Liquid System	Daily Spread	Solid storage/Deep litter ^b	Pasture Range and Paddock	Poultry without bedding	Poultry with bedding	Incineration
Cattle	Dairy cows	37.3	4.1	10.4	48.2	NA	NA	NA
	Beef cows	6.3	0.7	28.0	65.0	NA	NA	NA

Animal Type		Liquid System	Daily Spread	Solid storage/Deep litter ^b	Pasture Range and Paddock	Poultry without bedding	Poultry with bedding	Incineration
	Dairy heifers	5.3	0.6	23.5	70.6	NA	NA	NA
	Beef heifers	6.3	0.7	28.0	65.0	NA	NA	NA
	Dairy replacements >1 year	5.3	0.6	23.5	70.6	NA	NA	NA
	Beef all others >1 year	6.3	0.7	28.0	65.0	NA	NA	NA
	Dairy calves <1 year	0.0	0.0	45.8	54.2	NA	NA	NA
	Beef calves <1 year	0.0	0.0	45.8	54.2	NA	NA	NA
Pigs	Sows	37.7	0.0	20.3	42.0	NA	NA	NA
	Gilts	37.7	0.0	20.3	42.0	NA	NA	NA
	Boars	37.7	0.0	20.3	42.0	NA	NA	NA
	Fattening & other pigs 80 - >110 kg	35.3	0.0	62.7	2.0	NA	NA	NA
	Fattening & other pigs 50-80 kg	35.3	0.0	62.7	2.0	NA	NA	NA
	Other pigs 20-50 kg	35.3	0.0	62.7	2.0	NA	NA	NA
	Pigs <20 kg	45.0	0.0	34.0	21.0	NA	NA	NA
Sheep	Breeding sheep	0.0	0.0	4.2	95.8	NA	NA	NA
	Other sheep	0.0	0.0	4.2	95.8	NA	NA	NA
	Lambs < 1 year	0.0	0.0	4.2	95.8	NA	NA	NA
Goats		0.0	0.0	8.2	91.8	NA	NA	NA

Animal Type		Liquid System	Daily Spread	Solid storage/Deep litter ^b	Pasture Range and Paddock	Poultry without bedding	Poultry with bedding	Incineration
Deer		0.0	0.0	24.9	75.1	NA	NA	NA
Horses		0.0	0.0	0.0	100.0	NA	NA	NA
Poultry	Growing pullets	NA	0.0	NA	1.2	0.0	98.8	0.0
	Laying fowls	NA	0.0	NA	8.8	91.2	0.0	0.0
	Breeding flock	NA	0.0	NA	0.2	0.0	99.8	0.0
	Table chicken	NA	0.0	NA	1.0	0.0	68.4	30.7
	Turkeys	NA	0.0	NA	3.6	0.0	96.4	0.0
	Total other poultry	NA	0.0	NA	2.0	0.0	98.0	0.0

^aMisselbrook et al., 2011

^bFarmyard manure

Table A 3.3.13 Nitrous Oxide Emission Factors for Animal Waste Handling Systems^a

Waste Handling System	Emission Factor (EF ₃), kg N ₂ O per kg N excreted
Liquid - with natural crust cover ^d	0.005
Liquid - without natural crust cover ^d	0
Daily Spread ^b	0
Deep bedding	0.01
Grazing (prp) - for cattle (dairy, non-dairy), poultry and pigs	0.02
Grazing (prp) - for sheep and other animals	0.01
Poultry manure - with bedding	0.001
Poultry manure - without bedding	0.001

^a2006 IPCC guidelines

^bReported under Agricultural Soils

^cFraction of cattle and pig liquid MMS with a natural crust cover is 80% and 20%, respectively.

A 3.3.3 Agricultural Soils (3D)

A 3.3.3.1 Inorganic Fertiliser

Table A 3.3.14 Areas of UK Crops and quantities of fertiliser applied for 2013^a

Crop Type	Crop area, ha	Fertiliser, ktN
Winter wheat	1,614,511	295.9
Spring barley	903,171	97.2
Winter barley	309,825	44.2
Oats	176,541	17.4
Rye, triticale & mixed corn	24,214	1.1
Maize	194,477	10.9
Main crop potatoes	139,293	23.6
Sugar beet	116,973	11.0
Oilseed rape	715,218	126.9
Peas (green)	32,005	0.0
Peas (dry)	28,984	0.0

Crop Type	Crop area, ha	Fertiliser, ktN
Beans (human consumption)	4,965	0.0
Beans (animal consumption)	118,315	0.1
Root crops for stock feed	50,656	3.7
Leafy forage crops	4,088	0.2
Other forage crops	21,334	0.9
Vegetable (brassicae)	2,842	0.4
Vegetables (other)	76,134	6.3
Soft fruit	9,580	0.6
Top fruit	23,398	1.4
Hops	0	0
Linseed	34,498	2.9
Other tillage	49,679	0.8
Grass under 5 years	1,389,665	135.2
Permanent grass	5,941,798	338.3

^aData includes England, Wales, Scotland and Northern Ireland, June survey results: England: <https://www.gov.uk/government/statistical-data-sets/structure-of-the-agricultural-industry-in-england-and-the-uk-at-june> and Jennie Blackburn (DEFRA); Scotland: <http://www.scotland.gov.uk/Resource/0045/00455080.xls>; <http://www.scotland.gov.uk/Resource/0043/00436132.xls> and Graeme Kerr (The Scottish Government); Wales: <http://wales.gov.uk/docs/statistics/2013/131128-survey-agricultural-horticulture-june-2013-en1.xls> and John Bleasdale (Welsh Government); Northern Ireland: <http://www.dardni.gov.uk/index/statistics/crops-livestock-and-labour-numbers/crop-areas-and-production-1981-on-wards.htm>, Conor McCormack and Paul Caskie (DARDNI); BSFP (2013). British Survey of Fertiliser Practice: Fertiliser Use on Farm Crops for Crop Year 2013, the BSFP Authority, Peterborough. Data for preceding years comes from earlier versions of the same publication.

Table A 3.3.15 Trends in area grown ('000 ha) and N fertiliser applied (kg/ha) for the major UK crops, 1990-2013^a

Year	Winter wheat		Spring barley		Winter barley		Main crop potatoes		Oilseed rape		Grass leys (<5yrs)		Permanent grassland	
	'000 ha	kg/ha N	'000 ha	kg/ha N	'000 ha	kg/ha N	'000 ha	kg/ha N	'000 ha	kg/ha N	'000 ha	kg/ha N	'000 ha	kg/ha N
1990	2,014	180	635	85	882	146	177	168	390	221	1,606	151	5,503	108
1991	1,980	186	552	88	841	143	177	160	440	205	1,603	156	5,520	107
1992	2,066	183	515	88	784	146	181	162	421	205	1,573	155	5,466	107
1993	1,759	186	518	90	650	150	171	160	377	205	1,567	157	5,455	108
1994	1,811	188	483	89	628	150	165	157	404	205	1,456	159	5,552	111
1995	1,859	196	504	92	689	153	172	156	354	189	1,407	165	5,546	111
1996	1,977	188	518	90	749	147	177	160	357	187	1,396	155	5,516	108
1997	2,034	188	518	90	839	145	166	164	445	187	1,394	152	5,454	105
1998	2,045	181	484	87	769	142	164	173	506	185	1,302	145	5,524	105
1999	1,847	184	631	95	548	148	178	137	417	192	1,226	169	5,608	104
2000	2,086	182	539	101	589	146	166	135	332	187	1,226	140	5,520	96
2001	1,635	193	783	102	462	152	165	131	404	193	1,205	131	5,738	93
2002	1,996	192	555	103	546	157	158	146	357	203	1,243	133	5,670	87
2003	1,836	191	621	104	455	154	145	130	460	201	1,200	126	5,836	84
2004	1,990	187	587	100	420	148	148	131	498	204	1,246	114	5,771	79
2005	1,870	184	553	98	384	148	137	148	588	207	1,193	112	5,860	73
2006	1,836	186	494	98	388	141	140	123	568	185	1,137	101	6,119	72

Year	Winter wheat		Spring barley		Winter barley		Main crop potatoes		Oilseed rape		Grass leys (<5yrs)		Permanent grassland	
	'000 ha	kg/ha N	'000 ha	kg/ha N	'000 ha	kg/ha N	'000 ha	kg/ha N	'000 ha	kg/ha N	'000 ha	kg/ha N	'000 ha	kg/ha N
2007	1,830	181	515	96	383	139	140	118	674	183	1,176	95	6,111	64
2008	2,080	173	616	92	416	137	144	144	598	178	1,141	90	6,183	53
2009	1,814	183	749	100	411	142	147	152	581	176	1,262	86	6,223	57
2010	1,939	178	539	96	382	144	138	123	642	184	1,231	97	6,066	61
2011	1,969	181	611	96	359	144	146	142	705	186	1,278	90	6,018	56
2012	1,992	179	618	96	385	149	149	130	756	183	1,357	91	5,938	55
2013	1,615	177	903	105	310	145	139	159	715	180	1,390	96	5,942	62

^aData includes England, Wales, Scotland and Northern Ireland, June survey results: England: <https://www.gov.uk/government/statistical-data-sets/structure-of-the-agricultural-industry-in-england-and-the-uk-at-june> and Jennie Blackburn (DEFRA); Scotland: <http://www.scotland.gov.uk/Resource/0045/00455080.xls>; <http://www.scotland.gov.uk/Resource/0043/00436132.xls> and Graeme Kerr (The Scottish Government); Wales: <http://wales.gov.uk/docs/statistics/2013/131128-survey-agricultural-horticulture-june-2013-en1.xls> and John Bleasdale (Welsh Government); Northern Ireland: <http://www.dardni.gov.uk/index/statistics/crops-livestock-and-labour-numbers/crop-areas-and-production-1981-onwards.htm>, Conor McCormack and Paul Caskie (DARDNI); BSFP (2013). British Survey of Fertiliser Practice: Fertiliser Use on Farm Crops for Crop Year 2013, the BSFP Authority, Peterborough. Data for preceding years comes from earlier versions of the same publication.

A 3.3.3.2 Organic Fertilizers

See **Table A 3.3.1**, **Table A 3.3.8** and **Table A 3.3.10** for activity data.

A 3.3.3.3 Application of sewage sludge to land**Table A 3.3.16 Nitrous oxide emissions from sewage sludge (kt N₂O/yr)^a**

Year	Application of sewage sludge to land (t DM/yr)	Direct N ₂ O	Indirect N ₂ O from atmospheric deposition	Indirect N ₂ O from leaching and runoff
1990	507,855	0.287	0.057	0.065
1995	546,746	0.309	0.062	0.070
2000	590,160	0.334	0.067	0.075
2001	831,133	0.470	0.094	0.106
2002	896,885	0.507	0.101	0.114
2003	1,059,808	0.600	0.120	0.135
2004	1,118,583	0.633	0.127	0.142
2005	1,216,378	0.688	0.138	0.155
2006	1,288,025	0.729	0.146	0.164
2007	1,333,078	0.754	0.151	0.170
2008	1,360,893	0.770	0.154	0.173
2009	1,309,479	0.741	0.148	0.167
2010	1,281,602	0.725	0.145	0.163
2011	1,259,683	0.713	0.143	0.160
2012	1,269,713	0.718	0.144	0.162
2013	1,286,915	0.728	0.146	0.164

^aData includes England, Wales, Scotland and Northern Ireland, see data sources in Waste sector **Section A 3.5.3**

A 3.3.3.4 Grazing Animals

See **Table A 3.3.1**, **Table A 3.3.9** and **Table A 3.3.10** for parameters used in calculations.

A 3.3.3.5 Crop Residues**Table A 3.3.17 Dry Mass Content and Residue Fraction of UK Crops for 2013**

Crop Type ^d	Fraction dry mass ^a	Residue/Crop (R _{AG(T)})	N content of above-ground residues (N _{AG})	N content of below-ground residues (N _{BG})	Ratio of below ground residues to above-ground biomass (RBG-BIO)
Field Beans ^c	0.86	1.1	0.01	0.01	0.19

Crop Type ^d	Fraction dry mass ^a	Residue/Crop (R _{AG(T)})	N content of above-ground residues (N _{AG})	N content of below-ground residues (N _{BG})	Ratio of below ground residues to above-ground biomass (RBG-BIO)
Peas (green)	0.08	1.1	0.008	0.008	0.19
Beans (human consumption)	0.08	1.2	0.008	0.008	0.19
All peas harvested dry	0.86	1.1	0.008	0.008	0.19
Rye, mixed corn, triticale	0.855	1.6	0.005	0.011	0.22
Wheat	0.855	1.3	0.006	0.009	0.24
Oats	0.855	1.3	0.007	0.008	0.25
Barley	0.855	1.2	0.007	0.014	0.22
OSR	0.91	1.2	0.006	0.009	0.22
Linseed	0.91	1.2	0.006	0.009	0.22
Maize	0.5	1	0.006	0.007	0.22
Sugar beet	0.1	0.2	0.016	0.014	0.20
Hops ^b	0.2	1.2	0.015	0.012	0.54
Potatoes	0.2	0.4	0.019	0.014	0.20
Total roots & onions	0.07	1.2	0.016	0.014	0.20
Total brassicas	0.06	1.2	0.015	0.012	0.54
Total others	0.05	1.2	0.015	0.012	0.54

^aBurton (1982), Nix (1997) or AUK

^bHops dry mass from Brewers Licensed Retail Association (1998)

^cField beans dry mass from PGRE (1998)

^dDefault factors for estimation of N added to soils from crop residues taken from Table 11.2, 2006 IPCC guidelines

Table A 3.3.18 Production of UK Crops for 2013^a

Crop Type	Crop production, kt
Wheat	11,921
Barley	7,092
Oats	964
Rye, mixed corn, triticale	114
Maize	5,834

Crop Type	Crop production, kt
Potatoes	5,618
Sugar beet	8,430
Oilseed rape	2,128
Peas (green)	158
Peas (dry)	48
Beans (human consumption)	26
Beans (animal cons)	378
Vegetables (brassicae)	428
Vegetables (other)	1,565
Hops	0
Linseed	62
Other tillage	42

^aData includes England, Wales, Scotland and Northern Ireland (Lindsay Holmes, DEFRA (England & Wales), Nicola Kerr, The Scottish Government and Alison Lambert, Conor McCormack, DARDNI).

A 3.3.3.6 Mineralisation

Mineralised N is reported in **Table A 3.3.19**.

Table A 3.3.19 Mineralised N from soils

N in mineral soils that is mineralized in association with loss of soil C	1990	1995	2000	2005	2009	2010	2011	2012	2013
kt N/yr	39	44	52	73	86	86	87	87	88

A 3.3.3.7 Histosols

Total area 2857 km².

A 3.3.4 Field Burning of Agricultural Residues (3F)**Table A 3.3.20 Constants for crop residue calculations (Tables 2.5, 2.6 & p.11.14 2006GL)**

Description	Value
C _f (value for wheat applied to wheat, barley, oats, linseed)	0.9
EF for burning of agricultural residues - CH ₄ (g/kg/DM burnt)	2.7
EF for burning of agricultural residues - N ₂ O (g/kg/DM burnt)	0.07
FraC _{Renew} (annual crops)	1
FraC _{Renew} (grass <5 yrs)	0.25
FraC _{Remove}	0

A 3.3.4.1 CO₂ emissions from Lime and Urea Application**Table A 3.3.21 CO₂ emissions from the application of lime**

Year	Application of Limestone CaCO ₃ (t /yr)	CO ₂ emissions from limestone (kt)	Application of Dolomite CaMg(CO ₃) ₂ (t /yr)	CO ₂ emissions from dolomite (kt)
1990	2,219,000	976	1,259,000	600
1995	2,441,000	1,074	1,300,000	620
2000	1,338,000	589	670,000	319
2001	1,281,000	564	560,000	267
2002	1,273,000	560	590,000	281
2003	1,619,000	712	690,000	329
2004	1,430,000	629	600,000	286
2005	1,247,000	549	570,000	272
2006	1,358,000	598	580,000	276
2007	1,421,000	625	620,000	296
2008	1,301,000	572	550,000	262
2009	1,591,000	700	580,000	276
2010	1,558,000	686	530,000	253
2011	1,571,000	691	716,000	341
2012	1,362,000	599	485,000	231
2013	1,230,000	541	485,000	231

Table A 3.3.22 CO₂ emissions from the application of urea

Year	Application of urea (kt /yr)	CO ₂ emissions from urea application (t)
1990	526	385,461
1995	282	207,157
2000	176	129,354
2001	230	168,635
2002	318	232,872
2003	234	171,696
2004	344	252,120
2005	311	227,938
2006	307	225,226
2007	445	326,219
2008	276	202,145
2009	404	296,364
2010	418	306,615
2011	478	350,795
2012	444	325,681
2013	334	245,223

A 3.4 LAND USE, LAND USE CHANGE AND FORESTRY (CRF SECTOR 4)

The following section describes in detail the methodology used in the LULUCF sector described in **Chapter 6**.

A 3.4.1 Carbon stock changes due to afforestation and forest management (4A)

A 3.4.1.1 The Forest carbon accounting model CARBINE

Carbon uptake by the forests planted in the UK is calculated by a carbon accounting model, CARBINE, as gains and losses in pools of carbon in standing trees, litter and soil in conifer and broadleaf forests and in harvested wood products. Restocking is assumed in all forests. The method is Tier 3, as defined by IPCC (2006). Matthews *et al.* (2014) gives an overview of the CARBINE model and a comparison of its use in the 1990-2012 LULUCF inventory with the C-Flow model previously used model forest carbon stock changes.

CARBINE uses as input data estimates of stand structure and growth obtained from yield tables that are applied at the stand level (Edwards and Christie, 1981). Stand-level carbon estimates are combined with area/age-class information to estimate forest and national carbon

stocks. CARBINE can be used to estimate historical forest carbon stocks, as well as current and future carbon stocks under different forest area and management scenarios. Using one set of yield tables assumes the same growth rates/patterns occur at any time: historic, current or future. This means that changes that might affect growth rate or form are excluded, such as the improvement of planting material or better site quality. Carbon stock changes are inferred from differences in carbon stock estimates at different times. The model can represent all of the introduced and native plantation and naturally-occurring species relevant to the UK.

The model as used for this inventory consists of three sub-models or ‘compartments’ which estimate carbon stocks in the forest, soil, and wood products. The forest carbon sub-model is further compartmentalised to represent fractions due to tree stems, branches, foliage, and roots. The impact of different forest management regimes can be assessed for the range of tree species, yield classes and management regimes represented in published yield tables (Edwards and Christie, 1981). At present not all of these are implemented in CARBINE. Currently the model contains the tables for 19 different tree species (Norway spruce, Sitka spruce, Scots pine, Corsican pine, Lodgepole pine, European larch, Japanese larch, Douglas fir, Grand fir, Noble fir, Western Red cedar, Western hemlock, Oak, Beech, Nothofagus, Poplar and a combined model table that covers Sycamore, Ash and Birch). Yield tables were extrapolated where necessary to cover longer rotations and management and yield in non-clearfell and un-thinned forests. All areas for a species are assumed to have been planted at the same spacing.

Increases in stemwood volume were based on standard yield tables, with an extrapolation function for stemwood volumes prior to the first table age. The mass of carbon in a forest was calculated from volume by multiplying by species-specific wood density, stem:branch and stem:root mass ratios and the fraction of carbon in wood (0.5 assumed). The values used for these parameters for Sitka spruce (*P. Sitchensis*) are given in **Table A 3.4.1**. Sitka spruce is the most common species in UK forests (c. 30%); parameters for other tree species are given in Matthews *et al.* (2014).

Table A 3.4.1 Main parameters for forest carbon flow model used to estimate carbon uptake by planting of forests of Sitka spruce (*P. Sitchensis*), yield class 12.

	<i>P. sitchensis</i>
Time of maximum mean annual increment (years)	60
Initial spacing (m)	2
First table age	20
Year of first thinning	25
Stemwood density (oven dried tonnes m ⁻³)	0.33
Stemwood conversion loss	10%
% Branchwood left in forest	100%
% Branchwood harvested for fuel	0%
% fuel from bark	30%

	<i>P. sitchensis</i>
% non-fuel products from bark	70%
% small roundwood (underbark) used as fuel	20%
% Pallets and fencing from small roundwood (under bark)	20%
% Paper from small roundwood (under bark)	35%
% Particleboard etc. from small roundwood (under bark)	25%
% Fuel from sawlogs (under bark)	30%
% Pallets and fencing from sawlogs (under bark)	0%
% Particleboard from sawlogs (under bark)	40%
% Structural timber from sawlogs (under bark)	30%
Root:Stem ratio	0.49
Crown:Stem ratio	0.32
Underbark/overbark ratio at 15cm DBH (varies with DBH)	0.9
Ratio of thinned stem volume that is sawlog at 15cm DBH (varies with DBH)	0.05

The forest carbon sub-model also represents accumulation and loss of carbon in dead wood and litter. Tree mortality is accounted for implicitly in the standard Forestry Commission growth and yield tables (Edwards and Christie, 1981), and explicit estimates are included in models for unthinned stands, where mortality levels are high. The annual deadwood volume estimates need to be accumulated over an appropriate period to give the total stem volume in dead wood for a given stand age, to allow for the time taken for dead trees to decay. In the current version of CARBINE, the carbon in standing dead wood at any time step is calculated as a weighted sum of the carbon in trees that have died in the current year and the preceding 33 years. The weighting function has the exponential form

$$w_D(T) = -0.10554 + 1.10554 \times 0.93148T$$

where $w_D(T)$ is the fraction of dead wood remaining and T is the time in years since the material entered the dead wood pool. If $T > 33$ years then $w(T)$ is set to zero.

Root and branch wood volume associated with dead trees is estimated in the same way as for living stem wood.

Standing dead wood is regarded as distinct from other forms of dead wood, which effectively form part of the litter pool. An assumption is made that 5% of the carbon in standing dead wood is transferred to the litter pool; implicitly all other losses as standing deadwood degrades involve oxidation of carbon to the atmosphere as CO_2 .

The other significant input of carbon to the dead wood and litter pool is due to harvesting operations (as part of either thinning or clearfelling). The carbon in roots of harvested trees is assumed to all enter the litter pool. The harvesting of stem wood is assumed to involve a conversion loss equivalent to 10% of standing stem volume, which also enters the litter pool.

It is difficult to make robust assumptions about the fate of branch wood and foliage at time of harvesting. In many situations, this material will be left on site to deteriorate and decay. Sometimes it is possible that branch wood remaining after clearfelling may be deliberately burned. There is also an increasing trend towards active harvesting of branch wood (or at least some proportion of it) to supply biomass to the Energy sector. For this inventory the assumption has been made that none of branch wood is harvested but is left to degrade and decay on site as part of the litter pool.

It is assumed in the CARBINE model that harvested material from thinning and felling is made into wood products. This is described further in **Section A 3.4.11**. The net change in the carbon in this pool of wood products is reported in Category 4G.

Carbon entering the litter pool is assumed to be transferred to the soil or to be released to the atmosphere as CO₂. Emissions to the atmosphere are assumed to follow a Weibull 'hazard' function, such that 50% of the material has been lost after 20 years and 95% of the material has been lost after 40 years. The Weibull function has the form

$$w_L(T) = \exp(-(T/B)^C)$$

where $w_L(T)$ is the fraction of material remaining, T is the time in years since the material entered the litter pool and B and C are constant parameters of the equation. Transfer from the litter pool to soil occurs with 1.6% of the remaining litter being added to the soil carbon pool at each time step.

CARBINE contains a basic soil sub-model to estimate carbon stocks and stock changes in this pool which runs independently of the forest sub-model. Initial soil carbon is estimated based on land use/cover and soil texture (sand, loam, clay and peat). Changes in soil carbon are assumed to take place in response to land-use change and the magnitude and time course are estimated according to soil type (texture) and major land use category. This information is based on RothC, a UK soil carbon model, and published literature (Coleman et al., 1997). The estimates for these soil texture classes are then combined to give estimates for an organic and a mineral soil for conifers. This combination of soil texture classes, and the CARBINE soil carbon model more generally, was parameterised for this inventory to give similar results to the CFlow soil carbon model (UK Greenhouse Gas Inventory, 1990-2011, Annex 3.6).

The wood products sub-model represents wood products as long-lived and short-lived sawn timber, particleboard, paper and fuel (**Section A 3.4.11**). Carbon in harvested stemwood is allocated to these wood product categories using an assortment forecasting model that accounts for variation in product out-turn due to tree species and tree size class distribution at time of harvest (Rollinson and Gay, 1983). Wood products in primary use are assumed to decay over time with no account taken of carbon stocks in landfill or greenhouse gas emissions (due to wood products) from landfill (these are taken into account in the Waste sector).

A 3.4.1.2 Forest activity data: management

The forest data for the inventory have been estimated by using data from the Forestry Commission planting statistics and the National Inventory of Woodlands and Trees. Information from the Sub-compartment database (the Forestry Commission database of information on the growth rate and management of all GB public forest estate; see **Section 7.2.4**) was used to create a distribution of species and yield class. It was assumed that the private sector forests would follow the same distribution.

Management of forests are represented as one of four options: Clearfell with thinnings, clearfell without thinnings, managed but not clearfelled, and not used for timber production. The forests used for timber production have a rotation length assigned. For the clearfell forests restocking occurs after the rotation length. For non-clearfell productive woodlands it is assumed there is a 30 year overlap of restocking and non-restocked trees. It was assumed that the private sector distribution of managed forests would follow the same pattern as for the public forest estate. The percentage of private sector woodlands that are not managed for timber production was estimated separately for conifer and broadleaves using information from the woodland grant scheme (to give areas definitely in production), and by comparing the CARBINE timber production estimates to the timber production statistics. **Table A 3.4.2** shows the forest area by management prescription and **Table A 3.4.3** by rotation for conifers and broadleaves in the 1990-2012 inventory; a breakdown by tree species is given in Matthews *et al.* (2014).

Table A 3.4.2 Forest area by management prescription in the UK (from 1990-2012 inventory)

Type	Forest area (kha) by management prescription			
	No thin, clear-felling	Thinning, clear-felling	Continuous cover	No thinning, no felling (i.e. not under management for production)
Total conifer	520.4	490.8	94.1	433.6
Total broadleaf	85.7	128.2	46.0	751.7
Total forest	606.1	619.1	140.1	1185.2

Table A 3.4.3 Forest area by rotation length for forests managed for production (from 1990-2012 inventory)

Type	Rotation range (years)								
	<30	30-39	40-49	50-59	60-69	70-99	100-119	120-149	150+
Total conifer	37.4	190.4	81.9	289.5	210.4	201.9	40.6	34.0	19.2
Total broadleaf	0.0	0.3	61.3	79.7	0.6	20.9	22.1	0.1	13.3
Total forest	37.4	190.8	143.2	369.2	211.0	222.8	62.7	34.0	32.5

The rotation lengths are based on the time of maximum mean annual stem volume increment. A range of rotations lengths were generated around this value to even out felling events. An assumption was also made that managed but un-thinned Sitka spruce would be on a considerably shorter rotation, as the most likely reason for managed Sitka spruce not being thinned would be the threat of windblow (as Sitka was widely planted on upland sites). This

was necessary for the algorithm to be able to successfully assign areas of forest to planting years.

A 3.4.1.3 Forestry activity data: historical and current afforestation rates

Irrespective of species assumptions, the variation in removals from 1990 to the present is determined by the afforestation rate in earlier decades, the effect this has on the age structure in the present forest estate, and hence the average growth rate. At the current rate of forest expansion removals of atmospheric carbon increased until 2005 and have now started to decrease gradually, reflecting the reduction in afforestation rate after the 1970s. This afforestation is all on ground that has not been wooded for many decades.

A comparison of historical forest census data and the historical annual planting rates has been undertaken. Forest censuses were taken in 1924, 1947, 1965, 1980 and the late 1990s. The latest census (National Forest Inventory) will not be completed until 2014/15. The comparison of data sources showed that discrepancies in annual planting rates and inferred planting/establishment date (from woodland age in the forest census) are due to restocking of older (pre-1920) woodland areas and variations in the harvesting rotations. However, there is also evidence of shortened conifer rotations in some decades and transfer of woodland between broadleaved categories (e.g. between coppice and high forest). It is difficult to incorporate non-standard management in older conifer forests and broadleaved forests into the Inventory because it is not known whether these forests are on their first rotation or subsequent rotations (which would affect carbon stock changes, particularly in soils). For this inventory submission the assumption was made that any forest area in the inventory planted post-1920 that is not in the new planting record must be restocking. Forests planted pre-1920 were assumed to have been on land that has been forest since time immemorial. In both cases the forest area was assumed to have been restocked twice and had been managed in the same fashion and on the same rotation. The only exception to this was where the data indicated that some of the forest area that had been felled in the past was no longer used for timber production. It was assumed that forest had previously been felled on a rotation was harvested at the time of maximum mean annual increment.

Where the annual area of new planting was greater than the area implied by the inventory, it was assumed to have been restocked. This gave an indication of the maximum length of some of the rotations that had been applied. As an example, if the area in the inventory for 35 years before the base year is 2kha and the new planting record indicates that 3kha were afforested that year, then 1kha of the new planting must have been restocked. The rotation length for this area must also be a maximum of 35 years, otherwise it would not have been felled.

The planting data used as input to the CARBINE model comes from national planting statistics from 1921 to the present (provided by the Forestry Commission for England, Scotland and Wales and from 1900 to the present by the Northern Ireland Forest Service), and estimates of historical planting year (from analysis of the 2000 National Inventory of Woodland and Trees). We assumed that the National Inventory of Woodlands and Trees survey gives a distribution of all the forest area by broad age classes for a base year of 2000, classified by conifers/broadleaves and by country. An algorithm was used to obtain the area of woodland planted pre-1920 by removing the area of new planting from the age class distribution. The species were then allocated to this "residual distribution" by starting in the base year of 2000 and allocating the shortest rotations first. Forests planted pre-1920 were assumed to have been on land that has been forest since time immemorial. The planting years for all dates pre-1921 are purely notional, and were determined from a combination of the age of the forest

areas and the assumed rotation. We assume that the trees were planted (or regenerated) one rotation ago, two rotations ago, or more. In practice, we assume they were planted (or regenerated) several rotations ago. This approach was undertaken to spin up the model in terms of soil and litter. The new planting records were assigned based on the percentage of area previous allocated to each species and management (by country). Conifer planting on organic soil is a subset of total conifer planting. All broadleaf planting is assumed to be on non-organic soil. The afforestation rates for each planting type in the UK have been calculated from the planting data and are shown in **Table A 3.4.4**.

Table A 3.4.4 Afforestation rate of conifers and broadleaves in the United Kingdom since 1500 (using pre-1920 estimates from woodland age and post-1921 planting data)

Period	Planting rate (k ha a ⁻¹)		
	Conifers on all soil types	Conifers on organic soil	Broadleaves
1501-1600	0.01	0.00	0.00
1601-1700	0.14	0.00	0.41
1701-1750	0.50	0.00	3.50
1751-1800	0.91	0.00	6.29
1801-1850	1.04	0.00	2.55
1851-1900	0.88	0.00	1.92
1901-1910	0.58	0.00	0.79
1911-1920	0.34	0.00	0.25
1921-1930	5.45	0.53	2.44
1931-1940	7.46	0.73	2.13
1941-1950	7.43	0.91	2.22
1951-1960	21.66	3.29	3.09
1961-1970	30.08	5.57	2.55
1971-1980	31.92	6.94	1.12
1981-1990	22.69	5.31	2.12
1991	13.46	3.27	6.84
1992	11.56	2.75	6.60
1993	10.08	2.47	9.08
1994	7.35	1.83	11.34


Period	Planting rate (k ha a ⁻¹)		
	Conifers on all soil types	Conifers on organic soil	Broadleaves
1995	9.45	2.19	10.80
1996	7.42	1.78	9.26
1997	7.72	1.75	9.87
1998	6.98	1.50	9.92
1999	6.63	1.39	10.36
2000	6.52	1.32	11.37
2001	4.90	0.94	13.84
2002	3.89	0.73	10.51
2003	3.75	0.69	9.93
2004	2.92	0.54	9.44
2005	2.10	0.35	9.87
2006	1.14	0.19	7.66
2007	2.13	0.33	8.69
2008	0.85	0.14	6.67
2009	1.21	0.17	5.22
2010	0.54	0.07	4.90
2011	1.55	0.19	6.64
2012	3.45	0.46	9.25
2013	1.86	0.35	8.92

The proportion of forest planting on mineral and organic soils was re-assessed in 2012, as part of the work to estimate N₂O emissions due to drainage on forest soils (Yamulki *et al.* 2012). This work is described below.

A 3.4.1.4 Allocation of CARBINE outputs to UNFCCC inventory sub-categories

The CARBINE model output was post-processed using the IPCC default 20-year transition period for land converted to forest to move into the Forest remaining Forest category. The area within the Land converted to Forest Land sub-category is split between cropland, pasture grassland, semi-natural grassland, settlement and other areas. Pasture grassland and semi-natural grassland are then combined for Grassland reporting in the CRF. This split is based on the relative proportions of historical land use change from these categories to forest. The proportions for each country change over time because the 20-year transition period has a different start date for each inventory year (**Figure A 3.4.1**).

Figure A 3.4.1 Proportional land use contribution to forest conversion in the UK in the previous 20 years, 1990-2013



The area and carbon stock changes in the Forest remaining Forest category are adjusted to take account of losses of forest converted to other land use categories, as these losses, in their entirety, are not reflected in the statistics published by the Forestry Commission. Implied carbon stock changes per unit area are calculated using the unadjusted forest area and carbon stock changes. The forest area is then adjusted to reflect losses due to forest conversion and multiplied by the implied carbon stock change to obtain the adjusted carbon stock change.

The CARBINE model has not yet been implemented for forest in the Isle of Man and Guernsey (Crown Dependencies of the UK) and instead the CFlow model is used as it was in previous submissions (UK Greenhouse Gas Inventory, 1990-2011, Annex 3.6).

A 3.4.1.5 Nitrogen fertilization of forest land

Nitrogen fertilization of forest land is assumed to occur only when absolutely necessary, i.e. new planting on ‘poor’ soils (slag heaps, impoverished brown field sites, or upland organic soils). In terms of the inventory, this means that N fertilisation is assumed for Settlement converted to Forest land and Grassland converted to Forest Land on organic soils. The areas of new planting with these conditions were taken from the same dataset used in the CARBINE model (see **Table A 3.4.4**) for 4.A.2. Land converted to Forest land.

Where fertilisation occurs, an application rate of 150 kg N ha⁻¹ is assumed based on Forestry Commission fertilisation guidelines (Taylor 1991). The guidelines recommend applying fertiliser on a three-year cycle until canopy closure (at c. 10 years), but this is thought to be rather high (Skiba 2007) and unlikely to occur in reality, so two applications are adopted as a compromise. These applications occur in year 1 and year 4 after planting. The emission factor for N₂O of applied nitrogen fertiliser is the default value of 1% used in the IPCC 2006


Guidelines. Emissions of N₂O from N fertilisation of forests have fallen since 1990 due to reduced rates of new forest planting. The GWP of N₂O has been changed to 298 in line with the 2006 Guidelines.

A 3.4.1.6 Emissions from drainage on forest soils

Work on developing this method was undertaken by Forest Research in 2012 (Yamulki *et al.* 2012), using new GIS data on forest planting in England, Wales and Scotland. Comparable data were not available for Northern Ireland. This method was described in the 1990-2012 National Inventory Report (Webb *et al.* 2014).

For the 1990-2013 inventory the method has been modified to use the same data on forest planting on mineral and organic soils as is used by the CARBINE model for the calculation of carbon stock changes. It is assumed that only forests planted since 1920 have been drained. The areas of forest planted on mineral soil, nutrient-rich organic soil, nutrient-poor organic soils, nutrient-rich organo-mineral soils and nutrient-poor organo-mineral soils (Figure A 3.4.2) are estimated based on the proportion of forest cover on different soil types (Yamulki *et al.* 2012), adjusted by the amount of forest planted since 1920. The area of forest on mineral soil is adjusted further by splitting it between free-draining mineral soils and imperfectly draining (easily waterlogged) mineral soils, which require artificial drainage (based on the current guidance and policy for forest operations and management). The proportion of mineral soils requiring artificial drainage is: 34% in England, 24% in Scotland, 3% in Wales, 68% in Northern Ireland and 26% in the UK as a whole. We assumed all forest on organic and organo-mineral soils is cultivated prior to planting and therefore effectively drained.

Figure A 3.4.2 Area of forest on different drained soil types in the UK 1990-2013



N₂O emissions are estimated using the Tier 1 methodology and the IPCC default emission factors for drained mineral (0.06 kg N₂O-N ha⁻¹ yr⁻¹), nutrient-rich organic (0.6 kg N₂O-N ha⁻¹ yr⁻¹) and nutrient-poor organic soils (0.1 kg N₂O-N ha⁻¹ yr⁻¹) (IPCC, 2006).

A 3.4.2 Land Use Change and Soils (4B, 4C, 4E)

Changes in soil carbon due to land use change are modelled with a dynamic model of carbon stock change which is driven by matrices of change calculated from land surveys.

A 3.4.2.1 Land Use Change Matrices

For Great Britain (England, Scotland and Wales), matrices from the Monitoring Landscape Change (MLC) data from 1947 & 1980 (MLC 1986) and the Countryside Surveys (CS) of 1984, 1990, 1998 (Haines-Young *et al.* 2000) and 2007 (Smart *et al.* 2009) are used.

In Northern Ireland, matrices were calculated from the Northern Ireland Countryside Surveys of 1990, 1998 (Cooper and McCann 2002) and 2007 (Cooper, McCann and Rogers 2009). The only data available for Northern Ireland pre-1990 is land use areas from The Agricultural Census and The Forest Service (Cruickshank and Tomlinson 2000). Matrices of land use change were estimated for 1970-79 and 1980-89 using area data. The relationship between the matrix of land use transitions and initial area from recent Countryside Surveys is assumed to be the same as the relationship between the matrix and area data for each of the earlier periods – 1970-79 and 1980-89. The matrices developed in this approach were used to extrapolate areas of land use transition back to 1950 to match the start year in the rest of the UK.

The AFOLU Guidance (IPCC 2006) recommends use of six types of land for descriptive purposes: Forest, Grassland, Cropland, Settlements, Wetlands and Other Land. Areas undergoing active commercial peat extraction and areas of inland water and flooded land are reported under Wetlands in the current inventory, and the remaining land in the UK has been placed into the five other types. The more detailed habitats for the two surveys in Great Britain were combined as shown in **Table A 3.4.5** for the Monitoring Landscape Change dataset and **Table A 3.4.6** for the Countryside Survey Broad Habitats (Jackson, 2000).

Table A 3.4.5 Grouping of MLC land cover types for soil carbon change modelling

CROPLAND	GRASSLAND	FORESTLAND	SETTLEMENTS (URBAN)	OTHER
Crops	Upland heath	Broadleaved wood	Built up	Bare rock
Market garden	Upland smooth grass	Conifer wood	Urban open	Sand/shingle
	Upland coarse grass	Mixed wood	Transport	Inland water
	Blanket bog	Orchards	Mineral workings	Coastal water
	Bracken		Derelict	
	Lowland rough grass			

CROPLAND	GRASSLAND	FORESTLAND	SETTLEMENTS (URBAN)	OTHER
	Lowland heather			
	Gorse			
	Neglected grassland			
	Marsh			
	Improved grassland			
	Rough pasture			
	Peat bog			
	Fresh Marsh			
	Salt Marsh			

Table A 3.4.6 Grouping of Countryside Survey Broad Habitat types (Jackson, 2000) for soil carbon change modelling

CROPLAND	GRASSLAND	FORESTLAND	SETTLEMENTS (URBAN)	OTHER
Arable and horticulture	Improved grassland	Broadleaved, mixed and yew woodland	Built up areas and gardens	Inland rock
	Neutral grassland	Coniferous woodland	Unsurveyed urban land	Supra littoral rock
	Calcareous grassland		Boundary and linear features	Littoral rock
	Acid grassland			Standing open water and canals
	Bracken			Rivers and streams
	Dwarf shrub heath			Sea
	Fen, marsh, swamp			
	Bogs			
	Montane			
	Supra littoral sediment			
	Littoral sediment			

The area data used between 1947 and 2007 are shown in **Table A 3.4.7** and **Table A 3.4.8**. The land use change data over the different periods were used to estimate annual changes by

assuming that these were uniform across the measurement period. The full set of annual land use change matrices 1990-2013 are provided in Table 4.1 in the Common Reporting Format tables.

Table A 3.4.7 Sources of land use change data in Great Britain for different periods in estimation of changes in soil carbon

Year or Period	Method	Change matrix data
1950-1979	Measured LUC matrix	MLC 1947->MLC1980
1980 - 1984	Interpolated	CS1984->CS1990
1984 - 1989	Measured LUC matrix	CS1984->CS1990
1990 - 1998	Measured LUC matrix	CS1990->CS1998
1999-2007	Measured LUC matrix	CS1998->CS2007
2008-2013	<i>Extrapolated</i>	CS1998->CS2007

Table A 3.4.8 Sources of land use change data in Northern Ireland for different periods in estimation of changes in soil carbon.

Year or Period	Method	Change matrix data
1950 – 1969	Extrapolation and ratio method	NICS1990->NICS1998
1970 – 1989	Land use areas and ratio method	NICS1990->NICS1998
1990 – 1998	Measured LUC matrix	NICS1990->NICS1998
1999-2007	Measured LUC matrix	NICS1998->NICS2007
2008-2013	<i>Extrapolated</i>	NICS1998->NICS2007


NICS = Northern Ireland Countryside Survey

The transitions between habitat types in the Countryside Surveys for the latest survey (2007) were calculated with Geographical Information System software (ArcGIS). We identified 544 Countryside Survey squares of Great Britain that coincided between the 1998 and 2007 surveys. Survey square locations are confidential. For each coincident square, we calculated the area that changed from one habitat type in 1998 to another in 2007. There are 47 broad habitats described by the Countryside Survey. Individual surveyed squares contain a subset of these habitats and changes between habitats are called transitions. Each coincident survey square also has a ‘land class’ assigned to it that does not change between survey years. There are currently 45 land classes in the Land Classification of Great Britain. Land classes represent the stratification of environments across the UK. A simplified picture of the stratification is shown in **Figure A 3.4.3**. Northern Ireland is treated as a single uniform unit as its smaller area means that it does not display the climatic or elevation variations evident across the rest of the UK.

Transitions between broad habitats were grouped by land class. The ratio of the total area of each land class to the total area sampled within each land class is calculated so that the transitions can be up-scaled to the land class areas. Transitions can then be extracted at

various scales i.e. UK or Devolved Authorities scale or 20 km by 20 km squares. These scales are required by the soil carbon and non-forest biomass models.

Figure A 3.4.3 Stratification of environments across the UK with areas 1 to 6 based on the underlying Land Classification (45 classes).



A 3.4.2.2 Soils modelling

A database of soil carbon density for the UK (Milne & Brown 1997, Cruickshank *et al.* 1998, Bradley *et al.* 2005) is used in conjunction with the land use change matrices. There are three soil survey groups covering the UK and the field data, soil classifications and laboratory methods have been harmonized to reduce uncertainty in the final joint database. The depth of soil considered was also restricted to 1 m at maximum as part of this process. **Table A 3.4.9** shows total stock of soil carbon (1990) for different land types in the four devolved areas of the UK.

Table A 3.4.9 Soil carbon stock (TgC = MtC) for depths to 1 m in different land types in the UK

Region Type	England	Scotland	Wales	N. Ireland	UK
Forestland	108	295	45	20	467
Grassland	995	2,349	283	242	3,870
Cropland	583	114	8	33	738
Settlements	54	10	3	1	69
Other	0	0	0	0	-
TOTAL	1,740	2,768	340	296	5,144

The dynamic model of carbon stock change requires the change in equilibrium carbon density from the initial to the final land use. The core equation describing changes in soil carbon with time for any land use transition is:

$$C_t = C_f - (C_f - C_0)e^{-kt}$$

where

C_t is carbon density at time t

C_0 is carbon density initial land use

C_f is carbon density after change to new land use

k is time constant of change

By differentiating we obtain the equation for flux f_t (emission or removal) per unit area:

$$f_t = k(C_f - C_0)e^{-kt}$$

From this equation we obtain, for any inventory year, the land use change effects from any specific year in the past. If A_T is area in a particular land use transition in year T considered from 1950 onwards then total carbon lost or gained in an inventory year, e.g. 1990, is given by:

$$F_{1990} = \sum_{T=1950}^{t=1990} kA_T (C_f - C_0)(e^{-k(1990-T)})$$

This equation is used with k , A_T and $(C_f - C_0)$ chosen by Monte Carlo methods within ranges set by prior knowledge, e.g. literature, soil carbon database, agricultural census, LUC matrices.

In the model, we calculate the change in equilibrium carbon density from the initial to the final land use during a transition. These are calculated for each land use category as averages for Scotland, England, Wales and Northern Ireland. These averages are weighted by the area of Land Use Change occurring in four broad soil groups (organic, organo-mineral, mineral, unclassified) in order to account for the actual carbon density where change has occurred. In

the UK land use change other than afforestation generally does not occur on organic soils. Changes on soil carbon stock on afforested land are modelled using the CARBINE model rather than the exponential loss model above. Other areas of land use change on organic soils are believed to be very small and are currently not separated out from change on mineral soils.

Hence mean soil carbon density change is calculated as:

$$\bar{C}_{ijc} = \frac{\sum_{s=1}^6 (C_{sijc} L_{sijc})}{\sum_{s=1}^6 L_{sijc}}$$

This is the weighted mean, for each country, of change in equilibrium soil carbon when land use changes, where:

i = initial land use (Forestland, Grassland, Cropland, Settlements)

j = new land use (Forestland, Grassland, Cropland, Settlements)

c = country (Scotland, England, N. Ireland & Wales)

s = soil group (organic, organo-mineral, mineral, unclassified)

C_{sijc} is change in equilibrium soil carbon for a specific land use transition

The land use data (1990 to 1998) is used in the weighting (this will be updated). The averages calculated are presented in **Table A 3.4.10-Table A 3.4.13**.

Table A 3.4.10 Weighted average change in equilibrium soil carbon density (t ha⁻¹) to 1 m deep for changes between different land types in England

To \ From	Forestland	Grassland	Cropland	Settlements
Forestland	0	25	32	83
Grassland	-21	0	23	79
Cropland	-31	-23	0	52
Settlements	-87	-76	-54	0

Table A 3.4.11 Weighted average change in equilibrium soil carbon density (t ha⁻¹) to 1 m deep for changes between different land types in Scotland

To \ From	Forestland	Grassland	Cropland	Settlements
Forestland	0	47	158	246
Grassland	-52	0	88	189
Cropland	-165	-90	0	96
Settlements	-253	-187	-67	0

Table A 3.4.12 Weighted average change in equilibrium soil carbon density (t ha⁻¹) to 1 m deep for changes between different land types in Wales

To \ From	Forestland	Grassland	Cropland	Settlements
Forestland	0	23	57	114
Grassland	-18	0	36	101
Cropland	-53	-38	0	48
Settlements	-110	-95	-73	0

Table A 3.4.13 Weighted average change in equilibrium soil carbon density (t ha⁻¹) to 1 m deep for changes between different land types in Northern Ireland

To \ From	Forestland	Grassland	Cropland	Settlements
Forestland	0	94	168	244
Grassland	-94	0	74	150
Cropland	-168	-74	0	76
Settlements	-244	-150	-76	0

The rate of loss or gain of carbon is dependent on the type of land use transition (**Table A 3.4.14**). For transitions where carbon is lost e.g. transition from Grassland to Cropland, a ‘fast’ rate is applied whilst a transition that gains carbon occurs much more slowly. A literature search for information on measured rates of changes of soil carbon due to land use was carried out and ranges of possible times for completion of different transitions were selected, in combination with expert judgement. These are shown in **Table A 3.4.15**.

Table A 3.4.14 Rates of change of soil carbon for land use change transitions.

		Initial			
		Forestland	Grassland	Cropland	Settlement
Final	Forestland		<i>slow</i>	<i>slow</i>	<i>slow</i>
	Grassland	<i>fast</i>		<i>slow</i>	<i>slow</i>
	Cropland	<i>fast</i>	<i>fast</i>		<i>slow</i>
	Settlement	<i>fast</i>	<i>fast</i>	<i>fast</i>	

(“Fast” & “Slow” refer to 99% of change occurring in times shown in **Table A 3.4.15**)

Table A 3.4.15 Range of times for soil carbon to reach 99% of a new value after a change in land use in England (E), Scotland (S) and Wales (W)

	Low (years)	High (years)
Carbon loss ("fast") E, S, W	50	150
Carbon gain ("slow") E, W	100	300
Carbon gain ("slow") S	300	750

Changes in soil carbon from equilibrium to equilibrium ($C_r - C_0$) were assumed to fall within ranges based on 2005 database values for each transition and the uncertainty indicated by this source (up to $\pm 11\%$ of mean). The areas of land use change for each transition were assumed to fall a range of uncertainty of $\pm 30\%$ of mean.

A Monte Carlo approach is used to vary the rate of change, the area activity data and the values for soil carbon equilibrium (under initial and final land use) for all countries in the UK. The model of change was run 1000 times using parameters selected from within the ranges described above. The mean carbon flux for each region resulting from this imposed random variation is reported as the estimate for the Inventory. An adjustment was made to these calculations for each country to remove increases in soil carbon due to afforestation, as the CARBINE model provides a better estimate of these fluxes in the Land Converted to Forest Land category. Variations from year to year in the reported net emissions reflect the trend in land use change as described by the matrices of change.

Change in soil carbon stock due to Cropland Management.

A new methodology has been developed to assess change in soil carbon stocks due to Cropland Management. This implements the findings of Defra project SP1113 (Moxley *et al*, 2014) which reviewed of UK relevant literature on the effects of Cropland management practices on soil carbon stocks and attempted to modelling UK specific emission factors.

Increases in inputs of fertiliser, manure and crop residues were found to increase soil carbon stocks of tillage land, but changes in the tillage regime from conventional tillage to reduced or zero tillage were found to have no significant effect in a UK context.


Using this methodology, tillage crops are divided into Medium and Low residue groups based on the data on total crop biomass. Where land receives inputs of fertiliser or manure the inputs moved up a class (e.g. Cropland producing a Low residue crop which receives manure is considered to receive Medium inputs, while land producing a Medium residue Crop which received manure inputs is considered to receive High inputs). If crop residues are removed from land the input level drops. A decision tree for assess the effect of Cropland management on soil carbon stocks is shown in **Figure A 3.4.4**.

For most Cropland management activities there was insufficient UK field data to develop reliable Tier 2 emission factors, and so Tier 1 factors have been used (for manure and residue inputs, and for differentiating perennial crops, annual crops and set-aside). However, for tillage reduction both a literature review and modelling work suggested that it did not have a significant effect on soil carbon stocks, and that the Tier 1 emission factors over-estimated its effect under UK conditions. Therefore a stock change factor of 1 has been used for tillage reduction.

As change in soil carbon stocks due to changes in Cropland Management are smaller than changes due to land use change the IPCC default transition time of 20 years is used.

Data on the areas under the main crop types is obtained from the annual June Agricultural Censuses carried out by each UK administration (Defra, 2013a; Welsh Government, 2013; Scottish Government 2013; DARDNI, 2013). Data on the areas of Cropland receiving inputs of manure, fertiliser and crop residues is obtained from the annual British Survey of Fertiliser Practice (Defra, 2013b and previous editions).

Figure A 3.4.4 Decision tree for assessing the effects of Cropland Management on soil carbon stocks.



Change in soil carbon stock due to Grassland Management.

Defra project SP1113 attempted to develop a methodology to allow reporting of changes in soil carbon stocks resulting from Grassland Management. Although there is reasonable data

on the effects of management practices such as liming, reseeded and drainage on improved Grassland on mineral soils there is little data on the effect of many management practices if applied to semi-natural grassland or those on organo-mineral or organic soils where there is a risk that more intensive management could increase carbon losses. As semi-natural Grassland makes up a large proportion of Grassland in the UK the lack of field data makes it impossible to reliably report changes soil carbon stocks from Grassland Management. If improved UK-specific field data becomes available it will be possible to include reporting of Grassland Management in the LULUCF inventory.

A 3.4.2.3 Future development

A new vector based approach to tracking land use change is being developed. The starting point will be CARBINE land cover maps from 1990, 2000 and 2007 overlain with the CEH Countryside Survey data and the Forestry Commission National Forest Estate and Woodlands ground-based data. These data together will be used to produce a set of 100 x 100 m resolution maps, where each 100 m square has an associated vector of land use over time. The maps can be aggregated into the set of distinct representative vectors with their corresponding areas. The vector areas can then be calibrated to match the observed time series of the areas of Forest, Urban, Arable, and Grassland reported in national statistics. This will improve the modelling of change in soil carbon stocks resulting from land use change compared to the current statistically based land use change matrices.

By adding spatial data on crop type from the Intergrated Administration and Control System(IACS) dataset used to administer Common Agricultural Policy payments it will be possible to track changes in Cropland Management more fully, and to improve data on grass/crop rotation patterns. This will improve modelling of soil carbon stocks as the current approach assumes that all change is from an equilibrium soil carbon stock, which will not be the case for rotational Grassland.

Although the intention was to use this approach in the 1990 – 2013 inventory, this was delayed (on the advice of the LULUCF scientific steering committee) to allow the inventory team time to implement the changes required for the 2006 AFOLU Guidelines and the move to the new Common Reporting Format software.

A 3.4.3 Changes in stocks of carbon in non-forest biomass due to land use change (4B2, 4C2, 4E2)

Change in biomass carbon stock due to land use change.

Changes in stocks of carbon in biomass due to land use change are based on the same area matrices used for estimating changes in carbon stocks in soils (see previous section). The biomass carbon density for each land type other than Forest is assigned by expert judgement based on the work of Milne and Brown (1997) and these are shown in **Table A 3.4.16**. Five basic land uses were assigned initial biomass carbon densities, and then the relative occurrences of these land uses in the four countries of the UK were used to calculate mean densities for each of the IPCC types, Cropland, Grassland and Settlements.

Living biomass carbon stocks and Dead Organic Matter (DOM) stocks on Forest Land are modelled using CARBINE and used to calculate changes in carbon stocks due to conversions to and from Forest Land. When land is deforested to another land use, it is assumed that all living biomass and DOM is either converted to Harvested Wood Products or burnt on site in

the year in which deforestation takes place. (Under KP-LULUCF reporting all HWP from Article 3.3 Deforestation are assumed to be instantaneously oxidised). Increase in biomass carbon and DOM stocks on afforested land is modelled in CARBINE. Full details of CARBINE modelling of carbon stocks on Forest Land are given in **Annex A 3.4.1.1**.

The mean biomass carbon densities for each land type were further weighted by the relative proportions of change occurring between non-Forest land types (**Table A 3.4.17-Table A 3.4.20**), in the same way as the calculations for changes in soil carbon densities. Changes between these equilibrium biomass carbon densities were assumed to happen in a single year.

Table A 3.4.16 Equilibrium biomass carbon density (kg m⁻²) for different land types

Density (kg m ⁻²)	Scotland	England	Wales	N. Ireland
Arable	0.15	0.15	0.15	0.15
Gardens	0.35	0.35	0.35	0.35
Natural	0.20	0.20	0.20	0.20
Pasture	0.10	0.10	0.10	0.10
Urban	0	0	0	0
Equilibrium biomass carbon density of IPCC types weighted by occurrence (kg m⁻²)				
Cropland	0.15	0.15	0.15	0.15
Grassland	0.18	0.12	0.13	0.12
Settlements	0.29	0.28	0.28	0.26

Biomass carbon stock for the IPCC land use categories were estimated using land cover data from the 2007 Land Cover Map (LCM) and the biomass carbon stocks for each LCM land cover type developed in Milne and Brown (1997). For Settlements LCM was used to assess the proportion of gardens, pasture-type grass (including sports pitches, golf courses and parks) and urban (built over) area within areas identified as Settlements. The biomass carbon estimates for Grassland and Cropland do not include explicit estimates of hedgerow biomass, this will be included in the 1990 – 2014 inventory as a planned improvement to the inventory.

Table A 3.4.17 Weighted average change in equilibrium biomass carbon density (kg m⁻²) for changes between different land types in England

To \ From	Forestland	Grassland	Cropland	Settlements
Forestland				
Grassland		0	0.08	-0.08
Cropland		-0.08	0	-0.13
Settlements		0.08	0.13	0

(Transitions to and from Forestland are considered elsewhere)

Table A 3.4.18 Weighted average change in equilibrium biomass carbon density (kg m⁻²) for changes between different land types in Scotland.

To \ From	Forestland	Grassland	Cropland	Settlements
Forestland				
Grassland		0	0.02	-0.09
Cropland		-0.02	0	-0.14
Settlements		0.09	0.14	0

(Transitions to and from Forestland are considered elsewhere)

Table A 3.4.19 Weighted average change in equilibrium biomass carbon density (kg m⁻²) for changes between different land types in Wales.

To \ From	Forestland	Grassland	Cropland	Settlements
Forestland				
Grassland		0	0.07	-0.08
Cropland		-0.07	0	-0.13
Settlements		0.08	0.13	0

(Transitions to and from Forestland are considered elsewhere)

Table A 3.4.20 Weighted average change in equilibrium biomass carbon density (kg m⁻²) for changes between different land types in Northern Ireland.

To \ From	Forestland	Grassland	Cropland	Settlements
Forestland				
Grassland		0	0.08	-0.06
Cropland		-0.08	0	-0.11
Settlements		0.06	0.11	0

(Transitions to and from Forestland are considered elsewhere)

A 3.4.3.1 Future development

A new vector based approach to tracking land use change is being developed, as described in section 29.1.1.3. This will improve modelling of biomass carbon stock changes resulting from land use change.

An approach to reporting the effect of Cropland and Grassland Management on biomass carbon stocks has been developed which will use Tier 2 stock change factors and should allow the effect of these activities to be included in the 1990 - 2014 inventory.

A 3.4.4 Carbon stock changes and biomass burning emissions due to Deforestation (4B, 4C, 4E, 4G)

Deforestation is an activity that cuts across LULUCF categories, affecting net emissions and removals in all the land use categories except 4D Wetlands. The process of land use change affects carbon stock changes in biomass and soil, and the woody material left after felling either moves into the harvested wood products pool or is assumed to be burnt on-site, resulting in immediate biomass burning emissions.

Levy and Milne (2004) discuss methods for estimating deforestation since 1990 using a number of data sources. Their approach of combining Forestry Commission felling licence data for rural areas with Ordnance Survey data for non-rural areas has been expanded to include new sources of information and to improve coverage of all countries in the UK. Deforestation before 1990 (which contributes to soil carbon stock changes from historical land use change) is estimated from the land use change matrices described in **Section A 3.4.2**. In the 1990-2013 inventory the forestry agencies Forest Research and Forestry Commission have revised the estimates of deforestation for Great Britain under different land types from 2000 onwards.

A 3.4.4.1 Sources of deforestation activity in the UK

In Great Britain, some activities that involve tree felling require permission from the Forestry Commission, in the form of a felling licence, or a felling application within the various forestry grant schemes. Under the Forestry Act 1967, there is a presumption that the felled areas will be restocked, usually by replanting but sometimes by natural regeneration. Thus, in the 1990s, around 14,000 ha yr⁻¹ were felled and restocked. However, some licences are granted without the requirement to restock, where there is good reason – so-called unconditional felling licences. A felling licence is not required only under certain conditions (<http://www.forestry.gov.uk/forestry/INFD-6DFKW6>), e.g. if felling is allowed as part of planning permission (for building work) or for service maintenance (for gas, water, electricity). Most unconditional felling licence applications are for small areas (6.8 ±19.2 ha). Larger area of deforestation may be undertaken for the purposes of renewable energy development (wind farms) or habitat restoration (e.g. peatbog rewetting).

Felling for urban development (with no requirement to restock) can be allowed under planning permission but only local planning authorities hold documentation for this, and the need for collation makes estimating the national total difficult. However, in England, the Ordnance Survey (national mapping agency) makes an annual assessment of land use change from the data it collects for map updating and provides this assessment to the Department of Communities and Local Government (DCLG) (<http://www.communities.gov.uk/planningandbuilding/planningbuilding/planningstatistics/landusechange/>). DCLG provide an extract of this dataset, listing annual land use change from Forest to developed land uses (1990-2008 in the latest submission). This dataset comes from a continuous rolling survey programme, both on the ground and from aerial photography. The changes reported each year may have actually occurred in any of the preceding 1-5 years (the survey frequency varies among areas, and can be up to 10 years for moorland/mountain areas). Consequently, a five-year moving average is applied to the data to smooth out the between-year variation appropriately, to give a suitable estimate with annual resolution.

The Countryside Survey land use change matrix (**Section A 3.4.2.1**) gives estimates of forest conversion to other land use categories for all countries in the UK for 1990-1998 and 1999-2007. There are known issues with Countryside Survey over-estimating the extent of Forest

conversion compared with the extent estimated by the Forestry Commission. This is due to differences in Forest definitions, amongst other causes.

A 3.4.4.2 Compilation of activity datasets

For 1990-1999 the deforestation activity dataset is compiled from the felling licence and DCLG datasets as far as possible, using Countryside Survey (CS) data to fill gaps in the time series, to estimate deforestation in Northern Ireland (for which no direct data is available) and to estimate the conversion to different land use categories⁶. The DCLG data is used to estimate the area of Forest Land converted to Settlement (4.E.2.1). The unconditional felling licence data is used to estimate the area of Forest Land converted to Cropland (4.B.2.1) and of Forest Land converted to Grassland (4.C.2.1). The split between the Cropland and Grassland categories is based on the proportional split between forest to grassland conversion and forest to cropland conversion in the most recent Countryside Surveys. Only England has any post-1990 forest to cropland conversion: the estimated areas in Scotland, Wales and Northern Ireland are so small that they are thought to be due to survey classification error than genuine land use change.

The CS data is used to estimate the relative split of Forest conversion between Grassland, Cropland and Settlements (**Table A 3.4.21**), using other known data (e.g. felling licences) to correct the CS areas where datasets overlap in time (**Table A 3.4.22**). This correction ratio is used to adjust the estimated deforestation areas (as Countryside Survey is known to over-estimate deforestation as described in the section above). There is no non-CS data for Northern Ireland so the correction ratios for England or Wales are used, depending on availability. The 1990-98 correction ratios are also applied to the pre-1990 CS land use change estimates.

The annual area of forest converted to other land uses is removed from the area of 4A1 Forest Land remaining Forest Land to maintain consistency in the land area matrix.

⁶ Discussion with Northern Ireland experts confirmed that there are no direct and comprehensive datasets on woodland loss available. 127 ha of deforestation between 2000 and 2006 is recorded in Environmental Impact

Table A 3.4.21 Countryside Survey data for Forest conversion

Countryside Survey land use change		Annual rate of change, kha/yr				Grassland/Cropland fractional split		
		England	Scotland	Wales	N Ireland	England	Scotland	Wales
1990-1998	Forest to Natural Grassland	5.600	4.418	1.099	0.171	0.61	0.86	0.72
	Forest to Pasture Grassland	3.081	0.608	0.418	0.086	0.33	0.14	0.28
	Forest to Cropland	0.545	0.097	0.019	0.008	0.06	0.00	0.00
	Forest to Settlements	1.242	0.293	0.132	0.072			
	Forest to Other Land	0.169	0.231	0.058	0.025			
1999-2007	Forest to Natural Grassland	2.656	10.327	0.120	0.209	0.86	0.98	0.42
	Forest to Pasture Grassland	0.277	0.186	0.162	0.102	0.09	0.02	0.58
	Forest to Cropland	0.141	0.006	0.001	0.001	0.05	0.00	0.00
	Forests to Settlements	0.617	0.098	0.095	0.142			
	Forest to Other Land	0.430	0.695	0.374	0.027			

Table A 3.4.22 Corrected Forest conversion rates

		Correction ratio			Estimated annual rate of change, kha/yr			
		England	Scotland	Wales	England	Scotland	Wales	N Ireland
1990-1998	Grassland & Cropland	2% ^a			0.159	0.088 ^c	0.026 ^c	0.005 ^c
	Settlements & Other Land	28% ^b			0.390	0.145 ^c	0.052 ^c	0.027 ^c
1999-2000	Grassland & Cropland	20% ^a	2% ^a	15% ^a	0.602	0.262	0.041	0.045 ^d

		Correction ratio			Estimated annual rate of change, kha/yr			
		England	Scotland	Wales	England	Scotland	Wales	N Ireland
	Settlements & Other Land	28% ^b			0.296	0.224 ^c	0.133 ^c	0.048 ^c

^a Unconditional felling licence data used for correction

^b Land Use Change Statistics used for correction

^c England correction ratio used


^d Wales correction ratio used

For 2000 onward, the area and subsequent land-use of deforestation were estimated based on a combination of data sources:

- observations on forest loss by the National Forest Inventory (internal Forestry Commission analysis) by IPCC category;
- unconditional felling licences granted (assumed all converted to Grassland);
- analysis of the FC Sub-Compartment Database for restoration of Forest land to open habitats (assumed all converted to Grassland); and
- conversion to non-forest on private sector forest covered by long-term forest plans rather than felling licences (internal Forestry Commission report, assumed all converted to Grassland).

The changes in the overall deforestation area are shown in **Figure A 3.4.5**. The revision in deforestation was only done from 2000 onwards, partly because there was no suitable data on which to base adjustments for 1990-1999, but also because a number of policy developments came into play in 2000 or shortly beforehand, which affected deforestation to restore open habitats or develop wind-farms. These include the introduction of the UK's climate change policy (2000), and the diversification in relevant forest policies in England, Scotland and Wales following the devolution of forest policy to countries in the late 1990s (Matthews et al. 2014).

Figure A 3.4.5 Comparison of estimated areas of deforestation in the UK between the 1990-2012 and 1990-2013 inventories




A 3.4.4.3 Estimation of emissions

Soil carbon stock changes associated with deforestation are estimated using the dynamic soil carbon model described in **Section A 3.4.2**. When deforestation occurs it is assumed that 60% of the standing biomass is removed as timber products and the remainder is burnt (in the UNFCCC inventory, KP-LULUCF reporting of deforestation assumes instantaneous oxidation of HWP). Country-specific forest biomass densities for living and dead organic matter from CARBINE are used (**Figure A 3.4.6**). Biomass losses are reported in the relevant carbon stock change tables (assuming a carbon fraction of 0.5). The carbon removed as timber is reported as Harvested Wood Products (HWP) in 4G, using CARBINE to model emission from HWP (described in **Section A 3.4.11**).

Direct and indirect greenhouse gas emissions from associated biomass burning is estimated using the Tier 1 methodology described in the IPCC 2006 guidelines (IPCC 1997 a, b, c) and the emission ratios for CH₄, CO, N₂O and NO_x from Table 3A.1.15 in the IPCC 2003 GPG for LULUCF. Only immediate losses are considered because sites are normally completely cleared, leaving no debris to decay.

Figure A 3.4.6 Country-specific biomass densities for biomass burning, 1990-2013. Living densities (top) and dead organic matter – litter and dead wood (bottom).



A 3.4.5 Biomass Burning – Forest and Non-Forest Wildfires (4A, 4B, 4C)

A 3.4.5.1 Activity dataset

Until 2010 only wildfires on Forest land were reported due to a lack of activity data for wildfires on other land use categories. Data on Forest wildfires prior to 2010 come from the Forestry Commission and the Forest Service of Northern Ireland.

In 2010 the Fire and Rescue Service began recording wildfires in England, Scotland and Wales on a new Incidence and Reporting Systems (IRS) which includes wildfires on all land use categories. The IRS database contains 30 attributes for each fire to which a fire appliance was called, including date, spatial location, property type description (e.g. heathland and moorland, standing crop) and an estimate of the area burnt. This dataset is available from 1st April 2009. The original dataset had >126,000 fire records but 99% of these fires were less than 1 ha in size. The IRS database is manually completed by fire service personnel and requires some subjective judgement. This is likely to lead to non-systematic differences in the accuracy and precision of the data. The accuracy of the locations is variable, but an assessment of a number of the larger fires suggests that the land cover type attribute is reliable. The accuracy of the burnt area estimates could not be validated using aerial photography as the available imagery was not recent enough. Landsat images were used, however, it was still difficult to find cloud-free, pre- and post-fire images for fires in 2010. In addition Landsat has been affected by image 'striping' since 2003, which affects the quality of the images and causes some data loss. There are issues with re-ignited fires or additional fires in the same area being logged in the database as separate events. Overall, the uncertainty associated with this dataset is high but should be re-assessed once a longer time series is available.

To provide data on non-Forest wildfires prior to 2010, thermal anomaly data for 2010 from the NASA-operated MODerate Resolution Imaging Spectroradiometer (MODIS) was obtained from the Fire Information Resource Management System (FIRMS) and allocated to land uses using the proportions of fire on each land use type from the Fire and Rescue Service IRS data. The correlation between MODIS data and IRS data breaks down below 25 ha, so for consistency a 25 ha threshold was set for reporting wildfires logged on the IRS.

Thermal anomalies usually represent active fires, but may detect industrial heat sources, although these are typically masked out by the thermal anomaly processing chain. The IRS data set records 89 fires > 25ha occurring in 2010. The FIRMS data set records 335 fire detections for the same period, however, the FIRMS detections may contain multiple detections for a single fire event and the FIRMS detections are for a single 1km pixel, and do not have a straightforward conversion to burnt area. Searching the IRS and FIRMS data sets for temporally and spatially coincident events (using a 2km buffer around the IRS data) suggests that 22 fires were recorded by both the IRS and FIRMS systems. There are wide discrepancies between the two datasets, reflecting their different natures. The IRS data set records fires where a fire service response was required, so does not record controlled burning, unless the fire gets out of control. The FIRMS data set however, responds to anomalous heat signatures, so records controlled and uncontrolled fires, however, it is only able to detect fires under cloud-free or light cloud conditions. It is also only able to detect fires alight at the time of the satellite overpass. The FIRMS data is more likely to detect larger fires than smaller ones, probably due to the stronger heat signature and the longer burn time that larger fires tend to exhibit.

The IRS and FIRMS thermal anomalies give a very different perspective on the extent, timing and duration of fire events in the UK. However, the datasets did show correlation ($R^2= 70-81\%$) for fires larger than 25 ha, which enabled an empirical relationship to be derived to extend the burnt area record back to 2001. A burnt area threshold of 25 hectares was used to extract a subset of the IRS database: this captured 84% of the IRS wildfire-burnt area in England, 94% in Scotland and 66% in Wales.

As more IRS data becomes available it will be possible to increase confidence in the relationship between fires detected by FIRMS and fires logged in the IRS. This may allow FIRMS data to be extrapolated to fires covering less than 25 ha the inventory in future. However, there would need to be reasonable confidence that the ratio of large to small fires used was valid and also some investigation of whether the distribution of small fires across land use classes was the same as that of larger fires.

It was assumed that all fires in the IRS database were wildfires: even if they started as controlled burning, the need for a fire appliance call-out indicates that they are no longer under control. The IRS property type descriptions were assigned to LULUCF sub-categories (**Table A 3.4.23**). There is a very small area of wildfires that occur on Settlement types, and these are included in the Grassland category as the IRS land type classification suggests that they occur on grassy areas within Settlements and there is not a separate reporting field for wildfires in Settlements in the CRF.

Table A 3.4.23 IRS database property type descriptions by LULUCF sub-category

LULUCF sub-category	Forest	Cropland	Grassland	Settlement
IRS property type description	Woodland/forest - conifers/softwood	Straw/stubble burning	Heathland or moorland	Domestic garden (vegetation not equipment)
	Woodland/forest - broadleaf/hardwood	Stacked/baled crop	Grassland, pasture, grazing etc.	Park
		Nurseries, market garden	Scrub land	Roadside vegetation
		Standing crop	Tree scrub	Railway trackside vegetation
				Wasteland
				Canal/riverbank vegetation

A time series of wildfire-burnt areas for each non-forest land use type was constructed for 1990-2013 (**Figure A 3.4.8**). For non-forest wildfires for England, Scotland and Wales the IRS burnt areas were used for 2010-2013 and the burnt area estimated from thermal anomalies (using equation 1) from 2000 to 2010. For 1990-2000 the average annual burnt area 2001-2011 was used.

In Northern Ireland, where no IRS data were available, it was assumed that the heathland and grassland burning rates were in the same proportions as the Scottish burning rates, using the area of heathland and grassland from the 2007 Northern Ireland Countryside Survey.

Estimates of the forest area burnt in wildfires 1990-2004 are published in different locations (FAO/ECE 2002; Forestry Commission 2004; FAO 2005) but all originate from either the Forestry Commission (Great Britain) or the Forest Service (Northern Ireland). There is a gap in the time series 2005-2010 for Great Britain but areas of forest wildfires are reported annually for Northern Ireland. The gap was filled using the annual average areas burnt 1995-2005. These areas refer only to fire damage in state forests; no information is collected on fire damage in privately owned forests. The proportion of private-owned forest that was burnt each year was assumed to be the same as the percentage of the state forest that was burnt each year.

Figure A 3.4.7 Annual area of FIRMS thermal anomalies for GB for 2001-2013 (thermal anomalies were filtered to exclude those recorded over urban/industrial areas).





Figure A 3.4.7 shows the temporal pattern of FIRMS thermal anomalies, with peaks in hot dry years such as 2003. The FIRMS data used only includes thermal anomalies for March – August for each year, as these are the months where the IRS database recorded fires greater than 25ha. Some FIRMS thermal anomalies were recorded outwith these months due to FIRMS detecting both controlled burns and some fires less than 25ha in size which are not included in the IRS data.

Figure A 3.4.8 Time series of wildfire burnt areas in the UK 1990-2013**A 3.4.5.2 Estimation of emissions**

The IPCC Tier 1 method is used for estimating emissions of CO₂ and non-CO₂ gases from wildfires (IPCC 2006). The *Calluna* heath fuel biomass consumption factor and grassland emission factors are used for heathland and moorland fires, the agricultural residues EFs for cropland and the “savannah and grassland” EFS for other grassland and settlements.

Country-specific biomass and Dead Organic Matter densities from the CARBINE model are used for estimating fuel consumption in forest fires (as discussed in the deforestation methodology section) and the ‘extra tropical forest’ EFs in the 2006 Guidelines.

Emissions from all wildfires are reported under the ‘Land remaining Land’ categories (i.e. 4A1, 4B1 and 4C1) and IE reporting under 4A2, 4B2 and 4C2.

A 3.4.6 Emissions from drainage (organic soils) (4B1, 4C1)

Some Wetlands in the UK were drained many years ago for agricultural purposes and continue to emit carbon from the soil. Previous inventories have used the methodology described by Bradley (1997) to assess these emissions. However, because no data existed on the extent of drained organic soils elsewhere only emissions from drained organic soils under Cropland in England were included in the inventory. A new assessment of areas of drained organic soils under Cropland and improved Grassland throughout the UK recently became available (Steve Anthony, ADAS pers. comm) from work on the UK Agricultural Greenhouse Gas Platform project (Defra project AC0114). This work gives more complete information on the area of Cropland on drained organic soils, and also attempted to assess the area of Grassland on drained organic soils. It was also possible to estimate the area of improved Grassland on drained organic soils, but lack of data drainage of semi-natural Grassland meant that the area of semi-natural Grassland on drained organic soils could not be estimated. These areas of drained organic soils have also been used in the 1990 – 2013 inventory for the Agricultural Sector, so there is consistency within the UK Greenhouse Gas Inventory. Work to implement the Wetlands Supplement (IPCC 2013) guidance will develop a methodology to estimate the

area of semi-natural Grassland on drained organic soils. Previous inventories have used a model which gives different emissions rates to thick and thin peat. The drained organic soil area data do not include information on the depth of the organic soil. Therefore in this inventory emissions have been estimated using emissions factors for drained organic soils under Cropland and Grassland taken from the AFOLU Guidelines (IPCC, 2006) applied to all depths of drained organic soil. Work to implement the Wetland Supplement guidance may allow more detailed emissions trajectories to be developed in future inventories.

It is assumed area of drained organic soils has remained constant as no drainage or rewetting has occurred since 1990. It is also assumed that land on organic soils has not been converted between land uses.

A 3.4.7 Changes in Stocks of Carbon in Non-Forest Biomass due to Yield Improvements (4B1)

Up until the 1990-2012 inventory an annual increase in the biomass of cropland vegetation in the UK that is due to yield improvements (from improved species strains or management, rather than fertilization or nitrogen deposition) was included in estimates of carbon stock change in non-forest biomass. This was based on the assumption that the annual average standing biomass of cereals has increased linearly with increase in yield between 1980 and 2000 (Sylvester-Bradley *et al.* 2002). This assumption was re-examined, using annual crop yield data used by the Agriculture Inventory sector. No common crops in the UK showed any sustained increase in yield over time since 1990 so the decision was taken to cease reporting of this activity.

A 3.4.8 Emissions of N₂O due to disturbance associated with land use conversion and land management changes (4(III))

Methodological coverage of this activity has expanded in the IPCC 2006 Guidelines, as previously only N₂O emissions due to soil disturbance associated with land use conversion to Cropland were included. All land use conversions or land management changes that will result in a loss of soil carbon, leading to N mineralization and N₂O emissions are now reported. The UK now includes emissions resulting from the land use conversions: 4B2 Forest to Cropland, 4B2 Grassland to Cropland, 4C1 Grassland remaining Grassland, 4C2 Forest to Grassland, 4E1 Settlement remaining Settlement, 4E2 Forest to Settlement, 4E2 Cropland to Settlement and 4E2 Grassland to Settlement. Emissions of N₂O from 4C1 and 4E1 arise from land use change over 20 years before the inventory reporting year where soil carbon stock losses are still included in the LULUCF inventory estimates. Emissions of N₂O from 4B1 Cropland remaining Cropland (resulting from land use change over 20 years before the inventory year) are calculated in the same way by the LULUCF inventory team but are included in the Agriculture sector (category 3D1).

The Tier 1 methodology described in the IPCC 2006 Guidelines is used. The activity data are the areas and soil carbon stock changes reported in the relevant categories in 4B, 4C and 4E. Some C:N ratios for UK soil/vegetation combinations are published in the Countryside Survey (with values of 11.7 to 13.4) but only for the top 15cm of soil. However, the soil carbon stock changes reported in the inventory are from the top 1m of soil, so these C:N ratios were not felt to be applicable. Therefore, the IPCC default C:N ratio of 15 is used for estimating mineralised

N. The emission factor of 1% in the 2006 Guidelines was used to estimate N₂O emissions from mineralised N.

A 3.4.9 On-site and off-site emissions from peat extraction (4D)

On-site emissions of CO₂ and N₂O from peat extraction activities (for energy and horticultural use) and off-site emissions of CO₂ from the decomposition of horticultural peat are reported in category 4D.

A 3.4.9.1 Activity datasets


Separate activity datasets have been compiled for Northern Ireland and for Great Britain (England, Scotland and Wales). Information for Northern Ireland is taken from papers by Cruikshank and Tomlinson (1997) and Tomlinson (2010). These provide estimates of the extent of peat extraction in 1990-1991 and 2007-2008 by different methods (mechanical extraction, sod-cutting and hand-cutting) and by different end uses (fuel or horticultural peat) (**Table A 3.4.24**). Estimates for 1992-2006 were interpolated and the estimate for 2012-2013 was assumed to be the same as that for 2008-2011.

Table A 3.4.24 Activity data for peat extraction sites in Northern Ireland

End use	Method	Area in 1990-1991, ha	Area in 2007-2008, ha
Fuel	Mechanical	3855	329
Fuel	Hand-cutting	107	16
Horticultural	57% vacuum harvesting, 22% mechanical extraction, 18% sod cutting, 3% turfs	576	
Horticultural	95% vacuum harvesting, 5% mechanical extraction		689

For Great Britain areas undergoing peat extraction in 1991 were calculated using the GB area of peat with planning permission (7598 ha) and splitting it between the three countries in proportion to their production volume in 1991 (for both horticultural and fuel peat). Peat extraction sites were estimated using the Directories of Mines and Quarries published in 2002, 2005, 2010 and 2014 (Cameron et al. 2014). These give the location of active commercial extraction sites, but do not contain information on extraction area.

The areas of peat extraction at the site identified from Directory of Mines and Quarries can be clearly seen on Google Earth satellite imagery as shown in the figure below. The areas undergoing peat extraction were measured from the Google Earth imagery. The imagery has been taken at varying (but known) dates. Coverage is not consistent across the UK. For the 1990-2013 inventory the Google imagery was checked for updates against the 2013 data supplied by BGS. The Google imagery for extraction sites reporting changes since 2010 was found to be from approximately 2010, the same age as at previous assessment. This gave the peat extraction areas in **Table A 3.4.25**.

Figure A 3.4.9 Peat extraction site visible on Google Earth imagery

This method was repeated to check for updates to the data in 2014 for the 1990-2013 inventory and no changes to extraction site status or updated imagery were found since the 2010 data.

Any sites abandoned since 2002 (where a change of land use cannot be identified) are still estimated to be producing on-site emissions, in line with good practice guidance. A time series was constructed using linear interpolation. The extraction area (active and abandoned) declined between 1991 and 2009 by 18% in England and 60% on fuel sites in Scotland but increased by 16% on horticultural sites in Scotland. This area was assumed to be converted to Grassland.

The Directory of Mines and Quarries lists two peat extraction sites in Wales, but the Mineral Extraction in Great Britain report for 2012 does not report any peat production in Wales, so it is assumed that the registered sites are currently inactive. In a few cases sites straddle to the Wales-England border and so sites may be registered in one administration but have part of the extraction area in the other. A small area of land conversion to Wetland (<0.14 kha) was recorded which was assumed to be all from Grassland.

Table A 3.4.25 Activity data for peat extraction sites in England, Scotland and Wales

Country	Area in 1991, ha	Area in 2002, ha	Area in 2005, ha	Area in 2010, ha
England	5854	4785	4785	4794
Scotland	1734	1471	1471	1585
<i>Horticultural</i>	1174	1285	1285	1362
<i>Fuel</i>	560	186	186	223
Wales	482	482	482	482

Annual peat production in Great Britain is inferred from extractor sales by volume as published in the annual Mineral Extraction in Great Britain report (formerly known as the Minerals Raised Inquiry) (ONS 2014). This gives a breakdown for horticultural and other uses of peat (assumed to be fuel) for English regions and for Scotland (no peat extraction is reported in Wales) (Table A 3.4.26). Annual production is highly variable because extraction methods depend on suitable summer weather for drying peat.

Table A 3.4.26 Annual peat production, m³ for England and Scotland (from Annual Minerals Raised Inquiry/Mineral Extraction in Great Britain reports)

Year	England		Scotland	
	Horticultural	Fuel	Horticultural	Fuel
1990	1,116,940	2,727	293,170	93,163
1991	1,202,000	2,000	241,000	115,000
1992	1,079,000	4,000	332,000	91,000
1993	1,069,820	2,180	306,511	73,489
1994	1,375,000	1,000	498,000	108,000
1995	1,578,000	2,000	657,000	44,000
1996	1,313,000	2,000	517,000	53,000
1997	1,227,000	2,000	332,000	59,000
1998	936,000	0	107,000	32,000
1999	1,224,000	0	392,000	37,000
2000	1,258,000	1,000	336,000	31,000

Year	England		Scotland	
	Horticultural	Fuel	Horticultural	Fuel
2001	1,459,000	1,000	325,000	30,000
2002	856,000	1,000	107,000	10,000
2003	1,227,000	1,000	741,000	38,000
2004	902,000	1,000	338,000	21,000
2005	927,000	1,000	556,000	21,000
2006	856,000	1,000	712,000	24,000
2007	654,000	0	221,000	10,000
2008	455,000	41,000	243,000	21,000
2009	476,000	0	390,000	21,000
2010	456,000	1,000	527,000	21,000
2011	429,000	0	369,000	26,000
2012	422,000	0	126,000	20,000
2013	422,000	0	126,000	20,000

* The latest statistics were not published in time for inclusion in this submission, so the volumes for 2013 were carried forward from 2012

A 3.4.9.2 Estimation of emissions

Default on-site emission factors for Tier 1 reporting (IPCC 2006) are used to estimate emissions. Peat extracted for horticultural use is inferred to be from oligotrophic (nutrient-poor) bogs. Peat for fuel is inferred to be from minerotrophic (nutrient-rich) fens or bogs. On-site emissions of CO₂ and N₂O from drainage are reported.

A value of 0.0641 tonnes C m⁻³ is used for Great Britain to estimate emissions from extracted horticultural peat volumes based on previous work (Thomson et al, 2011). This is higher than the previously used factor of 0.0557 (Cruikshank and Tomlinson 1997) but slightly lower than the default emission factor of 0.07 tonnes C m⁻³ air-dry peat for nutrient-poor peats.

Tomlinson (2010) gives production estimates of horticultural peat production for Northern Ireland for 1990/91 and 2007/2008. These have been interpolated to produce a time series. The total emission from horticultural peat production is the sum of emissions from vacuum harvesting production, sod extraction production and mechanical extraction production.

Emissions from vacuum harvesting production =

$$\text{area} * \text{annual depth of extraction} * \text{carbon fraction by volume}$$

where

$$\text{Annual depth of extraction by vacuum harvesting, m/ha} = 0.1$$

$$\text{Carbon fraction of air-dry peat by volume, tonnes C/m}^3 \text{ air-dry peat} = 0.0641$$

Emissions from sod extraction production =

$$\text{area} * \text{sod extraction rate} * \% \text{ dry matter for sods} * \text{mean \% C}$$

where

$$\text{Sod extraction rate, tonnes/ha/yr} = 200$$

$$\text{Sod extraction, mean \% dry matter} = 35\%$$

$$\text{Mean \% carbon} = 49\%$$

Emissions from mechanical extraction production =

$$\text{area} * \text{extraction rate} * \% \text{ dry matter for mechanical extraction} * \text{mean \% C}$$

where

The mechanical extraction rate was estimated to be 206.45 tonnes/ha in 1990/91 and 243.06 tonnes/ha in 2007/08 (Tomlinson 2010).

$$\text{Mechanical extraction, mean \% dry matter} = 67\%$$

$$\text{Mean \% carbon} = 49\%$$

A 3.4.10 Flooded Lands (4D)

Carbon stock changes on land converted to Flooded Land (reservoirs) were included in the inventory for the first time this year, based on the IPCC 2006 Guidance. Data on all reservoirs over 1 km² were compiled but only reservoirs established since 1990 were reported (areas of inland water under 1 km² are reported under 4F Other Land). Activity data were compiled for England and Wales from the Public Register of Large Raised Reservoirs provided by the Environment Agency, which listed location, surface area and year built. Activity data for Scotland was compiled from the SEPA Water Body Classification database (of water bodies > 0.5 km²) and the associated Water Body data sheets. Additional information on the year of building was obtained from:

- the Gazetteer for Scotland <http://www.scottish-places.info>
- hydro-electric power generators <http://sse.com/whatwedo/ourprojectsandassets/> <http://www.power-technology.com/projects/glendoehydropowerpla/> , and
- local authorities <http://www.argyll-bute.gov.uk>.

It was established through discussion with local experts that no new large reservoirs had been built in Northern Ireland since the 1950s.

Only five large reservoirs have been established in the UK since 1990, three in England and one each in Scotland and Wales (another five in England are sacrificial floodplains and do not fit the criteria of permanent conversion to Flooded Land). These cover a total of 1.895 kha.

The location of each reservoir was examined using the www.magic.gov.uk geographic information portal. All reservoirs were in upland locations and were assumed to be Grassland prior to their conversion to Flooded Land. (Any forest removed as part of the land conversion will have been captured under the deforestation activity methodology). A Tier 1 methodology was followed, so carbon stock changes in living biomass stock in the year of flooding were

estimated, but not carbon stock change in soils. A living biomass density of 2 t dry matter/ha was used to estimate carbon stock losses.

A 3.4.11 Harvested Wood Products (4G)

The activity data used for calculating this activity is the annual forest planting rates. CARBINE then applies a forest management regime as given in input to the model. For a given forest stand, carbon enters the HWP pool when thinning is undertaken (depending on the species first thinning occurs c. 20 years after planting) and when harvesting takes place.

At thinning and harvest, the CARBINE model allocates merchantable stem volume to various wood products, while the remainder is transferred to the waste pool. The 'end-use' wood products represented are:

- Long-lived sawn timber, e.g. timber used for construction
- Short-lived sawn timber, e.g. timber used for fencing
- Particleboard
- Paper
- Fuel.

For reporting purposes the long-lived and short-lived sawn timber are reported together in the Sawnwood category and treated the same.

During wood processing, conversion losses are assumed and enter the waste stream and decay within a year. The allocation of carbon to wood product categories is estimated by inputting the merchantable stem carbon (from the forest yield model) to a stand volume assortment forecasting model which estimates the volume allocated to sawn timber, roundwood and waste. This is implemented in CARBINE as a set of functions derived from the output of a more general and flexible assortment forecasting program known as ASORT (Rollinson and Gay, 1983). The proportions allocated to different product categories depends on the species harvested, with waste being either left in the forest or used for fuel. This information is based on expert opinion rather than data or scientific research. A carbon retention curve is used to estimate product decay and return of carbon to the atmosphere. Each wood product category has its own carbon retention curve using the default half-lives in the IPCC 2006 Guidelines (from the Durban KP decision, paragraph 29), taking into account the decay rate of wood products and the service life as influenced by socio-economic factors (sawn timber: 35 years, particleboard: 25 years, paper: 2 years, fuel: instantaneous oxidation).

In implementing the 2006 IPCC guidelines for HWP the UK has elected to report using the production approach B2, which requires disaggregation of HWP into those produced and consumed domestically and those produced and exported. In the annual Forestry Statistics publication there is data on the apparent consumption of wood products in the UK. A consistent dataset is available at the product level (i.e. sawnwood, wood panels and paper & paperboard) for 2002 onwards. The ten year average of 2002-2011 was calculated for each product type and those values were used for the years 1990-2001. This dataset was used to assign the HWP output from the CARBINE model into either consumed domestically or exported.

A 3.4.12 Methods for the Overseas Territories (OTs) and Crown Dependencies (CDs)

A web search of statistical publications was undertaken for any updates in datasets in 2014. This work builds on an MSc project to calculate LULUCF net emissions/removals for the OTs and CDs undertaken during 2007 (Ruddock 2007).

The availability of data for the different OTs and CDs is very variable, so that emission estimates can only be made for the Isle of Man, Guernsey, Jersey and the Falkland Islands. These four comprise over 95% of the area in all the OTs and CDs. Gibraltar wished to produce their own inventory: their LULUCF net emissions/removals are likely to be extremely small, given the size of the country (6 km²), and will have little impact on overall numbers. A lack of suitable data for the Caribbean territories (discussed in the 1990-2006 NIR) makes it impossible to create inventories for them at the present time.

Information on the area of each IPCC land category, dominant management practices, land use change, soil types and climate types were compiled for each OT/CD from statistics and personal communications from their government departments (**Table A 3.4.27**). This allowed Tier 1 level inventories to be constructed for the four OT/CDs already mentioned, and a Tier 3 approach for Forest Land on the Isle of Man and Guernsey (using the C-Flow model, for information on CFlow model please refer to 1990-2011 NIR). The assumptions and factors used for the estimation of emissions are given in **Table A 3.4.28** and **Table A 3.4.29**. The estimates have high uncertainty and may not capture all relevant activities, but given the size of the territories any missing sources are likely to be small.

Table A 3.4.27 Information sources for estimating LULUCF emissions from the Overseas Territories and Crown Dependencies

Territory	LULUCF category	Time period	Reference
Isle of Man	4A	1970-2011	Personal communication from Isle of Man Department of Agriculture, Fisheries and Forestry (Peter Williamson) FAO (2010) Global Forest Resources Assessment: Isle of Man
	4B, 4C	2002-2013	Isle of Man Agricultural and Horticultural Census: completed by all farmland occupiers on an annual basis Isle of Man Digest of Economic and Social Statistics
	4E	1991-1994	Isle of Man Ecological Habitat Survey, Phase 1 Report (Sayle et al, 1995)
Guernsey	4A	1990-2010	FAO Global Forest Resources Assessment 2010: Guernsey

Territory	LULUCF category	Time period	Reference
	4A, 4B, 4C, 4E	1998/9, 2005, 2010	Guernsey Habitat Survey Sustainable Guernsey 2005, 2009, Guernsey Facts and Figures 2011
Jersey	4A	1990-2010	FAO Global Forest Resources Assessment 2010: Jersey
	4A, 4B, 4C, 4E	2006, 2008-2012	Jersey In Figures 2006/2008/2009/2010/2011/2012
Falkland Islands	4A	1990-2011	Department of Mineral Resources, personal communication FAO Global Forest Resources Assessment 2010: Falkland Islands
	4B, 4C	1991-2013	Falkland Islands Agricultural Statistics
	4E	1990-2005	Falkland Islands Environment and Planning Department, personal communication

Table A 3.4.28 Assumptions used in applying the Tier 1 methodology to the Overseas Territories and Crown Dependencies

Land Use category	Sub-category	Isle of Man	Guernsey	Jersey	Falkland Islands
Forest land fluxes	Living biomass, DOM, Mineral soils, Organic soils	From C-Flow model	From C-Flow model	Assumed in equilibrium	No forest on Falklands
Crop remaining crop	Living biomass	N/A. Only for perennial crops	N/A. Only for perennial crops	N/A. Only for perennial crops	N/A. Only for perennial crops
	Dead organic matter	N/A	N/A	N/A	N/A
	Mineral soils	No change in SOC	No change in SOC	No change in SOC	N/A
	Organic soils	N/A	N/A	N/A	Default (-5 tC/ha)

Land Use category	Sub-category	Isle of Man	Guernsey	Jersey	Falkland Islands
Land converted to Crop	Living biomass	Use Wales values, grass to crop (-0.5 tC/ha)	Use England values, grass to crop (-0.5 tC/ha)	Use England values, grass to crop (-0.5 tC/ha)	Use Wales values, grass to crop (-0.5 tC/ha)
	Dead organic matter	N/A	N/A	N/A	N/A
	Mineral soils	Default . SOC = 95 tC/ha, assume conversion from natural grassland (-2.603 tC/ha)	Default . SOC = 95 tC/ha, assume conversion from natural grassland (-0.95 tC/ha)	Default . SOC = 95 tC/ha, assume conversion from natural grassland (-0.95 tC/ha)	N/A
	Organic soils	N/A	N/A	N/A	Default (-5 tC/ha)
	N ₂ O emissions	Default (0.002727 t N ₂ O/ha)	Default (0.000995 t N ₂ O/ha)	Default (0.000995 t N ₂ O/ha)	Default (0.012571 t N ₂ O/ha)
Grass remaining grass	Living biomass	N/A	N/A	N/A	N/A
	Dead organic matter	N/A	N/A	N/A	N/A
	Mineral soils	No change in SOC	No change in SOC	No change in SOC	N/A
	Organic soils	N/A	N/A	N/A	Assume no soil C stock change
Land converted to grass	Living biomass	Use Wales values, crop to pasture grass (-0.5 tC/ha)	Use England values, settlement to pasture grass (-1.8 tC/ha)	Use England values, crop to pasture grass (-0.5 tC/ha)	Use Wales values, crop to pasture grass (-0.5 tC/ha)
	Dead organic matter	N/A	N/A	N/A	N/A
	Mineral soils	Default . SOC = 95 tC/ha, assume conversion from cropland (2.603 tC/ha)	Default . SOC = 95 tC/ha, assume conversion from settlement, assume same soil C as for cropland (0.95 tC/ha)	Default . SOC = 95 tC/ha, assume conversion from cropland (0.95 tC/ha)	N/A
	Organic soils	N/A	N/A	N/A	Default (-0.25 tC/ha)

Land Use category	Sub-category	Isle of Man	Guernsey	Jersey	Falkland Islands
Settlements remaining Settlements	Living biomass, DOM, Mineral soils, Organic soils	N/A	N/A	N/A	N/A
Land converted to Settlements	Living biomass	Use Wales values, grass to settlement (-0.8 tC/ha)	Use England values, grass to settlement (-0.8 tC/ha)	Use England values, grass to settlement (-0.8 tC/ha)	Use Wales values, grass to settlement (-0.8 tC/ha)
	Dead organic matter	N/A	N/A	N/A	N/A
	Mineral soils	Default . SOC = 95 tC/ha, assume conversion from grassland and all soil C lost (-6.65 tC/ha)	Default . SOC = 95 tC/ha, assume 30% of land is paved over and the rest is turf grass (-1.14 tC/ha)	Default . SOC = 95 tC/ha, assume 30% of land is paved over and the rest is turf grass (-1.14 tC/ha)	N/A
	Organic soils	N/A	N/A	N/A	Default - assume cropland (-5 tC/ha)
	N ₂ O emissions	Default (0.002727 t N ₂ O/ha)	Default (0.00119 t N ₂ O/ha)	Default (0.00119 t N ₂ O/ha)	N/A
Other land remaining other land	Living biomass, DOM, Mineral soils, Organic soils	N/A	N/A	N/A	N/A
Land converted to other land	Living biomass	N/A	Assume loss of grassland to standing water or cliff (-2 tC/ha)	Assumed loss of grassland to standing water (-2 tC/ha)	N/A
	Dead organic matter	N/A	N/A	N/A	N/A
	Mineral soils	N/A	Assume no change in soil stocks	N/A	N/A
	Organic soils	N/A	N/A	N/A	N/A
	N ₂ O emissions	N/A	0	N/A	N/A
Harvested wood products		From C-Flow model	From C-Flow model	N/A	N/A

Table A 3.4.29 Tier 1 factors used for estimating LULUCF emissions from Overseas Territories and Crown Dependencies

	Factor	Isle of Man/ Guernsey/ Jersey	Falkland Islands
Biomass carbon densities, tC/ha	Cropland	1.5	1.5
	Grassland	2	2
	Pasture Grassland	1	1
	Settlements	2.8	2.8
	Soil C density	95	87
	Grass Flu	1	1
	Grass Fmg	1	1
	Grass Fi	1	1
	Crop Flu	0.8	0.69
	Crop Fmg	1	1
	Crop Fi	1	0.92
	C/N ratio kg N ₂ O-N/kg N	15	15
	N ₂ O EF	0.01	0.01
	Cropland Organic soils EF, tC/ha/yr		-5
	Grassland Organic soils EF, tC/ha/yr		-0.25
	EF2 for temperate organic crop and grassland soils, kg N ₂ O-N/ha		8

A 3.4.13 Uncertainty analysis of the LULUCF sector

The purpose of carrying out uncertainty analysis within the LULUCF inventory is to quantify where the largest sources of errors lie, and to identify areas to be targeted in future work so as to reduce the uncertainties. In the 1990-2010 inventory report a sensitivity analysis of the whole of the existing inventory methodology was undertaken, applying uncertainty quantification more widely and rigorously to all model parameters and empirical conversion factors, in order to quantify the impact of those uncertainties on the inventory. Although this analysis was carried out for the CFlow model, which is no longer used, it is likely to be applicable to the CARBINE model as both are similar forest carbon accounting models (based on the same underlying yield tables and input data). A more rigorous uncertainty analysis of the CARBINE model will be performed in the future.

The results of the simulations, including both input and parameter uncertainty, are that the area undergoing land use change is the single biggest uncertainty in the inventory, followed by uncertainty in the forest model parameters and the choice of model for the change in soil carbon following land use change (**Figure A 3.4.10**). The next five terms are all of a similar magnitude. Full details of the methodology and results are in the 1990-2010 inventory report. The combined uncertainties for the IPCC sub-categories for the 1990-2013 inventory are given in **Table A 3.4.30**.

The uncertainty in the land use change areas is being addressed by the development of a new vector-based approach (see **Chapter 7, Section 1.1**), combining multiple sources of land use data.

Figure A 3.4.10 The largest uncertainties in the LULUCF inventory, in terms of standard deviation in the output distributions




Table A 3.4.30 Combined uncertainties for the LULUCF sector 1990-2013 (rounded to 5%)

Sub-category	CO ₂		CH ₄		N ₂ O	
	1990	2013	1990	2013	1990	2013
4A Forest Land	30%	30%	55%	55%	40%	40%
4B Cropland	45%	45%	55%	55%	55%	55%

Sub-category	CO ₂		CH ₄		N ₂ O	
	1990	2013	1990	2013	1990	2013
4C Grassland	45%	45%	55%	55%	35%	35%
4D Wetland	100%	50%	-	-	100%	100%
4E Settlements	50%	50%	20%	20%	20%	20%
4F Other Land	-	-	-	-	-	-
4G HWP	45%	45%	-	-	-	-

Parameterisation of the forest model is the second largest source of uncertainty. This has been addressed with the move to CARBINE, as 19 tree species are now modelled instead of the two used in previous submissions. Results from the latest National Forest Inventory (NFI) will also provide additional information on carbon stocks in trees (e.g. Forestry Commission 2014⁷). The choice of soil carbon model and its parameters are also important, because the time course of the flux following land use change may be quite different, depending on the equations used to represent this, and how carbon is distributed between fast- and slow-turnover pools. The choice of forest model is less important, largely because all the UK forest models are based on the same yield table data.

A 3.5 WASTE (CRF SECTOR 5)

A 3.5.1 Solid Waste Disposal on Land (5A)

The assumed waste composition is set out in **Table A 3.5.1**. This table also sets out the assumed DDOC content of the waste.

The model allocates waste to two types of landfill – old, closed sites which last received waste in 1979, and modern engineered landfills that came into operation from 1980. Only these latter sites have gas management systems. The old closed sites have no gas control. The distribution of waste between these types of site is the same as used for compiling the previous NIR.

The quantities of Local Authority controlled and Commercial & Industrial waste sent to landfill are shown in **Table A 3.5.2**. The amounts of methane generated, recovered, used for power generation, flared, oxidised and emitted to the atmosphere from 1990 to 2013 are also shown.

For the overseas territories and crown dependencies, the IPCC landfill model is used, apart from for the Isle of Man where insufficient information is currently available. Where available, country-specific waste generation and composition data have been applied and appropriate

⁷ This survey is preliminary and the carbon stocks have been estimated using the same relationships and calculation parameters that underlie CARBINE; they are therefore not an independent validation of the LULUCF estimates.

defaults chosen e.g. taking into account climatic variation. There are no landfill emissions for Gibraltar as waste is exported. **Table A 3.5.3** below gives the parameters used.

Table A 3.5.1 Waste composition data

Local Authority controlled waste					C&I				
Material	Waste composition (2013)	DDOC			Material	Waste composition (2013)	DDOC		
		RDO	MDO	SDO			RDO	MDO	SDO
Paper	10.6%	0.00%	15.70%	0.42%	Paper and Card	19.4%	0.00%	15.70%	0.42%
Card	7.7%	0.00%	14.78%	0.39%	Food and Abattoir	8.2%	6.45%	1.95%	0.15%
Nappies	3.1%	0.00%	4.30%	0.00%	Food effluent	0.0%	0.00%	6.76%	0.00%
Textiles (and footwear)	5.6%	0.00%	6.67%	0.00%	Misc. Comb	2.7%	0.00%	11.00%	0.00%
Misc. Combustible	1.4%	0.00%	11.00%	0.00%	Furniture	3.5%	0.00%	5.18%	0.04%
Wood	5.3%	0.00%	11.84%	0.69%	Garden	3.1%	3.74%	4.40%	0.58%
Food	21.3%	6.74%	2.60%	0.17%	Sewage sludge	0.04%	2.31%	0.00%	0.00%
Garden	3.5%	3.74%	4.40%	0.58%	Textiles / Carpet and Underlay	9.7%	0.00%	6.67%	0.00%
Soil and other organic	2.1%	0.00%	0.27%	0.00%	Wood	10.0%	0.00%	11.84%	0.69%
Furniture	5.0%	0.00%	5.18%	0.04%	Sanitary	1.3%	0.00%	4.30%	0.00%
Non-inert fines	1.8%	0.00%	6.35%	0.00%	Other (as 100% inert)	42.5%	0.00%	0.00%	0.00%
Other (as 100% inert)	32.7%	0.00%	0.00%	0.00%					

Notes:

- a. DDOC values for Local Authority controlled waste and Commercial & Industrial waste since 1997
- b. Furniture in LA-managed waste is assumed to be 62% wood and 5% textile on fresh mass basis.

- c. "Other" is assumed to be 100% inert – i.e. non-biodegradable.
- d. Furniture in C&I waste is assumed to be 50% wood on fresh mass basis.

A 3.5.1.1 Methane emissions

The right-most column of **Table A 3.5.2** shows the current NIR estimate of methane emitted from UK landfills, according to the approach outlined in **Chapter 7**, taking account of recovery and oxidation.

Table A 3.5.2 Amount of waste landfilled and methane generated, captured, utilised, flared, oxidised and emitted

Year	Waste Landfilled (Mt)			Waste reported in CRF ⁸	Methane generated	Methane captured		Methane used for power generation		Methane flared		Residual methane oxidised		Methane emitted	
	MSW	C&I	MSW+ C&I	Mt	kt	kt	%	kt	%	kt	%	kt	%	kt	%
1990	18.19	74.69	92.88	58.35	2,810	33	1%	33	1%	0	0%	278	10%	2499	89%
1991	18.84	74.02	92.86	56.54	2,846	50	2%	50	2%	0	0%	280	10%	2517	88%
1992	19.47	73.35	92.82	54.72	2,879	90	3%	90	3%	0	0%	279	10%	2510	87%
1993	20.09	72.65	92.74	52.86	2,910	107	4%	107	4%	0	0%	280	10%	2522	87%
1994	20.71	72.00	92.71	51.04	2,938	124	4%	124	4%	0	0%	281	10%	2532	86%
1995	26.46	71.32	97.78	54.33	3,025	135	4%	135	4%	0	0%	289	10%	2601	86%
1996	25.75	70.64	96.39	51.16	3,076	170	6%	170	6%	0	0%	291	9%	2615	85%
1997	26.98	69.94	96.92	49.90	3,121	218	7%	218	7%	0	0%	290	9%	2613	84%
1998	26.67	67.53	94.20	49.31	3,149	278	9%	278	9%	0	0%	287	9%	2584	82%
1999	27.56	66.44	94.00	49.59	3,173	394	12%	394	12%	0	0%	278	9%	2502	79%
2000	27.57	65.44	93.01	50.06	3,198	500	16%	500	16%	0	0%	270	8%	2428	76%
2001	28.06	61.86	89.92	50.40	3,213	566	18%	566	18%	0	0%	265	8%	2382	74%

⁸ Waste reported in the CRF does not include inert C&I waste.

Year	Waste Landfilled (Mt)			Waste reported in CRF ⁸	Methane generated	Methane captured		Methane used for power generation		Methane flared		Residual methane oxidised		Methane emitted	
	MSW	C&I	MSW+C&I	Mt	kt	kt	%	kt	%	kt	%	kt	%	kt	%
2002	27.63	64.23	91.86	49.42	3,211	599	19%	598	19%	1	0%	261	8%	2351	73%
2003	26.24	65.65	91.89	46.54	3,172	723	23%	723	23%	0	0%	245	8%	2204	69%
2004	25.05	63.18	88.23	43.83	3,112	874	28%	874	28%	0	0%	224	7%	2015	65%
2005	22.66	60.22	82.88	39.94	3,022	926	31%	926	31%	0	0%	210	7%	1886	62%
2006	21.33	55.71	77.04	36.85	2,915	950	33%	944	32%	6	0%	196	7%	1768	61%
2007	19.72	51.56	71.28	34.42	2,806	989	35%	987	35%	2	0%	182	6%	1635	58%
2008	17.63	47.59	65.22	30.98	2,681	1080	40%	987	37%	93	3%	160	6%	1441	54%
2009	16.35	14.87	31.22	24.93	2,471	1118	45%	1018	41%	100	4%	135	5%	1218	49%
2010	14.73	14.94	29.66	23.34	2,280	1144	50%	1029	45%	115	5%	114	5%	1023	45%
2011	13.13	14.80	27.93	21.65	2,129	1115	52%	1029	48%	86	4%	101	5%	913	43%
2012	11.59	14.78	26.37	20.10	2,001	1099	55%	1031	52%	69	3%	90	5%	811	41%
2013	10.76	14.77	25.54	19.27	1,892	1158	61%	1034	55%	125	7%	73	4%	660	35%

Notes

- a. Methane generated is based on the MELMod model.
- b. Methane captured is the sum of methane used for power generation and methane flared.
- c. Methane used for power generation is calculated from official figures on landfill gas electricity generation (Digest of UK Energy Statistics (DUKES, 2013)), in MWh/year, assuming a net calorific value for methane of 50 GJ/tonnes and a conversion efficiency between methane use and electricity export of 30% rising to 36%, which includes parasitic losses and on-site use of electricity, e.g. for gas blowers, leachate treatment and site offices.
- d. Methane flared is calculated from site-specific data provided by the Environment Agency at regulated sites for 2009 to 2013, from SEPA for 2013, and from a study carried out during 2014 for other years.
- e. Methane oxidised is based on the IPCC default oxidation factor of 10%, applied to methane remaining after subtraction of the amount captured.
- f. Methane emitted = (methane generated – methane captured) x (1-oxidation factor).

Table A 3.5.3 Parameters used in landfill emission estimates for overseas territories and crown dependencies

Region	Methodology	Activity data	MCF	DOC
Guernsey	IPCC Landfill Model	2005 onwards: total MSW to landfill data and percentage that is plastics, other inert. Prior to 2005: flat-lined 2005 data	IPCC default values; waste management type is unmanaged, deep (results from expert consultation, 2014)	IPCC default values
Jersey	N/A, all MSW is incinerated for energy from waste			
Gibraltar	N/A, all MSW used to be incinerated, now all waste is exported to be landfilled in Spain.			
Isle of Man	Scale UK emissions by population			
Bermuda	IPCC Landfill Model	Total MSW to landfill (Environmental Statistics Compendium)	IPCC default values; no information on management system so assume unmanaged deep	IPCC default values
Cayman Islands	IPCC Landfill Model	2000 onwards: Total MSW to landfill (Department of Environmental Health). Prior to 2000: flat-line 2000 data	IPCC default values; landfill sites are lined and managed to some degree, but with limited information, "Uncategorised" considered appropriate	IPCC default values

Region	Methodology	Activity data	MCF	DOC
Falkland Islands	IPCC Landfill Model	1998: Halcrow Report. Other years: flat-lined after advice in personal communication from environmental officer	IPCC default values; waste management type is unmanaged, shallow (results from expert consultation, 2014)	IPCC default values
Montserrat	IPCC Landfill Model	Population data; IPCC default waste per capita for the Caribbean region	IPCC default values; no information on management system so assume uncategorised	IPCC default values

A 3.5.2 Waste Incineration (5C)**Table A 3.5.4 Activity Data: UK Waste Incineration 1990-2013**

Year	Municipal Waste Incineration ^a (Mt)	Clinical Waste Incineration (Mt)	Chemical Waste Incineration (Mt)	Sewage Sludge Incineration (Mt)
1990	2.194	0.350	0.290	0.075
1991	2.172	0.350	0.290	0.069
1992	2.083	0.330	0.290	0.072
1993	1.841	0.310	0.290	0.084
1994	1.341	0.290	0.289	0.072
1995	1.223	0.270	0.289	0.082
1996	1.340	0.250	0.288	0.088
1997	-	0.230	0.287	0.081
1998	-	0.236	0.287	0.185
1999	-	0.242	0.286	0.186
2000	-	0.248	0.285	0.188
2001	-	0.254	0.285	0.189
2002	-	0.260	0.284	0.191
2003	-	0.224	0.257	0.192
2004	-	0.188	0.231	0.194
2005	-	0.152	0.204	0.195
2006	-	0.115	0.177	0.196
2007	-	0.124	0.163	0.191
2008	-	0.131	0.136	0.168
2009	-	0.121	0.126	0.175
2010	-	0.125	0.140	0.183
2011	-	0.113	0.138	0.176
2012	-	0.096	0.134	0.161
2013	-	0.095	0.134	0.165

a Note that msw incinerators were closed or converted to extract power by 1997, so is considered to be power generation

Table A 3.5.5 Emissions Data: UK Waste Incineration 1990-2013

Year	Chemical Waste Incineration	Accidental Fires	MSW Incineration ^a	Clinical Waste Incineration	Sewage Sludge Incineration	Total
Carbon Dioxide (kt CO ₂)						
1990	326.8	NE	667.8	292.6	NA	1,287.2
1995	325.3	NE	322.8	225.7	NA	873.8
2000	277.7	NE	NO	207.3	NA	485.0
2005	251.2	NE	NO	126.7	NA	377.9
2009	168.4	NE	NO	101.5	NA	269.9
2010	169.4	NE	NO	104.5	NA	273.9
2011	174.0	NE	NO	94.3	NA	268.3
2012	167.1	NE	NO	80.5	NA	247.6
2013	170.6	NE	NO	79.7	NA	250.3
Methane (kt CH ₄)						
1990	NE	1.009	4.411	0.007	0.029	5.456
1995	NE	0.984	2.458	0.005	0.032	3.479
2000	NE	0.772	NO	0.005	0.073	0.850
2005	NE	0.704	NO	0.003	0.076	0.783
2009	NE	0.519	NO	0.002	0.068	0.590
2010	NE	0.345	NO	0.002	0.071	0.419
2011	NE	0.350	NO	0.002	0.069	0.421
2012	NE	0.332	NO	0.002	0.063	0.396
2013	NE	0.332	NO	0.002	0.064	0.398

Other Detailed Methodological Descriptions **A3**

Year	Chemical Waste Incineration	Accidental Fires	MSW Incineration ^a	Clinical Waste Incineration	Sewage Sludge Incineration	Total
Nitrous oxide (kt N ₂ O)						
1990	0.029	NE	0.001	0.011	0.060	0.100
1995	0.029	NE	0.000	0.008	0.066	0.103
2000	0.029	NE	NO	0.007	0.150	0.186
2005	0.020	NE	NO	0.005	0.156	0.181
2009	0.013	NE	NO	0.004	0.140	0.156
2010	0.014	NE	NO	0.004	0.146	0.164
2011	0.014	NE	NO	0.003	0.141	0.158
2012	0.013	NE	NO	0.003	0.129	0.145
2013	0.013	NE	NO	0.003	0.132	0.149

a Note that msw incinerators were closed or converted to extract power by 1997, so is considered to be power generation

A 3.5.3 Wastewater Handling (5D)

A 3.5.3.1 5D1 Domestic and Commercial Waste Water Handling and Sludge Disposal

Table A 3.5.6 UK Domestic and Commercial Waste Water Treatment (5D1) Activity Data

Treatment/disposal route		unit	1990	1995	1998	1999	2000	2001	2002	2005	2010	2011	2012	2013
Total Sludge		kt tds	1,423	1,470	1,429	1,437	1,444	1,534	1,589	1,768	1,666	1,635	1,589	1,598
Population Equivalent		million	68.3	69.2	69.7	70.0	70.2	70.5	70.8	70.5	69.2	70.2	71.2	72.9
Additional Treatment	Digested	kt tds	409	441	406	441	477	649	685	824	787	765	717	694
	Advanced Digested	kt tds	93	99	92	99	106	130	249	312	350	351	385	295
	Composted	kt tds	8	8	8	8	9	9	13	17	25	32	43	48
Disposal route	Farmland	kt tds	508	547	504	547	590	831	897	1,216	1,282	1,260	1,270	1,287
	Landfill	kt tds	160	153	130	120	110	159	107	131	35	26	14	6
	Incineration	kt tds	68	80	185	198	211	221	287	252	238	248	237	252
	Sea	kt tds	570	534	476	424	372	-	-	-	-	-	-	-
	Composted	kt tds	2	2	2	2	2	8	7	13	23	30	42	53
	Land Reclamation	kt tds	31	30	30	30	30	32	151	96	44	41	10	-
	Other	kt tds	84	124	101	115	129	283	140	61	44	30	16	-

Where tds is total dissolvable solids

Table A 3.5.7 UK Domestic and Commercial Waste Water Treatment (5D1) Implied Emission Factors

Treatment/disposal route		unit	1990	1995	1998	1999	2000	2001	2002	2005	2010	2011	2012	2013
Mechanical treatment and storage ¹		kt/Mt tds	2.70	2.70	2.70	2.70	2.70	2.70	2.70	2.70	2.70	2.70	2.70	2.70
Additional Treatment	Digested ²	kt/Mt tds	16.56	16.31	16.08	16.15	16.49	16.92	16.94	16.96	16.44	16.23	16.20	16.01
	Advanced Digested	kt/Mt tds	4.56	4.56	4.56	4.56	4.56	4.56	4.56	4.56	4.56	4.56	4.56	4.56
	Composted	kt/Mt tds	1.46	1.46	1.46	1.46	1.46	1.46	1.46	1.46	1.46	1.46	1.46	1.46
Disposal route	Farmland ³	kt/Mt tds	1.39	1.39	1.39	1.39	1.40	1.59	1.47	1.46	1.43	1.42	1.38	1.45
	Landfill	kt/Mt tds	3.79	3.79	3.79	3.79	3.79	3.79	3.79	3.79	3.79	3.79	3.79	3.79
	Incineration	kt/Mt tds	-	-	-	-	-	-	-	-	-	-	-	-
	Sea ⁴	kt/Mt tds	60.00	60.00	60.00	60.00	60.00	60.00	60.00	60.00	60.00	60.00	60.00	60.00
	Composted	kt/Mt tds	0.63	0.63	0.63	0.63	0.63	0.63	0.63	0.63	0.63	0.63	0.63	0.63
	Land Reclamation ⁵	kt/Mt tds	1.41	1.41	1.41	1.41	1.41	1.41	1.41	1.41	1.41	1.41	1.41	1.41
	Other ⁶	kt/Mt tds	1.20	1.20	1.20	1.20	1.20	1.20	1.20	1.20	1.20	1.20	1.20	1.20
Total ⁷		kt/Mt tds	32.84	30.76	28.51	26.68	24.96	11.76	12.05	12.83	12.70	12.52	12.33	11.74

1. All waste is mechanically treated and stored, so the emission factor is applied to total sludge.

2. Implied emission factor after methane capture.

3. Emission factor varies depending on how the waste is treated.

4. Not an IEF, this is the default IPCC factor for sea, river and lake discharge.

5. Land reclamation hasn't got associated reported emissions, so the factor is based on a weighted average of other waste to land (farmland, composting) IEFs.

6. Other hasn't got associated reported emissions, the factor is based on a weighted average of all other disposal IEFs.

7. For information, IEF when dividing total emissions by total activity.

Table A 3.5.8 UK Domestic and Commercial Waste Water Treatment (5D1) Emission Estimates (kt CH₄)

Treatment/disposal route		1990	1995	1998	1999	2000	2001	2002	2005	2009	2011	2012	2013
Mechanical treatment and storage		3.84	3.97	3.86	3.88	3.90	4.14	4.29	4.78	4.61	4.42	4.29	4.31
Additional Treatment	Digested	6.77	7.19	6.53	7.13	7.86	10.98	11.60	13.98	13.66	12.41	11.62	11.11
	Advanced Digested	0.43	0.45	0.42	0.45	0.49	0.59	1.14	1.42	1.58	1.60	1.76	1.34
	Composted	0.01	0.01	0.01	0.01	0.01	0.01	0.02	0.03	0.02	0.05	0.06	0.07
Disposal route	Farmland	0.71	0.76	0.70	0.76	0.82	1.32	1.32	1.77	1.89	1.79	1.75	1.86
	Landfill	0.61	0.58	0.49	0.46	0.42	0.60	0.41	0.50	0.17	0.10	0.05	0.02
	Incineration	-	-	-	-	-	-	-	-	-	-	-	-
	Sea	34.22	32.05	28.57	25.46	22.34	-	-	-	-	-	-	-
	Composted	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.01	0.01	0.02	0.03	0.03
	Land Reclamation	0.04	0.04	0.04	0.04	0.04	0.05	0.21	0.14	0.07	0.06	0.01	-
	Other	0.10	0.15	0.12	0.14	0.15	0.34	0.17	0.07	0.07	0.04	0.02	-
Total		46.73	45.22	40.75	38.33	36.03	18.05	19.15	22.69	22.08	20.47	19.59	18.75

A 3.5.3.2 5D2 Industrial Waste Water Handling and Sludge Disposal**Table A 3.5.9 UK Industrial Waste Water Treatment Activity Data (5D2) (1990-2013)**

Sector	Unit	1990	1995	2000	2005	2010	2011	2012	2013
Organic chemical production	Mt	1.617	1.617	1.751	1.752	1.487	1.590	1.539	1.536
Milk-processing	million PE	1.464	1.464	1.464	0.626	0.624	0.601	0.609	0.613
Manufacture of fruit and vegetable products	million PE	1.145	1.145	1.145	1.473	1.495	1.630	1.658	1.734

Sector	Unit	1990	1995	2000	2005	2010	2011	2012	2013
Potato-processing	million PE	0.302	0.302	0.302	0.389	0.395	0.430	0.438	0.458
Meat industry	million PE	0.623	0.623	0.623	0.619	0.638	0.662	0.664	0.633
Breweries	million PE	0.094	0.094	0.104	0.109	0.109	0.108	0.100	0.100
Production of alcohol and alcoholic beverages	million PE	1.931	1.931	1.931	2.022	2.010	2.002	1.858	1.856
Manufacture of animal feed from plant products	million PE	0.476	0.476	0.476	0.302	0.377	0.503	0.435	0.451
Manufacture of gelatine and of glue from hides, skin and bones	million PE	0.013	0.013	0.013	0.013	0.013	0.013	0.013	0.013
Malt-houses	million PE	0.207	0.207	0.207	0.216	0.215	0.214	0.199	0.199
Fish-processing industry	million PE	0.018	0.018	0.018	0.006	0.006	0.005	0.005	0.005
Total Food and Drink	million PE	6.273	6.273	6.283	5.774	5.883	6.169	5.980	6.062

Where PE is population equivalent

A 3.6 DATA FOR THE UK'S CROWN DEPENDENCIES AND OVERSEAS TERRITORIES

Table A 3.6.1 Isle of Man, Guernsey and Jersey – Emissions of Direct GHGs (Mt CO₂ equivalent)

Sector	1990	1995	2000	2005	2009	2010	2011	2012	2013
1. Energy	1.47	1.63	1.64	1.44	1.46	1.44	1.37	1.49	1.46
2. Industrial Processes and Product Use	0.00	0.01	0.03	0.06	0.08	0.08	0.08	0.08	0.07
3. Agriculture	0.20	0.20	0.21	0.14	0.18	0.17	0.17	0.13	0.13
4. LULUCF	-0.02	-0.03	-0.03	-	-	-	-	-	-
5. Waste	0.21	0.20	0.19	0.14	0.14	0.14	0.14	0.13	0.13
7. Other	-	-	-	-	-	-	-	-	-
Total	1.86	2.01	2.04	1.75	1.84	1.81	1.75	1.81	1.79

Table A 3.6.2 Isle of Man, Guernsey and Jersey – Fuel use data

Fuel	Fuel Unit	1990	1995	2000	2005	2009	2010	2011	2012	2013
Aviation spirit	Mt	0.01	0.01	0.01	0.01	0.01	0.01	0.01	0.01	0.01
Aviation turbine fuel	Mt	0.08	0.08	0.10	0.11	0.09	0.09	0.09	0.09	0.09
Burning oil	Mt	0.22	0.26	0.35	0.37	0.38	0.40	0.38	0.42	0.44
Coal	Mt	0.12	0.10	0.04	0.02	0.01	0.01	0.01	0.01	0.01
DERV	Mt	0.07	0.09	0.13	0.12	0.12	0.12	0.13	0.13	0.13
Fuel oil	Mt	0.48	0.59	0.45	0.09	0.15	0.11	0.10	0.18	0.17
Gas oil	Mt	0.12	0.13	0.12	0.14	0.11	0.09	0.08	0.07	0.05
LPG	Mth	25.65	25.37	60.57	38.72	34.31	34.90	30.99	28.22	27.09
Natural gas	Mth	0.00	0.00	0.00	101.94	119.88	135.02	135.54	136.05	136.56
Petrol	Mt	0.24	0.23	0.22	0.22	0.20	0.19	0.19	0.18	0.17
Wood	Mt	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00

Table A 3.6.3 Isle of Man, Guernsey and Jersey – Animal numbers

Livestock Category	1990	1995	2000	2005	2009	2010	2011	2012	2013
Dairy	15,888	15,729	16,186	13,127	11,658	11,455	10,944	9,136	9,163
Non dairy	28,663	28,333	29,176	16,770	30,147	28,615	27,137	27,475	27,586

Livestock Category	1990	1995	2000	2005	2009	2010	2011	2012	2013
Sheep	151,764	160,228	176,259	87,537	145,962	138,251	134,963	59,558	55,017
Goats	333	347	376	196	304	288	301	190	183
Horses	1,928	1,928	1,928	1,965	2,301	2,379	2,346	2,376	2,408
Pigs	4,854	5,411	4,609	1,148	1,420	1,114	920	872	918
Poultry	84,048	46,481	46,448	58,160	55,092	54,400	52,152	57,193	57,808

Table A 3.6.4 Isle of Man, Guernsey and Jersey – Total emissions from Agricultural Soils (kg N₂O-N)

Territory	1990	1995	2000	2005	2009	2010	2011	2012	2013
Isle of Man	17,440	17,673	18,741	10,872	16,354	15,612	14,886	10,905	10,767
Guernsey	1,318	1,318	1,093	962	948	952	942	946	947
Jersey	2,330	2,323	2,369	1,768	1,703	1,749	1,721	1,735	1,750

Table A 3.6.5 Isle of Man, Guernsey and Jersey – Amount of synthetic fertilizer applied

Country	kg N applied
Isle of Man	3,921,100
Guernsey	297,200
Jersey	594,400

Table A 3.6.6 Cayman Islands, Falklands Islands, Bermuda and Montserrat – Emissions of Direct GHGs (Mt CO₂ equivalent)

Sector	1990	1995	2000	2005	2009	2010	2011	2012	2013
1. Energy	1.09	1.10	1.18	1.39	1.39	1.38	1.43	1.31	1.32
2. Industrial Processes and Product Use	0.00	0.01	0.02	0.03	0.04	0.04	0.04	0.04	0.04
3. Agriculture	0.27	0.26	0.25	0.22	0.20	0.21	0.21	0.20	0.20
4. LULUCF	0.01	0.01	0.02	0.02	0.02	0.02	0.01	0.02	0.02
5. Waste	0.12	0.10	0.09	0.09	0.13	0.13	0.13	0.12	0.12
7. Other	-	-	-	-	-	-	-	-	-
Total	1.49	1.48	1.54	1.75	1.79	1.78	1.81	1.69	1.70

Table A 3.6.7 Cayman Islands, Falklands Islands, Bermuda and Montserrat – Fuel use data

Fuel	Fuel Unit	1990	1995	2000	2005	2009	2010	2011	2012	2013
Aviation spirit	Mt	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
Aviation turbine fuel	Mt	0.28	0.20	0.21	0.18	0.20	0.20	0.18	0.18	0.19
Burning oil	Mt	0.00	0.01	0.01	0.01	0.01	0.01	0.01	0.01	0.01
Clinical waste	Mt	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
DERV	Mt	0.16	0.13	0.11	0.24	0.10	0.10	0.14	0.09	0.09
Fuel oil	Mt	0.21	0.22	0.37	0.41	0.42	0.43	0.43	0.45	0.45
Gas oil	Mt	0.41	0.46	0.43	0.48	0.55	0.55	0.60	0.58	0.59
LPG	Mth	7.59	7.79	7.26	12.54	12.41	12.68	12.86	13.27	13.53
MSW	Mt	0.00	0.20	0.20	0.20	0.19	0.19	0.16	0.17	0.17
Natural gas	Mth	0.84	1.04	1.20	1.58	1.27	1.87	1.62	1.73	1.74
Petrol	Mt	0.20	0.21	0.22	0.18	0.21	0.20	0.18	0.14	0.14

Table A 3.6.8 Cayman Islands, Falklands Islands, Bermuda and Montserrat – Animal numbers

Livestock Category	1990	1995	2000	2005	2009	2010	2011	2012	2013
Dairy Cattle	11,529	11,507	11,054	10,999	10,668	10,833	10,628	10,765	10,753
Non-dairy Cattle	4,978	4,710	5,748	6,988	6,585	6,145	6,272	5,775	5,823
Sheep	733,949	690,456	647,525	538,251	483,425	492,865	491,575	490,837	490,837
Goats	7,507	8,342	8,891	9,440	9,800	9,347	9,512	9,426	9,313
Horses	2,500	2,194	1,744	1,594	1,419	1,394	1,373	1,342	1,342
Swine	2,116	2,232	2,055	1,880	2,619	2,429	2,370	2,198	2,348
Poultry	45,319	49,555	50,714	50,525	49,593	49,041	49,498	49,937	49,873
Deer	124	124	124	124	184	184	243	243	243

Table A 3.6.9 Cayman Islands, Falklands Islands, Bermuda and Montserrat – Total emissions from Agricultural Soils (kg N₂O-N)

Territory	1990	1995	2000	2005	2010	2011	2012	2013
Bermuda	257	257	257	257	257	257	257	257
Cayman Islands	386	477	529	581	724	770	688	685
Falkland Islands	27,596	25,828	24,378	20,743	18,675	18,581	18,483	18,486
Montserrat	3,540	3,669	3,669	3,665	3,712	3,712	3,763	3,763

Table A 3.6.10 Cayman Islands, Falklands Islands, Bermuda and Montserrat - Amount of synthetic fertilizer applied

Country	kg N applied
Cayman Islands	5,400
Falklands	0
Bermuda	1,480
Montserrat	6,000

Table A 3.6.11 Cayman Islands, Falklands Islands, Bermuda and Montserrat – Production of non-N-fixing crops (tonnes)

Territory	1990	1995	2000	2005	2009	2010	2011	2012	2013
Montserrat	1,231	1,404	1,843	2,104	2,015	1,931	2,106	2,113	2,113
Cayman Islands	382	629	669	445	447	459	486	509	509
Bermuda	4,256	4,505	4,419	3,876	5,029	5,240	5,161	5,413	5,413

Table A 3.6.12 Montserrat – Production of N-fixing crops (tonnes)

Production of N-fixing crops (tonnes)	1990	1995	2000	2005	2009	2010	2011	2012	2013
Montserrat	0.26	0.42	0.42	0.36	0.40	0.34	0.36	0.40	0.40

Table A 3.6.13 Gibraltar – Emissions of Direct GHGs (Mt CO₂ equivalent)

Sector	1990	1995	2000	2005	2009	2010	2011	2012	2013
1. Energy	0.174	0.179	0.201	0.233	0.250	0.245	0.238	0.240	0.286
2. Industrial Processes and Other Product Use	0.001	0.001	0.004	0.007	0.009	0.010	0.010	0.011	0.011
3. Agriculture	-	-	-	-	-	-	-	-	-
4. LULUCF	-	-	-	-	-	-	-	-	-
5. Waste	0.006	0.006	0.006	-	0.001	0.002	0.000	0.000	0.002
6. Other	-	-	-	-	-	-	-	-	-
Total	0.180	0.186	0.211	0.240	0.260	0.256	0.249	0.251	0.298

Table A 3.6.14 Gibraltar – Fuel use data

Fuel	Fuel Unit	1990	1995	2000	2005	2009	2010	2011	2012	2013
Aviation turbine fuel	Mt	0.009	0.007	0.006	0.009	0.008	0.007	0.009	0.009	0.009

Other Detailed Methodological Descriptions **A3**

Fuel	Fuel Unit	1990	1995	2000	2005	2009	2010	2011	2012	2013
Charcoal	Mt	-	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000
Road fuels	Mt	0.001	0.002	0.005	0.008	0.008	0.008	0.008	0.008	0.008
Fuel oil	Mt	0.023	0.021	0.024	0.021	0.020	0.018	0.018	0.017	0.013
Gasoil	Mt	0.020	0.024	0.026	0.033	0.040	0.041	0.038	0.041	0.058
MSW	Mt	0.016	0.019	0.024	-	-	-	-	-	-
Natural gas	Mth	1.121	0.925	0.914	0.898	0.769	0.860	0.642	0.750	0.735
Clinical waste	Mt	-	-	-	-	0.001	0.002	0.000	0.000	0.002

ANNEX 4: National Energy Balance for the Most Recent Inventory Year

A 4.1 UK ENERGY BALANCE

The UK energy balance is produced and published annually by the Department of Energy and Climate Change, in the Digest of UK Energy Statistics – DUKES. This is available online from:

<https://www.gov.uk/government/collections/digest-of-uk-energy-statistics-dukes>

The aggregate energy balance for the latest year is presented below. The following sections explain how the energy balance is used for the UK inventory for individual fuel types, and how the data are supplemented with other statistics that may lead to deviations from the DUKES statistics.

The UK energy statistics (detailed breakdown) are presented on a mass basis for liquid and solid fuels, and on a gross energy basis for gaseous fuels (including derived gases). The UK inventory is calculated using these data directly, and for the purposes of reporting in the CRF and NIR, activity data and emission factors are converted to energy units, on a net basis.

Table A 4.1.1 UK Energy Balance (thousand tonnes of oil equivalent, gross energy basis)

	Coal	Manufactured fuel(1)	Primary oils	Petroleum products	Natural gas(2)	Bioenergy & waste(3)	Primary electricity	Electricity	Heat sold	Total
Supply										
Indigenous production	8,025	-	44,468	-	36,523	6,883	18,467	-	-	114,366
Imports	32,122	593	64,675	30,908	46,011	2,167	-	1,508	-	177,984
Exports	-447	-83	-37,009	-28,698	-9,429	-247	-	-267	-	-76,182
Marine bunkers	-	-	-	-2,691	-	-	-	-	-	-2,691
Stock change(4)	-789	-87	+791	+84	+53	-	-	-	-	+53
Primary supply	38,911	423	72,926	-397	73,157	8,803	18,467	1,241	-	213,530
Statistical difference(5)	-111	-4	-104	-131	-19	-	-	-15	-	-384
Primary demand	39,022	426	73,030	-266	73,177	8,803	18,467	1,256	-	213,914
Transfers	-	+5	-2,023	+2,026	-5	-	-3,024	+3,024	-	+3
Transformation	-37,053	1,514	-71,007	70,181	-19,388	-5,934	-15,442	27,608	1,451	-48,071
Electricity generation	-31,432	-939	-	-588	-17,397	-5,834	-15,442	27,608	-	-44,024
Major power producers	-31,308	-	-	-246	-14,751	-2,403	-15,442	25,225	-	-38,925

	Coal	Manufactured fuel(1)	Primary oils	Petroleum products	Natural gas(2)	Bioenergy & waste(3)	Primary electricity	Electricity	Heat sold	Total
Autogenerators	-124	-939	-	-342	-2,646	-3,430	-	2,383	-	-5,098
Heat generation	-378	-51	-	-68	-1,991	-101	-	-	1,451	-1,138
Petroleum refineries	-	-	-71,007	70,932	-	-	-	-	-	-75
Coke manufacture	-4,020	3,574	-	-	-	-	-	-	-	-446
Blast furnaces	-1,073	-1,308	-	-	-	-	-	-	-	-2,381
Patent fuel manufacture	-150	239	-	-95	-	-	-	-	-	-6
Other	-	-	-	-	-	-	-	-	-	-
Energy industry use	2	777	-	4,657	4,718	-	-	2,283	160	12,597
Electricity generation	-	-	-	-	-	-	-	1,538	-	1,538
Oil and gas extraction	-	-	-	672	4,003	-	-	49	-	4,725
Petroleum refineries	-	-	-	3,984	99	-	-	361	160	4,604
Coal extraction	2	-	-	-	14	-	-	68	-	84
Coke manufacture	-	378	-	-	-	-	-	7	-	385
Blast furnaces	-	400	-	-	31	-	-	38	-	469

	Coal	Manufactured fuel(1)	Primary oils	Petroleum products	Natural gas(2)	Bioenergy & waste(3)	Primary electricity	Electricity	Heat sold	Total
Patent fuel manufacture	-	-	-	-	-	-	-	-	-	-
Pumped storage	-	-	-	-	-	-	-	89	-	89
Other	-	-	-	-	571	-	-	133	-	705
Losses	-	215	-	-	643	-	-	2,322	-	3,179
Final consumption	1,967	953	-	67,285	48,423	2,868	-	27,283	1,292	150,069
Industry	1,441	592	-	4,351	8,023	550	-	8,427	847	24,231
Unclassified	-	74	-	3,512	1	550	-	-	-	4,137
Iron and steel	38	518	-	4	459	-	-	327	-	1,346
Non-ferrous metals	14	-	-	0	165	-	-	381	-	560
Mineral products	776	-	-	163	1,305	-	-	578	-	2,822
Chemicals	55	-	-	103	1,330	-	-	1,486	419	3,392
Mechanical engineering etc.	8	-	-	-	486	-	-	607	-	1,102
Electrical engineering etc.	4	-	-	1	225	-	-	531	-	760
Vehicles	37	-	-	175	386	-	-	436	-	1,033

	Coal	Manufactured fuel(1)	Primary oils	Petroleum products	Natural gas(2)	Bioenergy & waste(3)	Primary electricity	Electricity	Heat sold	Total
Food, beverages etc.	31	-	-	134	1,750	-	-	951	0	2,866
Textiles, leather etc.	42	-	-	43	443	-	-	249	-	778
Paper, printing etc.	70	-	-	29	645	-	-	948	-	1,692
Other industries	362	-	-	31	449	-	-	1,808	427	3,078
Construction	5	-	-	156	377	-	-	126	-	664
Transport (6)	10	-	-	51,964	-	1,091	-	353	-	53,418
Air	-	-	-	12,258	-	-	-	-	-	12,258
Rail	10	-	-	700	-	-	-	350	-	1,060
Road	-	-	-	38,177	-	1,091	-	3	-	39,271
National navigation	-	-	-	828	-	-	-	-	-	828
Pipelines	-	-	-	-	-	-	-	-	-	-
Other	516	221	-	3,982	39,918	1,227	-	18,502	445	64,811
Domestic	492	221	-	2,769	29,622	884	-	9,755	52	43,794
Public administration	16	-	-	290	3,826	98	-	1,618	381	6,230
Commercial	4	-	-	386	5,184	33	-	6,796	11	12,414
Agriculture	-	-	-	294	94	211	-	333	-	932
Miscellaneous	5	-	-	243	1,193	0	-	-	-	1,441

	Coal	Manufactured fuel(1)	Primary oils	Petroleum products	Natural gas(2)	Bioenergy & waste(3)	Primary electricity	Electricity	Heat sold	Total
Non energy use	-	140	-	6,988	481	-	-	-	-	7,609

(1) Includes all manufactured solid fuels, benzole, tars, coke oven gas and blast furnace gas.

(2) Includes colliery methane.

(3) Includes geothermal and solar heat.

(4) Stock fall (+), stock rise (-).

(5) Primary supply minus primary demand.

(6) See paragraphs 5.12 regarding electricity use in transport and 6.40 regarding renewables use in transport.

A 4.2 FUELS DATA

The fuels data are taken from DUKES - the Digest of UK Energy Statistics (DECC, 2014), so the fuel definitions and the source categories used in the NAEI reflect those in DUKES.

IPCC Guidelines (IPCC, 2006) lists fuels that should be considered when reporting emissions. **Table A 4.2.1** lists the fuels that are used in the GHGI (based on DUKES) and indicates how they relate to the fuels listed in the IPCC Guidelines. In most cases the mapping is obvious but there are a few cases where some explanation is required.

Table A 4.2.1 Mapping of fuels used in IPCC and the NAEI

	IPCC	NAEI
Category	Subcategory	Subcategory
Liquid	Motor Gasoline	Petrol
	Aviation Gasoline	Aviation Spirit
	Jet Kerosene	Aviation Turbine Fuel ¹ (ATF)
	Other Kerosene	Burning Oil
	Gas/Diesel Oil	Gas Oil/ DERV
	Residual Fuel Oil	Fuel Oil
	Orimulsion	Orimulsion
	Liquefied Petroleum Gases	Liquefied Petroleum Gas (LPG)
	Naphtha	Naphtha
	Petroleum Coke	Petroleum Coke
	Refinery Gas	Other Petroleum Gas (OPG)
	Other Oil: Other Petroleum Products	Refinery Miscellaneous
	Lubricants	Lubricants
Solid	Anthracite	Anthracite
	Coking Coal	Coal ²
	Other Bituminous Coal	Coal
		Slurry ³
	Coke Oven Coke	Coke
	Patent Fuel	Solid Smokeless Fuel (SSF)
	Coke Oven Gas	Coke Oven Gas
	Blast Furnace Gas	Blast Furnace Gas

	IPCC	NAEI
Category	Subcategory	Subcategory
Gas	Natural Gas	Natural Gas
		Sour Gas ⁴
		Colliery Methane ⁵
Other Fuels	Municipal Solid Waste	Municipal Solid Waste
	Industrial Waste: Scrap Tyres	Scrap Tyres
	Waste Oils	Waste Oil
Peat	Peat	Peat
Biomass	Wood/Wood Waste	Wood
	Other Primary Solid Biomass	Straw
		Poultry Litter, Meat & bone meal
	Landfill Gas	Landfill Gas
	Sludge Gas	Sewage Gas
	Charcoal	Charcoal
	Other liquid biofuels	Liquid Biofuels
	Other biogas	Biogas

- 1 Includes fuel that is correctly termed jet gasoline.
- 2 Used in coke ovens.
- 3 Coal-water slurry used in some power stations
- 4 Unrefined natural gas used on offshore platforms and some power stations
- 5 IPCC Guidelines (IPCC, 2006) specifies coal seam methane is included in Natural Gas.

A 4.2.1 Reallocations of energy data and differences from UK energy statistics

The main source of energy consumption data used in the UK inventory is the Digest of UK Energy Statistics (DECC, 2014). This annual publication gives detailed sectoral energy consumption broken down by fuel type, and covering the entire time period covered by the inventory. In many cases, these data are used directly in the inventory without modification. However there are instances where the activity data used in the inventory are not based directly on DUKES data, but where alternative data sources provide supplementary data to inform energy use and emission estimates.

The reasons for any deviations from use of DUKES data in the inventory are discussed within the source category methodological descriptions in Section 3 of the main report. The main reasons for reallocations or modifications are:

- To account for differences in geographical scope

- To make best use of EU ETS data (this currently does not directly feed into the UK energy statistics)
- To utilise other operator reported data (e.g. direct to the Inventory Agency, or via the regulator's inventories).

The fuel reconciliation tables (**Table A 4.2.2 – Table A 4.2.6**) show how the deviations are applied and how the energy data for the major fuels in the UK inventory are reconciled against the energy demand data from DUKES. The tables show:

1. where fuels are re-allocated between sectors, but the overall annual fuel consumption across all UK sectors is kept consistent with the data in DUKES, and
2. where deviations are made to DUKES figures for total UK consumption of a given fuel, and in which source categories these deviations are made.

The Inventory Agency presents data below for the fuel allocations for coal, natural gas, fuel oil, gas oil (including DERV) and petroleum gases (LPG, OPG) for the latest inventory year. Together these fuels constitute the majority of the UK inventory 1A sector emissions total.

Deviations to the energy balance are made in consultation with the authors of the energy statistics.

A 4.2.1.1 Coal

Total industrial coal use within the GHG inventory is consistent with the DUKES total and in the most part, coal use at the sectoral level is consistent with the DUKES data. However, there is an apparent step change in the amount of fuel allocated to other industries within DUKES between 1999 and 2000. In addition, between 1997 and 1999, the total coal use allocated to 1A2f is less than the independent estimates for cement and lime production used within the inventory. Cement and lime production would fall into the 1A2f category for IPCC reporting. Therefore Inventory Agency estimates have been made to construct a consistent time series for coal use. **Table A 4.2.2** below compares inventory estimates with DUKES estimates for the latest inventory year.

Table A 4.2.2 Fuel reconciliation - coal use in the latest year (Mtonnes)

DUKES Category	DUKES	GHGI	Difference	GHGI category	CRF	Comment
Major power producers	49.842	49.842	0.000	Power stations	1A1a	
Blast furnaces	1.411	1.411	0.000	Blast furnaces	2C1	
Coal extraction	0.003	0.003	0.000	Collieries - combustion	1A1c	
		0.112		Autogenerators	1A2b	
		0.084		Autogenerators - exported to grid	1A2b	
Autogenerators	0.200	0.200	0.000			
Patent fuel manufacture etc.	0.212	0.212	0.000	Solid smokeless fuel production	1B1b	

DUKES Category	DUKES	GHGI	Difference	GHGI category	CRF	Comment
Coke manufacture	5.288	5.107		Coke Production	1B1b	Operator data used in place of DUKES
Iron and steel	0.053	0.053		Iron & steel - combustion plant	1A2a	
Non-ferrous metals	0.040	0.040		Non-Ferrous Metal	1A2b	
Chemicals	0.502	0.502		Chemicals	1A2c	
Paper, printing etc.	0.128	0.128		Pulp, Paper and Print	1A2d	
Food, beverages etc.	0.068	0.068		Food & drink, tobacco	1A2e	
Other industry	1.834					
		1.484		Other industrial combustion	1A2g	Includes re-allocation to balance change to coke manufacture
		0.484		Cement production - combustion	1A2f	Operator data
		0.046		Lime production - non decarbonising	1A2f	EU ETS
Industry + Coke total	7.913	7.913	0.000			
Rail	0.014	0.014	0.000	Rail	1A3c	
Domestic - anthracite	0.263	0.263	0.000	Domestic combustion - anthracite	1A4b	
		0.380		Domestic combustion - UK	1A4b	
		0.003		Domestic combustion - crown dependencies	1A4b	
Domestic - coal	0.383	0.383	0.000			
Agriculture	0.000	0.000	0.000	Agriculture - stationary combustion	1A4c	
Commercial	0.006					
Miscellaneous	0.007					
	0.012	0.012	0.000	Miscellaneous combustion	1A4a	
Public administration	0.152	0.152	0.000	Public sector combustion	1A4a	

DUKES Category	DUKES	GHGI	Difference	GHGI category	CRF	Comment
TOTAL	60.405	60.405	0.000			

Notes: Sequences of shaded rows indicate categories which are grouped for purposes of data reconciliation, and should be considered together.

A 4.2.1.2 Natural Gas

Data for natural gas use is largely taken directly from DUKES and the national total is consistent between the inventory and the energy statistics, other than a small additional use of gas on the Isle of Man which is added to the inventory. Operator estimates for ammonia production (both fuel and feedstock), and ETS data for gas separation plant lead to minor reallocations of the DUKES data, these are summarised below in **Table A 4.2.3**.

Table A 4.2.3 Fuel reconciliation – natural gas use in 2013 (Mtherms)

DUKES Category	DUKES	GHGI	Difference	GHGI Category	CRF	Comment
Major power producers	5854	5890	+37	Power stations	1A1a	GHGI includes Isle of Man
Autogeneration	1033					Some gas re-allocated to refineries in GHGI
		445		Autogenerators	1A2g	
		326		Autogenerators - exported to grid	1A2g	
		0		Railways - stationary combustion	1A4a	
Petroleum refineries	124	385		Refineries - combustion	1A1b	Gas re-allocated from autogenerators
	1157	1157	0			
Coal extraction	2	2	0	Collieries - combustion	1A1c	
		1087		Upstream oil production	1A1c	
		502		Upstream gas production	1A1c	
Oil and gas extraction	1589	1589	0			
Blast furnaces	12	12	0	Blast furnaces	1A2a	
		227		Gas production	1A1c	EU ETS
		0		Nuclear fuel production	1A1c	
Other energy industries	227	227	0			
Non-ferrous metals	73	73		Non-Ferrous Metal	1A2b	
Chemicals	715	715		Chemicals	1A2c	

DUKES Category	DUKES	GHGI	Difference	GHGI Category	CRF	Comment
Paper, printing, etc.	305	305		Pulp, Paper and Print	1A2d	
Food, beverages, etc.	739	739		Food & drink, tobacco	1A2e	
Other industry	1574					
		1365		Other industrial combustion	1A2g	
		50		Lime production	1A2f	EU ETS
		2		Cement production	1A2f	Operator's data
		97		Ammonia production - fuel	2B1	
		159		Ammonia production – feedstock	2B1	
		93		Other non-energy use (stored carbon)	-	
Non-energy use	191					
	1765	1765	0			
Iron and steel	196	196	0	Iron and steel - combustion plant	1A2a	
Domestic	11755	11760	+6	Domestic combustion	1A4b	GHGI includes Isle of Man
Public administration	1664	1664	0	Public sector combustion	1A4a	
Commercial	2199					
Miscellaneous	473					
	2673	2676	+3	Miscellaneous combustion	1A4a	GHGI includes Isle of Man
Agriculture	37	37	0	Agriculture - stationary	1A4c	
Autogenerators (colliery methane)	17					
Coal extraction (colliery methane)	3					
	20	20	0	Collieries – combustion (colliery methane)	1A1c	
Total	28782	28828	+46			Gas use in Isle of Man

Notes: Sequences of shaded rows indicate categories which are grouped for purposes of data reconciliation, and should be considered together.

1 Mtherm = 105.51 TJ

A 4.2.1.3 Fuel Oil

A revised domestic/international split for shipping means that the UK inventory deviates from the national statistics total. Additional sectoral deviations are also made to account for known use of fuel oil in power stations, and the Crown Dependencies.

Table A 4.2.4 Fuel reconciliation – Fuel oils use in 2013 (Mtonnes)

DUKES Category	DUKES	GHGI	Difference	GHGI Category	CRF	Comment
Major power producers	0.158	0.165		Power Stations - UK	1A1a	EU ETS data
		0.049		Power Stations - crown dependencies	1A1a	Local data sets
Autogenerators	0.041					
Iron and steel	0.007	0.020		Iron and steel - combustion plant	1A2a	Includes autogenerators
Other industry	0.046	0.037		Other industrial combustion	1A2g	Reduced to offset increase in 1A1a; includes autogenerators
		0.000		Cement production - combustion	1A2f	Operator's data
Commercial	0.047					
Miscellaneous	0.010					
		0.028		Miscellaneous combustion - UK	1A4a	Reduced to offset increase in 1A1a
		0.008		Miscellaneous combustion - crown dependencies	1A4a	Local data sets
	0.309	0.309	0.000			
Non-ferrous metals	0.001	0.001	0.000	Non-Ferrous Metal	1A2b	
Chemicals	0.041	0.041	0.000	Chemicals	1A2c	
Paper, printing etc.	0.004	0.004	0.000	Pulp, Paper and Print	1A2d	
Food, beverages etc.	0.104	0.104	0.000	Food & drink, tobacco	1A2e	
Petroleum refineries	0.331	0.331	0.000	Refineries - combustion	1A1b	
Agriculture	0.014	0.014	0.000	Agriculture - stationary combustion	1A4c	
Domestic	0.000	0.000	0.000	Domestic combustion	1A4b	
Public administration	0.019	0.019	0.000	Public sector combustion	1A4a	

DUKES Category	DUKES	GHGI	Difference	GHGI Category	CRF	Comment
National navigation	0.089	0.073		Shipping - coastal	1A3d	Revised UK/international split
		0.008		Shipping between UK and Gibraltar	1A3d	Revised UK/international split
		0.000		Shipping between UK and other overseas territories	1A3d	Revised UK/international split
Marine bunkers	1.292	1.301		Shipping - international IPCC definition		Revised UK/international split
	1.382	1.382	0.000			
Total	2.055	2.055	0.000			

Notes: Sequences of shaded rows indicate categories which are grouped for purposes of data reconciliation, and should be considered together. "Total" includes marine bunkers.

A 4.2.1.4 Gas Oil

Gas oil is used in both off-road transport and machinery diesel engines, and as a fuel for stationary combustion. The varied use of this fuel complicates the means of allocating consumption across the wide range of sectors that use the fuel in the inventory. DUKES provides a breakdown of gas oil consumption in different industry and other sectors, but the data resolution in DUKES does not distinguish between use of the fuel for stationary combustion and off-road machinery, a distinction which is necessary for the inventory.

The GHGI estimates consumption of gas oil and emissions for off-road machinery using a bottom-up method based on estimates of population and usage of different types of machinery. However, this has led to a situation where the total amount of gas oil consumption across sectors exceeds that which is available as given in DUKES. Therefore consumption figures, mainly for stationary combustion in industry sectors, have had to be adjusted to obtain a total fuel balance.

The problem is extended when new sources of gas oil consumption are found. For example, the recent development of an inventory for the UK's inland waterways requires the allocation of gas oil to this sector (Walker et al, 2011). During the process of compiling the inland waterways inventory, it became clear that not all vessels with diesel engines use gas oil, but use road diesel and that this may also be the case for other off-road machinery sources, especially those that consume small amounts of fuel on an irregular basis, e.g. for private or recreational use rather than commercial use. There are also inconsistencies in terminology used to define types of fuel; it became apparent that the terms "gas oil", "red diesel" and "diesel" are used interchangeably by fuel suppliers and consumers and this confuses the situation when considering fuel allocations across different sectors.

In light of this, Task 5 of the 2011 UK GHG Inventory Improvement Programme aimed to address the allocation of gas oil and DERV in the GHGI (Murrells et al., 2011). The methodology outlined in Murrells et al. (2011) was used in the compilation of the 2011 inventory, and is summarised here. The same approach has been used in this inventory.

Several fuel suppliers and experts in the petroleum industry and at the Department for Transport were consulted to understand terminologies used, the physical differences between

gas oil and DERV, and to gauge opinions on what determines where the fuels are mainly used where it is possible to use either gas oil or DERV. The study concluded that while the majority of agricultural and industrial machinery will be using low tax gas oil (red diesel), a small amount of DERV is likely to be used by private recreational boat users and by equipment with small engines used for private or small-scale commercial use on an irregular basis and the gas oil fuel supply infrastructure makes it more convenient to use DERV.

The study provided new estimates of the amount of DERV and petrol consumed by non-road transport sources with small internal combustion engines. This reduces the overestimation of gas oil consumption and relieves the pressure on how much gas oil consumption by other sources has to be adjusted to match the total amount available as given in DUKES.

The study also considered the allocation of gas oil given in DUKES to different industry and other sectors and how these can be mapped to inventory reporting categories. The detailed bottom-up method is used to estimate gas oil consumption by different off-road machinery and marine vessel types. Independent sources were used to estimate gas oil used by the rail sector while data provided by industrial sites reporting under emission trading schemes (EU ETS) were used to derive an allocation of gas oil consumption by stationary combustion sources in different industry, commercial and other sectors. Also, the UK energy statistics now include an allocation of gas oil for consumption by the oil and gas sector, but since only a partial time series was made available, the study included making estimates of gas oil for this category back to 1990.

A method of re-allocation was developed using an over-arching condition that the total sum of gas oil consumption across all sectors was consistent with the total consumption figures given in DUKES across all years. The method allowed the consumption estimates for industrial off-road machinery and stationary combustion by industry, commercial and public sector activities to vary in order to align the total consumption estimates with DUKES on the basis that the estimates for these sources are the most uncertain.

Details of the methodology were given in Murrells et al (2011). The report considers the uncertainties in the sector allocations and makes recommendations on how these can be improved based on current activities known to be taking place in the UK to understand the allocation of gas oil across some sectors. The method was first applied to the UK inventory reported in 2012 and the application of the method to define the allocation of gas oil consumption for all source categories across the time series from 1990-2010 was described in the 2012 UK National Inventory Report. The same allocation method has been applied across the current time series for this version of the inventory.

Table A 4.2.5 below summarised the DUKES and GHGI allocations for the latest inventory year.

Table A 4.2.5 Fuel reconciliation – Gas oil use in 2013 (Mtonnes)

DUKES Category	DUKES	GHGI	Difference	GHGI Category	CRF	Comment
Refinery	0.000	0.000	0.000	Refineries	1A1b	
Major power producers	0.042					
		0.042		Power stations - UK	1A1a	EU ETS

DUKES Category	DUKES	GHGI	Difference	GHGI Category	CRF	Comment
		0.005		Power stations - CDs	1A1a	Local data sets
Autogenerators	0.058					
Iron and steel	0.000	0.003		Iron and steel	1A2a	Includes autogenerators
Non-ferrous metals	0.000	0.000		Non-Ferrous Metal	1A2b	Reduced to offset higher consumption elsewhere
Chemicals	0.075	0.003		Chemicals	1A2c	
Paper, printing etc.	0.027	0.001		Pulp, Paper and Print	1A2d	
Food, beverages etc.	0.032	0.002		Food, drink, tobacco	1A2e	
Other industry	1.391					
		0.044		Other industry UK	1A2f	Reduced to offset higher consumption elsewhere
		0.000		Other industry CDs	1A2f	Local data sets
		0.009		Cement production	1A2f	Operator's data
		0.144		Aircraft - support vehicles	1A3e	Bottom up model
		1.119		Industrial off-road	1A2fii	Bottom up model
Commercial	0.312					
Miscellaneous	0.215					
		0.038		Miscellaneous (UK)	1A4a	Reduced to offset higher consumption elsewhere
		0.007		Miscellaneous (CDs)	1A4a	
Public	0.249	0.024		Public sector	1A4a	Reduced to offset higher consumption elsewhere
Agriculture	0.146					
		1.147		Agriculture - mobile	1A4c	Bottom up model
Rail	0.645					
		0.604		Railways	1A3c	Bottom up model
	3.191	3.191	0.000			
		0.061		Upstream gas	1A1c	
		0.558		Upstream oil	1A1c	

DUKES Category	DUKES	GHGI	Difference	GHGI Category	CRF	Comment
Oil & gas	0.619	0.619	0.000			
National navigation	0.678	0.295		Shipping - coastal	1A3d	Revised UK/int'l split
		0.029		Fishing vessels	1A4c	
		0.209		Shipping - naval	1A3d	From MoD
		0.098		Motorboats	1A3d	Bottom up model
		0.002		Inland goods-carrying vessels	1A3d	Bottom up model
Marine bunkers	1.248	1.292		Shipping - int'l IPCC definition		Revised UK/int'l split
	1.926	1.926	0.000			
		0.010		House and garden machinery - DERV	1A4b	Bottom up model
		0.252		Industrial off-road mobile machinery - DERV	1A2fii	Bottom up model
		0.002		Sailing boats with auxiliary engines	1A3d	Bottom up model
		0.097		Motorboats / workboats	1A3d	Bottom up model
		21.522		Road transport - UK	1A3b	Reduced to offset data for off-road and other sources
		0.043		Road transport - CDs	1A3b	Local data sets
Road	21.926	21.926	0.000			
		0.123		Domestic (UK)	1A4b	
		0.006		Domestic (CDs)	1A4b	Local data sets
Domestic	0.129	0.129	0.000			
Total	27.791	27.791	0.000			

Notes: Sequences of shaded rows indicate categories which are grouped for purposes of data reconciliation, and should be considered together. "Total" includes marine bunkers.

A 4.2.1.5 Petroleum gases

For petroleum gases (LPG, OPG), a number of gaps in the UK energy statistics have been identified and as such, the total fuel use in the inventory is greater than the national statistics. These modifications to the energy balance are set out in **Table A 4.2.6 4.2.6**. They mostly

relate to refineries, use of feedstock as fuel in the petrochemicals sector, and fuel use for offshore oil and gas production.

Table A 4.2.6 Fuel reconciliation – Use of petroleum gases in 2013 (Mtherms)

DUKES Sector	DUKES	GHGI	Difference	GHGI sector	CRF	Comment
Petroleum refineries, other gases	996					
Autogenerators, other gases	94					
	1090	1177	+87	Refineries, OPG	1A1b	EU ETS higher than DUKES
Petroleum refineries, propane	3	3	0	Refineries, LPG	1A1b	
Iron & steel industry, propane	0	0	0	Iron & steel combustion, LPG	1A2a	
Industry (excluding iron & steel), propane	99					
Industry, butane	31					
Agriculture, propane	48					
Agriculture, butane	0					
		177		Industrial combustion, LPG - UK	1A2f	
		1		Industrial combustion, LPG - crown dependencies	1A2f	
	177	177	0			
Industry, ethane & other gases	0	539	+539	Petrochemical industry, use of off-gases	2B8g	Process off-gas use reported in EU ETS, outside scope of DUKES
Road, propane	44	44	0	Road transport - all vehicles LPG use	1A3b	
Domestic, propane	128					
Domestic, butane	13					
		132		Domestic combustion, LPG - UK	1A4b	
		8		Domestic combustion, LPG - crown dependencies	1A4b	
	141	141	0			

DUKES Sector	DUKES	GHGI	Difference	GHGI sector	CRF	Comment
(excluded from DUKES)		105	+105	Gas separation plant, OPG	1A1c	EEMS. Outside scope of DUKES
(excluded from DUKES)		5	+5	Gas separation plant, LPG	1A1c	EEMS. Outside scope of DUKES
Total	1455	2238	+782			LPG and OPG data from EEMS, EU ETS

Notes: Sequences of shaded rows indicate categories which are grouped for purposes of data reconciliation, and should be considered together.

1 Mtherm = 105.51 TJ

A 4.2.1.6 Modifications for other years

The previous subsections are focussed on deviations from the National Energy Balance for the latest inventory year. Many of the deviations outlined above are relevant for all of the time series, or more than one year.

For the 2013 inventory, some reallocations have been introduced for earlier years in the time series, which do not affect the latest year. This is because, where research is carried out and revisions are made to DUKES, the modifications are typically only made for the latest 5 years. For the 2013 inventory, modifications have been made to the allocation of fuels between energy and non-energy use. These have been incorporated into the inventory directly, however in order to ensure time series consistency, data have been estimated for earlier years in the time series for the inventory, leading to deviations from the energy balance in earlier years that are not highlighted in the tables above.

ANNEX 5: Additional Information to be Considered as Part of the Annual Inventory Submission and the Supplementary Information Required Under Article 7, paragraph 1, of the Kyoto Protocol Other Useful Reference Information.

A 5.1 ANNUAL INVENTORY SUBMISSION

No additional information.

A 5.2 SUPPLEMENTARY INFORMATION UNDER ARTICLE 7, PARAGRAPH 1

No additional information.

ANNEX 6: Verification

This Annex discusses the verification of the UK estimates of the Kyoto Gases.

A 6.1 MODELLING APPROACH USED FOR THE VERIFICATION OF THE UK GHGI

In order to provide verification of the UK Greenhouse Gas Inventory (GHGI), DECC (Department of Energy and Climate Change) have established and maintained a high-quality remote observation station at Mace Head (MHD) on the west coast of Ireland. The station reports high-frequency concentrations of the key greenhouse gases and is under the supervision of Prof. Simon O'Doherty of the University of Bristol (O'Doherty *et al.* 2004). DECC extended the measurement programme in 2012 with three new tall tower stations across the UK (UK DECC network): Tacolneston (TAC) near Norwich; Ridge Hill (RGL) near Hereford; Tall Tower Angus (TTA) near Dundee, Scotland.

The Met Office, under contract to DECC, employs the Lagrangian dispersion model NAME (Numerical Atmospheric dispersion Modelling Environment) (Ryall *et al.* 1998, Jones *et al.* 2007) driven by three-dimensional modelled meteorology to interpret the observations. NAME determines the history of the air arriving at Mace Head at the time of each observation. By estimating and removing the underlying *baseline* trends (Northern Hemisphere mid-latitude atmospheric concentrations where the short-term impact of regional pollution have been removed from the data) from the observations and by modelling where the air has passed over on route to Mace Head on a regional scale, estimates of UK emissions are made. A methodology called Inversion Technique for Emission Modelling (InTEM) has been developed that uses an iterative best-fit technique which searches a set of random emission maps to determine the one that most accurately mimics the Mace Head observations [Manning *et al.* 2003, 2011].

In this Chapter both the NAME *baseline* trends and the UK emission estimates are presented. InTEM estimates using only Mace Head (MHD) data are presented along with the estimates made using the full UK DECC network. When only MHD data are used a three-year inversion window is assumed (calculated up until 2013 for most gases), however with the additional data from the other UK stations the inversion time window has been shortened to one-year. The geographical spread of the UK DECC network allows the spatial distribution of the emissions across the UK to be better constrained within InTEM. The 'top-down' InTEM estimates of UK emissions are compared to the 'bottom-up' GHGI estimates.

A 6.2 METHANE

Figure A 6.2.1 shows the baseline atmospheric concentration of methane from 1990 onwards. The underlying trend is positive but there is strong year-to-year variability and a strong seasonal cycle. The growth rate over the last 8 years has been consistently positive.

In **Figure A 6.2.2** the emission estimates made for the UK with the InTEM methodology are compared to the GHGI emission estimates for the period 1990 onwards.

Methane has a natural (biogenic) component and it is estimated that 22% of the annual global emission is released from wetlands (Nilsson *et al.* 2001). Usually natural emissions are strongly

dependent on a range of meteorological factors such as temperature and diurnal, annual, growth and decay cycles. Such non-uniform emissions will add to the uncertainties in the modelling, although in North West Europe the natural emissions are thought to be small compared to the anthropogenic emissions (<5%, Bergamaschi *et al* 2005). Due to the relatively strong local (within 20km) influence of biogenic emissions at Mace Head, a peat bog area, the influence of observations taken when local emissions are significant (low wind speeds and low boundary layer heights) has been reduced within InTEM

The GHGI trend is monotonically downwards whereas the median of the InTEM estimates, after a rapid fall, shows only a very modest decline from 1993 onwards (**Figure A 6.2.2**).

The GHGI and InTEM emissions estimates are in significant disagreement in the early part of the time series. Recent Inventory research has investigated the historical data available to estimate emissions from landfill methane early in the time series and made a series of improvements. Nevertheless robust data remains scarce. It is considered probable that the discrepancy principally results from limited landfill methane data availability, specifically relating to volumes of gas flared, and this situation is unlikely to ever be improved upon. Note that the Inventory estimates are higher than InTEM during this period and are therefore conservative.

Figure A 6.2.1 Monthly Northern Hemisphere trend in methane estimated from Mace Head observations (blue line). Red line denotes the de-seasonalised long-term trend. Data under grey shading are not yet ratified.





Figure A 6.2.2 Verification of the UK emission inventory estimates for methane in Gg yr⁻¹ from 1990. GHGI estimates are shown in orange (uncertainty in red). InTEM (MHD-only, 3-year) estimates are shown in blue (uncertainty in light blue). InTEM (all data, 1-year) estimates are shown in black (uncertainty in grey).



A 6.3 NITROUS OXIDE

Figure A 6.3.1 shows the baseline atmospheric concentration of nitrous oxide from 1990 onwards. The annual trend is monotonic and positive at ~0.7 ppb yr⁻¹.

The main activities in Europe resulting in the release of nitrous oxide are agricultural practices resulting in emissions from soils (~60%), chemical industry (~20%) and combustion (~15%) (UNFCCC 1998 figures). The amount emitted from soils has significant uncertainty and has a diurnal and seasonal release cycle. It is driven by the availability of nitrogen, temperature and the soil moisture content.

Figure A 6.3.2 shows the InTEM and GHGI emission estimates for the UK for nitrous oxide for the period 1990 onwards. The median InTEM estimates are approximately 10-40 kt lower than the GHGI estimates up to 2011. The trends in the time-series are in good agreement up to 2011 with both show declining UK totals. The GHGI estimates show a sharp decline (40 Gg) between 1998 and 1999 in line with the introduction of the clean technology at an adipic acid plant in Wilton, north east England. It is estimated to have cut its emissions of N₂O by 90%, from 46 thousand tonne yr⁻¹ to around 6 thousand tonne yr⁻¹ (Defra, 2000). The InTEM estimates, with a longer averaging period, show a more gradual decline from 1998 to 2003 but the overall reduction is similar. In 2012 and 2013 InTEM emissions have risen, this is seen both when using only MHD observations and when observations from the whole UK DECC network are used. Although the emission uncertainties of both InTEM and GHGI are significant, the latter are considerably larger.

The nature of the nitrous oxide emissions challenges the InTEM assumption of uniformity of release both in time and space. Also the point of release to the atmosphere may not be coincident with the activity generating the nitrous oxide e.g. the nitrous oxide may be

transported from its source, for example by rivers to an ocean, prior to its release to the atmosphere.

Figure A 6.3.1 Monthly Northern Hemisphere trend in nitrous oxide estimated from Mace Head observations (blue line). Red line denotes the de-seasonalised long-term trend. Data under grey shading are not yet ratified.





Figure A 6.3.2 Verification of the UK emission inventory estimates for nitrous oxide in Gg yr⁻¹ from 1990. GHGI estimates are shown in orange (uncertainty in red). InTEM (MHD-only, 3-year) estimates are shown in blue (uncertainty in light blue). InTEM (all data, 1-year) estimates are shown in black (uncertainty in grey).



A 6.4 HYDROFLUOROCARBONS

A 6.4.1 HFC-134a

Figure A 6.4.1 shows the baseline atmospheric concentration of HFC-134a from 1995 onwards. The annual trend is monotonic and positive with a growth rate of over 4 ppt yr⁻¹.

Figure A 6.4.1 Monthly Northern Hemisphere trend in HFC-134a estimated from Mace Head observations (blue line). Red line denotes the de-seasonalised long-term trend. Data under grey shading are not yet ratified.





Figure A 6.4.2 shows the InTEM and GHGI emission estimates for the UK for HFC-134a for the period 1990 onwards. The GHGI shows a stronger increase in emission compared to the InTEM estimates. The InTEM estimates have risen at about 50% of the rate of the GHGI. From the 1997 onwards there is poor agreement between the GHGI and InTEM, with the InTEM estimates being about 60% of the GHGI estimates and well outside both uncertainty ranges. A similar result is obtained when the Tacolneston observations are included within InTEM. GHGI and InTEM both show the UK reducing its emissions from a peak in 2009.

Figure A 6.4.2 Verification of the UK emission inventory estimates for HFC-134a in Gg yr⁻¹ from 1990. GHGI estimates are shown in orange (uncertainty in red). InTEM (MHD-only, 3-year) estimates are shown in blue (uncertainty in light blue). InTEM (all data, 1-year) estimates are shown in black (uncertainty in grey).



A 6.4.2 HFC-152a


Figure A 6.4.3 shows the baseline atmospheric concentration of HFC-152a from 1995 onwards. The annual trend shows a strong rise from the mid-1990s until 2008, then a much reduced annual increase, from 2012 onwards no annual growth has been observed.

Figure A 6.4.3 Monthly Northern Hemisphere trend in HFC-152a estimated from Mace Head observations (blue line). Red line denotes the de-seasonalised long-term trend. Data under grey shading are not yet ratified.



Figure A 6.4.4 shows the InTEM and the GHGI emission estimates for the UK for HFC-152a for the period 1990 onwards. Between 2002 and 2008 the GHGI estimates are significantly larger than those estimated through the inversion modelling. Before 2002 and after 2008, the agreement is better with the median InTEM estimate falling within the GHGI uncertainty albeit consistently on the lower side. The InTEM uncertainties when only MHD data are used are large; they are significantly improved when the Tacolnaston data are included.

Figure A 6.4.4 Verification of the UK emission inventory estimates for HFC-152a in Gg yr⁻¹ from 1990. GHGI estimates are shown in orange (uncertainty in red). InTEM (MHD-only, 3-year) estimates are shown in blue (uncertainty in light blue). InTEM (all data, 1-year) estimates are shown in black (uncertainty in grey).



A 6.4.3 HFC-125


Figure A 6.4.5 shows the baseline atmospheric concentration of HFC-125 from 1998 onwards. The annual trend is monotonic and exponentially increasing. The growth rate is now over 2 ppt yr⁻¹.

InTEM emission estimates for the UK for HFC-125 for the period 1999 onwards are shown in **Figure A 6.4.6**. Both estimates suggest that the emissions of HFC-125 from the UK have increased significantly from the mid-1990s. The agreement between the 2 methods is excellent up until 2009 when InTEM reaches its peak. From 2010 the InTEM estimates show a modest decline in sharp contrast to the strongly increasing GHGI. By 2012 the difference between the GHGI and InTEM estimates is approximately 0.2 Gg yr⁻¹ (20%).

Figure A 6.4.5 Monthly Northern Hemisphere trend in HFC-125 estimated from Mace Head observations (blue line). Red line denotes the de-seasonalised long-term trend. Data under grey shading are not yet ratified.



Figure A 6.4.6 Verification of the UK emission inventory estimates for HFC-125 in Gg yr⁻¹ from 1990. GHGI estimates are shown in orange (uncertainty in red). InTEM (MHD-only, 3-year) estimates are shown in blue (uncertainty in light blue).



A 6.4.4 HFC-143a

Figure A 6.4.7 shows the baseline atmospheric concentration of HFC-143a from 2004 onwards. The annual trend is monotonic and positive at more than 1 ppt yr⁻¹.

InTEM emission estimates for the UK for HFC-143a for the period 2004 onwards are shown below in **Figure A 6.4.8** and are compared to the GHGI estimates. UK emissions, as estimated through the GHGI, are increasing year on year from the early 1990s. The InTEM estimates show a rise 2004-2009 and then a decline. The InTEM estimates are consistently higher than the GHGI estimates, with the uncertainty ranges not overlapping until 2010.

Figure A 6.4.7 Monthly Northern Hemisphere trend in HFC-143a estimated from Mace Head observations (blue line). Red line denotes the de-seasonalised long-term trend. Data under grey shading are not yet ratified.





Figure A 6.4.8 Verification of the UK emission inventory estimates for HFC-143a in Gg yr⁻¹ from 1990. GHGI estimates are shown in orange (uncertainty in red). InTEM (MHD-only, 3-year) estimates are shown in blue (uncertainty in light blue). InTEM (all data, 1-year) estimates are shown in black (uncertainty in grey).



A 6.4.5 HFC-23

Figure A 6.4.9 shows the baseline atmospheric concentration of HFC-23 from 2008 onwards. The annual trend is monotonic and positive, currently the growth rate is over 1 ppt yr⁻¹.

InTEM emission estimates for the UK for HFC-23 for 2008-2010 agree, within the uncertainty range, with the recent low emissions estimated by the GHGI (Figure A 6.4.10 and Figure A 6.4.11).

Figure A 6.4.9 Monthly Northern Hemisphere trend in HFC-23 estimated from Mace Head observations (blue line). Red line denotes the de-seasonalised long-term trend. Data under grey shading are not yet ratified.




Figure A 6.4.10 Verification of the UK emission inventory estimates for HFC-23 in Gg yr⁻¹ from 1990. GHGI estimates are shown in orange (uncertainty in red). InTEM (MHD-only, 3-year) estimates are shown in blue (uncertainty in light blue).





Figure A 6.4.11 Verification of the UK emission inventory estimates for HFC-23 in Gg yr⁻¹ zoomed in for 2004-2013. GHGI estimates are shown in orange (uncertainty in red). InTEM (MHD-only, 3-year) estimates are shown in blue (uncertainty in light blue).



A 6.4.6 HFC-32

Figure A 6.4.12 shows the baseline atmospheric concentration of HFC-32 from 2004 onwards. The annual trend is monotonic and positive. The growth rate is currently over 1 ppt yr⁻¹.

InTEM emission estimates for the UK for HFC-32 for 2004 onwards are shown in **Figure A 6.4.13**. The InTEM emission estimates are lower than the GHGI estimates. Both trends are positive however the rate of increase of the GHGI is larger than the InTEM. By 2012 the difference in estimated emissions is significant at more than 30%.

Figure A 6.4.12 Monthly Northern Hemisphere trend in HFC-32 estimated from Mace Head observations (blue line). Red line denotes the de-seasonalised long-term trend. Data under grey shading are not yet ratified.





Figure A 6.4.13 Verification of the UK emission inventory estimates for HFC-32 in Gg yr⁻¹ from 1990. GHGI estimates are shown in orange (uncertainty in red). InTEM (MHD-only, 3-year) estimates are shown in blue (uncertainty in light blue).



A 6.4.7 HFC-43-10mee

Figure A 6.4.14 shows the baseline atmospheric concentration of HFC-43-10mee from 2011 onwards. There is a slight positive trend with a growth rate of ~0.01 ppt yr⁻¹. The UK emissions of this gas are small. The GHGI estimates are higher than InTem but the agreement improves substantially in 2013.

Figure A 6.4.14 Monthly Northern Hemisphere trend in HFC-43-10mee estimated from Mace Head observations (blue line). Red line denotes the de-seasonalised long-term trend. Data under grey shading are not yet ratified.





Figure A 6.4.15 Verification of the UK emission inventory estimates for HFC-43-10mee in Gg yr⁻¹ from 1990. GHGI estimates are shown in orange (uncertainty in red). InTEM (MHD-only, 3-year) estimates are shown in blue (uncertainty in light blue). InTEM (all data, 1-year) estimates are shown in black (uncertainty in grey).



A 6.4.8 HFC-227ea

Figure A 6.4.16 shows the baseline atmospheric concentration of HFC-227ea from 2007 onwards. There is positive trend with a growth rate of ~0.1 ppt yr⁻¹. The GHGI estimates are more than double those obtained through inversion modelling.

Figure A 6.4.16 Monthly Northern Hemisphere trend in HFC-227ea estimated from Mace Head observations (blue line). Red line denotes the de-seasonalised long-term trend. Data under grey shading are not yet ratified.





Figure A 6.4.17 Verification of the UK emission inventory estimates for HFC-227ea in Gg yr⁻¹ from 1990. GHGI estimates are shown in orange (uncertainty in red). InTEM (MHD-only, 3-year) estimates are shown in blue (uncertainty in light blue). InTEM (all data, 1-year) estimates are shown in black (uncertainty in grey).



A 6.4.9 HFC-365mfc

Figure A 6.4.18 shows the baseline atmospheric concentration of HFC-365mfc from 2005 onwards. There is positive trend with a growth rate of ~0.07 ppt yr⁻¹. InTEM and the GHGI agree, within the uncertainties, for the first part of the atmospheric record 2005-2008. Both estimates then show a decline in UK emissions, however the GHGI estimates have a slower rate of reduction and are ~50% higher than InTEM 2009-2013.

Figure A 6.4.18 Monthly Northern Hemisphere trend in HFC-365mfc estimated from Mace Head observations (blue line). Red line denotes the de-seasonalised long-term trend. Data under grey shading are not yet ratified.





Figure A 6.4.19 Verification of the UK emission inventory estimates for HFC-365mfc in Gg yr⁻¹ from 1990. GHGI estimates are shown in orange (uncertainty in red). InTEM (MHD-only, 3-year) estimates are shown in blue (uncertainty in light blue). InTEM (all data, 1-year) estimates are shown in black (uncertainty in grey).



A 6.4.10 HFC-245fa

Figure A 6.4.20 shows the baseline atmospheric concentration of HFC-245fa from 2007 onwards. There is positive trend with a growth rate of ~0.2 ppt yr⁻¹. The InTEM estimates are higher than the GHGI estimates however they agree within the inversion uncertainties. No observations are available from Tacolneston.

Figure A 6.4.20 Monthly Northern Hemisphere trend in HFC-245fa estimated from Mace Head observations (blue line). Red line denotes the de-seasonalised long-term trend. Data under grey shading are not yet ratified.





Figure A 6.4.21 Verification of the UK emission inventory estimates for HFC-245fa in Gg yr⁻¹ from 1990. GHGI estimates are shown in orange (uncertainty in red). InTEM (MHD-only, 3-year) estimates are shown in blue (uncertainty in light blue).



A 6.5 PERFLUOROCARBONS

A 6.5.1 PFC-14

Figure A 6.5.1 shows the baseline atmospheric concentration of PFC-14 from 2004 onwards. The annual trend is monotonic and positive with a growth rate of around 0.7 ppt yr⁻¹. Within the uncertainty ranges of the InTEM and GHGI estimates, the UK emissions agree. The sharp drop in emissions in 2012 in the GHGI reflects the closure of the last significant aluminium production plant in the UK. The InTEM uncertainty ranges for PFC-14 are large because the overwhelming majority of emissions come from a few point sources (smelters), which are not well captured in the large area averages within InTEM. If prior knowledge of the point sources

is included within InTEM the uncertainty ranges are considerably reduced. InTEM estimates a further significant reduction in UK emissions in 2014.

Figure A 6.5.1 Monthly Northern Hemisphere trend in PFC-14 estimated from Mace Head observations (blue line). Red line denotes the de-seasonalised long-term trend. Data under grey shading are not yet ratified.





Figure A 6.5.2 Verification of the UK emission inventory estimates for PFC-14 in Gg yr⁻¹ from 1990. GHGI estimates are shown in orange (uncertainty in red). InTEM (MHD-only, 3-year) estimates are shown in blue (uncertainty in light blue). InTEM (all data, 1-year) estimates are shown in black (uncertainty in grey).



A 6.5.2 PFC-116

Figure A 6.5.3 shows the baseline atmospheric concentration of PFC-116 from 2004 onwards. The annual trend is monotonic and positive at around 0.1 ppt yr⁻¹.

The UK InTEM estimates are consistent with those reported in the GHGI (**Figure A 6.5.4**) given the significant uncertainties in the InTEM solutions.

Figure A 6.5.3 Monthly Northern Hemisphere trend in PFC-116 estimated from Mace Head observations (blue line). Red line denotes the de-seasonalised long-term trend. Data under grey shading are not yet ratified.





Figure A 6.5.4 Verification of the UK emission inventory estimates for PFC-116 in Gg yr⁻¹ from 1990. GHGI estimates are shown in orange (uncertainty in red). InTEM (MHD-only, 3-year) estimates are shown in blue (uncertainty in light blue). InTEM (all data, 1-year) estimates are shown in black (uncertainty in grey).



A 6.5.3 PFC-218

Figure A 6.5.5 shows the baseline atmospheric concentration of PFC-218 from 2004 onwards. The annual trend is monotonic and positive at around 0.02 ppt yr⁻¹.

The median UK InTEM estimates are higher than those reported in the GHGI (Figure A 6.5.6). However within the uncertainty ranges of the InTEM results, the two methods agree. Similar to PFC-14, without prior knowledge of the point source nature of the PFC-218 emissions, InTEM struggles to constrain the emissions.

Figure A 6.5.5 Monthly Northern Hemisphere trend in PFC-218 estimated from Mace Head observations (blue line). Red line denotes the de-seasonalised long-term trend. Data under grey shading are not yet ratified.





Figure A 6.5.6 Verification of the UK emission inventory estimates for PFC-218 in Gg yr⁻¹ from 1990. GHGI estimates are shown in orange (uncertainty in red). InTEM (MHD-only, 3-year) estimates are shown in blue (uncertainty in light blue). InTEM (all data, 1-year) estimates are shown in black (uncertainty in grey).



A 6.5.4 PFC-318

Figure A 6.5.7 shows the baseline atmospheric concentration of PFC-318 from 2010 onwards. The annual trend is monotonic and positive at around 0.05 ppt yr⁻¹.

The median UK InTEM estimates are significantly higher than those reported in the GHGI (Figure A 6.5.8). However the uncertainty ranges of the InTEM results are large and the estimated quantities are very small.

Figure A 6.5.7 Monthly Northern Hemisphere trend in PFC-318 estimated from Mace Head observations (blue line). Red line denotes the de-seasonalised long-term trend. Data under grey shading are not yet ratified.





Figure A 6.5.8 Verification of the UK emission inventory estimates for PFC-318 in Gg yr⁻¹ from 1990. GHGI estimates are shown in orange (uncertainty in red). InTEM (MHD-only, 3-year) estimates are shown in blue (uncertainty in light blue). InTEM (all data, 1-year) estimates are shown in black (uncertainty in grey).



A 6.6 SULPHUR HEXAFLUORIDE

Figure A 6.6.1 shows the baseline atmospheric concentration of SF₆ from 2004 onwards. The annual trend is monotonic and positive, the current growth rate is now 0.35 ppt yr⁻¹.

The median UK InTEM estimates are higher than those reported in the GHGI by ~0.015 Gg yr⁻¹ (~20%) (**Figure A 6.6.2**). However, the InTEM uncertainty ranges incorporate the GHGI estimates entirely and the trends in emissions are very similar. When the TAC data are included the agreement between the GHGI and InTEM is excellent.

Figure A 6.6.1 Monthly Northern Hemisphere trend in SF₆ estimated from Mace Head observations (blue line). Red line denotes the de-seasonalised long-term trend. Data under grey shading are not yet ratified.





Figure A 6.6.2 Verification of the UK emission inventory estimates for SF₆ in Gg yr⁻¹ from 1990. GHGI estimates are shown in orange (uncertainty in red). InTEM (MHD-only, 3-year) estimates are shown in blue (uncertainty in light blue). InTEM (all data, 1-year) estimates are shown in black (uncertainty in grey).




A 6.7 NITROGEN TRIFLUORIDE

Figure A 6.7.1 shows the baseline atmospheric concentration of NF₃ from 2013 onwards. The annual trend is monotonic and positive, the current growth rate is estimated to be greater than 0.1 ppt yr⁻¹.

NF₃ is only measured at MHD, and measurements only started in May 2013. The InTEM emission estimates for the UK are insignificant. The GHGI estimate for 2013 is 21 kg.

Figure A 6.7.1 Monthly Northern Hemisphere trend in NF_3 estimated from Mace Head observations (blue line). Red line denotes the de-seasonalised long-term trend. Data under grey shading are not yet ratified.



A 6.8 CARBON DIOXIDE

High precision, high frequency measurements of CO_2 are made across the UK DECC network. The CO_2 observed has three principle components:

1. Northern hemisphere baseline (**Figure A 6.8.1**).
2. Anthropogenic (man-made)
3. Biogenic (natural)

Figure A 6.8.1 Monthly Northern Hemisphere trend in CO_2 estimated from Mace Head observations (blue line). Red line denotes the de-seasonalised long-term trend. Data under grey shading are not yet ratified.





Figure A 6.8.2 Verification of the UK emission inventory estimates for CO₂ in Gg yr⁻¹ for 1990-2011. GHGI estimates are shown in orange (uncertainty in red). InTEM (MHD-only, 3-year, using CO ratio) estimates are shown in blue (uncertainty in light blue). InTEM (all network, winter & daytime data only) estimate shown in black (uncertainty in black).



Plants both respire CO₂ and absorb it through photosynthesis. Therefore the CO₂ flux from vegetation has a diurnal and seasonal cycle and switches from positive to negative on a daily basis. This unknown natural (biogenic) component of the observed CO₂ is significant when compared to the anthropogenic (man-made) component and cannot be assumed negligible (except maybe during the winter months). From the CO₂ observations it is not possible to distinguish between biogenic and anthropogenic CO₂. Therefore it is difficult to use the CO₂ observations directly in an inversion to estimate anthropogenic emissions. This is because the diurnally varying biogenic CO₂ flux is at odds with a key assumption of the inversion method, namely that emissions do not strongly vary in time over the inversion time-window (one or three years in the standard analysis). Methods are under development to attempt to overcome these challenges, such as: the use of isotopic observations; through ratios with respect to anthropogenic CO; and through the use of just the winter-time observations. The uncertainties associated with each of these methods are predicted to be significant.

Figure A 6.8.2 are the preliminary results for UK emissions of CO₂ using the InTEM inversion results for CO. The InTEM CO emission maps have been scaled by the annually varying UK inventory ratio of CO₂:CO emissions (after removal of the CO₂ emissions estimated to be released from power stations – these are assumed to emit little CO due to abatement technologies). The InTEM uncertainties have been arbitrarily increased to a minimum of ±200,000 Gg yr⁻¹ to reflect the fact that the CO₂:CO ratio is variable across applications across the UK.

Also plotted is the InTEM estimate using the UK DECC network CO₂ observations (2012-2014). Only daytime (10am – 4pm), winter-time (15 Nov – 15 Feb) observations were included. The estimates will include some (unquantifiable) contribution from biogenic emissions and so should be larger than the reported anthropogenic-only inventory estimates. By choosing only

daytime, winter-time observations the impact of biogenic emissions is minimised but they are not eliminated entirely.

The estimated uncertainties in the inventory are also presented. It can be noted that the uncertainties in the InTEM results are considerably larger than the inventory uncertainties. Work is on-going to seek to improve our methods of verifying inventory CO₂ emission estimates.

ANNEX 7: Analysis of EU ETS Data

A 7.1 INTRODUCTION

This annex summarises the analysis of the 2013 European Union Emissions Trading System (EU ETS) energy and emissions data that is used within the compilation of the UK GHG inventory. The EU ETS data are used to inform activity data estimates for heavy industry sectors, carbon dioxide emission factors of UK fuels within those sectors, and for comparison of fuel allocations to specific economic sectors against data presented in the *Digest of UK Energy Statistics* (DUKES), published by the Department of Energy and Climate Change (DECC).

The EU ETS data are used in the UK GHGI compilation as follows:

- EU ETS raw data on energy and emission estimates are processed and checked to enable integration of the activity data, implied emission factors and installation emission estimates as far as practicable within the UK GHG inventory compilation. Emission sources reported in EU ETS are allocated to inventory fuels and source codes, outliers are identified and clarifications of data inconsistencies are sought with the regulatory agencies;
- The verified EU ETS data provides up to date high quality fuel compositional analysis of UK fuels, and these data are used to improve inventory emission estimates across the highly energy intensive sources such as power stations, refineries, cement kilns, and oil and gas sources;
- EU ETS activity data are closely compared against the UK national energy balance (DUKES) published by DECC, and any inconsistencies are researched, seeking to resolve these through consultation with DECC wherever possible;
- The EU ETS dataset for offshore oil and gas installations are checked to assess data consistency in emissions reporting between the EU ETS and the (more comprehensive) EEMS dataset that is used within the UK GHGI compilation;
- Overall, the Inventory Agency approach seeks to minimise data discrepancies between EU ETS and the GHGI as far as practicable, in order that the derivation of traded and non-traded emission estimates from the UK GHGI are as accurate as possible. Close consistency between the EU ETS and GHGI is an important aspect of the development of a complete and consistent evidence base for policy development and tracking progress towards UK GHG reduction targets in the non-traded sector under the EU Effort Sharing Decision.

The scope of reporting under EU ETS has increased from the 2013 dataset onwards. Phase II of the EU ETS ran from 2008-2012 inclusive. Phase III reporting began in 2013, with some new emission sources and new installations now reporting for the first time on their GHG emissions; in particular, the definition of combustion has now been extended to cover installations such as furnaces, driers, and other plant where heat is used directly. A handful of industrial process sources of CO₂ are also included for the first time, such as soda ash production. The UK Inventory Agency has analysed the 2013 dataset in order to use, where possible, any new data on emissions from the first year of EU ETS Phase III reporting. In the UK, the changes in reporting in Phase III are most significant for the chemicals sector, where the scope of reporting is larger than previously and now encompasses both new industrial

process emission sources, and additional energy use. There is also a notable shift towards estimation methods that are based on mass balance calculations (e.g. for chemical manufacturing) within the UK operator reporting to EU ETS. Other sectors with significant increases in reporting are food and drink manufacture, where installations such as driers, ovens etc. were included for the first time thus adding to the emissions from boilers and CHP plant that were reported in previous years, and roadstone coating, a sector which has not been present in the dataset before. Analysis of the 2013 dataset has enabled the Inventory Agency to improve estimates of emissions from the combustion of waste residues and process off-gases within the chemical and petrochemical sectors (which are all reported under IPPU sector 2B10), as well as estimate the IPPU component of several specific manufacturing processes, such as for soda ash (2B7), and titanium dioxide (2B6). In addition, following a review of methodology for all IPPU sources, EU ETS data for phase II onwards has been used to improve emission estimates for glass production (2A3), brickmaking (2A4) and reductant use in electric arc furnaces (2C1).

The key findings from the analysis and use of the EU ETS data include:

- In the 2013 EU ETS dataset, a very high coverage of Tier 3 emissions data is evident for all fuel use in the power sector, coal use in the lime sector, and refinery fuel oil and OPG use. All of the fuel quality data for these sources and fuels are therefore used within the UK GHGI, as the EU ETS fuel quality data is the most representative dataset available to inform UK carbon dioxide emission factors in the inventory;
- EU ETS emissions data from refineries are higher than estimates derived from DUKES activity data, with a discrepancy evident in OPG emissions. Consultation with the industry trade association, UKPIA, and cross-checking with their data shows that the EU ETS data are a true reflection of industry estimates of CO₂, and therefore UK GHGI estimates are based on EU ETS data rather than refinery fuel use data reported in the UK energy balance;
- There are a range of other activity data discrepancies within the oil & gas, cement and lime, other industry and iron and steel sectors. Revisions to fuel allocations within the UK GHGI have been implemented for a number of sources, whilst further research is needed in some instances to clarify the issues where the reporting format of EU ETS does not map explicitly to energy balance and GHG inventory reporting requirements;
- EU ETS data for fuel use at chemical and petrochemical production facilities has helped to identify and quantify under-reports within the UK energy statistics for the combustion of gases that are derived from Natural Gas Liquid (NGL) feedstock to petrochemical production processes, and from combustion of carbon-containing process residues. Analysis of “fuel gas” calorific values and carbon content has helped to inform the calculations to estimate emissions from NGL-derived gases and other residues, to address the under-report in UK energy statistics.

The use of EU ETS data in the UK GHG inventory is summarised in **Table A 7.1.1**.

Table A 7.1.1 Summary of the use of EU ETS data in the UK inventory

Category	Sub-categories	EU ETS data used			Comments
		Factors	Activity	Emission _s	
1A1a	Power stations - coal, fuel oil natural gas, sour gas	✓			
1A1a	Power stations – pet coke			✓	Some additional data is sourced from process operators.
1A1b	Refineries – pet coke & OPG			✓	EU ETS figures only used where higher than DUKES-based emissions.
1A1b	Refineries – natural gas	✓			
1A1c	Gas industry – natural gas		✓		
1A1c 1B1b 1A2a 2C1	Integrated steelworks	✓	✓		Use of various EU ETS data in complex carbon balance – factors for some fuels, activity data for others
1A1c	Collieries – Colliery methane	✓			
1A2b	Autogenerators - coal	✓			
1A2f	Lime - coal			✓	
1A2f	Lime – natural gas		✓		
1A2g	Industry - pet coke & waste solvents			✓	No alternative data available for this emission source.
1A2g	Industry – colliery methane	✓			
2A1	Cement			✓	Data used is actually from industry trade-association, but this is based on EU ETS returns
2A2	Lime			✓	
2A3	Glass			✓	
2A4	Bricks			✓	
2B7	Soda ash			✓	
2B8g	Ethylene & other petrochemicals			✓	
2C1	Electric arc furnaces - reductants			✓	

A 7.2 BACKGROUND

A 7.2.1 EU ETS Data and GHG Inventories

The European Union Emissions Trading System (EU ETS) data provides annual estimates of fuel use and fuel quality data from the most energy intensive sites in the UK, and provides a source of data, or can be used to cross-check data held in the UK Greenhouse Gas Inventory (GHGI), and to inform the carbon contents of current UK fuels. The EU ETS has operated since 2005, and there are now 9 years' worth of data on fuel use and emissions across major UK industrial plant, for 2005-2013.

The data reported under the EU ETS includes quantities of fuels consumed (or other activity data for process sources of CO₂), carbon contents of fuels and other inputs, calorific values (fuels only) and emissions of carbon dioxide, all presented by installation and by emission source. Activity data are also given for many biofuels, although emissions of CO₂ from these fuels are not included in the emissions data. This is useful though, since PI/SPRI/NIPI emissions data for CO₂ often include biocarbon as well as fossil carbon, and the EU ETS data on biofuels helps to explain differences between CO₂ emissions reported in EU ETS and in the PI/SPRI/NIPI. EU ETS data for individual installations are treated as commercially confidential by the UK regulatory authorities and so only aggregated emissions data are reported in inventory outputs.

As part of the UK's annual reporting requirements to the MMR and UNFCCC, the UK must include a comparison of the EU ETS data against the national inventory dataset within the National Inventory Report. Furthermore, the analysis of the inventory against the EU ETS dataset is coming under increasing scrutiny due to the development of domestic GHG reduction targets that are based on non-traded⁹ emissions data only, and the growing need to understand the UK non-traded sector emissions for future reporting under the Effort Sharing Decision.

The EU ETS dataset helps to improve the UK GHG inventory in a number of ways:

- Identifying new sources, therefore improving completeness;
- Helping assess true levels of uncertainty in fuel- and sector-specific data;
- Providing fuel quality data and oxidation factors for complex processes;
- Providing information on process-specific emissions that are not apparent from the national energy balances;
- Reducing uncertainty in the GHGI; and
- Acting as a source of quality assurance to inventory data.

In the 1990-2013 inventory cycle, the Inventory Agency has updated and extended the EU ETS analysis conducted for inventory compilation, using the 2013 EU ETS dataset, which is the first year of reporting under the Phase III EU ETS scope and includes approximately 180 new installations in the UK compared to the Phase II reporting scope (2008-2012), with increased scope of reporting at many more. This annex presents a comprehensive review of

⁹ All GHG emissions that are regulated within the EU ETS are defined as "traded" emissions, whilst all other GHG emissions are defined as "non-traded". The EU Effort Sharing Decision will lead to the UK adopting a new target for GHG reductions by 2020 for all of the non-traded emissions (i.e. everything outside of EU ETS), and progress towards this target will be monitored through the UK GHG inventory.

the nine years' of EU ETS data, indicating where the data have already been used in the improvement of the GHGI, as well as highlighting outstanding issues which could be investigated further, with potential for further revision and improvement of the GHGI.

The Inventory Agency has also been provided with full details of the 2011-2013 EU ETS data for all offshore oil and gas installations, which are regulated by the DECC Offshore Inspectorate. Access to these detailed data has enabled a more thorough review of the fuel/gas quality and reported emissions from combustion and flaring sources at offshore installations, and has directly improved the completeness and accuracy of the sector estimates within the UK GHGI.

The analysis of the EU ETS data for use in the UK GHGI necessitates a detailed review of the available data, in order to ensure correct interpretation and application of the available data. The study team prioritises effort to the sources and sites that are the most significant in UK GHGI terms, and/or where data reporting discrepancies have been identified from previous work. For those sectors where EU ETS data are used in the GHGI, it is important to review emission factors from all major installations to ensure that any outliers are identified and checked prior to their inclusion in inventory calculations.

Wherever possible, consistent assumptions are made when interpreting data across all years of the EU ETS. For instance ensuring that each site is allocated to the same inventory sector in each year (unless there is reason to change it – some industrial combustion plant in recent years have been converted into power stations, and so these sites do need to be allocated to different sectors in different years), and that there is consistency in the way in which site-specific names for fuels are interpreted across the entire period. The information on the EU ETS method “Tier” used for each of the data dictates whether they are used in inventory compilation. The highest tier EU ETS data are assumed to be subject to the lowest level of uncertainty, and so only tier 3 and tier 4 data are used. Occasionally there are internal inconsistencies in the EU ETS data between the data on consumption of a given fuel and emissions from the use of that fuel. These need to be resolved before the data can be used in the UK GHGI. As emissions data are verified, we cross-check the detailed emissions data against the final verified emissions for each site. As a general rule it is found that the most appropriate solution to inconsistencies is to assume that the EU ETS emissions data are correct as EU ETS reporting requirements are well regulated, and that it is the activity data that need to be amended instead.

A 7.2.2 Scope of the UK EU ETS and Implications for the GHG Inventory

There are a number of limitations to the EU ETS data that affect the data usefulness in GHG inventory compilation, including:

- The EU ETS data are only available from 2005 onwards, whilst the UK GHG inventory reports emission trends back to 1990. The additional information that EU ETS provides (e.g. year-specific emission factors for many fuels in energy intensive sectors) helps to reduce the uncertainties in inventory emission estimates for the later years, but care is needed where revisions to the time series are made back to 2005. A consistent approach to inventory compilation across the time series is a key tenet of IPCC good practice guidance, and care is needed to ensure that the use of EU ETS data does not introduce a systematic reporting step-change in the UK GHGI;

- Further to this point, it is important to note that the scope of EU ETS reporting has evolved through the years, from Phase I (2005 to 2007) into Phase II (2008 to 2012 data) and now to Phase III (2013 onwards). The comparability of EU ETS data for many sectors is poor between these three phases. For example, many cement kilns did not report to EU ETS until Phase II: several sectors including cement were reporting under Climate Change Agreements and were opted-out of EU ETS during Phase I. Therefore in several sectors, more complete coverage of EU ETS reporting is evident in Phase II and data from 2008 onwards are therefore much more useful for UK GHGI reporting. The scope of coverage of chemical industry emissions has gone through two step changes – in 2008, and again in 2013, and some sectors (such as roadstone coating) only appear for the first time in the 2013 EU ETS data. Less significantly for the GHGI, many small installations, mostly in the public sector, were removed from EU ETS at the end of phase I. It is vital that the GHGI takes full account of such changes and that UK inventory data do not include trends that merely result from the increase (or decrease) in scope of EU ETS. The changes in EU ETS scope have made the data set increasingly useful, and there are now five years' worth of Phase II data and one year of data under Phase III, hence the EU ETS dataset is now an important source of information for the UK inventory;
- In the UK during EU ETS Phases I and II, the regulators adopted a “medium” definition of the term “combustion”, and there are many sectors where fuel use in specific types of combustion unit have not been included in the EU ETS reporting scope until the start of Phase III (2013 onwards). Examples of this include flaring on chemical sites, fuel use in heaters, dryers, fryers and stenters in industry sectors such as: chemicals, food and drink, textiles, paper and pulp. Hence the total fuel use and GHG emissions from these sectors have typically been under-reported within the EU ETS historically, with many sites and sources excluded from the scope of EU ETS. However, the EU ETS data for these sectors is also incomplete both in phase II and phase III because small installations are not covered by EU ETS. Therefore, while the change in scope for combustion installations is a positive step, it has relatively little impact on the data used in GHGI compilation. Some phase III data has been used to improve the estimates of combustion of process wastes / off-gases in the chemical and petrochemical sector in the 2015 submission.

Phase III also brought an increased scope for industrial process sources of CO₂, and data appeared for the first time for soda ash production, and titanium dioxide manufacture. The data for soda ash production were used in the GHGI compilation, as were the partial data set available for titanium dioxide manufacture. Full data for the latter will be obtained in time for the next submission and the UK inventory updated if necessary.
- When using the EU ETS data, assumptions and interpretations are required to be made regarding the fuel types used by operators; assumptions are made on a case by case basis depending on knowledge of the site or industry and expert judgement. Operators are free to describe fuels as they wish in their returns, rather than choosing from a specific list of fuels, and so assumptions occasionally need to be made where the fuel type used is not clear from the operator's description of it. This issue was more significant in the earlier years of EU ETS reporting, with operators often using terms such as “Fuel 1”. The assumption then made about fuel type was based both on the other data the operator provided on the fuel such as calorific value, but also by

comparison with later data for the same site, since operators now tend to use standard fuel names.

Note that:

- The direct use of EU ETS data (e.g. fuel use data by sector) to inform UK GHGI estimates is limited to where the EU ETS is known to cover close to 100% of sector installations. For example, the EU ETS is regarded as representative and almost 100% comprehensive in coverage of refineries, power stations, cement and lime kilns; for many other industrial sectors (such as chemicals, non-ferrous metals, food and drink, engineering) the EU ETS is not comprehensive and therefore the data are of more limited use, mainly providing a de-minimis fuel consumption figure for these sectors;
- EU ETS Implied Emissions Factors (IEFs) can be used within the UK GHGI, but only where the evidence indicates that EU ETS data are representative of sector as whole and provides more comprehensive and accurate data than alternative sources. The key criteria to consider in the assessment of EU ETS IEF usefulness is the percentage of annual fuel use by sector where operator estimates use Tier 3 emission factors.
- Review of the EU ETS IEFs for different fuels across different sites provides a useful insight into the level of Tier 3 reporting within different sectors, the progression of higher-Tier reporting within EU ETS through the time series and the level of variability in fuel quality for the different major fuels in the UK. As a general rule, those energy-intensive sectors with near 100% coverage in EU ETS also report a very high proportion of emission factor data at Tier 3. Those sectors with incomplete coverage tend to report most emission factor data below Tier 3. As a result, in all cases where the level of sectoral coverage is high, the quality of reported data is also sufficiently high to be used with confidence in the UK inventory.

A 7.2.3 Limitations of EU ETS Data Integration with GHG Inventory: Autogeneration

Despite detailed research there remain some fundamental limitations in the use of EU ETS data within national inventories where the sector allocation of energy use and emissions cannot be resolved against the national energy statistics that underpin the GHG inventory compilation. One key example is that of the division between fuel use in autogeneration (or heat generation) and direct fuel use within a specific sector. For example, based on the data available from EU ETS, it is impossible to differentiate between gas use in autogeneration on a chemical installation, and gas used directly to heat chemical production processes. In this example, the allocation of EU ETS energy use and emissions between 1A2c (chemicals) and 1A2f (autogenerators) is uncertain, and therefore comparison of EU ETS and GHGI estimates is uncertain. The EU ETS data are not sufficiently detailed and transparent to enable accurate allocation, and so in all cases fuels and emissions are allocated to the industry sector, and not to autogeneration.

It is worth noting here that the UK energy statistics are also subject to some uncertainty, however small, and that there is likely to be more uncertainty in estimates at industrial sector-level, rather than at more aggregated levels. For example, while fuel producers and suppliers will be able to quantify total fuel demand with a high level of certainty, it would be far more difficult for them to estimate fuel use by specific industrial sectors. This will be reflected in the quality of UK energy statistics which are used to estimate emissions from 1A2c etc. We

consider that a high proportion of fossil fuel use by the UK chemical industry will be included in the EU ETS, on the basis that most industrial chemical processes will require sufficiently large combustion installations to exceed the threshold for EU ETS. Therefore, we consider that it is reasonable to assume that EU ETS emissions for the chemical sector should cover most of the sector and therefore be similar in magnitude to those estimated from UK energy statistics and even, given the uncertainty in fuel allocation for autogeneration, to exceed them. For other sectors such as metals, paper, and food and drink, we would assume that the level of sectoral coverage by the EU ETS would be lower, so that emission estimates based on EU ETS would probably be lower than those based on energy statistics, even taking into account the uncertainty regarding autogeneration.

A 7.3 DATA PROCESSING

DECC provided the detailed EU ETS regulator data from the Environment Agency, Scottish Environment Protection Agency and Northern Ireland Environment Agency during May 2014, and the Inventory Agency industrial emissions experts progressed the analysis, combining the datasets to generate a UK-wide EU ETS dataset. The work built on analysis conducted in previous years, as the EU ETS has been in place since 2005, but this latest analysis, while focussing on the 2013 submissions, did involve review of the complete 9-years of data, to ensure a consistent approach to the interpretation of energy and emissions data across the time series.

The initial step in the analysis is the allocation of all sites in the dataset to one of the economic sectors as reported within the DUKES Commodity Balance tables. Next, the reported fuels for every UK installation have to be allocated to one of the GHGI fuel names, which are also aligned with the fuel types reported within DUKES. This enables a direct comparison of EU ETS fuel totals against sector fuel allocations within DUKES and therefore used within the GHGI.

Most of the allocations have been made as part of previous years' work, and do not need to be revisited. However, the start of phase III, meant that there were approximately 180 new installations included in the 2013 EU ETS data which had to be allocated to DUKES' sectors, and all of the fuel data for 2013 also has had to be allocated to DUKES/GHGI fuel types. In a very small number of cases, we have revised data for earlier years, for example when it has become apparent that existing assumptions are likely to be incorrect. The allocation process does rely upon some expert judgement, with the Ricardo-AEA team using the reported EU ETS fuel names as well as the reported fuel quality data such as calorific values and carbon emission factors in order to make the fuel-type allocation for each entry in the EU ETS spreadsheet. The allocation is, occasionally, quite uncertain, particularly with the allocation of petroleum-based fuels such as the GHGI fuel categories LPG, OPG, gas oil and fuel oil, often because of the use of abbreviations or other ambiguous names for fuels within the EU ETS reporting system. Cross-checking of data across the time series for each installation has been used to ensure as much consistency in fuel allocations as possible, although in some cases, operators of installations use different fuel terminology in different years.

The quality checking and allocation process is an open-ended task for such a large dataset, and hence the Inventory Agency focuses on the highest emitters and the known "problem" sites and fuel types. Where uncertainties arise in allocations, the most important allocation decisions are copied across to the DECC DUKES team, for their information and input, as

ultimately the EU ETS analysis by the Inventory Agency is taken into account to some degree within the compilation of DUKES for the following year.

As a data verification step, the installation emissions (broken down by fuel) from the EU ETS regulator spreadsheets are then compared against the total installation emissions for 2013 on the European Union Transaction Log (EUTL) which is a central website that holds the verified EU ETS emissions totals for all EU installations in the scheme. Each year we have noted that for some sites the regulator data does not match the EUTL dataset, and therefore some “residual” emissions allocations are generated, from the difference between EUTL and regulator information. In cases where these residual emissions are large, then these are fed back to the regulator contacts, for their consideration and to request any insights into the likely fuels that the residual emissions should be allocated against. Minor residual emissions are ignored for the purposes of the analysis reported here.

A final data set is then available for fuel combustion emission sources, which includes the following data fields:

- GHGI Source Category;
- GHGI Fuel Category;
- Fuel Consumed;
- Fuel Calorific Value;
- Fuel Carbon Emission Factor; and
- Related Emissions of CO₂

The Inventory Agency then combines the data by sector and/or fuel category to provide data for comparison against GHGI emissions data, and energy statistics published in DUKES. In this way, the analysis can:

- provide improved CO₂ emission factors for highly energy-intensive industrial sectors covered by the GHGI through the use of verified data;
- provide a comparison with UK energy statistics, allowing the identification of inconsistencies between EU ETS and DUKES;
- Identify any emission sources that are not contained in the GHGI.

The analysis of the EU ETS data for all onshore facilities was completed by May 2014 and provided to the DECC team of energy statisticians, in time for them to consider the EU ETS dataset during compilation of the UK energy balance for 2013, as published within DUKES 2014.

The EU ETS data for offshore oil and gas installations was provided in May 2014 and were used directly in the compilation of emission estimates for the upstream oil and gas sector, after the UK energy balance had been compiled by DECC. Access to these EU ETS data for offshore facilities provided more fuel-specific information (GCV, carbon content) to help improve completeness and accuracy of the upstream oil and gas estimates in the UK GHGI, augmenting the EEMS dataset which is a more comprehensive dataset (i.e. EEMS covers more emission sources than EU ETS) but does not provide the same level of fuel-specific data.

A 7.4 EU ETS DATA COVERAGE

The coverage of the EU ETS data has changed over the 9 years for which data are available. Major changes have been outlined in **Section A 7.2.2**, and these changes in scope have an

impact on the usefulness of data for some sectors, with data generally being more complete for Phase II (2008-12) and Phase III (2013-) of EU ETS. In addition, smaller combustion installations in the industrial, commercial and public sectors are outside the scope of EU ETS, and in fact coverage was decreased after 2007 due to the exemption of certain 'small emitters' from the UK EU ETS. For some source sectors in the GHGI, the EU ETS data therefore only includes a small proportion of the sector and the EU ETS data are not useful to directly inform the GHGI.

The following GHGI source sectors are well represented in the EU ETS data sets in the UK, with all UK installations included:

- Power stations burning coal, gas, and fuel oil as the principal fuel;
- Oil refineries;
- Coke ovens & Integrated steelworks;
- Cement kilns (from Phase II onwards); and
- Lime kilns (from Phase II onwards, and excluding kilns used in the Soda Ash industry)
- Glassworks - container, flat, special & fibre subsectors only (from phase II onwards)
- Brickworks and other sites manufacturing heavy ceramic goods (from phase II onwards).

However, GHGI sectors such as industrial combustion, autogeneration, and public sector combustion are only partially represented in the EU ETS data. An indication of the actual level of coverage of the EU ETS data can be seen in **Table A 7.4.1** below. The number of sites in each sector which are included in the ETS dataset for 2005 and 2013 are given, together with the Inventory Agency's estimate of the total number of installations in that sector throughout the UK in those years.

Table A 7.4.1 Numbers of installations included in the EU ETS data

Sector	Number of installations			
	2005		2013	
	EU ETS	UK total	EU ETS	UK total
Power stations (fossil fuel, > 75MWe)	60	60	60	60
Power stations (fossil fuel, < 75MWe)	23	27	30	39
Power stations (nuclear)	12	12	9	9
Coke ovens	4	4	4	4
Sinter plant	3	3	3	3
Blast furnaces	3	3	3	3
Cement kilns	8	15	12	12
Lime kilns	4	17	15	15
Refineries	12	12	12	12
Combustion – iron & steel industry	11	200 ^a	26	200 ^a

Sector	Number of installations			
	2005		2013	
	EU ETS	UK total	EU ETS	UK total
Combustion – other industry	171	5000 ^a	~400	5000 ^a
Combustion – commercial sector	28	1000 ^a	55	1000 ^a
Combustion – public sector	169	1000 ^a	114	1000 ^a
Glassworks (flat, special, container & fibre)	6	32	24	24
Brickworks	18	80 ^b	52	52 ^b

^a These estimates are 'order of magnitude' figures, to show that the number of installations in the UK is likely to be considerably higher than the number of installations reporting in the EU ETS.

^b Numbers of brickworks are not certain in 2005 but will have been significantly higher than in 2008 (when there were about 70) since many brickworks were closed or mothballed in the second half of 2007. All brickworks are believed to be covered by EU ETS in 2013.

Data are included in EU ETS for all coke ovens, refineries, sinter plant and blast furnaces. Power stations are divided into three categories in the table in order to show that, although nine stations are not included in the EU ETS data for 2013 (4 in 2005), these are all small (in most cases, very small diesel-fired plant supplying electricity to Scottish islands). In comparison, coverage is quite poor in 2005 for cement and lime kilns (due to CCA participants opting out during Phase I) and for combustion processes (due to CCA/UKETS opt-outs and the fact that numerous combustion plant are too small to be required to join the EU ETS). All cement kilns and all lime kilns are included in 2013; until Phase III (2013) there were two lime kilns that were excluded on the basis that they are an integral part of installations for manufacturing soda ash, and these installations have only been included in the scope of EU ETS in the UK since 2013. Coverage of glassworks and brickworks was very limited during phase I, but since 2008 has been very good: all large glassworks have been included since 2008, and all but one brickworks were included in phase II, with that remaining site being added for phase III. UK totals for brickworks are subject to some uncertainty however, and may be revised in future should more data be obtained.

For most emission sources the level of detail given in the EU ETS data matches well with the structures of the GHGI, allowing comparison of like with like. Only in the case of coke ovens and integrated steelworks is this not the case, since the EU ETS reporting format does not provide a breakdown of emissions for the sectors reported within the GHGI: i.e. estimates of emissions from coke ovens, blast furnaces and sinter plants are not provided explicitly. However, for these sectors, additional detailed analysis, including the collection of other industry data, has allowed for far greater use of EU ETS data for the inventory.

A 7.5 EU ETS DATA USE IN THE UK GHGI

The use of EU ETS data in the UK GHGI may conveniently be divided into two classes:

- Instances where activity data and, in most cases, emission totals as well are taken from EU ETS;

- Instances where emission factors only are taken from EU ETS and then used in the UK GHG Inventory with activity data from other sources such as DUKES.

A 7.5.1 Activity and Emissions Data

A 7.5.1.1 Crude Oil Refineries

The comparison of EU ETS emissions data against GHGI data based on DUKES fuel use allocations for petcoke, natural gas, fuel oil and OPG use is inconsistent to varying degrees in different years. Previous EU ETS analysis indicated that petcoke data in DUKES were too low; the DECC energy statistics team have investigated this matter with the refinery operators and have revised data for a number of sites that had been misreporting through the DORS system used to compile DUKES. In recent years, therefore, the EU ETS and DUKES data are closely consistent for petcoke use by refineries.

Data inconsistencies between DUKES and EU ETS remain for other fuels, however. In some cases, this will be due to misallocation of fuel use data within the EU ETS analysis, where fuel names are unclear, e.g. “fuel gas” could be interpreted as refinery use of OPG or the use of natural gas as a support fuel within the refinery fuel gas system.

The fuel oil activity data in most years is around 10% higher in EU ETS than in DUKES. Natural gas is a relatively minor fuel in the sector; whilst the EU ETS allocations indicate an over-report in DUKES, there is considerable uncertainty over the allocations of gases in the EU ETS dataset, as noted above. For the latest inventory submission, it has been recognised that DUKES data for natural gas used in autogeneration includes some fuel burnt at refineries, thus the difference between refinery fuel use as given in EU ETS, and that derived from DUKES data has been reduced. Consumption of naphtha reported in DUKES as “unclassified industry” is allocated to refineries as the only known consumers in the UK. However, in the case of OPG, there is typically an under-report in DUKES, although the data in DUKES is higher in two years. **Table A 7.5.1** below presents the emissions allocated to OPG for those years (2004 onwards except 2005, 2012) where UKPIA and EU ETS data indicates that DUKES data are too low. Note that the GHGI estimates also include the assumption that all of the OPG allocation to “autogenerators” within the DUKES commodity balance tables (in the column “Other gases”) is used within the refinery sector. Consultation with the DECC DUKES team has indicated (Personal Communication, Evans, 2010) that the “Other gases” column in the Commodity Balance tables is the OPG on the refinery basis, with CHP plant on site allocated to the autogeneration line. We have therefore retained this assumption in the current analysis, including the autogenerator allocation of “other gases” within the refinery sector.

To resolve the refinery sector under-report, we have compared DUKES data against EU ETS data, and also considered the total carbon dioxide emissions for the refinery sector provided annually by UKPIA. At the installation level, the UKPIA and EU ETS data show very close consistency for recent years (typically within 1%). The close consistency of the EU ETS and UKPIA data further strengthens the case for using EU ETS data as the primary dataset to inform the UK GHG inventory, in preference to the DUKES energy statistics.

At the fuel-specific level, the greatest disparity is evident in the reporting of OPG use at refineries; the reporting disparity has therefore been resolved through a top-down emissions comparison between DUKES-derived data and the best available operator data from EU ETS (2005 onwards) and UKPIA (pre-2005), with the difference between the two then allocated to OPG use in the UK GHGI. UK inventory estimates of emissions for the sector are therefore

aligned with EU ETS totals back to 2005, and with UKPIA data prior to 2005, unless the estimates derived from DUKES data are higher than those from UKPIA or EU ETS (i.e. in 2005 and 2012).

No deviations from UK energy statistics have been made prior to 2004, as the data from UKPIA and GHGI estimates based on DUKES are closely consistent with the DUKES-derived data being slightly higher; therefore a conservative approach is adopted, using DUKES-derived GHG estimates.

The time series of emissions data and the additional OPG emissions data (where DUKES data are low) for the sector are shown below.

Table A 7.5.1 Refinery Emissions Data Comparison and Revision to OPG Activity

Year	Best Operator Data*	Refinery emissions total (if based on DUKES)	Data selection for the UK GHGI estimates	Additional emissions assumed from OPG
	kt C	kt C		ktC
2000	4599	4710	<i>Use DUKES as > UKPIA</i>	0
2001	4535	4658	<i>Use DUKES as > UKPIA</i>	0
2002	4767	5237	<i>Use DUKES as > UKPIA</i>	0
2003	4772	5079	<i>Use DUKES as > UKPIA</i>	0
2004	4999	4870	<u>Use UKPIA as > DUKES</u>	129
2005	5007	5376	<i>Use DUKES as > EU ETS</i>	0
2006	4910	4709	<u>Use EU ETS as > DUKES</u>	201
2007	4857	4725	<u>Use EU ETS as > DUKES</u>	131
2008	4709	4414	<u>Use EU ETS as > DUKES</u>	295
2009	4492	4068	<u>Use EU ETS as > DUKES</u>	423
2010	4632	4360	<u>Use EU ETS as > DUKES</u>	272
2011	4739	4448	<u>Use EU ETS as > DUKES</u>	291
2012	4287	4354	<i>Use DUKES as > EU ETS</i>	0

Year	Best Operator Data*	Refinery emissions total (if based on DUKES)	Data selection for the UK GHGI estimates	Additional emissions assumed from OPG
	kt C	kt C		ktC
2013	4003	3831	<u>Use EU ETS as > DUKES</u>	172

*For 2005 onwards, the EU ETS data are verified by third parties and regarded as the best available sector estimates; prior to 2005 the best available operator emissions data are from the trade association, UKPIA.

There is some level of uncertainty in the allocation of fuels in EU ETS to specific “DUKES” fuels, although the OPG use in refineries seems to be reported quite consistently as “Refinery Gas”, “Refinery Off-Gas”, or “OPG/RFG”. The DECC DUKES team have reviewed the year to year consistency of OPG use in refineries through the DORS system.

A 7.5.1.2 Oil & Gas Terminal OPG and LPG Use

The allocation of reported fuel use within EU ETS to UK energy balance fuel nomenclature is uncertain in some cases. Analysis of the EU ETS fuel use data does indicate that there are small amounts of these fuels being used in the upstream oil & gas sector that are not evident within DUKES.

The DECC DUKES team have noted previously (Personal communication, DECC, 2010) that some LPG and OPG fuels are abstracted from upstream oil and gas exploration and production sources, rather than purchased from other sources, and that no data have been collected for this source since DUKES last published data for these sources, for the year 2002.

Therefore, the data from the EU ETS from oil and gas processing terminals on LPG and OPG combustion are used directly within the UK GHG inventory for the Phase II and Phase III years of 2008 to 2013, with estimates for 2003 to 2007 derived by interpolation between the EU ETS 2008 data and the DUKES 2002 data.

A 7.5.1.3 Natural Gas Use by Downstream Gas Supply Installations

The EU ETS data includes natural gas use by large gas compressor and storage sites that operate on the UK gas transmission and distribution network, as well as the three operational LNG terminals and a small number of other downstream gas industry sites.

The gas use reported in EU ETS for these sites throughout Phase II and III has been notably higher than the allocation of gas within DUKES Commodity Balance table 4.2 (Energy Industry Use, Other). This has been evident in the traded / non-traded analysis for the gas supply sector in the UK and DA GHGI.

As this gas use arises from the downstream network, the Inventory Agency and the DECC DUKES team consider that the DUKES data indicate a small misallocation of gas use, rather than a gap in reported gas use. For 2005 to 2013, therefore, the EU ETS data for this source are used within the UK GHG inventory, and the overall gas use data are balanced by reducing the allocation of gas use to “other industrial combustion” (IPCC source 1A2g); the EU ETS data since 2005 shows good consistency with the data from DUKES for earlier years.

The increased gas use for this sector based on EU ETS data is expected to still be a small under-report for the sector as a whole, as the EU ETS scope only includes around 35 of the

larger gas compressor, LNG terminals and storage sites on the UK network, and it is likely that additional gas use on smaller sites also occurs. However, the Inventory Agency has no data to inform such estimates.

A 7.5.1.4 Other Industry OPG use

There are a number of “other industry” sites where OPG use has been allocated by the Inventory Agency from EU ETS data, where the fuel is defined as either a specific gas (e.g. ethane, propane, butane) or more generic terms such as “OPG”, “High Pressure Refinery Gas”, “Low Pressure Refinery gas”, “fuel gas” or “RFG/OPG/ROG” within the EU ETS forms. The direct reference to refinery gases and the locations of the installations raises questions about how the fuel consumption is tracked and then reported within DUKES.

For a small number of sites, consultation with the DUKES team, regulators and operators has clarified that there is an under-report within DUKES and that the EU ETS energy and emissions data are the more accurate dataset and should be used in the UK GHGI. At some sites, energy supplier data returns to DECC have been misinterpreted with gases allocated to non-energy uses in the UK energy balance, when in fact a higher proportion of petroleum-based gases are used in combustion.

Within the DUKES petroleum commodity balance tables, there is no allocation of OPG or other light hydrocarbons to these industrial combustion processes, but there is an allocation to non-energy use of these gases, as well as some OPG use reported in autogeneration. Based on the EU ETS evidence, some proportion of this non-energy use has been re-allocated to account for the reported GHG emissions from these facilities, to address this gap in the UK energy statistics.

In the 1990-2013 inventory cycle, EU ETS data for fuel use at petrochemical production facilities helped to identify and quantify under-reports within the UK energy statistics for the combustion of gases that are derived from Natural Gas Liquid (NGL) feedstock to petrochemical production processes. Analysis of “fuel gas” calorific values and carbon content has helped to inform the calculations to estimate emissions from NGL-derived gases, to address the under-report in UK energy statistics and fill a reporting gap in previous inventory submissions.

A 7.5.1.5 Industrial Processes

The EU ETS dataset contains data on a number of industrial processes for which alternative data sources are either unavailable or of low quality. The EU ETS data therefore constitute the most reliable set of emissions data for these processes and are used in the UK inventory. In almost all cases, the EU ETS activity data are difficult to use directly largely because different operators provide activity data and emission factors on a different basis (e.g. some may provide input material and emission factors on a consumption basis, others will provide production data and emission factors on a production basis). Therefore, for all of the industrial process sources, the EU ETS emissions data are adopted, and activity data are generally back-calculated from the emissions using a suitable IPCC emission factor. The industrial process sources where EU ETS data have been used to generate estimates of emissions included within the UK GHGI in this submission, include:

- Emissions from the manufacture of lime. UK activity data for limestone and dolomite consumption in lime production would yield much lower emission estimates than is suggested by EU ETS returns therefore, as a conservative approach, the EU ETS data

are used instead. Activity data are back-calculated using the IPCC default factor for lime production. See **Section 4.3** for further details.

- Emissions from the use of carbonates in the manufacture of glass. As with lime production, the available data on consumption of limestone and dolomite for glass production are suspect, being very inconsistent across the time series, and so EU ETS data are used in the generation of the inventory time series, as detailed in **Section 4.4**.
- Emissions from the use of clays, carbonate minerals and other additives in the manufacture of bricks and roofing tiles, as detailed in **Section 4.5**. The EU ETS data are very detailed, with separate lines for different input materials such as different types of clay, carbonate minerals used in the bricks or in scrubbers used to abate fluoride emissions, and coke oven coke/petroleum coke used as an additive in certain bricks. UK brick production data are used as activity data.
- Estimates for emissions from the use of limestone in flue-gas desulphurisation (FGD) plant for the years 2005-2013 are taken from EU ETS data, because UK activity data (for gypsum produced from the FGD plant) are incomplete for those years. Activity data for 1990-2004 are available from non-EU ETS sources, and are back-calculated from the EU ETS CO₂ emissions for 2005 onwards assuming an emission factor of 253 kg CO₂ per tonne gypsum produced (which is based on an assumed 100% conversion of limestone and SO₂ into gypsum and CO₂).
- EU ETS Phase III saw the introduction of data for soda ash manufacturing sites and EU ETS data, and CO₂ emissions reported for earlier years in the PI, are used as the basis of UK inventory emissions data for that sector. See **Section 4.12** for more details.
- Titanium dioxide production was also included in phase III of the EU ETS, but data for the UK plant was not included in the data set provided, and so was not used in the generation of emissions data for this submission. Emission estimates were generated using an alternative method, but it is anticipated that the next submission will use EU ETS data instead.
- Petroleum coke is added to some electric arc furnaces as a reductant, and emissions and activity data for the period 2005-2013 are taken from EU ETS data, with emissions in earlier years being extrapolated on the basis of plant production. See **Section 4.16** for further details.

A 7.5.2 Implied Emission Factors

A 7.5.2.1 Power Stations

Table A 7.5.2 summarises EU ETS data for fuels burnt by major power stations and coal burnt by autogenerators. The percentage of emissions based on Tier 3 emission factors is given (Tier 3 factors are based on fuel analysis, and are therefore more reliable than emission factors based on default values), as well as the average emission factor for EU ETS emissions based on Tier 3 factors.

Table A 7.5.2 EU ETS data for Fuels used at Power Stations and Autogenerators (Emission Factors in kt / Mt for Coal & Fuel Oil, kt / Mth for Gases)

Year	Fuel	% Tier 3	Average Carbon Emission Factor (Tier 3 only)
2005	Coal	99	615.3

Year	Fuel	% Tier 3	Average Carbon Emission Factor (Tier 3 only)
2006		100	615.0
2007		100	614.7
2008		100	612.4
2009		100	607.2
2010		100	609.0
2011		100	609.0
2012		100	611.5
2013		100	612.5
2005	Fuel oil / Waste oil ^a	59	860.3
2006		66	873.0
2007		68	871.1
2008		91	869.5
2009		94	872.7
2010		95	873.3
2011		94	873.9
2012		95	874.7
2013		93	871.3
2005	Natural gas	52	1.443
2006		76	1.465
2007		95	1.464
2008		97	1.467
2009		100	1.464
2010		99	1.460
2011		99	1.458
2012		100	1.461
2013		99	1.464
2005	Coal - autogenerators	100	594.3
2006		100	596.3
2007		100	594.5
2008		100	581.3
2009		100	600.6
2010		100	599.9

Year	Fuel	% Tier 3	Average Carbon Emission Factor (Tier 3 only)
2011		100	594.9
2012		0 ^b	N/A
2013		0 ^b	N/A

^a It is not possible to distinguish between fuel oil and waste oil in the EU ETS data, so all emissions have been reported under fuel oil.

^b Plant operated as a power station after 2011 and included in the figures for power stations burning coal

The EU ETS data shown are regarded as good quality data, since a high proportion of emissions are based on Tier 3 emission factors (i.e. verified emissions based on fuel analysis to ISO17025). The factors are also very consistent across the time-series, which would be expected for this sector. As shown in Section 3, the EU ETS data for power stations also cover almost all UK installations in this sector, and certainly cover all of the larger installations burning coal, fuel oil and natural gas.

A few power stations burn small quantities of petroleum coke as well as coal. One supplies data to ETS for coal/petroleum coke blends i.e. there are no separate emissions data or carbon factors for the coal and the petroleum coke at that site. We therefore back-calculate the coal IEF in those blends by using an assumed default for the petcoke carbon content and more detailed activity data on the constituents of the fuel blends, obtained directly from the operator.

The EU ETS based emission factors presented above for power stations are used directly as the emission factors in the GHGI, with the exception of the 2005 figure for gas, where Tier 3 factors were only used for about half of the sector's emissions reported in EU ETS. Small quantities of sour gas were burnt at one power station in 2005-2007 and 2009 and EU ETS Tier 3 emission factors are available and therefore used. [Due to the confidentiality of the data, the emission factors are not shown]. Prior to 2005, the emission factors for these sectors are based on the methodology established by Baggott *et al*, 2004, since it has been concluded that this represents the most reliable approach.

The EU ETS factors for coal-fired autogenerators are slightly different to the factors for the power stations in that, although the EU ETS data are exclusively Tier 3, they only represent about 80% of total fuel used by the sector.

A 7.5.2.2 Crude Oil Refineries

The tables below summarise the EU ETS data for the major fuels burnt by refineries in the UK.

The main fuels in refineries are fuel oil and OPG and emissions also occur due to the burning off of 'petroleum coke' deposits on catalysts used in processes such as catalytic cracking. In the latter case, emissions in the EU ETS are not generally based on activity data and emission factors but are instead based on direct measurement of carbon emitted. This is due to the technical difficulty in measuring the quantity of petroleum coke burnt and the carbon content. Refineries also use natural gas, although it is a relatively small source of emissions compared to other fuels.

Table A 7.5.3 Refinery EU ETS Data for Fuel Oil, OPG and Natural Gas (Emission Factors in kt / Mt for Fuel Oil and kt / Mth for OPG and Natural Gas)

Year	Fuel	% Tier 3	Average Carbon Emission Factor (Tier 3 sites only)
2005	Fuel Oil	25	861.0
2006		65	873.9
2007		79	877.4
2008		91	871.6
2009		91	876.2
2010		97	878.2
2011		85	877.6
2012		80	887.6
2013		98	874.3
2005	OPG	59	1.499
2006		57	1.475
2007		68	1.582
2008		80	1.483
2009		80	1.489
2010		80	1.501
2011		67	1.453
2012		64	1.470
2013		78	1.489
2005	Natural Gas	0	N/A
2006		43	1.460
2007		45	1.462
2008		98	1.475
2009		98	1.480
2010		97	1.465
2011		81	1.447
2012		63	1.442
2013		89	1.459

There has been a significant drop in the proportion of Tier 3 reporting for all three fuels since 2010, which will adversely affect the quality of the emission factors, although coverage is still in excess of 50% for all fuels.

Emission factors for **fuel oil** generated from EU ETS data have been adopted in the GHGI, with the exception of data for 2005, where Tier 3 methods were used for only 25% of fuel.

Carbon factors can be derived for **OPG** based on moderate levels of Tier 3 reporting for 2005-2007 and 2011-2013 but 80% for 2008-2010. There is some uncertainty regarding the allocation of EU ETS fuels to the OPG fuel category, and the derived emission factors do cover a wider spread of values than for many other fuels in EU ETS. However, this perhaps reflects the nature of this fuel, and the data for all years have been used in the inventory.

Carbon factors for natural gas are based on a low % of Tier 3 reporting until 2008; in 2008 to 2010 over 90% of gas use is reported at Tier 3 and 81% in 2011. Within the UK GHGI, the EU ETS factors for 2008 to 2012 are used directly, whilst emission factors for earlier years are derived from gas network operator gas compositional analysis.

EU ETS emission data for **petroleum coke** are higher in 2005-2010, when compared against the estimates derived from DUKES activity data and the industry-recommended emission factor. This is especially noticeable for 2005, where the petroleum coke consumption given in DUKES would have to be more than 100% carbon in order to generate the carbon emissions given in the EU ETS. Consultation with DECC energy statisticians has identified that the figures given in DUKES are subject to uncertainty and hence the EU ETS data are used directly within the UK GHGI for those years.

A 7.5.2.3 Integrated Steelworks & Coke Ovens

Table A 7.5.4 summarises EU ETS data for the major fuels burnt at integrated steelworks and coke ovens. The data exclude one independent coke oven which calculates emissions using a detailed mass balance approach which makes it more difficult to assess the data in the same way as the other installations.

Table A 7.5.4 EU ETS data for Fuels used at Integrated Steelworks and Coke Ovens (Emission Factors in kt / Mt for Solid & Liquid Fuels, kt / Mth for Gases)

Year	Fuel	% Tier 3	Average Carbon Emission Factor (Tier 3 sites only)
2005	Blast furnace gas	0	N/A
2006		100	6.871
2007		99	6.911
2008		97	6.903
2009		97	6.996
2010		100	6.920
2011		94	6.974
2012		96	6.811
2013		98	6.766

Year	Fuel	% Tier 3	Average Carbon Emission Factor (Tier 3 sites only)
2005	Coke oven gas	0	N/A
2006		0	N/A
2007		0	N/A
2008		54	1.094
2009		100	1.195
2010		100	1.150
2011		100	1.086
2012		100	1.097
2013		100	1.093
2005	Natural gas	0	N/A
2006		3	1.479
2007		2	1.478
2008		0	N/A
2009		58	1.425
2010		68	1.441
2011		64	1.441
2012		64	1.443
2013		27	1.447
2005	Fuel oil	0	N/A
2006		0	N/A
2007		0	N/A
2008		84	878.3
2009		89	884.7
2010		83	887.6
2011		87	888.7
2012		66	878.2
2013		0	N/A

Much of the ETS data for coke ovens and steelworks are now used in the GHGI, although not the emission factors shown above. Instead, the Inventory Agency have used the EU ETS data and other detailed, site-specific and fuel-specific data, provided by the process operators to refine the carbon balance model used to generate emission estimates for the sector. Details of the revisions to the carbon balance model can be found in the research report from the 2013-2014 inventory improvement programme (Ricardo-AEA, 2014)

A 7.5.2.4 Cement Kilns

Table A 7.5.5 summarises EU ETS data for the major fuels burnt at cement kilns.

Table A 7.5.5 EU ETS data for Fuels used at Cement Kilns (kt / Mt)

Year	Fuel	% Tier 3	Average Carbon Emission Factor (Tier 3 sites only)
2005	Coal	8	671.1
2006		100	546.2
2007		100	664.3
2008		100	655.8
2009		100	658.3
2010		100	637.7
2011		100	645.8
2012		100	662.4
2013		100	694.2
2005	Petroleum coke	-	N/A
2006		100	820.8
2007		100	830.2
2008		100	819.1
2009		100	796.8
2010		100	750.8
2011		100	738.4
2012		100	770.2
2013		100	811.1

The EU ETS dataset also provides a detailed breakdown of cement sector process emissions from the decarbonisation of raw materials during the clinker manufacturing process. These data are useful to compare against statistics provided by the Mineral Products Association (MPA) regarding clinker production and the non-combustion emissions associated with UK cement production. The MPA data on clinker production are commercially confidential.

The two data sets show significant differences for 2005-2007; however the EU ETS data cover only a fraction of the sector, so differences might be expected. The coal IEF data for 2008-2013 are closer, with a narrower (4%) range. Because of the good agreement in both activity data and emission factors for 2008 onwards, the industry-wide estimates provided by the MPA

and used within the GHGI show very close comparison with the EU ETS estimates, with the sum of the EU ETS data less than 1% lower than those reported to the GHGI, as outlined below in **Table A 7.5.6**.

Table A 7.5.6 Comparison of Cement Sector Carbon Dioxide Emissions* within the UK GHGI and the EU ETS for 2008-2012

	2008	2009	2010	2011	2012	2013
GHGI CO ₂ emissions (kt)	8294	5686	5788	6130	556 5	596 7
Sum of EU ETS CO ₂ emissions (kt)	8259	5647	5754	6087	555 6	597 2
EU ETS / GHGI	99.6%	99.3%	99.4%	99.3%	99.8%	100.1 %

*The data in this table include fuel combustion emissions (reported under IPCC 1A2f) and process emissions (reported under IPCC sector 2A1) from UK cement kilns.

A 7.5.2.5 Lime Kilns

Table A 7.5.7 summarises data given in the EU ETS datasets for the major fuels burnt at lime kilns. Unlike cement kilns, which often burn a variety of fuels, many lime kilns burn just a single fuel, often natural gas. The data below exclude coke oven coke used in lime kilns at soda ash plant since these kilns were not covered by EU ETS until phase III, and the small number of sites make the data confidential in any case.

Table A 7.5.7 EU ETS data for Fuels used at Lime Kilns (Emission Factors in kt / Mt for Solid Fuels and kt / Mth for Gases)

Year	Fuel	% Tier 3	Average Carbon Emission Factor
			(Tier 3 sites only)
2005	Coal*	-	N/A
2006		-	N/A
2007		34	846.9
2008		79	701.4
2009		100	698.9
2010		100	634.4
2011		100	703.9
2012		100	725.6
2013		100	689.1

*Coal used in the lime industry in the UK includes a proportion of anthracitic coal, and hence some of these IEFs are notably higher than for coal used in other sectors of UK industry.

The EU ETS data for lime kilns are variable across the time series, both in terms of the proportion of emissions based on Tier 3 factors, and in the emission factors themselves. EU

ETS based factors are currently used for coal and petroleum coke from 2008 onwards, as the EU ETS data do include all lime kilns burning those fuels and almost all of those data are Tier 3 and hence are regarded as highly reliable.

EU ETS data for natural gas use in the lime industry does cover all installations burning this fuel, however the proportion of emissions based on Tier 3 factors is very low. Therefore the EU ETS emission factors are not used in the UK GHGI, and the emission factors for natural gas continue to be based on the methodology given in Baggott *et al*, 2004.

Table A 7.5.8 shows implied emission factors for process-related emissions from lime kilns that are used within the UK GHG inventory. The lime industry can be sub-divided into those installations where lime is the primary product, and carbon dioxide is an unwanted by-product; and those installations where both lime and carbon dioxide are utilised. The latter include kilns in the sugar industry (where carbon dioxide is used in the purification stages) and soda ash production (where carbon dioxide is combined with other chemicals to produce sodium carbonate), and in these kilns, the carbon dioxide from decarbonisation of the limestone or dolomite feedstock is assumed to be fully consumed in the process, rather than emitted to atmosphere. **Table A 7.5.8** therefore does not cover these installations. None of the emission factors in EU ETS are Tier 3, so the table shows the overall emission factors for all tiers of data.

Table A 7.5.8 EU ETS emission factor data for production of lime (kt / Mt lime produced)

Year	Activity	EU ETS
2005	Lime production	200.4
2006		201.2
2007		201.3
2008		195.6
2009		195.0
2010		194.0
2011		195.6
2012		195.6
2013		194.4

These factors compare with a theoretical emission factor of 214 kt / Mt lime, assuming use of pure limestone. We note that the EU ETS factors are all lower than the theoretical emission factor based on the stoichiometry of the lime manufacturing process and this is despite some use of dolomitic limestone in the UK industry which would be expected to further increase the emission factor above the 214 kt/Mt lime factor. The EU ETS data are subject to third party verification, and therefore the emissions data are assumed to be accurate. It is assumed that the reason for this deviation from the theoretical emission factor is due to the production activity data being inflated by either the products containing some proportion of slaked lime (i.e. hydrated product and hence heavier than pure lime) and/or other additives to the lime product which increase the mass of the lime product.

A 7.5.2.6 Other Industrial Combustion

Table A 7.5.9 summarises EU ETS data for coal, fuel oil and natural gas used by industrial combustion installations.

At first sight, the data for coal looks like it should be reliable enough to be used in the GHGI with 92% or more of emissions based on Tier 3 factors in each year. However, it must be recalled that numerous smaller industrial consumers will not be represented in EU ETS and that the EU ETS data are not fully representative of UK fuels as a whole – see **Section A 7.4** for details. This is also true for EU ETS data for fuel oil and natural gas but here, in addition, very little of the EU ETS data are based on Tier 3 factors. Therefore, none of these data have been used directly in the compilation of the GHGI estimates.

Table A 7.5.9 EU ETS data for Coal, Fuel Oil and Natural Gas used by Industrial Combustion Plant (Emission Factors in kt / Mt for Coal & Fuel Oil, kt / Mth for Natural Gas)

Year	Fuel	% Tier 3	Average Carbon Emission Factor (Tier 3 sites only)	GHGI Carbon Emission Factor
2005	Coal	98	607.1	647.8
2006		98	603.0	647.8
2007		99	615.7	662.4
2008		94	598.6	656.6
2009		92	595.4	668.5
2010		92	589.0	674.3
2011		96	596.5	653.4
2012		96	605.8	653.7
2013		97	628.2	653.3
2005		Fuel oil	17	864.7
2006	27		865.3	879.0
2007	44		872.3	879.0
2008	24		871.4	879.0
2009	39		871.3	879.0
2010	40		873.0	879.0
2011	51		874.2	879.0
2012	48		875.1	879.0
2013	43		871.3	879.0

Year	Fuel	% Tier 3	Average Carbon Emission Factor (Tier 3 sites only)	GHGI Carbon Emission Factor
2005	Natural gas	14	1.593	1.478
2006		31	1.470	1.478
2007		42	1.466	1.477
2008		29	1.495	1.474
2009		43	1.499	1.474
2010		43	1.502	1.472
2011		41	1.468	1.470
2012		42	1.471	1.470
2013		44	1.474	1.474

Emission factors can also be derived from EU ETS where a high percentage of Tier 3 analysis is evident, for a number of other minor fuels. Due to the very low number of sites that report data for each fuel type, these EU ETS-derived emission factors are confidential and are not tabulated here. The source/activity combinations for which EU ETS emission factor data are used within the inventory are:

- Other industrial combustion / petroleum coke
- Other industrial combustion / waste solvents
- Other industrial combustion / colliery methane

The EU ETS-derived emission factors for colliery methane for each year (2005-2013) are also applied to all other sources using these fuels.

ANNEX 8: UK Domestic Emissions Reporting Requirements

In addition to the reporting requirements of the UNFCCC, Kyoto Protocol and EUMM, UK Greenhouse Gas Inventory statistics are published annually in a Department of Energy and Climate Change's National Statistics release¹⁰. The geographical coverage of these estimates differs from the UNFCCC and EUMM coverage, with the totals mainly including emissions from the UK and the UK's Crown Dependencies only, although progress towards the Kyoto Protocol is still reported. Summary tables of these data are presented below. The data are presented in the nine categories used for the UK's National Communications to the UNFCCC (NC Categories).

A 8.1 NATIONAL STATISTICS

Table A 8.1.1 Summary table of GHG (Including net emissions/removals from LULUCF) emissions by NC Category (Mt CO₂eq) – National Statistics coverage (UK and Crown Dependencies)

NC category	1990	1995	2000	2005	2009	2010	2011	2012	2013
Energy Supply	278.84	238.83	221.47	231.37	200.55	206.72	192.41	203.52	189.74
Transport	121.74	122.17	126.79	130.71	121.89	120.33	118.54	118.01	116.78
Residential	80.61	81.86	89.02	86.00	78.16	87.81	67.69	77.27	77.55
Business	115.43	113.70	117.24	109.52	90.75	94.42	88.65	88.42	90.94
Public	13.52	13.28	12.08	11.19	9.24	9.75	9.33	9.28	9.52
Industrial Process	60.03	50.93	27.22	20.41	11.62	12.48	10.98	10.45	12.75
Agriculture	65.95	65.11	61.41	57.30	54.17	54.59	54.53	53.96	53.75
Land Use Change	3.97	3.27	0.83	-2.93	-4.08	-4.29	-4.85	-4.97	-5.26
Waste Management	69.34	71.46	66.77	53.03	36.28	31.52	28.93	26.28	22.57
Total	809.41	760.61	722.83	696.60	598.58	613.33	566.22	582.21	568.35

Table A 8.1.2 Summary table of GHG emissions by Gas (Mt CO₂eq) – National Statistics coverage

Gas	1990	1995	2000	2005	2009	2010	2011	2012	2013
CO ₂	597.92	562.04	559.47	557.79	482.22	500.75	457.53	476.26	467.47

¹⁰ <https://www.gov.uk/government/publications/final-uk-emissions-estimates>

Gas	1990	1995	2000	2005	2009	2010	2011	2012	2013
CH ₄	136.94	129.93	113.89	92.08	71.88	67.00	63.95	61.20	56.19
N ₂ O	57.07	47.22	36.60	32.15	28.46	28.90	27.69	27.66	27.62
HFCs	14.55	19.55	10.45	13.13	15.17	15.66	15.98	16.19	16.21
PFCs	1.65	0.60	0.60	0.39	0.20	0.29	0.42	0.26	0.25
SF ₆	1.28	1.26	1.82	1.06	0.65	0.73	0.65	0.63	0.60
NF ₃	0.00042	0.00083	0.00169	0.00029	0.00026	0.00027	0.00030	0.00033	0.00036
Total	809.41	760.61	722.83	696.60	598.58	613.33	566.22	582.21	568.35

A 8.2 UK ONLY EMISSIONS STATISTICS

As part of the Climate Change Act 2008¹¹, the UK committed to reducing greenhouse gas emissions by at least 80 per cent by 2050 (relative to 1990), with an interim target of reducing greenhouse gas emissions by at least 34 per cent by 2020, also relative to 1990.

These targets along with the legally binding five-year carbon budgets (which set the trajectory to reaching the targets by placing a restriction on the total amount of greenhouse gases the UK can emit over the five-year period) are on a UK only basis, thus excluding Crown Dependencies and Overseas Territories.

Summary statistics for the UK only are presented below. Note that the scope of emissions used for calculating Carbon Budgets differs slightly from those presented here – for example Carbon Budgets exclude NF₃. The final 2013 UK GHG emissions statistical release included an update of the UK's performance against the first carbon budget. The release can be found at: <https://www.gov.uk/government/publications/final-uk-emissions-estimates>.

Table A 8.2.1 Summary table of GHG emissions by NC Category (Mt CO₂eq) – UK only

NC Sector	1990	1995	2000	2005	2009	2010	2011	2012	2013
Energy Supply	278.32	238.18	220.99	231.10	200.18	206.37	192.08	203.10	189.34
Transport	121.22	121.66	126.21	130.10	121.34	119.77	117.99	117.47	116.28
Residential	80.28	81.51	88.60	85.60	77.76	87.40	67.29	76.86	77.14
Business	115.21	113.46	116.93	109.16	90.40	94.08	88.34	88.09	90.63
Public	13.52	13.28	12.08	11.19	9.24	9.75	9.33	9.28	9.52
Industrial Process	60.03	50.93	27.22	20.41	11.62	12.48	10.98	10.45	12.75
Agriculture	65.75	64.91	61.19	57.16	53.99	54.42	54.37	53.82	53.61
Land Use Change	3.99	3.30	0.87	-2.91	-4.07	-4.28	-4.84	-4.96	-5.25

¹¹ <http://www.legislation.gov.uk/ukpga/2008/27/contents>

NC Sector	1990	1995	2000	2005	2009	2010	2011	2012	2013
Waste Management	69.15	71.27	66.60	52.90	36.15	31.40	28.80	26.15	22.45
Total	807.46	758.51	720.68	694.71	596.63	611.39	564.35	580.28	566.47

Table A 8.2.2 Summary table of GHG emissions by Gas (Mt CO₂eq) – UK Only

Gas	1990	1995	2000	2005	2009	2010	2011	2012	2013
CO ₂	596.39	560.36	557.76	556.24	480.66	499.20	456.04	474.67	465.93
CH ₄	136.59	129.59	113.57	91.86	71.63	66.76	63.72	61.00	55.98
N ₂ O	56.99	47.15	36.52	32.09	28.39	28.83	27.62	27.60	27.56
HFCs	14.55	19.55	10.42	13.08	15.09	15.58	15.90	16.11	16.14
PFCs	1.65	0.60	0.60	0.39	0.20	0.29	0.42	0.26	0.25
SF ₆	1.28	1.26	1.82	1.05	0.65	0.73	0.65	0.63	0.60
NF ₃	0.00042	0.00083	0.00169	0.00029	0.00026	0.00027	0.00030	0.00033	0.00036
Total	807.46	758.51	720.68	694.71	596.63	611.39	564.35	580.28	566.47

ANNEX 9: End User Emissions

A 9.1 INTRODUCTION

This Annex explains the concept of a final user or end user, summarises the final user calculation methodology with examples, and contains tables of greenhouse gas emissions according to final user from 1990 to 2013.

The final user sectoral categories used are consistent with those used in the National Communications (NC) to the UNFCCC. The sectoral categories in the NC are derived from the UNFCCC reporting guidelines on national communications¹².

The purpose of the final user calculations is to allocate emissions from fuel and electricity producers to the energy users - this allows the emission estimates for a consumer of energy to include the emissions from the production of the fuel or electricity they use.

The UNFCCC does not require final user data to be included in the UK's National Inventory Report. These data have been included to provide DECC with information for their policy support needs.

The tables in this Annex present summary data for UK greenhouse gas emissions for the years 1990-2013, inclusive. These data are updated annually to reflect revisions in the methods used to estimate emissions, and the availability of new information within the inventory. These recalculations are applied retrospectively to earlier years to ensure a consistent time series and this accounts for any differences in data published in previous reports.

Emissions from the UK Overseas Territories are not included in the calculations; there is not enough information available to reallocate emissions from energy supply. Emissions presented in this chapter show emissions from the UK and Crown Dependencies, consistent with the UK statistical release.

A 9.2 DEFINITION OF FINAL USERS

The final user¹³ or end user calculations allocate emissions from fuel producers to fuel users. The final user calculation therefore allows estimates to be made of emissions for a consumer of fuel, which also include the emissions from producing the fuel the consumer has used.

The emissions included in the final user categories can be illustrated with an example of two final users - the residential sector and road transport:

- Emissions in the **residential** final user category include:

¹² See page 84 of UNFCCC Guidelines contained in FCCC/CP/1999/7 available at: <http://unfccc.int/resource/docs/cop5/07.pdf>

¹³ A final user is a consumer of fuel for useful energy. A 'fuel producer' is someone who extracts, processes or converts fuels for the end use of final users. Clearly there can be some overlap of these categories but here the fuel uses categories of the UK DECC publication DUKES are used, which enable a distinction to be made.


1. Direct emissions from domestic premises, for example, from burning gas, coal or oil for space heating.
 2. Emissions from power stations generating the electricity used by domestic consumers; emissions from refineries including refining, storage, flaring and extraction; emissions from coal mines (including emissions due to fuel use in the mining industry itself and fugitive emissions of methane from the mines); and emissions from the extraction, storage and distribution of mains gas.
- Emissions in the **road transport** final user category include:
 1. Direct emissions from motor vehicle exhausts.
 2. Emissions from refineries producing motor fuels, including refining, storage, flaring and extraction of oil; and from the distribution and supply of motor fuels.

A 9.3 OVERVIEW OF THE FINAL USER CALCULATIONS

As fuel and electricity producers use energy from other producers, they are allocated emissions from each other and these have to then be reallocated to final users. This circularity results in an iterative approach being used to estimate emissions from categories of final users.

Figure A 9.3.1 shows a simplified view of the energy flows in the UK (the fuels used in the greenhouse gas inventory have hundreds of uses). This figure shows that while final users consuming electricity are responsible for a proportion of the emissions from power stations they are also responsible for emissions from collieries, and some of these emissions in turn come from electricity generated in power stations and from refineries.

Figure A 9.3.1 Simplified fuel flows for a final user calculation.



The approach for estimating end user emissions is summarised in the three steps below:

1. Emissions are calculated for each sector for each fuel.
2. Emissions from fuel and electricity producers are then distributed to those sectors that use the fuel according to the energy content¹⁴ of the fuel they use (these sectors can include other fuel producers). This distribution is based on inventory fuel consumption data and DUKES electricity consumption data.
3. By this stage in the calculation, emissions from final users will have increased and those from fuel and electricity producers will have decreased. The sum of emissions from fuel producers and power stations in a particular year as a percentage of the total emissions is then calculated. If this percentage, for any year, exceeds a predetermined value (In the model used to determine emissions from final users, the value of this percentage can be adjusted. The tables presented later in this Appendix were calculated for a convergence at 0.001%) the process continues at Step 2. If this percentage matches or is less than the predetermined value, the calculation is finished.

Convergence of this iterative approach is likely, as the fuel flows to the final users are much greater than fuel flows amongst the fuel producers.

While a direct solution could possibly be used it was decided to base the calculation on an iterative approach because:

¹⁴ If calorific data for the fuels is not available then the mass of fuel is used instead. This is the case for years prior to 1990.

- This can be implemented in the database structures already in existence for the UK greenhouse gas inventory;
- It can handle a wide range of flows and loops that occur without any of the limits that other approaches may incur; and
- The same code will cover all likely situations and will be driven by tabular data stored in the database.


A 9.4 EXAMPLE FINAL USER CALCULATION

The following example illustrates the methodology used to calculate emissions according to final users. The units in this example are arbitrary.

The example in **Figure A 9.4.1** has two fuel producers, *power stations* and *collieries*, and three final users, *residential*, *industry* and *commercial*. The following assumptions have been made for simplicity:

- The only fuels used are coal and electricity;
- Coal is the only source of carbon emissions (released from burning coal in power stations to produce electricity and from burning coal in the home for space heating); and
- Commerce uses no coal and so has zero 'direct' emissions.

Figure A 9.4.1 Fuel use in the example calculation



In **Figure A 9.4.1**, the tonnes refer to tonnes of coal burnt (black arrows), and the units refer to units of electricity consumed (blue arrows).

In this example the coal extracted by the colliery is burnt in the power station to produce electricity for the final users. Industrial and residential users also directly burn coal. Although the colliery uses electricity produced by the power station, it is not considered to be a final user. The colliery is a 'fuel producer' as it is part of the chain that extracts, processes and converts fuels for the final users.

Table A 9.4.1 summarises the outputs during this example final user calculation.

Table A 9.4.1 Example of the outputs during a final user calculation

		Sector					Unallocated emissions as percentage of total emission	Total emission of carbon (tonnes)	
		Colliery	Power Station	Residential	Industrial	Commercial			
Coal use (tonnes)	Mass	100	100,000	50,000	100,000	0			
	Energy content	25,000	25,000,000	12,500,000	25,000,000	0			
Electricity use (arbitrary units)	Energy units	10,000		50,000	100,000	100,000			
Emissions of carbon (tonnes)	Initial	70	70,000	35,000	70,000	0	40.02	175,070	
	Emissions after iteration step	1	2,692	28	48,476	96,951	26,923	1.55	175,070
		2	1	1077	49,020	98,039	26,934	0.62	175,070
		3	41	1	49,227	98,454	27,348	0.02	175,070
		4	0	17	49,235	98,470	27,348	0.01	175,070
		5	1	0	49,238	98,477	27,355	0	175,070
		6	0	0	49,239	98,477	27,355	0	175,070

The initial carbon emissions are 70% of the mass of coal burnt. The emissions from the power stations are distributed to the other sectors by using the factor:

- $(\text{Electricity used by that sector}) / (\text{total electricity used minus own use by power stations});$
- Similarly for the colliery emissions the following factor is used; and
- $(\text{Energy of coal used by that sector}) / (\text{total energy of coal consumed used minus own use by collieries}).$

At the end of iteration step one, the commerce sector has 26923 tonnes of carbon emissions allocated to it, mainly derived from power stations. Emissions allocated to the residential and industry sectors have also increased over their initial allocations. However collieries and power stations still have some emissions allocated to them (these come from each other) and so the reallocation process is repeated to reduce these allocations to zero – these two sectors are not final users. The total unallocated (in this example, equal to the total emissions from collieries and power stations) falls in each iteration until the emissions are consistently allocated across the sectors. In this example, six iterations are needed to achieve a consistent allocation across the sectors.

The sum of emissions allocated to the sectors (175,070 tonnes of carbon) remains unchanged from the initial allocation to the allocation in the sixth iteration. This check is an important quality control measure to ensure all emissions are accounted for during the final user calculations.

Figure A 9.4.2 Comparison of ‘direct’ and final user emissions of carbon according the sectors considered in the final user example




Figure A 9.4.2 compares the quantities of direct and final user carbon emitted from each sector at the end of the final user calculation. The direct emissions of carbon are from the combustion of coal in the sectors. The direct and final user emissions are from two distinct calculations and must be considered independently – in other words, the direct and final user emissions in each sector must not be summed. The sum of all the direct emissions and the sum of the final user emissions, are identical.

There are relatively large direct emissions of carbon from power stations, residential and industry sectors. The final user emissions from the power stations and the colliery are zero because these two sectors are not final users. The carbon emissions from these two sectors have been reallocated to the residential, industrial and commercial sectors. This reallocation means the final user emissions for the residential and industrial sectors are greater than their ‘direct’ emissions.

A 9.5 FINAL USER CALCULATION METHODOLOGY FOR THE UK GREENHOUSE GAS INVENTORY

The approach divides fuel user emissions into 8 categories (see column 1 of **Table A 9.5.1**). For each of these groups, source categories are distributed by the total energy consumption of a group of fuels. For example, for the coal group, the emissions of four source categories are distributed to final users according to the energy use of anthracite and coal combined.

Table A 9.5.1 Sources reallocated to final users and the fuels used

Final user group	Emission sources to be reallocated to final users	Fuels used for redistribution
1. Coke	Gasification processes	Coke
	Coke production	Blast furnace gas
	Iron and steel – flaring	
2. Coal	Closed Coal Mines	Coal
	Coal storage and transport	Anthracite
	Collieries – combustion	
	Deep-mined coal	
	Open-cast coal	
3. Natural gas	Gas leakage	Natural gas
	Gas production	
	Upstream Gas Production – flaring	
	Upstream Gas Production – fuel combustion	
	Upstream Gas Production – Gas terminal storage	
	Upstream Gas Production – Offshore Well Testing	
	Upstream Gas Production – process emissions	
	Upstream Gas Production – venting	
	Upstream Gas production – combustion at gas separation plant	
4. Electricity	Nuclear fuel production	Electricity
	Power stations	
	Autogeneration – exported to grid	
	Power stations – FGD	

Final user group	Emission sources to be reallocated to final users	Fuels used for redistribution
5. Petroleum	Upstream Oil Production – gas combustion	Naphtha
	Upstream Oil Production – gas flaring	Burning oil (premium)
	Upstream Oil Production – gas venting	Burning oil
	Upstream Oil Production – Offshore Oil Loading	Aviation turbine fuel
	Upstream Oil Production – Offshore Well Testing	Aviation spirit
	Upstream Oil Production – Oil terminal storage	Derv
	Upstream Oil Production – Onshore Oil Loading	Fuel oil
	Upstream Oil Production – process emissions	Gas oil
	Petrol stations – petrol delivery	OPG
	Petrol stations – vehicle refuelling	Refinery misc.
	Petrol terminals – storage	Petrol
	Petrol terminals – tanker loading	Petroleum coke
	Petroleum processes	Wide-cut gasoline
	Refineries – combustion	Vaporizing oil
	Refineries – drainage	LPG
	Refineries – flares	
	Refineries – general	
	Refineries – process	
	Refineries – road/rail loading	
	Refineries – tankage	
Sea going vessel loading		
Ship purging		
6. Solid Smokeless Fuels	Solid Smokeless fuel production	Solid Smokeless Fuels
7. Town gas	Town gas manufacture	Town gas
8. Charcoal	Charcoal production	Charcoal

Comments on the calculation methodology used to allocate emissions according final users are listed below:

- Emissions are allocated to final users on the basis of the proportion of the total energy produced used by a given sector. This approach is followed to allow for sectors such as petroleum where different products are made in a refinery;
- Some emissions are allocated to an “exports” category. This is for emissions within the UK from producing fuels, (for example from a refinery or coal mine), which are subsequently exported or sent to bunkers for use outside the UK. Therefore these emissions are part of the UK inventory even if the use of the fuel produces emissions that cannot be included in the UK inventory because it takes place outside the UK;
- No allowance is made for the emission from the production of fuels or electricity outside the UK that are subsequently imported;
- Some of the output of a refinery is not used as a fuel but used as feedstock or lubricants. This is not currently treated separately and the emissions from their production (which are small) are allocated to users of petroleum fuels. This is partly due to lack of data in the database used to calculate the inventory, and partly due to the lack of a clear, transparent way of separating emissions from the production of fuels and from the production of non-fuel petroleum products; and

Final user emissions are estimated for aviation in four categories: domestic take off and landing, international take off and landing, domestic cruise and international cruise. This enables both IPCC and UNECE categories to be estimated from the same final user calculation.

Our exact mapping of final user emissions to IPCC categories is shown in the following **Table A 9.5.2**. The NAEI source sectors and activity names are also shown, as it is necessary to subdivide some IPCC categories. This classification has been used to generate the final user tables for the greenhouse gases given in this section. As this table is for final users, no fuel producers are included in the table.

Table A 9.5.2 Final user category, IPCC sectors, and NAEI source names and activity names used in the emission calculation

NC Category	IPCC Category	Source Name	Activity Name
Agriculture	1A4ci_Agriculture/Forestry/Fishing:Stationary	Agriculture - stationary combustion	Coal Coke Fuel oil Natural gas Straw Vaporising oil
		Miscellaneous industrial/commercial combustion	Burning oil
	1A4cii_Agriculture/Forestry/Fishing:Off-road	Agriculture - mobile machinery	Gas oil Petrol
	2D1_Lubricant_Use	Agricultural engines	Lubricants
	3A1_Enteric_Fermentation_dairy_cattle	Agriculture livestock - dairy cattle enteric	Non-fuel combustion
	3A1_Enteric_Fermentation_non-dairy_cattle	Agriculture livestock - other cattle enteric	Non-fuel combustion
	3A2_Enteric_Fermentation_sheep	Agriculture livestock - sheep enteric	Non-fuel combustion
	3A3_Enteric_Fermentation_swine	Agriculture livestock - pigs enteric	Non-fuel combustion

NC Category	IPCC Category	Source Name	Activity Name
	3A4_Enteric_Fermentation_other:deer	Agriculture livestock - deer enteric	Non-fuel combustion
	3A4_Enteric_Fermentation_other:goats	Agriculture livestock - goats enteric	Non-fuel combustion
	3A4_Enteric_Fermentation_other:horses	Agriculture livestock - horses enteric	Non-fuel combustion
	3B1_Manure_Management_dairy_cattle	Agriculture livestock - dairy cattle wastes	Non-fuel combustion
	3B1_Manure_Management_non-dairy_cattle	Agriculture livestock - other cattle wastes	Non-fuel combustion
	3B2_Manure_Management_sheep	Agriculture livestock - sheep goats and deer wastes	Non-fuel combustion
	3B3_Manure_Management_swine	Agriculture livestock - pigs wastes	Non-fuel combustion
	3B4_Manure_Management_other:deer	Agriculture livestock - deer wastes	Non-fuel combustion
	3B4_Manure_Management_other:goats	Agriculture livestock - goats wastes	Non-fuel combustion
	3B4_Manure_Management_other:horses	Agriculture livestock - horses wastes	Non-fuel combustion
	3B4_Manure_Management_other:poultry	Agriculture livestock - broilers wastes	Non-fuel combustion
		Agriculture livestock - laying hens wastes	Non-fuel combustion

NC Category	IPCC Category	Source Name	Activity Name
		Agriculture livestock - other poultry wastes	Non-fuel combustion
		Agriculture livestock - all poultry wastes	Non-fuel combustion
	3B4_Other	Agriculture livestock - manure other	Non-fuel combustion
		Agriculture livestock - manure leaching (indirect)	Non-fuel combustion
		Agriculture livestock - manure liquid systems (indirect)	Non-fuel combustion
		Agriculture livestock - manure other (indirect)	Non-fuel combustion
		Agriculture livestock - manure solid storage and dry lot (indirect)	Non-fuel combustion
	3D_Agricultural_Soils	Agricultural soils	Non-fuel crops Non-fuel fertilizer
		OvTerr Agricultural Soils	Non-fuel combustion
	3D1_Agricultural_soils-Mineralization/Immobilization	Agricultural soils - Mineralization/Immobilization Associated with change in Soil Organic Matter	Non-fuel combustion
	3F_Field_burning	Field burning	Barley residue Linseed residue

NC Category	IPCC Category	Source Name	Activity Name
			Oats residue Wheat residue
	3G1_Liming - limestone	Liming	Limestone
	3G2_Liming - dolomite	Liming	Dolomite
	3H_Urea application	Agriculture - application of urea	Urea consumption
	non-IPCC	Agriculture - stationary combustion	Electricity
Business	1A2a_Iron_and_steel	Blast furnaces	Blast furnace gas Coke oven gas LPG Natural gas
		Iron and steel - combustion plant	Blast furnace gas Coal Coke Coke oven gas Fuel oil Gas oil LPG Natural gas Town gas

NC Category	IPCC Category	Source Name	Activity Name
			Blast Furnace Gas (Carbon)
	1A2b_Non-Ferrous_Metals	Autogeneration - exported to grid	Coal
		Autogenerators	Coal
		Non-Ferrous Metal (combustion)	Coal Fuel oil Gas oil Natural gas
	1A2c_Chemicals	Chemicals (combustion)	Coal Fuel oil Gas oil Natural gas
	1A2d_Pulp_Paper_Print	Pulp, Paper and Print (combustion)	Coal Fuel oil Gas oil Natural gas
	1A2e_food_processing_beverages_and_tobacco	Food & drink, tobacco (combustion)	Coal Fuel oil Gas oil

NC Category	IPCC Category	Source Name	Activity Name
			Natural gas
	1A2f_Non-metallic_minerals	Cement production - combustion	Coal Fuel oil Gas oil Natural gas Petroleum coke Scrap tyres Waste Waste oils Waste solvent
		Lime production - non decarbonising	Coal Coke Natural gas
	1A2gvii_Off-road_vehicles_and_other_machinery	Industrial off-road mobile machinery	DERV Gas oil Petrol
	1A2gviii_Other_manufacturing_industries_and_construction	Autogeneration - exported to grid	Natural gas
		Autogenerators	Biogas

NC Category	IPCC Category	Source Name	Activity Name
			Natural gas
		Other industrial combustion	Biomass Burning oil Coal Coke Coke oven gas Colliery methane Fuel oil Gas oil LPG Natural gas OPG Petroleum coke SSF Town gas Waste solvent Wood
	1A4ai_Commercial/Institutional	Miscellaneous industrial/commercial combustion	Coal Coke

NC Category	IPCC Category	Source Name	Activity Name
			Fuel oil Gas oil Landfill gas MSW Natural gas SSF Town gas
	2B1_Chemical_Industry:Ammonia_production	Ammonia production - combustion	Natural gas
	2B8g_Petrochemical_and_carbon_black_production:Other	Chemicals (combustion)	OPG
	2C1b_Pig_iron	Blast furnaces	Coal
	2D1_Lubricant_Use	Industrial engines	Lubricants
		Other industrial combustion	Lubricants
	2D4_Other_NEU	Non Energy Use: petroleum coke	Petroleum coke
	2E1_Integrated_circuit_or_semiconductor	Electronics - HFC	Non-fuel combustion
		Electronics - NF ₃	Non-fuel combustion
	2F1a_Commercial_refrigeration	Commercial Refrigeration	Refrigeration and Air Conditioning - Disposal
			Refrigeration and Air Conditioning - Lifetime

NC Category	IPCC Category	Source Name	Activity Name
			Refrigeration and Air Conditioning - Manufacture
	2F1b_Domestic_refrigeration	Domestic Refrigeration	Refrigeration and Air Conditioning - Disposal Refrigeration and Air Conditioning - Lifetime Refrigeration and Air Conditioning - Manufacture
	2F1c_Industrial_refrigeration	Industrial Refrigeration	Refrigeration and Air Conditioning - Disposal Refrigeration and Air Conditioning - Lifetime Refrigeration and Air Conditioning - Manufacture
	2F1d_Transport_refrigeration	Refrigerated Transport	Refrigeration and Air Conditioning - Disposal Refrigeration and Air Conditioning - Lifetime Refrigeration and Air Conditioning - Manufacture
	2F1e_Mobile_air_conditioning	Mobile Air Conditioning	Refrigeration and Air Conditioning - Disposal Refrigeration and Air Conditioning - Lifetime

NC Category	IPCC Category	Source Name	Activity Name
			Refrigeration and Air Conditioning - Manufacture
	2F1f_Stationary_air_conditioning	Stationary Air Conditioning	Refrigeration and Air Conditioning - Disposal Refrigeration and Air Conditioning - Lifetime Refrigeration and Air Conditioning - Manufacture
	2F2a_Closed_foam_blowing_agents	Foams	Non-fuel combustion
		Foams HFCs for the 2006 GLs	Non-fuel combustion
	2F2b_Open_foam_blowing_agents	One Component Foams	Non-fuel combustion
	2F3_Fire_Protection	Firefighting	Non-fuel combustion
	2F5_Solvents	Precision cleaning - HFC	Non-fuel combustion
	2F6b_Other_Applications:Contained-Refrigerant_containers	Refrigerant containers	Non-fuel combustion
	2G1_Electrical_equipment	Electrical insulation	Non-fuel combustion
	2G2_Military_applications	AWACS	Non-fuel combustion
	2G2_Particle_accelerators	Particle accelerators	Non-fuel combustion
	2G2e_Electronics_and_shoes	Electronics - PFC	Non-fuel combustion
		Electronics - SF ₆	Non-fuel combustion
		Sporting goods	Non-fuel combustion
	2G2e_Tracer_gas	SF ₆ used as a tracer gas	Non-fuel combustion

NC Category	IPCC Category	Source Name	Activity Name	
	2G3a_Medical applications	N ₂ O use as an anaesthetic	Population	
	5C2.2b_Non-biogenic:Other	Accidental fires - other buildings	Mass burnt	
	non-IPCC		Chemicals (combustion)	Electricity
			Food & drink, tobacco (combustion)	Electricity
			Iron and steel - combustion plant	Electricity
			Miscellaneous industrial/commercial combustion	Electricity
			Non-Ferrous Metal (combustion)	Electricity
			Other industrial combustion	Electricity
Energy Supply	1A1ai_Public_Electricity&Heat_Production	Power stations	Burning oil Coal Coke Fuel oil Gas oil LPG Natural gas OPG	

NC Category	IPCC Category	Source Name	Activity Name
			Petroleum coke
	1A1b_Petroleum_Refining	Refineries - combustion	Natural gas
	1A1ci_Manufacture_of_solid_fuels	Coke production	Natural gas
		Solid smokeless fuel production	Coke Natural gas
	1A1cii_Oil_and_gas_extraction	Upstream Gas Production - fuel combustion	Gas oil
		Upstream oil and gas production - combustion at gas separation plant	LPG OPG
		Upstream Oil Production - fuel combustion	Natural gas
	1A1ciii_Other_energy_industries	Collieries - combustion	Natural gas
		Nuclear fuel production	Natural gas
		Town gas manufacture	Burning oil Coal Coke LPG Natural gas
	1Aciii_other_energy_industries	Gas production	LPG

NC Category	IPCC Category	Source Name	Activity Name
			OPG Town gas
		1B1b_Solid_Fuel_Transformation	Coke production
	Solid smokeless fuel production	Coal Petroleum coke	
	non-IPCC	Collieries - combustion	Electricity
	Gas production	Electricity	
	Refineries - combustion	Electricity	
	Exports	Aviation_Bunkers	Aircraft - international cruise
Aircraft - international take off and landing			Aviation spirit Aviation turbine fuel
Aircraft between UK and Gibraltar - Cruise			Aviation spirit Aviation turbine fuel
Aircraft between UK and Gibraltar - TOL			Aviation spirit Aviation turbine fuel
Aircraft between UK and other Ots (excl Gib.) - Cruise			Aviation turbine fuel

NC Category	IPCC Category	Source Name	Activity Name
		Aircraft between UK and other OTs (excl Gib.) - TOL	Aviation turbine fuel
	Marine_Bunkers	Shipping - international IPCC definition	Fuel oil Gas oil
		Shipping between UK and Gibraltar	Fuel oil
		Shipping between UK and OTs (excl. Gib)	Fuel oil
	non-IPCC	Exports	Aviation turbine fuel Burning oil Coke DERV Fuel oil Petrol SSF Electricity
Industrial Process	2A1_Cement_Production	Cement - decarbonising	Clinker production
	2A2_Lime_Production	Lime production - decarbonising	Limestone
	2A3_Glass_production	Glass - general	Dolomite Limestone

NC Category	IPCC Category	Source Name	Activity Name
			Soda ash
	2A4d_Other_process_uses_of_carbonates	Brick manufacture - Fletton	Fletton bricks
		Brick manufacture - all types	Bricks
	2B1_Ammonia_Production	Ammonia production - feedstock use of gas	Natural gas
	2B10_Chemical_Industry:Other	Chemical industry - general	Process emission
	2B2_Nitric_Acid_Production	Nitric acid production	Acid production
	2B3_Adipic_Acid_Production	Adipic acid production	Adipic acid produced
	2B6_Titanium_dioxide_production	Chemical industry - titanium dioxide	Coke Petroleum coke
	2B7_Soda_Ash_Production	Chemical industry - soda ash	Soda ash produced
	2B8a_Methanol_production	Chemical industry - methanol	Methanol
	2B8b_Ethylene_Production	Chemical industry - ethylene	Ethylene
	2B8c_Ethylene_Dichloride_and_Vinyl_Chloride_Monomer	Chemical Industry – ethylene dichloride	Ethylene dichloride
	2B8d_Ethylene_Oxide	Chemical industry - ethylene oxide	Ethylene oxide
	2B8e_Acrylonitrile	Chemical industry - acrylonitrile	Acrylonitrile
	2B8f_Carbon_black_production	Chemical industry - carbon black	Carbon black capacity
	2B9a1_Fluorchemical_production:By-product_emissions	Halocarbons production - by-product	Non-fuel combustion

NC Category	IPCC Category	Source Name	Activity Name
	2B9b3_Fluorchemical_production:Fugitive_emissions	Halocarbons production - fugitive	Non-fuel combustion
	2C1a_Steel	Basic oxygen furnaces	Dolomite
		Electric arc furnaces	Petroleum coke Steel production (electric arc)
		Ladle arc furnaces	Steel production (electric arc) Steel production (oxygen converters)
	2C1b_Pig_iron	Blast furnaces	Coke Fuel oil
	2C1d_Sinter	Sinter production	Coke Dolomite Limestone
	2C3_Aluminium_Production	Primary aluminium production - general	Primary aluminium production
		Primary aluminium production - PFC emissions	Primary aluminium production
	2C4_Magnesium_production	Magnesium cover gas	Non-fuel combustion
	2C6_Zinc_Production	Non-ferrous metal processes	Coke
	non-IPCC	Blast furnaces	Electricity

NC Category	IPCC Category	Source Name	Activity Name
Land Use Change	4A_Forest Land (Biomass Burning - wildfires)	Forest Land - Biomass Burning\Wildfires	Biomass
	4A_Forest Land (organic soils drainage)	Forest Land - Drainage of Organic Soils	Non-fuel combustion
	4A1_Forest Land Remaining Forest Land	Forest Land remaining Forest Land	Non-fuel combustion
	4A2_Forest_Land-Direct_ N ₂ O _emission_from_N_fertilisation	Direct N ₂ O emission from N fertilisation of forest land	Non-fuel combustion
	4A2_Land Converted to Forest Land	Land converted to Forest Land	Non-fuel combustion
	4B_Cropland (Biomass Burning - controlled)	Cropland - Biomass Burning\Controlled Burning	Biomass
	4B_Cropland (Biomass Burning - wildfires)	Cropland - Biomass Burning\Wildfires	Biomass
	4B1_Cropland Remaining Cropland	Cropland remaining Cropland	Non-fuel combustion
	4B2_4(III) Direct N ₂ O Emissions from N Mineralization/Immobilization	Cropland - Drainage and rewetting and other management of organic and mineral soils	Non-fuel combustion
	4B2_Land Converted to Cropland	Land converted to Cropland	Non-fuel combustion
		Direct N ₂ O Emissions from N Mineralization/Immobilisation	Non-fuel combustion
	4C_Grassland (Biomass burning - controlled)	Grassland - Biomass Burning\Controlled Burning	Biomass
	4C_Grassland (Biomass Burning - wildfires)	Grassland - Biomass Burning\Wildfires	Biomass

NC Category	IPCC Category	Source Name	Activity Name
	4C_Grassland-Direct_ N ₂ O _emission_from_N_fertilisation	Grassland - Direct N ₂ O Emissions from N Mineralization/Immobilisation	Non-fuel combustion
	4C1_Grassland Remaining Grassland	Grassland remaining Grassland	Non-fuel combustion
	4C2_4(III) Direct N ₂ O Emissions from N Mineralization/Immobilization	Grassland - Drainage and rewetting and other management of organic and mineral soils	Non-fuel combustion
	4C2_Land converted to grassland	Land converted to Grassland	Non-fuel combustion
	4D1_Wetlands remaining wetlands	Wetlands remaining Wetland	Non-fuel combustion
	4D2_Land converted to wetlands	Land converted to Wetland	Non-fuel combustion
	4D2_Non- CO ₂ _emissions_from_drainage_of_soils_and_wetlands	Non- CO ₂ emissions from drainage of soils and wetlands	Non-fuel combustion
	4E_Settlements (Biomass burning - controlled)	Settlements - Biomass Burning\Controlled Burning	Biomass
	4E_Settlements-Direct_ N ₂ O _emission_from_N_fertilisation	Settlements - Direct N ₂ O Emissions from N Mineralization/Immobilisation	Non-fuel combustion
	4E1_Settlements remaining settlements	Settlements remaining Settlements	Non-fuel combustion
	4E2_Land converted to settlements	Land converted to Settlements	Non-fuel combustion
	4F2_Land converted to other land	Land converted to Other Land	Non-fuel combustion
	4G_Harvested wood products	Harvested Wood Products	Non-fuel combustion
Public	1A4ai_Commercial/Institutional	Public sector combustion	Burning oil

NC Category	IPCC Category	Source Name	Activity Name
			Coal Coke Fuel oil Gas oil Natural gas Sewage gas Town gas
	non-IPCC	Public sector combustion	Electricity
Residential	1A4bi_Residential_stationary	Domestic combustion	Anthracite Burning oil Charcoal Coal Coke Fuel oil Gas oil LPG Natural gas Peat Petroleum coke SSF

NC Category	IPCC Category	Source Name	Activity Name	
			Town gas	
			Wood	
		1A4bii_Residential:Off-road	House and garden machinery	DERV Petrol
		2D2 Non-energy_products_from_fuels_and_solvent_use:Paraffin_wax_use	Non-aerosol products - household products	Petroleum waxes
		2F4a_Metered_dose_inhalers	Metered dose inhalers	Non-fuel combustion
		2F4b_Aerosols:Other	Aerosols - halocarbons	Non-fuel combustion
		5B1a_composting_municipal_solid_waste	Composting (household)	Biological waste
		5C2.2b_Non-biogenic:Other	Small-scale waste burning	Waste
			Accidental fires - dwellings	Mass burnt
		5C2.2b_Non-biogenic:Other_Accidental fires (vehicles)	Accidental fires - vehicles	Mass burnt
non-IPCC	Domestic combustion	Electricity		
Transport	1A3a_Domestic_aviation	Aircraft - domestic cruise	Aviation spirit Aviation turbine fuel	
		Aircraft - domestic take off and landing	Aviation spirit Aviation turbine fuel	
		Aircraft between UK and CDs - Cruise	Aviation spirit Aviation turbine fuel	

NC Category	IPCC Category	Source Name	Activity Name
		Aircraft between UK and CDs - TOL	Aviation spirit Aviation turbine fuel
	1A3bi_Cars	Road transport - cars - cold start	DERV Petrol
		Road transport - cars - motorway driving	DERV Petrol
		Road transport - cars - rural driving	DERV Petrol
		Road transport - cars - urban driving	DERV Petrol
	1A3bii_Light_duty_trucks	Road transport - LGVs - cold start	DERV Petrol
		Road transport - LGVs - motorway driving	DERV Petrol
		Road transport - LGVs - rural driving	DERV Petrol

NC Category	IPCC Category	Source Name	Activity Name
		Road transport - LGVs - urban driving	DERV Petrol
	1A3biii_Heavy_duty_trucks_and_buses	Road transport - buses and coaches - motorway driving	DERV
	Road transport - buses and coaches - rural driving	DERV	
	Road transport - buses and coaches - urban driving	DERV	
	Road transport - HGV articulated - motorway driving	DERV	
	Road transport - HGV articulated - rural driving	DERV	
	Road transport - HGV articulated - urban driving	DERV	
	Road transport - HGV rigid - motorway driving	DERV	
	Road transport - HGV rigid - rural driving	DERV	
	Road transport - HGV rigid - urban driving	DERV	
	1A3biv_Motorcycles	Road transport - mopeds (<50cc 2st) - urban driving	Petrol
		Road transport - motorcycle (>50cc 2st) - urban driving	Petrol

NC Category	IPCC Category	Source Name	Activity Name
		Road transport - motorcycle (>50cc 4st) - motorway driving	Petrol
		Road transport - motorcycle (>50cc 4st) - rural driving	Petrol
		Road transport - motorcycle (>50cc 4st) - urban driving	Petrol
	1A3bv_Other_road_transport	Road transport - all vehicles LPG use	LPG
	1A3c_Railways	Rail - coal	Coal
		Railways - freight	Gas oil
		Railways - intercity	Gas oil
		Railways - regional	Gas oil
	1A3d_Domestic_navigation	Inland goods-carrying vessels	Gas oil
		Motorboats / workboats (e.g. canal boats, dredgers, service boats, tourist boats, river boats)	DERV
			Gas oil
			Petrol
		Personal watercraft e.g. jet ski	Petrol
		Sailing boats with auxiliary engines	DERV
		Shipping - coastal	Fuel oil

NC Category	IPCC Category	Source Name	Activity Name
			Gas oil
	1A3eii_Other_Transportation	Aircraft - support vehicles	Gas oil
	1A4ai_Commercial/Institutional	Railways - stationary combustion	Burning oil Coal Coke Fuel oil Natural gas
	1A4ciii_Fishing	Fishing vessels	Gas oil
	1A5b_Other:Mobile	Aircraft - military	Aviation spirit Aviation turbine fuel
		Shipping - naval	Gas oil
	2D1_Lubricant_Use	Marine engines	Lubricants
		Road vehicle engines	Lubricants
	2D3_Non-energy_products_from_fuels_and_solvent_use:Other	Road transport - urea	Urea consumption
	non-IPCC	Railways - regional	Electricity
		Road vehicle engines	Electricity
Waste Management	5A1a_Managed_Waste_Disposal_sites_anaerobic	Landfill	Non-fuel combustion
	5B1a_composting_municipal_solid_waste	Composting - NH ₃	Process emission

NC Category	IPCC Category	Source Name	Activity Name
		Total composting (non-household)	Biological waste
	5B2a_Anaerobic_digestion_municipal_solid_waste	Anaerobic Digestion (other)	Biological waste
		Mechanical Biological Treatment	Biological waste
	5C1.1b_Biogenic:Sewage_sludge	Incineration - sewage sludge	Sewage sludge combustion
	5C1.2a_Non-biogenic:municipal_solid_waste	Incineration	MSW
	5C1.2b_Non-biogenic:Clinical_waste	Incineration - clinical waste	Clinical waste
	5C1.2b_Non-biogenic:Other_Chemical_waste	Incineration - chemical waste	Chemical waste
	5D1_Domestic_wastewater_treatment	Sewage sludge decomposition	Non-fuel domestic
		Sewage sludge decomposition in private systems	Non-fuel domestic
	5D2_Industrial_wastewater_treatment	Industrial Waste Water Treatment	Non-fuel combustion

A 9.6 DETAILED EMISSIONS ACCORDING TO FINAL USER CATEGORIES

The final user categories in the data tables in this summary are those used in National Communications. The final user reallocation includes all emissions from the UK and Crown Dependencies, this is the coverage used for the UK statistical release¹⁵, where the final users data are presented in more detail.

The base year for hydrofluorocarbons, perfluorocarbons, nitrogen trifluoride and sulphur hexafluoride is 1995. For carbon dioxide, methane and nitrous oxide, the base year is 1990.

¹⁵ <https://www.gov.uk/government/publications/final-uk-emissions-estimates>

Table A 9.6.1 Final user emissions from all National Communication categories, MtCO₂ equivalent

Final user category	Base Year	1990	1995	2000	2005	2009	2010	2011	2012	2013
Agriculture	69.83	69.83	68.30	64.12	60.09	56.59	57.16	56.97	56.50	56.10
Business	251.19	250.09	220.55	218.91	212.50	178.19	186.62	175.38	180.68	177.77
Energy Supply	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
Industrial Process	65.29	63.12	53.49	29.23	21.11	12.64	13.29	11.76	11.22	13.32
Land Use Change	3.97	3.97	3.27	0.83	-2.93	-4.08	-4.29	-4.85	-4.97	-5.26
Public	31.62	31.62	29.04	24.37	22.43	19.06	19.33	18.30	19.01	18.25
Residential	172.96	172.30	158.00	158.70	162.96	145.47	156.12	128.86	145.33	139.89
Transport	139.81	139.81	143.18	146.83	150.46	138.98	137.41	135.03	133.86	131.79
Waste Management	69.34	69.34	71.46	66.77	53.03	36.28	31.52	28.93	26.28	22.57
Exports	9.33	9.33	13.32	13.06	16.94	15.46	16.17	15.84	14.30	13.93
Total greenhouse gas emissions	813.35	809.41	760.61	722.83	696.60	598.58	613.33	566.22	582.21	568.35

Table A 9.6.2 Final user CO₂ emissions from all National Communication categories, MtCO₂ equivalent

Final user category	Base Year	1990	1995	2000	2005	2009	2010	2011	2012	2013
Agriculture	10.66	10.66	10.12	8.32	8.27	7.54	7.72	7.75	7.60	7.21

Final user category	Base Year	1990	1995	2000	2005	2009	2010	2011	2012	2013
Business	231.12	231.12	204.68	203.52	194.88	159.61	167.50	156.05	161.20	158.88
Energy Supply	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
Industrial Process	20.88	20.88	18.95	18.13	16.73	10.70	11.14	10.66	10.50	12.63
Land Use Change	2.87	2.87	2.16	-0.23	-3.85	-4.90	-5.08	-5.62	-5.79	-6.01
Public	29.30	29.30	27.22	23.34	21.70	18.46	18.76	17.73	18.43	17.73
Residential	156.87	156.87	146.10	149.61	155.17	138.60	149.23	122.51	138.47	133.53
Transport	136.39	136.39	139.52	143.96	148.26	137.19	135.70	133.17	132.03	129.99
Waste Management	1.29	1.29	0.87	0.49	0.38	0.27	0.27	0.27	0.25	0.25
Exports	8.55	8.55	12.42	12.34	16.24	14.75	15.49	15.02	13.58	13.26
Total greenhouse gas emissions	597.92	597.92	562.04	559.47	557.79	482.22	500.75	457.53	476.26	467.47

Table A 9.6.3 Final user CH₄ emissions from all National Communication categories, MtCO₂ equivalent

Final user category	Base Year	1990	1995	2000	2005	2009	2010	2011	2012	2013
Agriculture	33.05	33.05	32.36	31.14	28.87	27.29	27.43	27.30	27.20	27.08
Business	15.54	15.54	11.68	7.38	4.72	3.73	3.63	3.61	3.39	2.89
Energy Supply	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00

Final user category	Base Year	1990	1995	2000	2005	2009	2010	2011	2012	2013
Industrial Process	2.03	2.03	1.67	1.08	0.46	0.39	0.35	0.32	0.30	0.28
Land Use Change	0.02	0.02	0.04	0.05	0.05	0.05	0.05	0.04	0.07	0.04
Public	2.13	2.13	1.69	0.94	0.65	0.53	0.50	0.51	0.51	0.44
Residential	14.53	14.53	10.54	6.55	4.75	4.16	4.23	3.71	4.08	3.62
Transport	2.02	2.02	1.77	1.20	0.84	0.74	0.66	0.76	0.69	0.65
Waste Management	66.92	66.92	69.40	64.97	51.20	34.42	29.61	27.01	24.37	20.63
Exports	0.71	0.71	0.78	0.59	0.54	0.57	0.55	0.68	0.59	0.56
Total greenhouse gas emissions	136.94	136.94	129.93	113.89	92.08	71.88	67.00	63.95	61.20	56.19

Table A 9.6.4 Final user N₂O emissions from all National Communication categories, MtCO₂ equivalent

Final user category	Base Year	1990	1995	2000	2005	2009	2010	2011	2012	2013
Agriculture	26.13	26.13	25.81	24.66	22.95	21.76	22.00	21.92	21.70	21.81
Business	2.30	2.30	1.95	1.73	1.79	1.40	1.43	1.35	1.53	1.41
Energy Supply	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
Industrial Process	23.86	23.86	14.36	5.39	2.89	1.19	1.30	0.23	0.08	0.09
Land Use Change	1.08	1.08	1.07	1.01	0.86	0.76	0.75	0.74	0.75	0.71

Final user category	Base Year	1990	1995	2000	2005	2009	2010	2011	2012	2013
Public	0.19	0.19	0.13	0.09	0.08	0.07	0.07	0.07	0.08	0.07
Residential	0.91	0.91	0.71	0.59	0.60	0.51	0.53	0.50	0.61	0.59
Transport	1.39	1.39	1.89	1.67	1.36	1.05	1.05	1.09	1.13	1.15
Waste Management	1.13	1.13	1.18	1.32	1.45	1.59	1.64	1.65	1.66	1.69
Exports	0.07	0.07	0.12	0.13	0.15	0.13	0.13	0.14	0.13	0.11
Total greenhouse gas emissions	57.07	57.07	47.22	36.60	32.15	28.46	28.90	27.69	27.66	27.62

