

Dr Clare Bayley  
Department for Environment, Food and Rural Affairs  
Ergon House,  
17 Smith Square  
SW1P 3JR

Telephone: 020 7238 3059  
Email: [clare.bayley@defra.gsi.gov.uk](mailto:clare.bayley@defra.gsi.gov.uk)  
Website: [www.defra.gov.uk](http://www.defra.gov.uk)



1<sup>st</sup> December 2011

## **RE: NOTIFICATION OF CHANGES TO THE AIR QUALITY INDEX**

The Committee on the Medical Effects of Air Pollutants (COMEAP) was asked by Defra and the Devolved Administrations to review the Air Quality Index to ensure that it is fit for purpose given developments in the field of air quality. The current index has been in operation essentially unchanged for a period of around 12 years. COMEAP published the report *Review of the UK Air Quality Index* in June 2011. Defra and the Devolved Administrations have considered the report recommendations and will implement several changes, following consultation with index users. The new index, to be renamed the Daily Air Quality Index (DAQI) is documented below.

The changes to the index are:

1. Updated health advice to accompany the DAQI to provide more focused and clearer information. The new index comes in three parts and includes additional advice for susceptible individuals, alongside advice for the general population:
  - A. Instructions on how the index should be used;**
  - B. The short-term health effects of air pollution and action that can be taken to reduce impacts;**
  - C. Health advice linked to each band to accompany the air quality index.**
2. Changes to the index bands (Low, Moderate, High, Very High) for particulate matter (PM<sub>10</sub>), nitrogen dioxide (NO<sub>2</sub>), and ozone (O<sub>3</sub>) to make them more stringent. See Table 1 for figures. Please note that the bands for ozone are slightly different to those recommended by COMEAP;
3. Inclusion of fine particulate matter (PM<sub>2.5</sub>) in the DAQI;
4. Removal of carbon monoxide (CO) from the DAQI in view of the dramatic reductions in outdoor concentrations of CO;
5. Changes to the presentation of the index, to be a 10-point scale with four bands of Low, Moderate, High, and Very High, with colour coding to aid the interpretation of the DAQI, see Table 1;

6. Inclusion of trigger values to allow for the prediction of episodes of elevated air pollution in real time as they emerge. With the averaging times<sup>1</sup> of 24 hours for particulate matter and 8 hours for ozone it is not possible to provide public information about an unexpected pollution episode until it is well established. 'Triggers' have been derived to provide information to the public to warn of likely exposure before it occurs and as a 'Moderate', 'High', or 'Very High' episode progresses. These 'triggers' can be used by organisations that operate real-time public information services. Further information on trigger values will follow;
7. Links to the long-term health effects of air pollution will be provided with the index, noting that the DAQI itself only addresses the health effects of short term exposure to elevated levels.

Defra recommend that public information services that use the Daily Air Quality Index implement these changes. Defra and the Devolved Administrations will be implementing these changes to national air quality information services from 1<sup>st</sup> January 2012.

#### **Actions on implementation for information providers:**

- Update the thresholds and definition of statistical calculations for the existing banding pollutants (O<sub>3</sub>, CO, SO<sub>2</sub>, NO<sub>2</sub> and PM<sub>10</sub>). In the case of ozone and PM<sub>10</sub> this will involve changing the averaging periods for the bandings as well as the thresholds.
- Add new bandings and statistical calculations for PM<sub>2.5</sub>.
- Change the format and presentation of air quality bulletins and other information provision methods to include PM<sub>2.5</sub> and drop CO.
- Update software for any data summaries and regional summaries where the index of "worst pollutant" is presented, so that PM<sub>2.5</sub> is included and CO is removed.
- Update any tables and descriptive pages on your website which present the bandings information, in particular to update the health information provided.
- Add the calculations and presentation of "Trigger Values" (further information on trigger values will follow).

Further information on air quality, including current and forecast levels of air pollution can be found at <http://uk-air.defra.gov.uk>. If you have any queries regarding the Daily Air Quality Index, please do not hesitate to contact me.

Yours sincerely,



Dr Clare Bayley  
Atmosphere and Local Environment

---

<sup>1</sup> The averaging time is the period of time which is used to smooth short term variations in pollutant concentrations. These differ from pollutant to pollutant reflecting two aspects of the evidence on health effects; firstly, the timescale of exposure over which adverse health effects might be caused and, secondly, the averaging times used in the studies on which the assessment is based.

## Daily Air Quality Index

The new bandings for the Daily Air quality Index are detailed in Table 1.

Band	Index	Ozone	Nitrogen Dioxide	Sulphur Dioxide	PM <sub>2.5</sub> Particles	PM <sub>10</sub> Particles
		Running 8 hourly mean	hourly mean	15 minute mean	24 hour mean	24 hour mean
		µgm <sup>-3</sup>	µgm <sup>-3</sup>	µgm <sup>-3</sup>	µgm <sup>-3</sup>	µgm <sup>-3</sup>
<b>LOW</b>						
	1	0-33	0-66	0-88	0-11	0-16
	2	34-65	67-133	89-176	12-23	17-33
	3	66-99	134-199	177-265	24-34	34-49
<b>MODERATE</b>						
	4	100-120	200-267	266-354	35-41	50-58
	5	121-140	268-334	355-442	42-46	59-66
	6	141-159	335-399	443-531	47-52	67-74
<b>HIGH</b>						
	7	160-187	400-467	532-708	53-58	75-83
	8	188-213	468-534	709-886	59-64	84-91
	9	214-239	535-599	887-1063	65-69	92-99
<b>VERY HIGH</b>						
	10	240 or more	600 or more	1064 or more	70 or more	100 or more

**Table 1: Daily Air Quality Index bands**

The new daily air quality index comes in three parts and includes additional advice for susceptible individuals, alongside advice for the general population:

- A. Instructions on how the index should be used;**
- B. The short-term health effects of air pollution and action that can be taken to reduce impacts;**
- C. Health advice linked to each band to accompany the air quality index.**

These are detailed below:

## A: HOW TO USE THE DAILY AIR QUALITY INDEX

**Step 1:** Determine whether you (or your children) are likely to be at-risk from air pollution. Information on groups who may be affected is on the 'Additional information on the effects of air pollution' page. Your doctor may also be able to give you advice.

**Step 2:** If you may be at-risk, and are planning strenuous activity outdoors, check the air pollution forecast.

**Step 3:** Use the health messages corresponding to the highest forecast level of pollution as a guide.

## B: ADDITIONAL INFORMATION ON THE SHORT-TERM EFFECTS OF AIR POLLUTION

The daily air quality index (DAQI) has been developed to provide advice on expected levels of air pollution. In addition, information on the short-term effects on health that might be expected to occur at the different bands of the index (Low, Moderate, High, Very High) is provided. It is possible that very sensitive individuals may experience health effects even on Low air pollution days. This advice applies to anyone experiencing symptoms.

### Short-term effects of air pollution on health

Air pollution has a range of effects on health. However, air pollution in the UK does not rise to levels at which people need to make major changes to their habits to avoid exposure; nobody need fear going outdoors.

Adults and Children with lung or heart conditions - It is known that, when levels of air pollutants rise, adults suffering from heart conditions, and adults and children with lung conditions, are at increased risk of becoming ill and needing treatment. Only a minority of those who suffer from these conditions are likely to be affected and it is not possible to predict in advance who will be affected. Some people are aware that air pollution affects their health: adults and children with asthma may notice that they need to increase their use of inhaled reliever medication on days when levels of air pollution are higher than average.

Older people are more likely to suffer from heart and lung conditions than young people and so it makes good sense for them to be aware of current air pollution conditions.

The general population - At Very High levels of air pollution, some people may experience a sore or dry throat, sore eyes or, in some cases, a tickly cough even in healthy individuals.

Children need not be kept from school or prevented from taking part in games. Children with asthma may notice that they need to increase their use of reliever medication on days when levels of air pollution are higher than average.

### Action that can be taken

When levels of air pollution increase it would be sensible for those who have noticed that they are affected to limit their exposure to air pollutants. This does not mean staying indoors, but reducing levels of exercise outdoors would be reasonable.

Older people and those with heart and lung conditions might avoid exertion on High pollution days.

Adults and children with asthma should check that they are taking their medication as advised by their health practitioner and may notice that they need to increase their use of inhaled reliever medication.

Adults with heart and circulatory conditions should **not** modify their treatment schedules on the basis of advice provided by the air quality index: such modification should only be made on a health practitioner's advice.

Some athletes, even if they are not asthmatic, may notice that they find their performance less good than expected when levels of a certain air pollutant (ground level ozone) are High, and they may notice that they find deep breathing causes some discomfort in the chest: This might be expected in summer on days when ground level ozone levels are raised. This does not mean that they are in danger but it would be sensible for them to limit their activities on such days.

**C: HEALTH ADVICE TO ACCOMPANY THE DAILY AIR QUALITY INDEX.**

Air pollution banding	Value	Accompanying health messages for at-risk groups and the general population	
		At-risk individuals *	General population
<b>Low</b>	<b>1 – 3</b>	<b>Enjoy</b> your usual outdoor activities.	<b>Enjoy</b> your usual outdoor activities.
<b>Moderate</b>	<b>4 – 6</b>	Adults and children with lung problems, and adults with heart problems, <b>who experience symptoms</b> , should <b>consider reducing</b> strenuous physical activity, particularly outdoors.	<b>Enjoy</b> your usual outdoor activities.
<b>High</b>	<b>7-9</b>	Adults and children with lung problems, and adults with heart problems, should <b>reduce</b> strenuous physical exertion, particularly outdoors, and particularly if they experience symptoms. People with asthma may find they need to use their reliever inhaler more often. Older people should also <b>reduce</b> physical exertion	Anyone experiencing discomfort such as sore eyes, cough or sore throat should <b>consider reducing</b> activity, particularly outdoors.
<b>Very High</b>	<b>10</b>	Adults and children with lung problems, adults with heart problems, and older people, should <b>avoid</b> strenuous physical activity. People with asthma may find they need to use their reliever inhaler more often.	<b>Reduce</b> physical exertion, particularly outdoors, especially if you experience symptoms such as cough or sore throat.

\* Adults and children with heart or lung problems are at greater risk of symptoms. Follow your doctor’s usual advice about exercising and managing your condition. It is possible that very sensitive individuals may experience health effects even on Low air pollution days. Anyone experiencing symptoms should follow the guidance provided in section B.

# Questions and answers on the Daily Air Quality Index (DAQI)

## 1. What is the Daily Air Quality Index?

The daily air quality index (DAQI) is used to communicate information about real-time and forecast levels of outdoor air pollution in the short-term. The DAQI uses a 1-10 index divided into four bands ('Low', 'Moderate', 'High' and 'Very High') to provide more detail about air pollution levels in a simple way, similar to the sun index or pollen index. The key index pollutants are particulate matter, ozone, sulphur dioxide, nitrogen dioxide, and carbon monoxide.

## 2. What is the DAQI used for?

Air quality information is communicated in terms of the DAQI and provides warning of potentially health-damaging air pollution events in the short-term. With advanced warning of poor air quality, individuals who are susceptible to the effects of air pollution, such as those with respiratory or cardio-vascular conditions, have the opportunity to manage their condition and reduce the severity of their symptoms.

## 3. Who uses the DAQI?

Individuals and organisations who want to know how good or bad air pollution is will look at air quality information which is described in terms of the DAQI. Defra, Local Authorities, and Universities use the DAQI to communicate air quality information, for example see:

- <http://uk-air.defra.gov.uk/air-pollution/bandings>
- <http://www.airtext.info/>
- <http://www.londonair.org.uk/london/asp/nowcast.asp>

## 4. How was the DAQI developed?

The DAQI was developed by Committee on the Medical Effects of Air Pollutants (COMEAP, <http://www.comeap.org.uk/>) and is based on the latest available health evidence. The 'Low' bands indicate air pollution levels where it is unlikely that anyone will suffer any adverse effects of short-term exposure, including people with lung or heart conditions who may be more susceptible to the effects of air pollution. The 'Moderate' band represents levels of air pollutants at which there are likely to be small effects that only affect susceptible people. Values for the 'High' bands are associated with significant effects in susceptible people. At 'Very High' levels of air pollution even healthy individuals may experience adverse effects of short-term exposure.

COMEAP reviewed the DAQI and published a report *Review of the UK Air Quality Index* in June 2011. See <http://comeap.org.uk/documents/reports/130-review-of-the-uk-air-quality-index.html>

## 5. Why was the DAQI reviewed?

The Committee on the Medical Effects of Air Pollutants (COMEAP) was asked by the Department of Environment, Food and Rural Affairs (Defra) to review the UK's air quality index (AQI) to ensure that it is fit for purpose. The current UK air quality index has now been in operation essentially unchanged for a period of around 12 years. Therefore, it was

appropriate to review the index to establish its suitability, given the developments in the field of air quality.

## **6. Why has Defra adopted the World Health Organisation guidelines for ozone instead of the advice recommended by COMEAP?**

COMEAP assessed the latest scientific evidence and provided Defra with their recommendations. Defra evaluated the recommendations in a wider context, taking other factors into consideration, including the purpose of the index and its users, to ensure that we are providing appropriate guidance to the public on daily levels of air pollution, and information on actions individuals can take to reduce the impacts on their health.

Defra will adopt a Low/Moderate breakpoint of  $100 \mu\text{g m}^{-3}$  for ozone. COMEAP recommended the breakpoint between the Low and Moderate band for ozone be set at  $80 \mu\text{g m}^{-3}$ . WHO recommend a guideline of  $100 \mu\text{g m}^{-3}$  but noted that some effects may be expected in susceptible individuals within the population at its recommended guideline of  $100 \mu\text{g m}^{-3}$ , and because of this COMEAP recommended the lower breakpoint of  $80 \mu\text{g m}^{-3}$ . However, the  $80 \mu\text{g m}^{-3}$  value is close to the hemispheric background level of ozone and it is likely that this level will be exceeded on the majority of days in the UK (up to 80% of days in a year) and elsewhere, and air pollution would be described as Moderate on most days.

As part of the review, customer insight research was conducted to investigate how people interpret air quality information, and how the provision of air quality information could be improved. A recommendation of the research was that users would like to see more variation in the reported levels of air pollution. The WHO guidelines provide for greater variation in the number of Low and Moderate air pollution days. If Moderate air pollution is reported on the majority of days then the importance of messages warning of elevated air pollution will be lost.

Additional text has been added to the index (in agreement with COMEAP) to ensure that individuals have the information they need to manage a health condition. The following statement has been added to the index information to inform of the risk:

*'It is possible that very sensitive individuals may experience health effects even on Low air pollution days. Anyone experiencing symptoms should follow the guidance provided.'*

COMEAP have been consulted on the changes to the Daily Air Quality Index and are supportive of the amendments.

Defra are committed to improving outdoor air quality. Reductions in the hemispheric background of ozone will require a global approach, as it is a trans-boundary pollutant.

## **7. When will the changes to the index be implemented?**

The changes to the DAQI will be implemented from 1<sup>st</sup> January 2012.

**8. There will be more moderate and higher days of air pollution reported with the revised index, does this mean that air pollution is getting worse?**

It may appear that air pollution is getting worse as more moderate days and higher will be reported. However, this not the case, air pollution is being reclassified. The revised index reflects updated scientific evidence and redresses the previous underestimation of the effects of short-term exposure to air pollution. The changes to the index will allow individuals who are susceptible to the effects of air pollution to manage their condition. This does not mean that levels of air pollution are increasing.