

Air Pollution in the UK 2016

September 2017





© Crown copyright 2017

You may re-use this information (excluding logos) free of charge in any format or medium, under the terms of the Open Government Licence v.3. To view this licence visit <u>www.nationalarchives.gov.uk/doc/open-government-licence/version/3/</u> or email <u>PSI@nationalarchives.gsi.gov.uk</u>

This publication is available at https://uk-air.defra.gov.uk/library/annualreport/index

Any enquiries regarding this publication should be sent to us at

air.quality@defra.gsi.gov.uk

Executive Summary

The UK is currently required to report air quality data on an annual basis under the following European Directives:

- The Council Directive on ambient air quality and cleaner air for Europe (2008/50/EC).
- The Fourth Daughter Directive (2004/107/EC) under the Air Quality Framework Directive (1996/62/EC).

This report provides background information on the pollutants covered by these Directives and the UK's Air Quality Strategy; their sources and effects, the UK's statutory monitoring networks, and the UK's modelling methodology. The report then summarises the UK's 2016 submission on ambient air quality to the European Commission, presenting air quality modelling data and measurements from national air pollution monitoring networks. The pollutants covered in this report are:

- Sulphur dioxide (SO₂)
- Nitrogen oxides (NO_x) comprising NO and NO₂
- PM₁₀ and PM_{2.5} particles
- Benzene
- 1,3-Butadiene
- Carbon Monoxide (CO)
- Metallic Pollutants
- Polycyclic aromatic hydrocarbons (PAH)
- Ozone (O₃)

These data are reported on behalf of Defra (the Department for Environment, Food and Rural Affairs) and the Devolved Administrations of Scotland, Wales and Northern Ireland.

For the purposes of air quality monitoring and assessment of compliance with the above Directives, the UK is divided into 43 zones. The 2016 results are detailed in section 4 of this report and summarised below:

- The UK met the limit value for hourly mean nitrogen dioxide (NO₂) in all but two zones.
- Six zones were compliant with the limit value for annual mean NO₂. The remaining 37 exceeded this limit value.
- Four zones exceeded the target value for benzo[a]pyrene.
- Three zones exceeded the target value for nickel.
- All zones met both the target values for ozone; the target value based on the maximum daily eight-hour mean, and the target value based on the AOT40 statistic.

- All zones except one exceeded the long-term objective for ozone, set for the protection of human health. This is based on the maximum daily eight-hour mean.
- Five zones exceeded the long-term objective for ozone, set for the protection of vegetation. This is based on the AOT40 statistic.
- All zones met the limit value for daily mean concentration of PM₁₀ particulate matter, without the need for subtraction of the contribution from natural sources.
- All zones met the limit value for annual mean concentration of PM₁₀ particulate matter, without the need for subtraction of the contribution from natural sources.
- All zones met the target value for annual mean concentration of PM_{2.5} particulate matter, the Stage 1 limit value, which came into force on 1st January 2015, and the Stage 2 limit value which must be met by 2020.
- All zones met the EU limit values for sulphur dioxide, carbon monoxide, lead and benzene.

A summary of the air quality assessment for 2016 with a comparison of the submissions carried out in the previous years (since 2008 when the Air Quality Directive came into force) can be found in section 4.3 of this report. Copies of those previous annual submissions can be found on the Commission website:

<u>http://cdr.eionet.europa.eu/gb/eu/annualair.</u> For more information on air quality in the UK visit the Defra website at <u>www.gov.uk/defra</u> and the UK Air Quality websites at <u>http://uk-air.defra.gov.uk/</u>, <u>www.scottishairquality.co.uk</u>, <u>www.welshairquality.co.uk</u> and <u>www.airqualityni.co.uk</u>.

Glossary

Air Quality Directive. The European Union's Directive 2008/50/EC of 21st May 2008, on Ambient Air Quality and Cleaner Air for Europe is often – as in this report - referred to as 'the Air Quality Directive'.

Air Quality Strategy. The United Kingdom's own National Air Quality Strategy, containing policies for assessment and management of air quality in the UK. This was first published in 1997, as a requirement of The Environment Act 1995.

Air Quality Strategy Objective. The Air Quality Strategy sets objectives for the maximum concentrations of eight pollutants. These are at least as stringent as the limit values of the Air Quality Directive.

Ambient Air. Outdoor air.

Benzene. A chemical compound that is harmful to human health. As an air pollutant, benzene can be emitted from domestic and industrial combustion processes, and road vehicles. Its chemical formula is $C_6 H_6$.

Benzo [a] Pyrene. One of a group of compounds called *polycyclic aromatic hydrocarbons (PAHs)* that can be air pollutants. The main sources of B[a]P in the UK are domestic coal and wood burning, fires, and industrial processes such as coke production.

1,3-Butadiene. This is an organic compound emitted into the atmosphere mainly from fuel combustion e.g. petrol and diesel vehicles. 1,3-butadiene is also an important chemical in certain industrial processes, particularly the manufacture of synthetic rubber. 1,3-butadiene is known to cause cancer in humans.

Carbon Monoxide (CO) a pollutant gas found released in road vehicle exhausts. When breathed in, carbon monoxide affects the blood's ability to carry oxygen around the body.

Episode (Air Pollution Episode). An 'air pollution episode' means a period of time (usually a day or several days) when air pollution is high (air quality is poor).

FDMS. This stands for 'Filter Dynamic Measurement System' and refers to a type of instrument for monitoring concentrations of particulate matter. The FDMS is a modified form of *TEOM*.

Fourth Daughter Directive. The European Union's Directive 2004/107/EC, which covers the four metallic elements cadmium, arsenic, nickel and mercury together with *polycyclic aromatic hydrocarbons (PAH)*. (Its name comes from its origin as one of four so-called Daughter Directives set up under an overarching 'framework Directive'.)

Limit value. The Air Quality Directive sets 'limit values' for ambient concentrations of pollutants. Limit values are legally binding and must not be exceeded. All Member States of the EU must make the limit values part of their own air quality legislation.

Long-Term Objectives. As well as limit values and target values, the Air Quality Directive sets 'long-term objectives' for ozone concentration. These are similar to limit values but are not legally mandatory. Member States must take all necessary measures not entailing disproportionate costs to meet the target values and long-term objectives.

Member States. Countries that are part of the European Union.

Microgramme per cubic metre (\mug m⁻³). Unit often used to express concentration of a pollutant in air. 1 μ g = 1 millionth of a gramme or 1 x 10⁻⁶ g.

Micrometre (µm). Unit of length often used for the size of particulate pollutants. $1 \mu m = 1$ millionth of a metre (1 x 10⁻⁶ m) or one thousandth of a millimetre.

Milligramme per cubic metre (mg m⁻³). Unit often used to express concentration of carbon monoxide in air. 1 mg = 1 thousandth of a gramme or 1×10^{-3} g.

Nickel (Ni) A toxic metallic element found in ambient air as a result of releases from oil and coal combustion, metal processes, manufacturing and other sources

Nitric oxide (NO). One of the oxides of nitrogen formed in combustion processes. NO is not harmful to human health but combines with oxygen to form nitrogen dioxide.

Nitrogen Dioxide (NO₂) One of the oxides of nitrogen formed in combustion processes. At high concentrations NO₂ is an irritant to the airways. NO₂ can also make people more likely to catch respiratory infections (such as flu), and to react to allergens.

Nitrogen Oxides. Compounds formed when nitrogen and oxygen combine. NOx, which comprises nitric oxide (NO) and nitrogen dioxide (NO₂), is emitted from combustion processes. Main sources include power generation, industrial combustion and road transport.

Ozone (O₃). A pollutant gas which is not emitted directly from any source in significant quantities, but is produced by reactions between other pollutants in the presence of sunlight. (This is what is known as a *'secondary pollutant'*.) Ozone concentrations are greatest in the summer. O₃ can travel long distances and reach high concentrations far away from the original pollutant sources. Ozone is an irritant to the airways of the lungs, throat and eyes: it can also harm vegetation.

Particulate Matter (PM). Small airborne particles. PM may contain many different materials such as soot, wind-blown dust or secondary components, which are formed

within the atmosphere as a result of chemical reactions. Some PM is natural and some is man-made. Particulate matter can be harmful to human health when inhaled, and research shows a range of health effects associated with PM. In general, the smaller the particle the deeper it can be inhaled into the lung.

PM₁₀. Particles which pass through a size-selective inlet with a 50 % efficiency cut-off at 10 µm aerodynamic diameter, as defined in ISO 7708:1995, Clause 6. This size fraction is important in the context of human health, as these particles are small enough to be inhaled into the airways of the lung – described as the 'thoracic convention' in the above ISO standard. PM₁₀ is often described as '*particles of less than 10 micrometres in diameter*' though this is not strictly correct.

PM_{2.5}. Particles which pass through a size-selective inlet with a 50 % efficiency cut-off at 2.5 µm aerodynamic diameter, as defined in ISO 7708:1995, Clause 7.1. This size fraction is important in the context of human health, as these particles are small enough to be inhaled very deep into the lung – described as the 'high risk respirable convention' in the above ISO standard. PM_{2.5} is often described as '*particles of less than 2.5 micrometres in diameter*' though this is not strictly correct.

Polycyclic Aromatic Hydrocarbons (PAH). PAHs are a large group of chemical compounds that are toxic and carcinogenic. Once formed, they can remain in the environment for a long time, and can be passed up the food chain. The main sources are domestic coal and wood burning, outdoor fires, and some industrial processes. The pollutant *benzo [a] pyrene* is a PAH, and because it is one of the more toxic PAH compounds it is measured as a 'marker' for this group of pollutants.

Secondary pollutant. A pollutant which is formed by chemical reactions from other pollutants in the atmosphere. Ozone, for example, is a secondary pollutant.

Sulphur dioxide (SO₂). An acid gas formed when fuels containing sulphur impurities are burned. SO₂ irritates the airways of the lung.

Target Value. As well as limit values, the *Air Quality Directive* and *Fourth Daughter Directive* set target values for some pollutants. These are similar to limit values but are not legally mandatory. *Member States* must take all necessary measures not entailing disproportionate costs to meet the target values.

TEOM. This stands for 'Tapered Element Oscillating Microbalance'. This is a type of instrument used to monitor concentrations of particulate matter.

TOMPs. This stands for 'Toxic Organic Micropollutants'. These are compounds that are present in the environment at very low concentrations, but are highly toxic and persistent. They include dioxins and dibenzofurans.

Contents

1	In	ntroduction1				
2	Le	gislative and Policy Framework3				
	2.1	European and International Background	3			
	2.2	The UK Perspective	7			
	2.3	Local Authority Air Quality Management	.13			
3	Tł	ne Evidence Base	.18			
	3.1	Pollutants of Concern	.18			
	3.2	Assessment of Air Quality in the UK	.27			
	3.3	Current UK Air Quality Monitoring	.27			
	3.4	Quality Assurance and Quality Control	.34			
	3.5	Modelling	.36			
	3.6	Access to Assessment Data	.38			
4	As	ssessment of Compliance	.39			
	4.1	Definition of Zones	.39			
	4.2	Air Quality Assessment for 2016	.41			
	4.3	Comparison with Previous Years	.51			
5	Sp	patial Variation and Changes Over Time	.56			
	5.1	Sulphur Dioxide	.57			
	5.2	Nitrogen Dioxide	.61			
	5.3	PM ₁₀ Particulate Matter	.66			
	5.4	PM _{2.5} Particulate Matter	.71			

	5.5	Benzene	75
	5.6	1,3-Butadiene	78
	5.7	Carbon Monoxide	80
	5.8	Ozone	83
	5.9	Metallic Elements	87
	5.10	Benzo [a] Pyrene	95
6	Pc	ollution Events in 2016	99
	6.1	Winter and Spring Particulate Pollution Episodes	99
	6.2	Summer Ozone and Particulate Matter Events	.105
	6.3	Bonfire Night Particulate Pollution Event	.107
	6.4	Forecasting Air Pollution Events	.111
7	W	here to Find Out More	.112
R	efere	nces	.114

This page is intentionally blank.

1 Introduction

A cleaner, healthier environment benefits people and the economy. Clean air is vital for people's health and the environment, essential for making sure our cities are welcoming places for people to live and work now and in the future, and to our prosperity. It is therefore important to monitor levels of air pollution. The broad objectives of monitoring air pollution in the UK are:

- To fulfil statutory air quality reporting requirements.
- To provide a sound scientific basis for the development of cost-effective control policies.
- To provide the public with open, reliable and up-to-date information on air pollution, enabling them to take appropriate action to minimise health impacts.
- To evaluate potential impacts on population, ecosystems and our natural environment.

Air quality standards are set in European Union (EU) Directive 2008/50/EC on Ambient Air Quality and Cleaner Air for Europe¹ and the Fourth Daughter Directive² (2004/107/EC). These Directives require all Member States to undertake air quality assessment, and to report the findings to the European Commission on an annual basis.

The UK has statutory monitoring networks in place to meet the requirements of these Directives, with air quality modelling used to supplement the monitored data. The results must be submitted to the European Commission each year. As of 2013, the air quality compliance assessment has been submitted to the Commission via e-Reporting (see Section 2.1.2). The UK's annual submission for 2016 can be found on the Commission website at http://cdr.eionet.europa.eu/gb/eu/aqd. All the compliance results are reported under 'Information on the Attainment of Environmental Objectives' in e-Reporting Data Flow G. Submissions for years up to and including 2012 (which were in the form of a standard questionnaire) can be found at http://cdr.eionet.europa.eu/gb/eu/aqd.

As well as reporting air quality data to the European Commission, the UK must also make the information available to the public. One way in which this is done is by the series of annual 'Air Pollution in the UK' reports. '*Air Pollution in the UK 201*6' continues this series, and this report has two aims:

• To provide a summary of the UK's 2016 air quality report to the Commission. A separate Compliance Assessment Summary document,

based upon Section 4 of this report, accompanies the UK's 2016 data submission to the Commission. This provides a concise summary aimed at the public.

• To act as a State of the Environment report, making information on the ambient air quality evidence base for the year publicly available. This includes an assessment of trends and spatial distribution, together with information on pollution events during the year.

This report:

- Outlines the air quality legislative and policy framework in Europe and the UK (Section 2).
- Describes the evidence base underpinning the UK's air quality assessment: the pollutants of concern, and where and how air pollution is measured and modelled *(Section 3)*.
- Presents an assessment of the UK's compliance with the limit values, target values and long term objectives set out in the Air Quality Directive and the Fourth Daughter Directive for 2016, and compares this with previous recent years. (Section 4).
- Explains the spatial distribution of the main pollutants of concern within the UK during 2016, and looks at how ambient concentrations have changed in recent years (*Section 5*).
- Explains pollution events 'episodes' of high pollution that occurred during 2016, (*Section 6*).

Further information on air quality in the UK can be found on Defra's online UK Air Information Resource (UK-AIR), at <u>http://uk-air.defra.gov.uk/</u>.

2 Legislative and Policy Framework

The UK air quality framework is derived from a mixture of domestic, EU and international legislation and consists of three main strands:

- Legislation regulating total emissions of air pollutants the UK is bound by both EU law (the National Emission Ceilings Directive) and international law (the Gothenburg Protocol to the UNECE Convention on Long-range Transboundary Air Pollution);
- 2) Legislation regulating concentrations of pollutants in the air; and
- 3) Legislation regulating emissions from specific sources such as legislation implementing the Industrial Emissions Directive and the Clean Air Act.

Reducing air pollution requires action to reduce domestic emissions as well as working closely with international partners to reduce transboundary emissions (pollutants blown over from other countries) which, at times, can account for a significant proportion of pollutant concentrations experienced in the UK. For example, a 2013 report³ prepared by the Air Quality Expert Group on behalf of Defra estimated that emission sources within the UK only accounted for 50-55% of measured annual average fine particulate matter (PM_{2.5}) concentrations, the remainder being formed or emitted elsewhere.

2.1 European and International Background

European Union (EU) air pollution legislation follows two complementary approaches;

- (i) controlling emissions at source, and
- (ii) setting of ambient air quality standards and long-term objectives.

All Member States must incorporate - or 'transpose' - the provisions of EU Directives into their own national law by a specified date and comply with legally binding implementing rules set out in the Decisions. The main Directives and Decisions are described below.

The UK has voted (in the referendum of 23rd June 2016) to leave the European Union, and in March 2017 the UK Government began the formal process of doing so. The UK has a long history of environmental protection and will safeguard and improve this. The Repeal Bill will ensure that the whole body of existing EU environmental law continues to have effect in UK law.

2.1.1 The Air Quality Directive and Fourth Daughter Directive

Directive 2008/50/EC of 21st May 2008, on Ambient Air Quality and Cleaner Air for Europe – referred to in this report as 'the Air Quality Directive'¹ - covers the following pollutants; sulphur dioxide, nitrogen oxides, particulate matter (as PM₁₀ and PM_{2.5}), lead, benzene, carbon monoxide and ozone. It revised and consolidated existing EU air quality legislation relating to the above pollutants.

Directive 2004/107/EC of 15th December 2004, relating to arsenic, cadmium, mercury, nickel and polycyclic aromatic hydrocarbons in ambient air - referred to in this report as 'the Fourth Daughter Directive'² - covers the four elements cadmium, arsenic, nickel and mercury, together with polycyclic aromatic hydrocarbons (PAH).

These two Directives set 'limit values', 'target values' and 'long-term objectives' for ambient concentrations of pollutants.

Limit values are legally binding and must not be exceeded. They are set for individual pollutants and comprise a concentration value, an averaging period for the concentration value, a number of exceedances allowed (per year) and a date by which it must be achieved. Some pollutants have more than one limit value, for example relating to short-term average concentrations (such as the hourly mean) and long-term average concentrations (such as the annual mean).

Target values and long-term objectives are set for some pollutants and are configured in the same way as limit values. Member States must take all necessary measures not entailing disproportionate costs to meet the target values and longterm objectives.

The Air Quality Directive and Fourth Daughter Directive include detailed provisions on monitoring and reporting air quality, including:

- The division of the UK into zones for the purposes of compliance reporting.
- The location and number of sampling points.
- The measurement methods to be used.
- Data quality objectives.
- Criteria each monitoring station must meet.
- Provision for reporting compliance.
- Provision of information to the public.

The Air Quality Directive also makes provisions for adapting standardised procedures to streamline the data provision, assessment and reporting of air quality, to electronically release information in line with the INSPIRE Directive (2007/2/EC). This led to the adoption of new implementing provisions for reporting (IPR) (Decision

2011/850/EC, referred to in this report as the Air Quality e-Reporting IPR). Section 2.1.2 provides a detailed description of the Air Quality e-Reporting IPR. The report of 2013 was the first in which e-Reporting was used to report emissions.

The provisions of the Air Quality Directive and Fourth Daughter Directive were transposed by the Air Quality Standards Regulations 2010⁴ in England, the Air Quality Standards (Scotland) Regulations 2010⁵ in Scotland, the Air Quality Standards (Wales) Regulations 2010 in Wales⁶ and the Air Quality Standards Regulations (Northern Ireland) 2010⁷. All the provisions made by the Directives are therefore incorporated into UK legislation.

The above Directives were amended by the Technical Directive 2015/1480/EC⁸ (which amended some details of the Directives' annexes relating to reference methods, data validation and locations of sampling points). The relevant regulations were amended accordingly in December 2016.

2.1.2 Air Quality e-Reporting

Defra is committed to the principles of Open Data. Air Quality e-Reporting is a process, developed by the European Commission, for reporting of compliance and provision of data under the Air Quality Directive. The development has been driven by the requirements of the INSPIRE Directive (which is concerned with the sharing of spatial data across EU Member States in a consistent and computer-readable format). Air Quality e-Reporting is a key tool to help ensure UK air quality data systems comply with the INSPIRE Directive and are available across Europe in a comparable form with other Member States. Operational Air Quality e-Reporting started on 1st January 2014; as of 21st October 2015, newly collected or extensively restructured spatial data sets have had to be available in INSPIRE conformant formats, and as of 10th December 2015 web based spatial data download services have been required to conform to INSPIRE Regulations.

European Commission Implementing Decision 2011/850/EU⁹ was introduced on 12th December 2011. This Decision laid down rules for the reciprocal exchange of information, and reporting on ambient air quality, in relation to the Air Quality Directive. The Decision provided an opportunity to modernise data reporting, improve data quality, facilitate information sharing and reduce the administrative burden of reporting. In adapting the procedures, Air Quality e-Reporting has embraced digital formats for reporting, and the internet as the core medium for reporting. Air Quality e-Reporting extends the core requirements of the INSPIRE Directive to meet the particular requirements of regulatory and informative air quality reporting.

The European Commission developed these procedures with assistance from the European Topic Centre on Air and Climate Change Mitigation and the European Environment Agency (EEA), and in close liaison with the European Environment Information and Observation Network (EIONET) air quality community. The e-Reporting system covers all regulatory and information reporting agreements set out by the Exchange of Information Decision (EoI) (Council Decision 97/101/EC¹⁰), the Air Quality Directive and the 4th Daughter Directive. By adopting data modelling approaches prescribed by INSPIRE, the e-Reporting data model is streamlined, internally consistent and meets modern standards for data encoding and data sharing. The data model is now organised into eight broad air quality data themes that service all reporting and information sharing needs of the air quality community.

2.1.3 The National Emission Ceilings Directive

The original National Emission Ceilings Directive (2001/81/EC) came into force in 2001, and was transposed into UK legislation by the National Emission Ceilings Regulations 2002. This Directive set national emission limits or 'ceilings' for the four main air pollutants responsible for the acidification and eutrophication (nutrient enrichment) of the natural environment, and the formation of ground level ozone which impacts both human health and the environment. The ceilings had to be met by 2010. They reflected the ceilings agreed internationally in the 1999 Gothenburg Protocol to the UNECE Convention on Long Range Transboundary Air Pollution (CLRTAP). Emissions of these pollutants can impact either locally or across national borders. The latter is known as transboundary air pollution. The 2001 Directive set emissions ceilings for: sulphur dioxide, oxides of nitrogen, volatile organic compounds, and ammonia which have applied since 2010.

The UK meets all current emission ceilings set under the National Emission Ceilings Directive. The National Emission Ceilings Directive report is available at https://www.eea.europa.eu/publications/nec-directive-status-report-2015.

The Gothenburg Protocol was revised in May 2012 to set emission reduction commitments (ERCs) for 2020 (from the 2005 baseline) for the same four pollutants and $PM_{2.5}$. The European Commission published a proposal to revise the National Emission Ceilings Directive in December 2013, to implement the new 2020 ERCs in the Protocol, and to set further commitments for 2030.

The revised National Emission Ceilings Directive (<u>Directive 2016/2284/EU</u>)¹¹ came into force on 31st December 2016. The Directive will be transposed into UK legislation by 1st July 2018. Under the new Directive and the amended Gothenburg Protocol, the UK has ambitious targets in place to reduce emissions of five damaging air pollutants (ammonia, nitrogen oxides, non-methane volatile organic compounds, fine particulate matter and sulphur dioxide) by 2020 and 2030. Defra will publish a Clean Air Strategy, setting out how we will work towards these goals, for consultation in 2018.

2.1.4 The Industrial Emissions Directive

The Industrial Emissions Directive (Directive 2010/75/EU) sets stringent provisions to reduce the emissions of pollutants from a diverse range of industrial sources - from intensive pig and poultry farms to chemical manufacturing sites and power stations – with the aim of achieving the environmental and human health benefits associated with a reduction in pollution. Under the Directive, industries must use best available techniques (BAT) to reduce their emissions. These techniques, and the emissions limits associated with the use of those techniques, are set out in best available technique reference documents (known as BREFs). BREFs are reviewed regularly, which will ensure an ongoing process of improvement to EU air quality achieved through the continuous reduction of pollution. In addition, the Directive sets emission limits for the emission of pollutants from particular sectors.

2.2 The UK Perspective

Domestic, EU and internationally-driven environmental legislation introduced over the past seventy years has provided a strong impetus to reduce the levels of harmful pollutants in the UK; as a result, current concentrations of many recognised pollutants are now at the lowest they have been since measurements began. Following the UK's Clean Air Act, the city smogs of the 1950s, caused by domestic and industrial coal burning, have now gone for good and significant progress has been made in improving air quality throughout recent decades. Between 2000 and 2015, UK estimated emissions of nitrogen oxides have fallen by 49%, and UK estimated emissions of PM₁₀ particulate matter have fallen by almost 26% (data from the National Atmospheric Emissions Inventory at <u>http://naei.beis.gov.uk/</u>).

Poor air quality is the greatest environmental risk to public health in the UK. It is known to exacerbate the impact of pre-existing health conditions, such as respiratory and cardio-vascular illnesses, especially for the elderly and infants.

Nitrogen dioxide (NO₂) is associated with adverse effects on human health. Although it has been more difficult to estimate the level of impact, there is enough evidence of such health effects to support the need to take action now.

It is likely that there is some overlap between the health impacts associated with ambient concentrations of particulate matter (PM) and NO₂. This is because of the uncertainty around causality between NO₂ and mortality and the potential overlap

between the health effects of PM and NO₂. Further work is being undertaken to understand and quantify this overlap, but the current recommendation is that between 0 and 33% of the effects associated with ambient concentrations of the two pollutants overlap.

2.2.1 The UK Air Quality Strategy

The Environment Act 1995 required that a National Air Quality Strategy be published, containing policies for assessment and management of air quality. The Air Quality Strategy¹² for England, Scotland, Wales and Northern Ireland was first published in March 1997. The overall objectives of the Strategy are to:

- Map out future ambient air quality policy in the United Kingdom in the medium term.
- Provide best practicable protection to human health by setting healthbased objectives for air pollutants.
- Contribute to the protection of the natural environment through objectives for the protection of vegetation and ecosystems.
- Describe current and future levels of air pollution.
- Establish a framework to help identify what we all can do to improve air quality.

The Strategy has established objectives for eight key air pollutants, based on the best available medical and scientific understanding of their effects on health, as well as taking into account relevant developments in Europe and the World Health Organisation. These Air Quality Objectives¹³ are at least as stringent as the limit values of the relevant EU Directives – in some cases, more so. The most recent review of the Strategy was carried out in 2007.

2.2.2 National Air Quality Statistics and Indicators

The UK reports on the following two indicators as National Air Quality Statistics for ambient air:

- Annual average concentrations of particles and ozone. These two types of air pollution are believed to have a significant impact on public health.
- Number of days in the year when air pollution is 'Moderate' or higher. This may relate to any one of five key air pollutants and is based on the UK's Daily Air Quality Index (see Section 2.2.4). From the 1st January 2012, PM_{2.5} particles replaced carbon monoxide in this suite of pollutants. The thresholds

used to define 'Moderate' and higher pollution levels in the air quality index were also revised at the beginning of 2012.

The National Air Quality Statistics summary for 2016 was released on 23rd April 2017 and is available from the Defra website¹⁴.

In August 2016, Defra published a revised edition of the England Natural Environment Indicators¹⁵. Indicator 11 for Environmental Quality and Health relates to air quality. This is based on:

- The average number of days per site when air pollution is 'Moderate' or higher for urban and for rural sites,
- Regional mortality due to anthropogenic particulate air pollution, compared to the England national average (5.6% in 2010, which is being taken as the baseline year for this indicator).

The UK Government's Public Health Outcomes Framework for England (published in 2012) recognises the burden of ill-health resulting from poor air quality as well as other public health concerns. This Framework sets out 60 health outcome indicators for England, and includes as an indicator:

• The fraction of annual all-cause adult mortality attributable to long-term exposure to current levels of anthropogenic particulate air pollution (measured as fine particulate matter, PM_{2.5})¹⁶.

This indicator is intended to enable Directors of Public Health to appropriately prioritise action on air quality in their local area. The indicator is calculated for each local authority in England based on modelled concentrations of fine particulate air pollution (PM_{2.5}). Estimates of the percentage of mortality attributable to long term exposure to particulate air pollution in local authority areas are available from the Public Health Outcomes Framework data tool at <u>http://www.phoutcomes.info/</u>. Current estimates at the time of writing, which are based on year 2013, range from around 3% in some rural areas to around 8% in some areas of London where pollution levels are highest.

The Defra document 'Air Quality: Public Health Impacts and Local Actions' can be found at <u>https://laqm.defra.gov.uk/documents/air_quality_note_v7a-(3).pdf</u>. A toolkit aimed at helping public health professionals appropriately prioritise assessment and action on $PM_{2.5}$ on a local level is available here:

http://randd.defra.gov.uk/Default.aspx?Menu=Menu&Module=More&Location=None &Completed=0&ProjectID=18580. Northern Ireland has a similar Public Health Strategy: *'Making Life Better – A Whole System Framework for Public Health 2013-2023'*. This document can be found at <u>https://www.health-ni.gov.uk/topics/public-health-policy-and-advice/making-life-better-whole-system-strategic-framework-public</u>, and also includes an air quality indicator.

Wales has a new national indicator under the Well-being of Future Generations (Wales) Act 2015 and the Welsh Public Health Outcomes Framework, which has now been published on StatsWales at

(https://statswales.gov.wales/Catalogue/Environment-and-Countryside/Air-Quality).

In 2015 the Scottish Government published '*Cleaner Air for Scotland – The Road to a Healthier Future*', Scotland's first separate air quality strategy. The first annual progress report, setting out progress towards delivering the actions in the strategy, was published in June 2017. This report is available for download at: <u>http://www.gov.scot/Publications/2017/06/2881/downloads#res521031</u>.

2.2.3 National Emissions Statistics

The UK reports annual emissions of the following pollutants via an annual National Statistics Release, available at <u>https://www.gov.uk/government/statistics/emissions-of-air-pollutants</u> :

- sulphur dioxide,
- oxides of nitrogen,
- non-methane volatile organic compounds (NMVOCs),
- ammonia (NH₃),
- particulate matter (as PM₁₀ and PM_{2.5}).

The most recent National Statistics Release covers 1970 to 2015 (the most recent year for which emission statistics are available). The main conclusions are as follows:

- 'There has been a long term decrease in the emissions of all of the air pollutants covered by this statistical release (ammonia, nitrogen oxides, non-methane volatile organic compounds, particulate matter (PM₁₀, PM_{2.5}) and sulphur dioxide).
- Emissions of sulphur dioxide decreased by 23 per cent from 2014 to 2015, dropping to the lowest level in the time series.
- Emissions of nitrogen oxides decreased in 2015 compared to 2014 by 4.0 per cent, dropping to the lowest level in the time series.

- Emissions of non-methane volatile organic compounds are continuing to decline, by 0.6 per cent between 2014 and 2015. The rate of decline was most pronounced in the 1990s and early 2000s and has slowed in recent years.
- *PM*₁₀ emissions have remained relatively static over the past five years decreasing by less than 0.1 kilotonnes from 2014 to 2015.
- *PM*_{2.5} emissions increased slightly by 0.9 per cent between 2014 and 2015. 2015 emissions are below the peak value of the last ten years, observed in 2010.
- There was an increase of 1.7 per cent in emissions of ammonia between 2014 and 2015. Increases in the past two years go against the trend of steady reduction observed from 1997 to 2013.
- The UK continues to meet current international and EU ceilings for emissions of ammonia, nitrogen oxides, non-methane volatile organic compounds and sulphur dioxide. The Gothenburg Protocol under the UNECE Convention on Long-range Trans-boundary Air Pollution, which the UK will shortly ratify, was revised in 2012 to set new emission ceilings to apply from 2020. The UK has also recently signed up to the revised National Emission Ceilings Directive (2016/2284/EC) which sets out stricter legally binding ceilings for emissions of five major air pollutants in 2020 and 2030.'

New emission statistics for 2016 are expected to be available in February 2018.

2.2.4 The Air Pollution Forecasting System

Daily UK air pollution forecasts are produced for five pollutants; nitrogen dioxide, sulphur dioxide, ozone, PM₁₀ particles and PM_{2.5} particles. The forecasts are communicated using the Daily Air Quality Index (<u>http://uk-air.defra.gov.uk/air-pollution/daqi</u>) which is a scale of one to ten divided into four bands. This allows the public to see at a glance whether the air pollution is low, moderate, high or very high and to look up any recommended actions to take.

The group of pollutants covered, and the thresholds between the various index bands, were updated by Defra as of 1st January 2012, in the light of recommendations by the Committee on the Medical Effects of Air Pollutants (COMEAP) in their 2011 review of the UK air quality index¹⁷.

The daily forecast is provided by the Met Office and is available from UK-AIR and from the Scottish, Welsh and Northern Ireland air quality websites (see Section 7), and is further disseminated via e-mail, Twitter and RSS feeds. Anyone may subscribe to the free air pollution bulletins at: <u>http://uk-air.defra.gov.uk/subscribe</u> Latest forecasts are issued daily, at: <u>http://uk-air.defra.gov.uk/forecasting/</u>. Defra also

provide automated updates on current and forecast air quality via Twitter @DefraUKAIR- see <u>http://uk-air.defra.gov.uk/twitter</u>.

2.2.5 The NO₂ Air Quality Plans

In July 2017, the Government launched '*The UK Plan for Tackling Roadside Nitrogen Dioxide Concentrations*' and announced £255m for local councils named in the plan as having persistent exceedances to accelerate their air quality plans. This is in addition to the £2.7bn already committed for tackling poor air quality, bringing total investment to £3bn. This includes the following commitments:

- £1bn to improve the infrastructure for ultra-low emission vehicles
- £290m to reduce transport emissions as part of the National Productivity Investment Fund
- £11m awarded to local authorities in the Air Quality Grant
- £89m for a Green bus fund
- £27m for the Clean Bus Technology Fund and Clean Vehicle Technology Fund
- £1.2bn for a Cycling and Walking Investment Strategy
- £100m for air quality as part of the Road Investment Strategy

The plan sets out how the UK will achieve compliance with EU limit values for NO₂ in the shortest possible time. To accelerate action, local areas have been asked to produce initial plans within eight months and final plans by the end of next year. Local authorities will also be able to bid for money from a new Clean Air Fund to support improvements which avoid the need for restrictions on polluting vehicles. This could include upgrading bus fleets, support for concessionary travel and more sustainable modes of transport such as cycling or infrastructure changes.

The plan includes confirmation that government is working with industry to end the sale of new conventional petrol and diesel cars and vans by 2040. The government will also issue a consultation in the autumn of 2017 to gather views on measures to support motorists, residents and businesses affected by the local plans – such as subsidised car club memberships, exemptions from any vehicle restrictions, or a targeted scrappage scheme for car and van drivers in most need of support.

Compliance is forecast to be achieved in all areas outside London by 2021, and in London by 2026 through the measures outlined in the plan.

The national air quality plan is available at https://www.gov.uk/government/publications/air-quality-plan-for-nitrogen-dioxide-no2-in-uk-2017

2.2.6 Measures to Address Target Value Exceedances of B[a]P and Nickel

EU Directive (2004/107/EC) sets target values for a number of metals and for benzo[a]pyrene. During 2013, 2014 and 2015 the UK exceeded target values for two pollutants B[a]P and nickel. These were reported in September of the following years as part of the UK's annual compliance assessment¹⁸.

The UK published reports at the end of 2015 and 2016, providing details of the assessment of these exceedances (2013 and 2014) and reporting the actions and measures that have already been taken or are planned that will help the UK meet the target values. An overview report was provided for each pollutant alongside more detailed information on any exceedances by zone. The reports on the 2014 and 2015 exceedances of the target values for B[a]P and nickel are available at: https://uk-air.defra.gov.uk/library/bap-nickel-measures.

2.3 Local Authority Air Quality Management

Requirements for local air quality management are set out in Part IV of the Environment Act 1995, and the Environment (Northern Ireland) Order 2002¹⁹. Authorities are required to carry out regular 'Review and Assessments' of air quality in their area and take action to improve air quality when the objectives set out in regulation cannot be met by the specified dates.

Local Authorities in England, Scotland, Wales and Northern Ireland have completed five rounds of review and assessment against the Strategy's objectives prescribed in the Air Quality (England) Regulations 2000²⁰, Air Quality (Scotland) Regulations 2000²¹, Air Quality (Wales) Regulations 2000²² and Air Quality (Northern Ireland) Regulations 2003²³, together with subsequent amendments^{24,25,26,27}. The sixth round began in 2015.

The Review and Assessment process was streamlined in England following a consultation in 2015, and in 2016 a new format, the Assessment Summary Review, was adopted. Scotland published new Local Air Quality Management Policy Guidance in 2016 (<u>http://www.gov.scot/Publications/2016/03/9717</u>). Wales undertook a consultation during the latter part of 2016 and the new Policy Guidance was published in June 2017 (downloadable from

<u>http://gov.wales/docs/desh/publications/170614-policy-guidance-cy.pdf</u> in Welsh and <u>http://gov.wales/topics/environmentcountryside/epq/airqualitypollution/airquality/guid</u> <u>ance/policy-guidance/?lang=en</u> in English).

When the Assessment Summary Review process identifies an exceedance of an Air Quality Strategy objective, the Local Authority must declare an 'Air Quality Management Area' (AQMA) and develop an Action Plan to tackle problems in the affected areas. Action Plans formally set out the measures the Local Authority proposes to take to work towards meeting the air quality objectives. They may include a variety of measures such as congestion charging, traffic management, planning and financial incentives. Advice for Local Authorities preparing an Action Plan is available from the Defra LAQM web pages at http://laqm.defra.gov.uk/action-planning/aqap-supporting-guidance.html.

Information on the UK's AQMAs is summarised in **Table 2-1** below. At present, 278 Local Authorities –71% of those in the UK – have one or more AQMAs. Some AQMAs are for more than one pollutant, and many Local Authorities have more than one AQMA.

Most Air Quality Management Areas in the UK are in urban areas and have been established to address the contribution to air pollution from traffic emissions of nitrogen dioxide or PM₁₀, or in some cases both. A small number are for SO₂. There are no longer any AQMAs for benzene.

Region	Total LAs	LAs with AQMAs	AQMAs for NO ₂	AQMAs for PM ₁₀	AQMAs for SO ₂
England (outside London)	293	209	508	44	6
London	33	33	33	29	0
Scotland	32	15	27	21	1
Wales	22	11	39	1	0
Northern Ireland	11	10	20	6	0
TOTAL	391	278	627	101	7

Table 2-1 Current UK-wide status of Air Quality Management Areas (AQMAs) and Action Plans (as of July 2017.)

Figure 2-1, **Figure 2-2** and **Figure 2-3** are pie charts illustrating the proportion of AQMAs declared as a result of various different emission sources, for the three pollutants NO₂, PM₁₀ and SO₂.

Road transport is specified as the main source in 96% of the AQMAs declared for NO₂. A further 2.7% of NO₂ AQMAs result from road transport mixed with industrial sources, 0.5% from a combination of road transport, industry and domestic sources, 0.2% from industrial sources alone, and the remaining 0.6% from other or unspecified sources.

Road transport is also the main source in the majority (76%) of AQMAs declared for PM_{10} , but with industry and domestic sources accounting for a larger proportion than is the case for NO₂.

Most of the seven AQMAs declared for SO₂ relate to industrial, domestic, or other non-traffic sources.

The locations of the UK's AQMAs are shown in **Figure 2-4**. Information on the UK's Air Quality Management Areas is published on the Defra LAQM web pages (link above). Information is provided on each AQMA, together with a map of the area, where available.





Figure 2-2 Proportions of AQMAs Resulting from Various Sources: PM₁₀







Sulphur Dioxide



Figure 2-4 Air Quality Management Areas in the UK, as of the end of 2016

3 The Evidence Base

A programme of air quality assessment and research is in place in the UK which delivers the evidential needs of Defra, associated with compliance with a range of European Directives, as well as the means to assess the effectiveness of air pollution mitigation policies.

This section explains Defra and the Devolved Administrations' evidence base for the annual assessment of compliance with the EU Directives on ambient air quality. It describes the air pollutants which are of concern and how these are monitored and modelled in the UK.

3.1 Pollutants of Concern

Table 3-1 below summarises the sources, effects and typical UK concentrations of the pollutants being assessed in relation to the Air Quality Directive and 4th Daughter Directive.

The information on sources has largely been summarised from the National Atmospheric Emission Inventory (NAEI) pollutant information pages²⁸ together with Table 1 of the Air Quality Strategy²⁹.

Information on health effects is summarised (and further information can be sought) from the following sources:

- The World Health Organization's Air Quality Guidelines Global Update (2005)³⁰ (which covers particulate matter, sulphur dioxide, nitrogen dioxide and ozone).
- The World Health Organization's '*Air Quality and Health*' factsheet (factsheet 313) at http://www.who.int/mediacentre/factsheets/fs313/en/index.html.
- Committee on the Medical Effects of Air Pollution COMEAP's "Statement on the Evidence for the Effects of Nitrogen Dioxide on Health" (COMEAP 2015)³¹ (referred to in the table as COMEAP 2015a).
- Reports by the Committee on the Medical Effects of Air Pollution (COMEAP):
 - \circ COMEAP's 2011 review of the air quality index³²,
 - COMEAP's 2009 report on long-term exposure to air pollution and its effect on mortality³³ (referred to in the table below as COMEAP 2009),
 - COMEAP's 2010 report on the mortality effects of long-term exposure to particulate air pollution in the United Kingdom³⁴ (referred to in the table as COMEAP 2010),

- COMEAP's 2015 report on quantification of effects associated with ozone³⁵ (referred to in the table as COMEAP 2015b)
- Expert Panel on Air Quality Standards (EPAQS) report 'Metals and Metalloids³⁶ (referred to as EPAQS 2009 in the table below).
- Public Health England's Compendium of Chemical Hazards web pages at https://www.gov.uk/government/collections/chemical-hazards-compendium
- World Health Organization's 2013 'Review of Evidence on Health Aspects of Air Pollution' (REVIHAAP) report³⁷.
- The Air Quality Strategy.

Information on typical ambient concentrations in the UK has been summarised from the Defra online air information resource, UK-AIR at <u>http://uk-air.defra.gov.uk/</u>, and a 2011 study by King's College London, the University of Leeds and AEA (now Ricardo Energy & Environment), which investigated the reasons why ambient concentrations of NO_x and NO₂ have decreased less than predicted on the basis of emissions estimates³⁸.

Table 3-1 Sources, Effects and Typical UK Concentrations

Pollutant and Sources	Health Effects	Environmental Effects	Typical Ambient Concentrations in the UK
Sulphur Dioxide (SO ₂): an acid gas formed when fuels containing sulphur impurities are burned. The largest UK source is currently power generation. Other important sources include industry, commercial fuel use, and residential fuel use in some areas.	A respiratory irritant that can cause constriction of the airways. People with asthma are considered to be particularly sensitive. Health effects can occur very rapidly, making short-term exposure to peak concentrations important. (Source: WHO AQG 2005)	Harmful to plants at high concentrations. Contributes to acidification of terrestrial and aquatic ecosystems, damaging habitats and leading to biodiversity loss. SO ₂ is also a precursor to the formation of secondary sulphate particles in the atmosphere.	Annual mean concentrations are typically less than 5 µg m ⁻³ except at sites in industrial locations or in residential areas with high use of solid fuel for heating.
Nitrogen Oxides (NOx) : NOx, which comprises nitric oxide (NO) and nitrogen dioxide (NO ₂), is emitted from combustion processes. Main sources include power generation, industrial combustion and road transport. According to the NAEI, road transport is now the largest single UK source of NOx, accounting for almost one third of UK emissions.	Short-term exposure to concentrations of NO ₂ higher than 200 µg m ⁻³ can cause inflammation of the airways. NO ₂ can also increase susceptibility to respiratory infections and to allergens. It has been difficult to identify the direct health effects of NO ₂ at ambient concentrations because it is emitted from the same sources as other pollutants such as particulate matter (PM). Studies have found that both day-to- day variations and long-term exposure to NO ₂ are associated with mortality and morbidity. Evidence from studies that have corrected for	In the presence of sunlight, nitrogen oxides can react with Volatile Organic Compounds to produce photochemical pollutants including ozone. NO _x contributes to the formation of secondary nitrate particles in the atmosphere. High levels of NO _x can harm plants. NO _x also contributes to acidification and eutrophication of terrestrial and aquatic ecosystems,	Annual mean concentrations of NO ₂ beside busy roads frequently exceed 40 µg m ⁻³ . This is not a UK-specific problem and is common in many other European countries. The main reasons why roadside NO ₂ concentrations have not decreased as expected is believed to be the failure of Euro vehicle emission standards for diesel vehicles to deliver the anticipated reductions in NO _x emissions in real world driving

Pollutant and Sources	Health Effects	Environmental Effects	Typical Ambient Concentrations in the UK
	the effects of PM is suggestive of a causal relationship, particularly for respiratory outcomes (Source: WHO 2013 REVIHAAP report, COMEAP 2015).	damaging habitats and leading to biodiversity loss.	conditions. At urban background locations, annual mean NO_2 concentrations are lower, typically 15-40 µg m ⁻³ . Peak hourly mean concentrations exceed 100 µg m ⁻³ at most urban locations, and occasionally exceed 300 µg m ⁻³ at congested urban roadside sites.
Particulate Matter: PM₁₀. This can be primary (emitted directly to the atmosphere) or secondary (formed by the chemical reaction of other pollutants in the air such as SO ₂ or NO ₂). The main source is combustion, e.g. vehicles, domestic burning and power stations. Other man-made sources include quarrying and mining, industrial processes and tyre and brake wear. Natural sources include wind- blown dust, sea salt, pollens and soil particles.	Research shows a range of health effects (including respiratory and cardiovascular illness and mortality) associated with PM ₁₀ . No threshold has been identified below which no adverse health effects occur. (Source: WHO AQG 2000)	Black carbon in PM is implicated in climate change. Secondary PM includes sulphate, nitrate and ammonium, formed from SO ₂ , NO _x and NH ₃ which are the main drivers for acidification and eutrophication.	Annual mean PM ₁₀ concentrations for urban AURN monitoring sites have been typically in the range 10-30 µg m ⁻³ in recent years.
Particulate Matter: PM_{2.5}. Like PM ₁₀ , the finer size fraction PM _{2.5} can be	Fine particulate matter can penetrate deep into the lungs and research in recent years	Secondary PM includes sulphate, nitrate and ammonium, formed from	Annual mean urban PM _{2.5} concentrations in the UK are

Pollutant and Sources	Health Effects	Environmental Effects	Typical Ambient Concentrations in the UK
primary or secondary, and has the same sources. Road transport becomes an increasingly important sector as the particle size decreases.	has strengthened the evidence that both short- term and long-term exposure to PM _{2.5} are linked with a range of health outcomes including (but not restricted to) respiratory and cardiovascular effects. COMEAP estimated that the burden of anthropogenic particulate air pollution in the UK in 2008 was an effect on mortality equivalent to nearly 29,000 deaths at typical ages and an associated loss of life across the population of 340,000 years. The burden can also be represented as a loss of life expectancy from birth of approximately six months. (Source: COMEAP 2010.)	SO ₂ , NO _x and NH ₃ which are the main drivers for acidification and eutrophication.	typically in the low teens of μg m ⁻³ but exceed 20 μg m ⁻³ at a few urban roadside locations.
Benzene: (C ₆ H ₆) is an organic chemical compound. Ambient benzene concentrations arise from domestic and industrial combustion processes, in addition to road transport. (Source: Air Quality Strategy).	Benzene is a recognised human carcinogen which causes changes in the genetic material (mutagenic effect) of the circulatory and immune systems. No absolutely safe level can be specified in ambient air. Acute exposure to high concentrations affects the central nervous system. (Source: WHO AQG 2000, PHE Compendium of Chemical Hazards)	Can also pollute soil and water, leading to exposure via these routes.	Annual mean concentrations of benzene are now low (consistently below 2 µg m ⁻³) due to the introduction of catalytic converters on car exhausts. The UK meets the benzene limit value of 5 µg m ⁻³ .

Pollutant and Sources	Health Effects	Environmental Effects	Typical Ambient Concentrations in the UK
Carbon Monoxide (CO) is produced when fuels containing carbon are burned with insufficient oxygen to convert all carbon inputs to carbon dioxide (CO ₂). Although CO emissions from petrol-engine road vehicles have been greatly reduced by the introduction of catalytic converters, road transport is still the most significant source of this pollutant (Source: NAEI).	CO affects the ability of the blood to take up oxygen from the lungs, and can lead to a range of symptoms. People are more likely to be exposed to dangerous concentrations of CO indoors, due to faulty or poorly ventilated cooking and heating appliances. Cigarette smoke is also a major source of exposure. (Source: NAEI, PHE Compendium of Chemical Hazards.)	Can contribute to the formation of ground-level ozone.	The UK is compliant with the European limit value for CO, with the 8-hour running mean concentration consistently below 10 mg m ⁻³ at all monitoring sites in recent years.
Ozone (O₃) is a secondary pollutant produced by the effect of sunlight on NO _x and VOCs from vehicles and industry. Ozone concentrations are greatest in the summer on hot, sunny, windless days. O ₃ can travel long distances, accumulate and reach high concentrations far away from the original sources.	A respiratory irritant: short-term exposure to high ambient concentrations can cause inflammation of the respiratory tract and irritation of the eyes, nose, and throat. High levels may exacerbate asthma or trigger asthma attacks in susceptible people and some non-asthmatic individuals may also experience chest discomfort whilst breathing. Evidence is also emerging of effects due to long-term exposure (WHO AQG 2000, WHO 2013 - REVIHAAP).	Ground level ozone can also cause damage to many plant species leading to loss of yield and quality of crops, damage to forests and impacts on biodiversity. Ozone is also a greenhouse gas implicated in climate change. In the upper atmosphere the ozone layer has a beneficial effect, absorbing harmful ultraviolet radiation from the sun.	In recent years, the annual mean daily maximum 8-hour running mean measured at AURN sites has been typically in the range 30-80 µg m ⁻³ . NO _X emitted in cities reduces local O ₃ concentrations as NO reacts with O ₃ to form NO ₂ and levels of O ₃ are often higher in rural areas than urban areas.

Pollutant and Sources	Health Effects	Environmental Effects	Typical Ambient Concentrations in the UK
Lead (Pb): a very toxic metallic element. Historically, lead was used as an additive in petrol, and road vehicles were the main source Lead's use in petrol was phased out in 1999, resulting in a 98% reduction of pre- 1999 UK emissions. Today, the main sources are metal production and industrial combustion of lubricants containing small amounts of lead. (Source: NAEI.)	Lead inhalation can affect red blood cell formation and have effects on the kidneys, circulatory system, gastrointestinal tract, the joints, reproductive systems, and can cause acute or chronic damage to the central nervous system (CNS). Long term low level exposure has been shown to affect intellectual development in young children (Source: EPAQS 2009). A threshold, below which the adverse effects of lead are not anticipated, has not been established (source: WHO AQG 2000, PHE Compendium of Chemical Hazards).	Can also pollute soil and surface waters. Exposure to contaminated soil and water may then become a health risk. Lead may accumulate in other organisms such as fish, and be passed up the food chain.	In recent years, UK annual mean concentrations of lead have typically ranged from less than 5 ng m ⁻³ at rural monitoring sites, to nearly 90 ng m ⁻³ at urban industrial sites. The EU limit value for Pb (0.5 μ g m ⁻³ or 500 ng m ⁻³) is met throughout the UK.
Nickel (Ni) is a toxic metallic element found in ambient air as a result of releases from oil and coal combustion, metal processes, manufacturing and other sources. Currently the main source is the combustion of heavy fuel oil, the use of coal having declined. (Source: NAEI.)	Nickel compounds are human carcinogens by inhalation exposure. Can cause irritation to the nose and sinuses and allergic responses and can lead to the loss of the sense of smell. Long-term exposure may lead to respiratory diseases and cancers. (Source: WHO AQG 2000, EPAQS 2009, PHE Compendium of Chemical Hazards.)	Can also pollute soil and water, leading to exposure via these routes.	Annual mean ambient particulate phase concentrations in the urban environment are typically of the order of 1 ng m ⁻³ with the exception of a few industrial areas, where higher annual means may occur, in some locations exceeding the 4 th

Pollutant and Sources	Health Effects	Environmental Effects	Typical Ambient Concentrations in the UK
			Daughter Directive target value of 20 ng m ⁻³ .
Arsenic (As) is a toxic element emitted into the atmosphere in the form of particulate matter. Historically the largest source was coal combustion, but as this has declined, the use of wood treated with preservatives containing As has become the most significant component of As emissions. (Source: NAEI.)	Acute inhalation exposure to high levels of arsenic primarily affects the respiratory system and can cause coughs, sore throat, breathlessness and wheezing. Long term inhalation exposure is associated with toxic effects on the respiratory tract and can cause lung cancer. (Source: WHO AQG 2000, EPAQS 2009, PHE Compendium of Chemical Hazards.)	Can also pollute soil and water, leading to exposure via these routes. Arsenic in water or soil can be taken up by plants or fish.	Measured UK annual mean concentrations in the particulate phase are now typically less than 1 ng m ⁻³ , meeting the 4 th Daughter Directive target value of 6 ng m ⁻³ .
Cadmium (Cd): a toxic metallic element whose main sources are energy production, non-ferrous metal production, iron and steel manufacture (as well as other forms of industrial combustion). (Source: NAEI.)	Acute inhalation exposure to cadmium causes effects on the lung such as pulmonary irritation. Chronic effects via inhalation can cause a build-up of cadmium in the kidneys that can lead to kidney disease and long term inhalation can lead to lung cancer. (Source: WHO AQG 2000, EPAQS 2009, PHE Compendium of Chemical Hazards.)	Can also pollute soil and water, leading to exposure via these routes.	Annual mean particulate phase concentrations in the UK in recent years are now typically < 2 ng m ⁻³ , and meet the 4 th Daughter Directive target value of 5 ng m ⁻³ .
Mercury (Hg): released to the air by human activities, such as fossil fuel combustion, iron and steel production	Acute exposure to high levels of Hg can cause chest pain and shortness of breath, and affect the central nervous system (CNS) and	Can also pollute soil, fresh water and sea water. Exposure to contaminated soil and water may then become a	There is no target value for mercury. Annual mean ambient concentrations (total of vapour

Pollutant and Sources	Health Effects	Environmental Effects	Typical Ambient Concentrations in the UK
processes, waste incineration, the manufacture of chlorine in mercury cells, and coal combustion. Emissions have declined in recent years as a result of improved controls on mercury cells, the reduction in coal use, and improved controls on waste incineration processes from 1997 onwards. (Source: NAEI.)	kidneys. Chronic exposure leads to CNS disorders, kidney damage and stomach upsets. (Source: WHO AQG 2000, PHE Compendium of Chemical Hazards.)	health risk. Mercury may accumulate in other organisms such as fish, and be passed up the food chain.	and particulate phases) are typically in the range 1-3 ng m ⁻³ , although higher concentrations (over 20 ng m ⁻³) have been measured at industrial sites in recent years.
Benzo[a]pyrene (B[a]P) is used as a 'marker' for a group of compounds known as polycyclic aromatic hydrocarbons (PAHs). The main sources of B[a]P in the UK are domestic coal and wood burning, fires (e.g. accidental fires, bonfires, forest fires, etc.), and industrial processes such as coke production. (Source: Air Quality Strategy).	PAHs are a large group of persistent, bio- accumulative, organic compounds with toxic and carcinogenic effects. Lung cancer is most obviously linked to exposure to PAHs through inhaled air. (Source: WHO AQG 2000, PHE Compendium of Chemical Hazards)	PAHs can bio-accumulate and be passed up the food chain.	Annual mean concentrations in most urban areas are below the EU target value of 1 ng m ⁻³ : the only exceptions are areas with specific local sources – such as industrial installations or domestic solid fuel burning.
3.2 Assessment of Air Quality in the UK

The evidence base for the annual assessment of compliance is based on a combination of information from the UK national monitoring networks and the results of modelling assessments. The use of models reduces the number of monitoring stations required. It has the added benefits of enabling air quality to be assessed at locations without monitoring sites and providing additional information on source apportionment and projections required for the development and implementation of air quality plans.

UK compliance assessment modelling is undertaken using national models known as the Pollution Climate Mapping (PCM) models. The PCM models have been designed to assess compliance with the limit values at locations defined within the Directives. Modelled compliance assessments are undertaken for 11 air pollutants each year. This assessment needs to be completed each year in the relatively short period between the time when the input data (including ratified monitoring data and emission inventories) become available and the reporting deadline at the end of September.

It is important to understand the differences between modelling carried out for compliance assessment purposes, and that carried out for Local Air Quality Management. National air quality modelling for the UK focuses on two components: pollutant concentrations at background locations, on a 1x1km grid square basis, and roadside pollutant concentrations, at four metres from the kerb of urban major road links. By contrast, Local Air Quality Management (LAQM) modelling is different in scope, purpose and methodology from the national modelling and will usually output contour plots showing dispersion away from the source, on a fine resolution grid. The level of detail and resolution of LAQM modelling is therefore much greater in order to focus on local exposure and hotspots. See Section 3.5 for more details on the modelling carried out for compliance assessment.

3.3 Current UK Air Quality Monitoring

During 2016 there were 284 national air quality monitoring sites across the UK, comprising several networks, each with different objectives, scope and coverage and these are operated on behalf of Defra by the Environment Agency (EA). This section provides a brief description of those used to monitor compliance with the Air Quality Directive and the 4th Daughter Directive. A summary of the UK national networks is provided in **Table 3-2** (the numbers of sites shown in this table add up to considerably more than 284 because some sites belong to more than one network). This table shows the numbers of sites in operation during part or all of 2016.

Network	Pollutants	Number of Sites operating in 2016
Automatic Urban and Rural Network (AURN)	CO, NO _x , NO ₂ , SO ₂ , O ₃ , PM ₁₀ , PM _{2.5} .	161
UK Metals Network	Metals in PM ₁₀ Including: As, Cd, Co, Cr, Cu, Fe, Mn, Ni, Pb, Se, V, Zn.	24
	Measured deposition. Including: Al, As, Ba, Be, Cd, Co, Cr, Cs, Cu, Fe, Li, Mn, Mo, Ni, Pb, Rb, Sb, Sc, Se, Sn, Sr, Ti, U, V, W, Zn.	6
	Hg deposition	4
	Total gaseous mercury	2
Non-Automatic Hydrocarbon	Benzene	35
Automatic Hydrocarbon	Range of volatile organic compounds (VOCs)	4
Polycyclic Aromatic Hydrocarbons (PAH).	21 PAH species including benzo[a]pyrene	30
European Monitoring and Evaluation Programme (EMEP)	Wide range of parameters relating to air quality, precipitation, meteorology and composition of aerosol in PM ₁₀ and PM _{2.5} .	2
Particle Concentrations and Numbers	Total particle number, concentration, size distribution, anions, EC/OC, speciation of PM ₁₀ , PM _{2.5} and PM ₁ .	4
Toxic Organic Micropollutants	Range of toxic organics including dioxins and dibenzofurans.	6
UK Eutrophying and Acidifying Pollutants: NO ₂ Net (rural diffusion tubes)	NO ₂ (rural)	25
UK Eutrophying and Acidifying Pollutants: AGANet	NO3, HCI, HNO3, HONO, SO2, SO4	30
UK Eutrophying and Acidifying Pollutants: NAMN	NH ₃ and/or NH ₄	93
UK Eutrophying and Acidifying Pollutants: PrecipNet	Major ions in rain water	47
Black Carbon	Black Carbon	15
Upland Waters Monitoring Network <i>(not funded by Defra)</i>	Chemical and biological species in water	26
Rural Mercury Network	Tekran analyser used to measure mercury in PM _{2.5} , reactive mercury and elemental mercury at Auchencorth Moss, and total gaseous mercury at Chilbolton Observatory.	2

3.3.1 The Automatic Urban and Rural Network (AURN)

The AURN is currently the largest automatic monitoring network in the UK and forms the bulk of the UK's statutory compliance monitoring evidence base. Data from the AURN are available on Defra's online UK Air Information Resource, UK-AIR at <u>http://uk-air.defra.gov.uk/</u>.

The techniques used for monitoring gaseous pollutants within the AURN are the reference measurement methods defined in the relevant EU Directives. For particulate matter the AURN uses methods which have demonstrated equivalence to the reference method, but which (unlike the reference method) allow continuous on-line monitoring. Details are provided in **Table 3-3**.

Pollutant	Method used, including details of CEN Standard Methods
O ₃	EN 14625:2012 'Ambient air quality – standard method for the measurement of the concentration of ozone by ultraviolet photometry' ³⁹
NO ₂ /NO _x	EN 14211:2012 'Ambient air quality - Standard method for the measurement of the concentration of nitrogen dioxide and nitrogen monoxide by chemiluminescence' ⁴⁰
SO ₂	EN 14212:2012 'Ambient air quality – Standard method for the measurement of the concentration of sulphur dioxide by UV fluorescence' ⁴¹
CO	EN 14626:2012 'Ambient air quality - Standard method for the measurement of the concentration of carbon monoxide by non-dispersive infrared spectroscopy' ⁴²
PM ₁₀ and PM _{2.5}	EN 12341:2014 'Ambient air quality - Standard gravimetric measurement method for the determination of the PM_{10} or $PM_{2.5}$ mass fraction of suspended particulate matter' ⁴³
	The AURN uses three methods which are equivalent for one or both pollutants: the Filter Dynamic Measurement System (FDMS), which determines particulate concentration by continuously weighing particles deposited on a filter: the Beta-Attenuation Monitor (BAM) which measures the attenuation of beta radiation passing through a paper filter on which particulate matter from sampled air has been collected, and the Partisol – a gravimetric sampler that collects daily samples onto a filter for subsequent weighing.

Table 3-3 AURN Measurement Techniques

3.3.2 The UK Metals Network

The UK Metals Network forms the basis of the UK's compliance monitoring for:

- The Air Quality Directive (for lead).
- The 4th Daughter Directive (for arsenic, cadmium, nickel and mercury).

At the end of 2013 Defra merged the existing Urban and Industrial Network with the Rural Network to form the UK Metals Network. The merged network monitors a

range of metallic elements at urban, industrial and rural sites, using a method equivalent to the CEN standard method⁴⁴. Metals (As, Cd, Co, Cr, Cu, Fe, Mn, Ni, Pb, Se, V and Zn) in PM₁₀ are measured at 24 sites. (The network stopped measuring mercury in PM₁₀ as of 2014.)

Metal deposition (Al, As, Ba, Be, Cd, Co, Cr, Cs, Cu, Fe, Li, Mn, Mo, Ni, Pb, Rb, Sb, Sc, Se, Sn, Sr, Ti, U, V, W and Zn) was measured at the following sites: Auchencorth Moss, Chilbolton Observatory, Heigham Holmes, Lough Navar and Yarner Wood. Rain water sampling for metals at Harwell stopped on 6th January 2016).

Hg deposition was measured at Auchencorth Moss, Chilbolton Observatory, Heigham Holmes and Yarner Wood.

Within the Metals Network total gaseous mercury is measured at London Westminster and Runcorn Weston Point.

3.3.3 Non-Automatic Hydrocarbon Network

In this network, ambient concentrations of benzene are measured by the CEN standard method⁴⁵. This involves pumping air through an adsorption tube to trap the compound, which is later analysed in a laboratory. This network monitors compliance with the Air Quality Directive's limit value for benzene. All sites in the Non-Automatic Hydrocarbon Network are co-located with AURN sites. In September 2016, the Birmingham Tyburn Roadside site was replaced by Birmingham A4540 Roadside.

3.3.4 Automatic Hydrocarbon Network

The Air Quality Directive also requires measurement and reporting of ozone precursor substances (29 species), which include volatile organic compounds (VOCs). Annex X (ten) of the Directive provides a list of compounds recommended for measurement. Ozone precursor measurement is carried out by the Automatic Hydrocarbon Network.

Automatic hourly measurements of a range of hydrocarbon species (including all those specified in Annex X of the Directive except formaldehyde and total non-methane hydrocarbons), are made at four sites using automated pumped sampling with *in-situ* gas chromatography. The VOCs monitored include benzene, which is covered by the Air Quality Directive as a pollutant in its own right.

3.3.5 PAH Network

The PAH Network monitors compliance with the 4th Daughter Directive, which includes a target value of 1 ng m⁻³ for the annual mean concentration of benzo[a]pyrene as a representative PAH, not to be exceeded after 31st December

2012. Samples are collected on filters using the PM_{10} 'Digitel' sampler. Samples are subsequently analysed in a laboratory. During 2016, there were 30 sites in this network measuring 23 PAH compounds.

3.3.6 EMEP

EMEP (European Monitoring and Evaluation Programme) is a programme set up to provide governments with qualified scientific information on air pollutants, under the UNECE Convention on Long-range Transboundary Air Pollution. There are currently two EMEP 'supersites', at Auchencorth Moss in Lothian (representing the north of the UK) and at Chilbolton Observatory in Hampshire (representing the south). The site at Chilbolton replaced the long running site at Harwell at the start of 2016. A representativeness analyses showed that both sites were similar in their rural background nature. A very wide range of measurements are taken at EMEP sites, supplemented by data from other UK networks which are co-located.

Monitoring includes:

- Hourly meteorological data,
- Soil and vegetation measurements,
- Metallic elements in PM₁₀ and precipitation,
- Deposition of inorganic ions,
- Major ions in PM_{2.5} and PM₁₀, as well as HCI, HNO₂, HNO₃, NH₃ and SO₂,
- Trace gases (ozone, NO_x and SO₂),
- Black carbon, organic carbon (OC) and elemental carbon (EC),
- Ammonia (monthly),
- Daily and hourly PM10 and PM2.5 mass,
- Volatile Organic Compounds,
- Carbonyls,
- CH₄ and N₂O fluxes.

3.3.7 Particle Concentrations and Numbers

The Air Quality Directive requires that the chemical composition of PM_{2.5} is characterised at background locations in the United Kingdom. The Particle Concentrations and Numbers Network contribute to this statutory requirement. During 2016, the network consisted of four measurement sites; two rural sites (Auchencorth Moss and Chilbolton Observatory), and two in London (London Marylebone Road and London North Kensington).

Among the parameters measured are:

- Total particle numbers per cubic centimetre of ambient air,
- Particle numbers in different particle size fractions,

- Major ions in PM₁₀,
- CI, NH4, NO3, OC, SO4 and organic carbon (OC) in PM1,
- Organic carbon (OC) and elemental carbon (EC) concentrations in PM_{2.5}.

As well as its statutory function, this network provides data on the chemical composition of particulate matter, primarily for the use of researchers of atmospheric processes, epidemiology and toxicology.

Measurements of elemental carbon (EC) and organic carbon (OC) began at Auchencorth Moss at the start of 2011 and Chilbolton Observatory at the start of 2016. EC and OC measurements were also made using a thermal/optical method involving both reflectance and transmission correction methods. Comparing both correction methods aims to provide valuable understanding of the measurement process for EC and OC.

3.3.8 TOMPs Network

This research-based network monitors a range of toxic organic micropollutants (compounds that are present in the environment at very low concentrations, but are highly toxic and persistent). These include dioxins, dibenzofurans and polychlorinated biphenyls. The TOMPs Network consists of six sites: Auchencorth Moss, Hazelrigg, High Muffles, London Nobel House, Manchester Law Courts and Weybourne.

The purpose of the TOMPs Network is to provide data on these air pollutants, and to support the development of policy to protect the environment and human health. Further information on the TOMPs Network can be found on UK-AIR at http://uk-air.defra.gov.uk/networks/network-info?view=tomps. However, this network is not used for compliance monitoring and will not be discussed further in subsequent sections of the report.

3.3.9 UK Eutrophying and Acidifying Pollutants Network

The UK Eutrophying and Acidifying Atmospheric Pollutants (UKEAP) network provides information on deposition of eutrophying and acidifying compounds in the UK and assessment of their potential impacts on ecosystems. The UKEAP network is an 'umbrella' project covering four groups of sites:

- The UKEAP rural NO₂ diffusion tube network (NO₂Net). This measures NO₂ concentrations as required as input to the rural NOx concentration field in the Pollution Climate Model.
- In 2016, the Acid Gas and Aerosol Network (AGANet) comprised a total of 30 sites, though sampling stopped at Barcombe Mills, Edinburgh St Leonards and Shetland in October 2016. The network measures a range of gases and aerosol components. Samples are collected monthly and are

analysed by either inductive coupled plasma optical emission spectrometry (ICP-OES) or ion chromatography.

- The National Ammonia Monitoring Network (NAMN) which characterizes ammonia and ammonium concentrations using both passive samplers (Alpha Samplers) and low volume denuders (Delta Samplers) at more than 90 locations in 2016. Monitoring also began at six sites (Ainsdale Dunes and Sands, Burnham Beeches, Ingleborough, May Moss, Monks Wood, Stiperstones) that had formed part of Natural England's Long Term Monitoring Network (LTMN).
- The Precipitation Network (PrecipNet) underwent a reconfiguration in 2016 with sampling stopping at the following sites: Barcombe Mills, Bottesford, Harwell, Llyn Llagi, Loch Chon, Lochnagar and Scoat Tarn. Sampling started at the former LTMN sites: Ainsdale Dunes and Sands, Bure Marshes, Fenn's, Whixall and Bettisfield Mosses, Ingleborough, Lullington Heath, Stiperstones and Thursley Common 2.
- The network allows estimates of sulphur and nitrogen deposition. Samples are collected fortnightly at all sites and daily at two sites.

3.3.10 Black Carbon Network

Black carbon is fine, dark carbonaceous particulate matter produced from the incomplete combustion of materials containing carbon (such as coal, oil, and biomass such as wood). It is of concern due to health effects, and also as a suspected contributor to climate change. In 2016, the Black Carbon Network measured black carbon at 14 sites using an automatic instrument called an Aethalometer[™]. During the year Birmingham Tyburn Roadside was replaced by Birmingham A4540 Roadside. The Aethalometer[™] measures black carbon directly, using a real-time optical transmission technique. The objectives of the network are as follows:

- To maintain coverage of black carbon measurements across the whole UK;
- To maintain continuity of historic datasets;
- To gather data for epidemiological studies of black carbon and health effects;
- To gather information about black carbon PM sources in the UK;
- To assess PM reductions from air quality management interventions;
- To quantify the contribution of wood burning to black carbon and ambient PM in the UK; and
- To gather data to address future policy considerations including black carbon and climate change.

3.3.11 UK Upland Waters Monitoring Network (UKUWMN)

The UK Upland Waters Monitoring Network (UWMN) was set up in 1988 (then called the Acid Waters Network) to assess the chemical and biological response of acidified lakes and streams in the UK to the planned reduction in emissions. It provides chemical and biological data on the extent and degree of surface water acidification in the UK uplands, in particular to underpin the science linking acid deposition to water quality and to monitor the response of aquatic ecosystems to reductions in air pollution. The sites making up the network were selected on the basis of acid deposition inputs being the only major sources of pollution, i.e. with no point sources of pollution or direct catchment disturbances other than traditional upland land use practices such as sheep grazing or forestry. There are 26 primary monitoring sites including 12 lakes and 14 streams across the UK, and a series of secondary sites, monitoring a range of parameters including water chemistry, algae and higher aquatic plants, invertebrates, fish, water temperature and sediment biology and chemistry.

3.3.12 Rural Mercury Monitoring

During 2016, in addition to the weekly monitoring of total gaseous mercury carried out at London Westminster and Runcorn Weston Point, and the monthly mercury deposition measurements at Auchencorth Moss, Harwell, Heigham Holmes and Yarner Wood as described in Section 3.3.2, speciated mercury monitoring is carried out using the Tekran automatic instrument. The Tekran instrument at Auchencorth Moss measures the mercury composition of PM_{2.5} as well as mercury in its elemental and reactive forms, whereas at Chilbolton Observatory it measured just total gaseous mercury. The Harwell monitoring station closed at the end of 2015 and was relocated to Chilbolton Observatory at the start of 2016.

3.4 Quality Assurance and Quality Control

Air quality monitoring in the UK is subject to rigorous procedures of validation and ratification. The well-established monitoring networks each have a robust and documented Quality Assurance and Quality Control (QA/QC) programme designed to ensure that measurements meet the defined standards of quality with a stated level of confidence. Essentially, each programme serves to ensure that the data obtained are:

- Representative of ambient concentrations existing in the various areas under investigation.
- Sufficiently accurate and precise to meet specified monitoring objectives.
- Comparable and reproducible. Results must be internally consistent and comparable with international or other accepted standards, if these exist.

- Consistent over time. This is particularly important if long-term trend analysis of the data is to be undertaken.
- Representative over the period of measurement; for most purposes, a yearly data capture rate of not less than 90% is usually required for determining compliance with EU limit values where applicable. An allowance of 5% is made in some cases for planned maintenance.
- Consistent with Data Quality Objectives⁴⁶. The uncertainty requirements of the EU Directives are specified as data quality objectives. In the UK, all air quality data meet the data quality requirements of the EU Directives in relation to uncertainty.
- Consistent with methodology guidance defined in EU Directives for relevant pollutants and measurement techniques. The use of tested and approved analysers that conform to Standard Method (or equivalent) requirements and harmonised on-going QA/QC procedures allows a reliable and consistent quantification of the uncertainties associated with measurements of air pollution.

Most UK networks use a system of regular detailed audits of all monitoring equipment at every site. These audits supplement more regular calibrations and filter changes and test all critical parameters of the measuring equipment including, where appropriate, linearity, converter efficiency (in the case of NO_x analysers) response time, flow rate etc.

Data verification is the process of checking and validating the data. (The term 'ratification' is used in some networks). Data entered on the Defra Air Information Resource (UK-AIR at <u>http://uk-air.defra.gov.uk</u>) in near real time are provided as provisional data. All these data are then carefully screened and checked via the verification process. The verified data then overwrite the provisional data on the website. It should however be noted that there are occasionally circumstances where data which have been flagged as 'Verified' could be subject to further revision. This may be for example where:

- A QA/QC audit has detected a problem which affects data back into an earlier verification period.
- Long-term analysis has detected an anomaly between expected and measured trends which requires further investigation and possible data correction. This was the case with 2000-2008 particulate monitoring data in the UK national network.
- Further research comes to light which indicates that new or tighter QA/QC criteria are required to meet the data quality objectives. This may require review and revision of historical data by applying the new criteria.

Only verified data are provided to the Commission in compliance with EU Directives.

Further details on the QA/QC procedures appropriate to each network can be obtained from the annual reports of the monitoring networks, and (in the case of the AURN) from the report 'QA/QC Procedures for the UK Automatic Urban and Rural *Air Quality Monitoring Network (AURN)*'⁴⁷ available from Defra's air quality web pages.

3.5 Modelling

3.5.1 Why Do Modelling?

The UK's monitoring programmes are supplemented by air quality modelling. There are several benefits of using modelling to complement the monitoring data gathered across the UK national monitoring networks:

- A reduction in the number of fixed continuous monitoring locations required for compliance with European air quality Directives freeing up resources and ensuring value for money.
- Coverage of the whole UK rather than specific locations where there is a monitoring site. A monitoring site might not fully represent the wider region in which it is located due to local characteristics such as buildings affecting dispersion, localised or temporary sources.
- Providing a framework within which to assess different air quality scenarios – for example projecting concentrations forward to assess levels in future years, representing potential changes to emissions in order to assess the impact of policy initiatives on air quality.

3.5.2 How the Models Work

The national modelling methodology varies between pollutants. The detailed methodology is explained in a technical report⁴⁸ (the latest versions of these can be found in the Library section of Defra's UK-AIR website⁴⁹).

Defra's air quality national modelling assessment for the UK consists of two components:

- Background concentrations on a 1x1km resolution, representing ambient air quality concentrations at background locations.
- Roadside concentrations concentrations at the roadside of urban major road links throughout the UK (i.e. motorways and major A-roads). There are approximately 10,000 of these urban major road links.

Roadside concentrations are not modelled for CO, SO₂, ozone, benzo[a]pyrene and metals as these are deemed not to have significant traffic-related sources.

The models have been designed to assess compliance at locations defined by the Directives as relevant for air quality assessment.

3.5.3 Background Air Quality

The 1x1 km background maps are made up of several components which are modelled separately and then added together to make the final grid. These individual components (supplemented by some additional components for various pollutants) are:

- Large point sources (e.g. power stations, steel works and oil refineries),
- Small point sources (e.g. boilers in town halls, schools or hospitals and crematoria),
- Distant sources (characterised by the rural background concentration),
- Local area sources (e.g. road traffic, domestic and commercial combustion and agriculture).

In order to ensure that these ambient concentrations from area sources are representative of the real world situation, they are validated against measurements taken from the national networks (including the AURN). After the validation has been completed the large points, small points, distant sources and area source components are added together to provide the final background map.

3.5.4 Roadside Air Quality

Roadside concentrations are determined by using a roadside increment model which attempts to estimate the contribution from road traffic sources and adds this on top of the modelled background concentrations discussed above.

For each of the road links that are modelled, there are emission estimates from the National Atmospheric Emissions Inventory (NAEI)⁵⁰ for each pollutant and road traffic counts. A measured roadside increment concentration is calculated for road links with a roadside monitoring station on them by subtracting the link's modelled background concentration (from the 1x1 km modelled maps) from the relevant measured roadside increment concentration for each road by applying the NAEI emissions and road traffic counts (annual average daily traffic flow) in a dispersion model. The RKM is calibrated by comparing the measured roadside increment concentrations with the modelled roadside increment concentrations with the modelled roadside increment concentrations at roadside monitoring stations with the modelled roadside increment concentrations on these road links. The application of the roads kernel model ensures that a process-based modelling approach is used to determine the local component of roadside concentrations, including factors influencing dispersion at the roadside e.g. road orientation, width, and additional vehicle induced turbulence.

3.6 Access to Assessment Data

Data from the UK's air quality monitoring networks and annual compliance modelling is available under the Open Government Licence <u>http://www.nationalarchives.gov.uk/doc/open-government-licence/version/2/</u> from UK-AIR.

Defra has produced a searchable online catalogue of air quality and emissions datasets which allows people to browse the extent of data available and access key metadata. This is available at <u>http://uk-air.defra.gov.uk/data/data-catalogue</u>.

Historical monitoring data can be accessed through the data selector tools in UK-AIR, at <u>http://uk-air.defra.gov.uk/data/</u>. Modelled data from the Pollution Climate Mapping model are available as .csv files for download from the modelled air quality data pages at <u>http://uk-air.defra.gov.uk/data/modelling-data</u> or can be accessed through the Ambient Air Quality Interactive Map at <u>http://uk-air.defra.gov.uk/data/gismapping</u> - a GIS tool which provides enhanced visualisation capability and access to roadside concentration data.

4 Assessment of Compliance

4.1 Definition of Zones

The UK is divided into 43 zones for air quality assessment. There are 28 agglomeration zones (large urban areas) and 15 non-agglomeration zones. Each zone has an identification code (**Table 4-1**). Zones are shown in **Figure 4-1**.

Zana zada Zana tura	
Zone code Zone type	
West Midlanda Urban Area UK0001 Aggiomeration	
West Midiands Orban Area OK0002 Aggiomeration	
Greater Manchester Orban Area UK0003 Aggiomeration	
West Yorkshire Urban Area UK0004 Aggiomeration	
Tyneside UK0005 Agglomeration	
Liverpool Urban Area UK0006 Agglomeration	
Sheffield Urban Area UK0007 Agglomeration	
Nottingham Urban Area UK0008 Agglomeration	
Bristol Urban Area UK0009 Agglomeration	
Brighton/Worthing/Littlehampton UK0010 Agglomeration	
Leicester Urban Area UK0011 Agglomeration	
Portsmouth Urban Area UK0012 Agglomeration	
Teesside Urban Area UK0013 Agglomeration	
The Potteries UK0014 Agglomeration	
Bournemouth Urban Area UK0015 Agglomeration	
Reading/Wokingham Urban Area UK0016 Agglomeration	
Coventry/Bedworth UK0017 Agglomeration	
Kingston upon Hull UK0018 Agglomeration	
Southampton Urban Area UK0019 Agglomeration	
Birkenhead Urban Area UK0020 Agglomeration	
Southend Urban Area UK0021 Agglomeration	
Blackpool Urban Area UK0022 Agglomeration	
Preston Urban Area UK0023 Agglomeration	
Glasgow Urban Area UK0024 Agglomeration	
Edinburgh Urban Area UK0025 Agglomeration	
Cardiff Urban Area UK0026 Agglomeration	
Swansea Urban Area UK0027 Agglomeration	
Belfast Metropolitan Urban Area UK0028 Agglomeration	
Eastern UK0029 Non-agglomeration	
South West UK0030 Non-agglomeration	
South East UK0031 Non-agglomeration	
East Midlands UK0032 Non-agglomeration	
North West & Merseyside UK0033 Non-agglomeration	
Yorkshire & Humberside UK0034 Non-agglomeration	
West Midlands UK0035 Non-agglomeration	
North East UK0036 Non-agglomeration	
Central Scotland UK0037 Non-agglomeration	
North East Scotland UK0038 Non-agglomeration	
Highland LIK0039 Non-agglomeration	
Scottish Borders UK0040 Non-agglomeration	
South Wales UK0041 Non-agglomeration	
North Wales UK0042 Non agglomeration	
Northern Ireland UIK0043 Non-agglomeration	

Table 4-1 UK Zones for Ambient Air Quality Reporting 2016

Figure 4-1 UK Zones for Ambient Air Quality Reporting 2016



© Crown copyright. All rights reserved Defra, Licence number 100022861 [2017]

4.2 Air Quality Assessment for 2016

The air quality assessment for each pollutant is derived from a combination of measured and modelled concentrations. Where both measurements and model results are available the assessment of compliance for each zone is based on the higher concentration of the two.

The air quality compliance assessment is submitted to the European Commission via e-Reporting. All the compliance results come under 'Information on the Attainment of Environmental Objectives' in e-Reporting Data Flow G.

The results of the air quality assessment submitted to the European Commission for 2016 are summarised in the tables below. The tables have been completed as follows:

- Where all measurements were within the relevant limit values in 2016, the table shows this as 'OK'.
- In the above cases, where compliance was determined by modelling or supplementary assessment, this is indicated by '(m)' – i.e. 'OK (m)'.
- Where locations were identified as exceeding a limit value, target value or long-term objective, this is identified as '>LV', '>TV' or '>LTO' as applicable.
- Where a non-compliance was determined by modelling or supplementary assessment, this is indicated by (m), as above.
- The abbreviation 'n/a' (not applicable) means that an assessment is not relevant for this zone, such as for the NO_X vegetation critical level in agglomeration zones.
- Zones that complied with the relevant limit values, targets or long-term objectives are shaded blue, while those that did not are shaded red. For ozone, zones that met the relevant target value but not the long-term objective are shaded purple.

There are no longer any zones where margins of tolerance apply.

4.2.1 Directive 2008/50/EC on Ambient Air Quality and Cleaner Air for Europe

Sulphur dioxide (SO₂): in 2016, all zones and agglomerations within the UK complied with the limit values for 1-hour mean and 24-hour mean SO₂ concentration, set for protection of human health.

All non-agglomeration zones within the UK also complied with the critical levels for annual mean and winter mean SO₂ concentration, set for protection of ecosystems. (These are not applicable to built-up areas).

Carbon monoxide (CO), benzene and lead: all zones and agglomerations were compliant with the limit values for these three pollutants in 2016. The 2016 compliance assessment for CO has been based on objective estimation (explained in Defra's technical report on UK air quality assessment⁴⁸), underpinned by NAEI emission trends, AURN measurement trends and historical modelling assessments.

Nitrogen dioxide (NO₂): in 2016 not all zones and agglomerations were compliant with the limit values. The results of the air quality assessment for nitrogen dioxide for each zone are summarised in **Table 4-2**.

Two zones had locations where the 1-hour limit value (200 μ g m⁻³) was exceeded on more than the permitted 18 occasions during 2016: Greater London Urban Area (UK0001) and South Wales (UK0041). The remaining 41 zones and agglomerations complied with the 1-hour mean NO₂ limit value.

Six zones *met* the annual mean limit value for NO₂ in 2016:

- Brighton/Worthing/Littlehampton (UK0010)
- Blackpool Urban Area (UK0022)
- Preston Urban Area (UK0023)
- Highland (UK0039)
- Scottish Borders (UK0040)
- Northern Ireland (UK0043).

The remaining 37 zones had locations with measured or modelled annual mean NO_2 concentrations higher than the annual mean limit value (40 µg m⁻³).

All non-agglomeration zones within the UK complied with the critical level for annual mean NO_X concentration, set for protection of vegetation.

PM₁₀ **Particulate Matter:** all zones and agglomerations were compliant with the annual mean limit value of 40 μ g m⁻³ for PM₁₀. All zones and agglomerations were compliant with the daily mean limit value. The results of the air quality assessment for PM₁₀ for each zone, with respect to the daily mean and annual mean limit values, are summarised in **Table 4-3**.

Under Section 20 of the Air Quality Directive, Member States are required to inform the Commission where exceedances of PM₁₀ limit values are due to natural sources (for example sea salt), and where this is the case, the exceedance does not count as non-compliance. Particulate matter from sea salt is modelled and has been used in the past to determine whether compliance with the limit values has been achieved after contribution from natural sources has been subtracted. However, in 2016 there were no modelled exceedances of either the 24-hr or annual mean limit values, so no subtraction of contribution from natural sources has been carried out.

Zone	Zone code	NO ₂ LV for health (1hr mean)	NO₂ LV for health (annual mean)	NO _x critical level for vegetation (annual mean)
Greater London Urban Area	UK0001	> LV	> LV	n/a
West Midlands Urban Area	UK0002	OK	> I V (m)	n/a
Greater Manchester Urban Area	UK0003	OK	> LV (m)	n/a
West Yorkshire Urban Area	UK0004	OK	> LV	n/a
Tyneside	UK0005	OK	> LV (m)	n/a
Liverpool Urban Area	UK0006	ОК	> LV (m)	n/a
Sheffield Urban Area	UK0007	ОК	> LV (m)	n/a
Nottingham Urban Area	UK0008	ОК	> LV (m)	n/a
Bristol Urban Area	UK0009	OK	> LV (m)	n/a
Brighton/Worthing/Littlehampton	UK0010	OK	OK	n/a
Leicester Urban Area	UK0011	OK	> LV	n/a
Portsmouth Urban Area	UK0012	OK	> LV (m)	n/a
Teesside Urban Area	UK0013	OK	> LV (m)	n/a
The Potteries	UK0014	OK	> LV	n/a
Bournemouth Urban Area	UK0015	OK	> LV (m)	n/a
Reading/Wokingham Urban Area	UK0016	OK	> LV (m)	n/a
Coventry/Bedworth	UK0017	OK	> LV (m)	n/a
Kingston upon Hull	UK0018	OK	> LV (m)	n/a
Southampton Urban Area	UK0019	OK	> LV	n/a
Birkenhead Urban Area	UK0020	OK	> LV (m)	n/a
Southend Urban Area	UK0021	OK	> LV (m)	n/a
Blackpool Urban Area	UK0022	OK	OK	n/a
Preston Urban Area	UK0023	OK	OK	n/a
Glasgow Urban Area	UK0024	OK	> LV	n/a
Edinburgh Urban Area	UK0025	OK (m)	> LV (m)	n/a
Cardiff Urban Area	UK0026	OK	> LV (m)	n/a
Swansea Urban Area	UK0027	OK	> LV (m)	n/a
Belfast Urban Area	UK0028	OK	> LV	n/a
Eastern	UK0029	OK	> LV	OK
South West	UK0030	OK	> LV	OK
South East	UK0031	OK	> LV	OK
East Midlands	UK0032	OK	> LV (m)	OK
North West & Merseyside	UK0033	OK	> LV	OK (m)
Yorkshire & Humberside	UK0034	OK	> LV (m)	OK
West Midlands	UK0035	OK	> LV (m)	OK (m)
North East	UK0036	OK	> LV (m)	OK (m)
Central Scotland	UK0037	OK	> LV (m)	OK (m)
North East Scotland	UK0038	OK	> LV	OK (m)
Highland	UK0039	OK	OK	OK (m)
Scottish Borders	UK0040	OK	OK	OK
South Wales	UK0041	> LV	> LV	OK (m)
North Wales	UK0042	OK	> LV (m)	OK
Northern Ireland	UK0043	OK	OK	OK (m)

Table 4-2 Results of Air Quality Assessment for Nitrogen Dioxide in 2016

LV = limit value, (m) indicates that the compliance or exceedance was determined by modelling.

Table 4-3 Results of Air Quality Assessment for PM_{10} in 2016

		PM ₁₀ LV	PM ₁₀ LV
Zone	Zone code	(daily mean)	(annual mean)
Greater London Urban Area	UK0001	OK	OK
West Midlands Urban Area	UK0002	OK	ОК
Greater Manchester Urban Area	UK0003	OK	OK
West Yorkshire Urban Area	UK0004	ОК	OK
Tyneside	UK0005	ОК	OK
Liverpool Urban Area	UK0006	ОК	OK
Sheffield Urban Area	UK0007	ОК	OK
Nottingham Urban Area	UK0008	ОК	OK
Bristol Urban Area	UK0009	ОК	ОК
Brighton/Worthing/Littlehampton	UK0010	OK (m)	OK (m)
Leicester Urban Area	UK0011	ОК	OK
Portsmouth Urban Area	UK0012	OK (m)	OK (m)
Teesside Urban Area	UK0013	ОК	ОК
The Potteries	UK0014	ОК	ОК
Bournemouth Urban Area	UK0015	OK (m)	OK (m)
Reading/Wokingham Urban Area	UK0016	ОК	ОК
Coventry/Bedworth	UK0017	OK (m)	OK (m)
Kingston upon Hull	UK0018	ОК	ОК
Southampton Urban Area	UK0019	OK (m)	OK (m)
Birkenhead Urban Area	UK0020	OK (m)	OK (m)
Southend Urban Area	UK0021	OK (m)	OK (m)
Blackpool Urban Area	UK0022	OK (m)	OK (m)
Preston Urban Area	UK0023	OK (m)	OK (m)
Glasgow Urban Area	UK0024	ОК	ОК
Edinburgh Urban Area	UK0025	ОК	OK
Cardiff Urban Area	UK0026	OK (m)	OK (m)
Swansea Urban Area	UK0027	ОК	OK
Belfast Metropolitan Urban Area	UK0028	ОК	OK
Eastern	UK0029	ОК	OK
South West	UK0030	OK	OK
South East	UK0031	ОК	ОК
East Midlands	UK0032	ОК	ОК
North West & Merseyside	UK0033	OK	OK
Yorkshire & Humberside	UK0034	ОК	OK
West Midlands	UK0035	OK	OK
North East	UK0036	OK (m)	OK (m)
Central Scotland	UK0037	ОК	OK
North East Scotland	UK0038	OK	OK
Highland	UK0039	ОК	OK
Scottish Borders	UK0040	OK (m)	OK (m)
South Wales	UK0041	OK	OK
North Wales	UK0042	OK	OK
Northern Ireland	UK0043	OK	OK

In Table 4-3, LV = limit value, (m) indicates that the compliance or exceedance was determined by modelling.

PM_{2.5} **Particulate Matter:** all zones met the target value for annual mean concentration of PM_{2.5} particulate matter (25 μ g m⁻³ to be achieved by 1st Jan 2010), the Stage 1 limit value (25 μ g m⁻³ to be achieved by 1st Jan 2015), which came into force on 1st January 2015, and the Stage 2 limit value (20 μ g m⁻³ to be achieved by 1st Jan 2020). All three apply to the calendar year mean.

The results of the air quality assessment for $PM_{2.5}$ for each zone are summarised in **Table 4-4**. Subtraction of $PM_{2.5}$ contributions due to natural events (1999/30/EC Article 5(4)) or natural contributions (2008/50/EC Article 20) was not necessary for any zone.

Under the Air Quality Directive, Member States will be required to achieve a national exposure reduction target for PM_{2.5}, over the period 2010 to 2020. This is based on the Average Exposure Indicator (AEI) statistic. The AEI for the UK is calculated as follows: the arithmetic mean PM_{2.5} concentration at appropriate UK urban background sites only is calculated for three consecutive calendar years, and the mean of these values taken as the AEI.

The AEI for the reference year (2010) was used to determine the National Exposure Reduction Target (NERT), to be achieved by 2020 (see Annex XIV of the Air Quality Directive). The UK's reference year AEI was 13 μ g m⁻³; on this basis, the Air Quality Directive sets an exposure reduction target of 15%. This equates to reducing the AEI to 11 μ g m⁻³ by 2020. (The detailed methodology and results of this calculation are presented in Defra's technical report on UK air quality assessment⁴⁸.)

The AEI for the reference year 2015 is set at 20 μ g m⁻³ as an Exposure Concentration Obligation (ECO) in the Air Quality Directive. The UK already meets this obligation. There are no obligations or target values for the years *between* 2010, 2015 and 2020, but the running AEIs for these intervening years give an indication of progress towards the 2020 target. The running year AEI for 2016 was calculated as follows:

- 2014: 12 µg m⁻³
- 2015: 10 µg m⁻³
- 2016: 10 µg m⁻³

The mean of these three values (to the nearest integer) is 11 μ g m⁻³. The exposure reduction target has therefore been met before 2020.

			PM _{2.5} Stage 1	PM _{2.5} Stage 2
		PM _{2.5} target	limit value	limit value
		value (annual	(annual mean,	(annual
		mean, for 1 st	for 1 st Jan	mean, for 1 st
Zone	Zone code	Jan 2010)	2015)	Jan 2020)
Greater London Urban Area	UK0001	OK	OK	OK
West Midlands Urban Area	UK0002	OK	OK	OK
Greater Manchester Urban Area	UK0003	OK	OK	OK
West Yorkshire Urban Area	UK0004	OK	OK	OK
Tyneside	UK0005	OK	OK	OK
Liverpool Urban Area	UK0006	OK	OK	OK
Sheffield Urban Area	UK0007	OK	OK	OK
Nottingham Urban Area	UK0008	OK	OK	OK
Bristol Urban Area	UK0009	OK	OK	OK
Brighton/Worthing/Littlehampton	UK0010	OK	OK	OK
Leicester Urban Area	UK0011	OK	OK	OK
Portsmouth Urban Area	UK0012	OK	OK	OK
Teesside Urban Area	UK0013	OK	OK	OK
The Potteries	UK0014	OK	OK	OK
Bournemouth Urban Area	UK0015	OK	OK	OK
Reading/Wokingham Urban Area	UK0016	OK	OK	OK
Coventry/Bedworth	UK0017	OK	OK	ОК
Kingston upon Hull	UK0018	OK	OK	ОК
Southampton Urban Area	UK0019	OK (m)	OK (m)	OK (m)
Birkenhead Urban Area	UK0020	OK	OK	OK
Southend Urban Area	UK0021	OK	OK	ОК
Blackpool Urban Area	UK0022	OK	OK	OK
Preston Urban Area	UK0023	OK	OK	OK
Glasgow Urban Area	UK0024	OK	OK	OK
Edinburgh Urban Area	UK0025	OK	OK	OK
Cardiff Urban Area	UK0026	OK	OK	OK
Swansea Urban Area	UK0027	OK	OK	OK
Belfast Metropolitan Urban Area	UK0028	OK	OK	OK
Fastern	UK0029	OK	OK	OK
South West		OK	OK	OK
South Fast		OK	OK	OK
East Midlands		OK	OK	OK
North West & Merseyside		OK	OK	OK
Vorkshiro & Humborsido		OK	OK	OK
West Midlands		OK	OK	OK
Nexth Fast		OK	OK	OK
North East		OK	OK	OK
Central Scotland		OK	OK	0K
Highland		OK	OK	OK
	UK0039			
Scottish Borders		OK (m)	OK (m)	
	UK0041	OK	OK	OK
North Wales	UK0042	OK	OK	OK
Northern Ireland	UK0043	OK (m)	OK (m)	OK (m)

Table 4-4 Results of Air Quality Assessment for PM_{2.5} in 2016.

Subtraction of natural source contribution was not carried out for any zones in 2016.

LV = limit value, (m) indicates that the compliance or exceedance was determined by modelling.

Ozone: all zones and agglomerations met the target values for health and for protection of vegetation. The results of the air quality assessment for ozone are summarised in **Table 4-5**.

For ozone, there is a target value based on the maximum daily 8-hour mean. All 43 zones and agglomerations were compliant with this target value. There is also a long-term objective for protection of human health, based on the maximum daily 8-hour mean. All but one of the 43 zones and agglomerations were *above* the long-term objective (LTO) for health in 2016, the exception being Edinburgh Urban Area (UK0025).

There is also a target value based on the AOT40 statistic. The AOT40 statistic (expressed in μ g m⁻³.hours) is the sum of the difference between hourly concentrations greater than 80 μ g m⁻³ (= 40 ppb) and 80 μ g m⁻³ over a given period using only the hourly mean values measured between 08:00 and 20:00 Central European Time each day. All 43 zones and agglomerations met the target value based on the AOT40 statistic. There is also a long-term objective, for protection of vegetation, based on this statistic. Five zones (Yorkshire and Humberside, the West Midlands, the North East, South Wales and North Wales) were above the long-term objective for vegetation in 2016.

Ozone concentrations – and hence the number of zones exceeding the LTOs fluctuate from year to year as ozone is a transboundary pollutant and its formation is influenced by meteorological factors.

Table 4-5 Results of Air Qua	lity Assessment	for Ozone in 2016
------------------------------	-----------------	-------------------

Zone		O ₃ TV and LTO for health	O ₃ TV and LTO for
	Zone code	(8hr mean)	vegetation (AOT40)
Greater London Urban Area	UK0001	Met TV, > LTO	OK
West Midlands Urban Area	UK0002	Met TV, > LTO	OK
Greater Manchester Urban Area	UK0003	Met TV, > LTO	ОК
West Yorkshire Urban Area	UK0004	Met TV, > LTO	OK
Tyneside	UK0005	Met TV, > LTO (m)	OK
Liverpool Urban Area	UK0006	Met TV, > LTO	OK
Sheffield Urban Area	UK0007	Met TV, > LTO	OK
Nottingham Urban Area	UK0008	Met TV, > LTO	OK
Bristol Urban Area	UK0009	Met TV, > LTO (m)	OK
Brighton/Worthing/Littlehampton	UK0010	Met TV, > LTO (m)	OK
Leicester Urban Area	UK0011	Met TV, > LTO	OK
Portsmouth Urban Area	UK0012	Met TV, > LTO (m)	OK
Teesside Urban Area	UK0013	Met TV, > LTO	OK
The Potteries	UK0014	Met TV, > LTO	OK
Bournemouth Urban Area	UK0015	Met TV, > LTO	OK
Reading/Wokingham Urban Area	UK0016	Met TV, > LTO	OK
Coventry/Bedworth	UK0017	Met TV, > LTO	OK
Kingston upon Hull	UK0018	Met TV, > LTO	OK
Southampton Urban Area	UK0019	Met TV, > LTO (m)	OK (m)
Birkenhead Urban Area	UK0020	Met TV, > LTO	OK
Southend Urban Area	UK0021	Met TV, > LTO	OK
Blackpool Urban Area	UK0022	Met TV, > LTO	OK
Preston Urban Area	UK0023	Met TV, > LTO	OK
Glasgow Urban Area	UK0024	Met TV, > LTO (m)	OK
Edinburgh Urban Area	UK0025	ОК	OK
Cardiff Urban Area	UK0026	Met TV, > LTO (m)	OK
Swansea Urban Area	UK0027	Met TV, > LTO	OK
Belfast Metropolitan Urban Area	UK0028	Met TV, > LTO (m)	OK
Eastern	UK0029	Met TV, > LTO	OK
South West	UK0030	Met TV, > LTO (m)	OK
South East	UK0031	Met TV, > LTO	OK
East Midlands	UK0032	Met TV, > LTO	OK
North West & Merseyside	UK0033	Met TV, > LTO	OK
Yorkshire & Humberside	UK0034	Met TV, > LTO	Met TV, > LTO
West Midlands	UK0035	Met TV, > LTO	Met TV, > LTO (m)
North East	UK0036	Met TV, > LTO (m)	Met TV, > LTO (m)
Central Scotland	UK0037	Met TV, > LTO (m)	OK
North East Scotland	UK0038	Met TV, > LTO	OK
Highland	UK0039	Met TV, > LTO	ОК
Scottish Borders	UK0040	Met TV, > LTO	ОК
South Wales	UK0041	Met TV, > LTO	Met TV, > LTO (m)
North Wales	UK0042	Met TV, > LTO	Met TV, > LTO
Northern Ireland	UK0043	Met TV, > LTO	OK

Footnote to Table 4-5: TV = target value, LTO = long-term objective, (m) indicates that the compliance or exceedance was determined by modelling.

In 2016 there were five measured exceedances of the ozone population information threshold (at four sites), but no exceedances of the population warning threshold. The information threshold exceedances are detailed in **Table 4-6**. All occurred between 19:00 and 20:00 on 19th July 2016.

Table 4-6 Measured Exceedances of the Ozone Information Threshold	Value in
2016	

Site name	Zone code	Number of 1-hour exceedances of information threshold	Maximum 1-hour concentration (µg m ⁻³)
Canterbury	UK0031	1	186
Sibton	UK0029	2	181
Southend-on-Sea	UK0021	1	181
St Osyth	UK0029	1	194

4.2.2 Fourth Daughter Directive 2004/107/EC

All zones met target values for arsenic and cadmium but some zones exceeded target values for nickel and benzo[a]pyrene. The results of the air quality assessment for arsenic (As), cadmium (Cd), nickel (Ni) and benzo[a]pyrene (B[a]P) for each zone are summarised in **Table 4-7**.

All zones and agglomerations met the target values for arsenic and cadmium. Three zones (Sheffield Urban Area, Swansea Urban Area and South Wales) exceeded the target value for nickel. In these zones, the exceedance has been attributed to industrial sources.

Concentrations of B[a]P were above the target value in four zones; Swansea Urban Area, Yorkshire and Humberside, South Wales and Northern Ireland. In the Swansea Urban Area, South Wales, and Yorkshire and Humberside zones, exceedances are attributed to emissions from industrial sources. In Northern Ireland, domestic combustion is the main source of B[a]P.

The remaining 39 zones were compliant with the target value for B[a]P, as shown in **Table 4-7**.

Zone	Zone code	As TV	Cd TV	Ni TV	B[a]P TV
Greater London Urban Area	UK0001	OK	OK	OK	OK
West Midlands Urban Area	UK0002	OK	OK	OK	OK
Greater Manchester Urban Area	UK0003	OK (m)	OK (m)	OK (m)	OK
West Yorkshire Urban Area	UK0004	OK (m)	OK (m)	OK (m)	OK
Tyneside	UK0005	OK (m)	OK (m)	OK (m)	OK
Liverpool Urban Area	UK0006	OK (m)	OK (m)	OK (m)	ОК
Sheffield Urban Area	UK0007	OK	OK	> TV	OK (m)
Nottingham Urban Area	UK0008	OK (m)	OK (m)	OK (m)	OK (m)
Bristol Urban Area	UK0009	OK (m)	OK (m)	OK (m)	OK (m)
Brighton/Worthing/Littlehampton	UK0010	OK (m)	OK (m)	OK (m)	OK (m)
Leicester Urban Area	UK0011	OK (m)	OK (m)	OK (m)	OK (m)
Portsmouth Urban Area	UK0012	OK (m)	OK (m)	OK (m)	OK (m)
Teesside Urban Area	UK0013	OK (m)	OK (m)	OK (m)	ОК
The Potteries	UK0014	OK (m)	OK (m)	OK (m)	OK (m)
Bournemouth Urban Area	UK0015	OK (m)	OK (m)	OK (m)	OK (m)
Reading/Wokingham Urban Area	UK0016	OK (m)	OK (m)	OK (m)	OK (m)
Coventry/Bedworth	UK0017	OK (m)	OK (m)	OK (m)	OK (m)
Kingston upon Hull	UK0018	OK (m)	OK (m)	OK (m)	OK (m)
Southampton Urban Area	UK0019	OK (m)	OK (m)	OK (m)	OK (m)
Birkenhead Urban Area	UK0020	OK (m)	OK (m)	OK (m)	OK (m)
Southend Urban Area	UK0021	OK (m)	OK (m)	OK (m)	OK (m)
Blackpool Urban Area	UK0022	OK (m)	OK (m)	OK (m)	OK (m)
Preston Urban Area	UK0023	OK (m)	OK (m)	OK (m)	OK (m)
Glasgow Urban Area	UK0024	OK (m)	OK (m)	OK (m)	ОК
Edinburgh Urban Area	UK0025	OK (m)	OK (m)	OK (m)	ОК
Cardiff Urban Area	UK0026	OK (m)	OK (m)	OK (m)	ОК
Swansea Urban Area	UK0027	OK	OK	> TV	> TV (m)
Belfast Urban Area	UK0028	OK	OK	OK	ОК
Eastern	UK0029	OK	OK	OK	ОК
South West	UK0030	OK	OK	OK	OK (m)
South East	UK0031	OK	OK	OK	ОК
East Midlands	UK0032	OK	OK	OK	ОК
North West & Merseyside	UK0033	OK	OK	OK	OK
Yorkshire & Humberside	UK0034	OK	OK	OK	> TV
West Midlands	UK0035	OK	OK	OK	OK (m)
North East	UK0036	OK (m)	OK (m)	OK (m)	ОК
Central Scotland	UK0037	OK	OK	OK	OK
North East Scotland	UK0038	OK (m)	OK (m)	OK (m)	OK (m)
Highland	UK0039	OK (m)	OK (m)	OK (m)	OK
Scottish Borders	UK0040	OK	OK	OK	OK (m)
South Wales	UK0041	OK	OK	> TV (m)	> TV (m)
North Wales	UK0042	OK (m)	OK (m)	OK (m)	OK (m)
Northern Ireland	UK0043	OK (m)	OK (m)	OK (m)	> TV

TV = target value, (m) indicates that the compliance or exceedance was determined by modelling.

4.3 Comparison with Previous Years

Table 4-8 to **Table 4-12** summarise the results of the air quality assessment for 2016 and provide a comparison with the results of the assessments carried out in previous years since 2008 (the year in which the Air Quality Directive came into force). For information on compliance with the 1st and 2nd Daughter Directives in earlier years, please see the 2012 or earlier reports in this series. **Table 4-8** shows the number of zones exceeding the limit value plus any agreed margin of tolerance applicable in the relevant year (i.e. the numbers of zones that were non-compliant). If any additional zones were within the limit value plus an agreed MOT (and therefore compliant), for example for NO₂, this is shown in the footnotes. As of 1st January 2015, there are no longer any margins of tolerance (MOT) in force for any pollutants.

Pollutant	Avg. time	2008	2009	2010	2011	2012	2013	2014	2015	2016
SO ₂	1-hour	None	None	None	None	None	None	None	None	None
SO ₂	24-hour	None	None	None	None	None	None	None	None	None
SO ₂	Annual ⁱ	None	None	None	None	None	None	None	None	None
SO ₂	Winter ⁱ	None	None	None	None	None	None	None	None	None
NO ₂	1-hour ⁱⁱ	3 zones	2 zones	3 zones	3 zones	2 zones	1 zone	2 zones	2 zones	2 zones
		(London,	(London,	(London,	(London,	(London,	(London)	(London,	(London,	(London,
		Glasgow, NE	Glasgow)	Teesside,	Glasgow,	South East)		South Wales)	South Wales)	South Wales)
		Scotland)		Glasgow)	South East)					
NO ₂	Annual	40 zones	40 zones	40 zones	35 zones ⁱⁱⁱ	34 zones ^{iv}	31 zones v	30 zones vi	37 zones ^{vii}	37 zones
NOx	Annual ⁱ	None	None	None	None	None	None	None	None	None

Table 4-8 Non-Compliances with the Limit Values of the Air Quality Directive

Pollutant	Avg. time	2008	2009	2010	2011	2012	2013	2014	2015	2016
PM10	Daily	2 zones (1 zone after subtraction of natural contribution)	3 zones (1 zone after subtraction of natural contribution)	None (after subtraction of natural contrib- ution) ^{viii}	None (after subtraction of natural contribution) ^{ix}	None (after subtraction of natural contribution. No time extension.)	None			
PM10	Annual	None	None	None	None	None	None	None	None	None
Lead	Annual	None	None	None	None	None	None	None	None	None
Benzene	Annual	None	None	None	None	None	None	None	None	None
CO	8-hour	None	None	None	None	None	None	None	None	None

Footnotes to Table 4-8:

¹ Applies to vegetation and ecosystem areas only. Critical Levels are already in force, no MOT.

" No modelling for 1-hour LV.

ii A further five zones exceeded the annual mean NO₂ LV in 2011 but were covered by time extensions and within the LV+ MOT, therefore compliant.

^{*iv*} A further four zones exceeded the annual mean NO₂ LV in 2012 but were covered by time extensions and within the LV+ MOT, therefore compliant.

^v A further seven zones exceeded the annual mean NO₂ LV in 2013 but were covered by time extensions and within the LV+ MOT, therefore compliant.

^{vi} A further eight zones exceeded the annual mean NO₂ LV in 2014 but were covered by time extensions and within the LV+ MOT, therefore compliant.

^{vii} 2015 was the first year with no time extensions for NO₂: this is the reason for the apparent increase in zones exceeding between 2014 and 2015. In 2014, 5 zones met the limit value and a further 8 zones were legally compliant due to the time extension. The time extensions ended on 1st January 2015. In 2015, 6 zones met the limit value, but the remaining zones were no longer covered by the time extension, bringing the total number of non-compliant zones from 30 in 2014 to 37 in 2015.

^{viii} One zone exceeded the daily mean PM₁₀ limit value more than the permitted 35 times in 2010, after subtraction of natural contribution. This zone was covered by a time extension, and was within the LV+MOT so was therefore compliant.

^{ix} One zone exceeded the daily mean PM₁₀ limit value more than the permitted 35 times in 2011, after subtraction of natural contribution. This zone was covered by a time extension, and was within the LV+MOT so was therefore compliant.

The UK has been compliant with the limit values for both lead and CO since 2003, and for benzene since 2007: these limit values are the same as those contained in the 1st and 2nd Daughter Directives, which the Air Quality Directive superseded.

Pollutant	Averaging time	2008	2009	2010	2011	2012	2013	2014	2015	2016
O ₃	8-hour	1 zone measured (Eastern)	None							
O ₃	AOT40	None	None	None	None	None	None	None	None	None

Table 4-9 Exceedances of Air Quality Directive Target Values for Ozone (Health)

Table 4-10 Exceedances of Air Quality Directive Long Term Objectives for Ozone

Pollutant	Averaging time	2008	2009	2010	2011	2012	2013	2014	2015	2016
O ₃	8-hour	43 zones	39 zones	41 zones	43 zones	41 zones	33 zones	32 zones	43 zones	42 zones
O ₃	AOT40	41 zones	10 zones	6 zones	3 zones	3 zones	8 zones	3 zones	1 zone	5 zones

Pollu -tant	Averaging time	2007	2008	2009	2010	2011	2012	2013	2014	2015	2016
As	Annual	None	None	None	None	None	None	None	None	None	None
Cd	Annual	None	None	None	None	None	None	None	None	None	None
Ni	Annual	1 zone (Swansea)	2 zones (Swansea, S Wales)	2 zones (Swansea, S Wales)	2 zones (Swansea, S Wales)	2 zones, (Swansea, S Wales)	2 zones, (Swansea, S Wales)	2 zones, (Swansea, S Wales)	3 zones, (Sheffield, Swansea, S Wales)	2 zones, (Swansea and S Wales)	3 zones, (Sheffield, Swansea, S Wales)
B[a]P	Annual	1 zone (Yorkshire & Humberside)	6 zones (Yorkshire & Humberside, Teesside, N Ireland, Swansea, S Wales, Belfast)	6 zones (Yorkshire & Humberside, N Ireland, Teesside, Swansea, North East, S Wales)	8 zones, (Yorkshire & Humberside, N Ireland, Teesside, Belfast, W Midlands, North East, S Wales, N Wales.)	7 zones (Yorkshire & Humberside, N Ireland, Teesside, Swansea, Belfast, North East, South Wales)	8 zones (Yorkshire & Humberside, Teesside, Swansea, Belfast, the North East, South Wales, North Wales, North Wales, Northern Ireland.)	6 zones (Yorkshire & Humberside, Teesside, Swansea, the East Midlands, the North East, South Wales.)	6 zones (Yorkshire & Humberside, Teesside, Swansea, the East Midlands, the North East and South Wales).	5 zones (Yorks. & Humber- side, Teesside, Swansea, the North East and South Wales).	4 zones (Yorks. & Humber- side, Swansea, South Wales and Northern Ireland).

Table 4-11 Exceedances of 4th Daughter Directive Target Values

Pollutant	Ave. time	2009	2010	2011	2012	2013	2014	2015	2016
PM _{2.5}	Annual	None							

5 Spatial Variation and Changes Over Time

This section looks at the spatial distribution of pollutants across the UK, based upon the modelled maps of ambient pollutant concentration discussed in Section 3.5 of this report, "Modelling". In the case of traffic-related pollutants such as NO₂, roadside and background concentrations are discussed separately.

For each pollutant, this section also discusses how ambient concentrations have changed over time, using data from the relevant ambient air quality monitoring networks: the Automatic Urban and Rural Network (AURN), the Automatic Hydrocarbon Network, the Non-Automatic Hydrocarbon Network, the UK Metals Network, and the PAH Network.

The AURN has been in operation since 1992. Since that time, it has grown considerably in size, and the proportion of urban traffic ('roadside') monitoring sites has increased. Therefore, we have in some cases based our investigation of trends on data from sub-sets of long-running AURN monitoring sites, rather than the whole network. This is intended to show changes over time, without introducing any spurious effects due to changes in the number and distribution of the sites.

For pollutants monitored using continuous automatic techniques, trend analysis has been carried out using Openair: a free, open-source software package of tools for analysis of air pollution data. Openair was developed by King's College London with the University of Leeds. The Openair project is currently led by Dr David Carslaw, of Ricardo Energy & Environment and the University of York. A range of Openair functions are available on UK-AIR: for more information on the functions and how to use them, please refer to: https://uk-air.defra.gov.uk/data/openair.

The Openair 'TheilSen' function has been used here. This uses the Theil-Sen statistical method to determine trends in pollutant concentrations over several years. The trend analysis is based on monthly mean pollutant concentrations. (At least 75% data capture is required for a valid monthly mean.) The 'TheilSen' function includes an option to 'de-seasonalise' the data (i.e. statistically modify the plotted data to remove the influence of seasonal cycles, thus providing a clearer indication of the overall trend). The de-seasonalise option has been used for most pollutants, particularly those that show a seasonal pattern; SO₂, NO₂, particulate matter and CO - but not 1,3-butadiene or ozone. Where the 'de-seasonalise' option has been used here, this is indicated in the graph title. In these trend graphs, the trend line is shown by a solid red line, with 95% confidence intervals for the trend shown by dotted red lines. The trend is given at the top of the graph in green, with confidence intervals shown in square brackets. The trend is given as units (e.g. μ g m⁻³) per year, over the period shown. This may be followed by a number of symbols, with + indicating that

the trend is significant at the 0.1 level, * indicating that the trend is statistically significant at the 0.05 level, ** indicating significance at the 0.01 level and *** indicating significance at the 0.001 level. The 'de-seasonalise' option also fills in gaps in the dataset using an interpolation method.

These changes over time are compared to changes in estimated total UK emissions where appropriate. Estimated UK emissions data are taken from the National Atmospheric Emissions Inventory (NAEI) website⁵⁰ at http://naei.defra.gov.uk/index.php. (The most recent year for which NAEI emission estimates are available is 2015). For some pollutants, the NAEI website allows bar charts to be created, showing emissions over a selected period of years, split between various emission source categories.

Please note that this section only aims to provide a general indication of changes in pollutant concentration over time, based in most cases on averages from groups of long-running sites. Patterns in specific regions or individual sites may be different.

5.1 Sulphur Dioxide

5.1.1 SO₂: Spatial Distribution in the UK

Figure 5-1 shows how the modelled 99.73rd percentile^a of hourly mean sulphur dioxide concentration varied across the UK during 2016. This statistic corresponds approximately to the 25th highest hourly mean (in the case of a full year's data); if greater than the hourly mean limit value it indicates that the limit value was exceeded on more than the 24 permitted occasions. There were no areas in which this statistic exceeded the limit value of 350 μ g m⁻³.

Figure 5-2 shows the modelled 99.18th percentile of 24-hour means (which corresponds to the 4th highest day in a full year). If greater than the 24-hourly mean limit value of 125 μ g m⁻³, this would indicate that there were more than the permitted three exceedances in the year. There were no areas of the UK where this was the case in 2016.

5.1.2 SO₂: Changes Over Time

Figure 5-3 shows how ambient concentrations have changed over the period 1992 to 2016, at the seven AURN monitoring stations that have monitored this pollutant for the longest time, and have remained in operation in 2016. All seven stations show a

^a Where the Directive allows exceedances on a number of occasions (i.e. limit value not to be exceeded more than a specified number of times per year), percentiles are used to illustrate this. These are simply the xth highest hourly mean divided by the number of hours in a year, or yth highest daily mean divided by the days in a year, expressed as a percentage.

downward trend that is statistically significant at the 0.001 level - highly significant – as denoted by the three asterisks (***) on the plots.

However, the decrease has not been linear. At most of these sites, the downward trend is steepest for the 1990s and early 2000s: there is a clear flattening-off in more recent years from around 2010.

The pattern observed in ambient SO₂ concentrations appears to reflect changes in national emissions. **Figure 5-4** is taken from the NAEI and shows the UK's estimated annual emissions of sulphur dioxide from 1992 (the first year of operation of the AURN) to 2015. The main source of this pollutant is fossil fuel combustion. SO₂ emissions in the UK have decreased substantially since 1992, due to reductions in the use of coal, gas and oil, and also to reductions in the sulphur content of fuel oils and DERV (diesel fuel used for road vehicles). The decrease in emissions over time shown here is the continuation of an on-going trend observed by the NAEI throughout the 1970s and 1980s, partly due to the decline of the UK's heavy industry.

However, around 2009, the graph flattens off, and shows a slight upturn in total SO₂ emissions in 2012. The NAEI pollutant information page for SO₂ (at http://naei.defra.gov.uk/overview/pollutants?pollutant_id=8) explains this as follows: "As a result of the economic downturn the drive to cut energy costs has resulted in an increase in solid fuel use, particularly in 2012 some coal-sensitive pollutants have seen a significant rise in coal burning emissions." Following 2012, the downward trend in SO₂ emissions continues. Figure 5-1 99.73rd Percentile of 1-hour Mean SO₂ Concentration, 2016 (µg m⁻³)



Figure 5-2 99.18th Percentile of 24-hour Mean SO₂ Concentration, 2016 (μ g m⁻³)



Figure 5-3 De-seasonalised Trends in SO₂ Concentration, 1992-2016 at 7 Longrunning AURN Sites



Figure 5-4 Estimated Annual UK Emissions of SO₂ (kt), 1992 – 2015 Source: NAEI



5.2 Nitrogen Dioxide

5.2.1 NO₂: Spatial Distribution in the UK

Figure 5-5 shows the modelled annual mean NO₂ concentrations for 2016, at *urban roadside* locations only. Although not every road link is clearly visible, it is possible to see that many are shaded yellow, orange and red - indicating that they had annual mean NO₂ concentrations above the limit value of 40 μ g m⁻³. These locations are widespread in London and also visible in urban areas elsewhere in the UK.

Figure 5-6 shows the modelled annual mean *background* NO₂ concentrations for 2016. Most background locations were within the limit value of 40 μ g m⁻³, but some small areas were not. These are shaded yellow, orange and red. These were largely confined to the major urban areas, and principal road links.

5.2.2 NO₂: Changes Over Time

Figure 5-7 and **Figure 5-8** show how ambient concentrations of nitrogen dioxide (as measured by the AURN) have decreased since 1992 (the Network's first year of operation). Time series of annual mean NO₂ concentrations are shown for the following sub-sets of long-running sites:

- Eight urban background sites operating since 1995 or earlier (**Figure 5-7**); Belfast Centre, Cardiff Centre, Leeds Centre, London Bloomsbury, Middlesbrough, Newcastle Centre, Sheffield Tinsley and Southampton Centre.
- Eight urban traffic sites operating since 1998 or earlier (**Figure 5-8**); Bath Roadside, Camden Kerbside, Exeter Roadside, Glasgow Kerbside, Haringey Roadside, London Marylebone Road, Oxford Centre Roadside and Tower Hamlets Roadside.

(As explained above, the use of sub-sets of long-running monitoring sites is intended to show changes over time, without introducing any spurious effects due to changes in the number and distribution of the sites.)

Figure 5-5 Urban Major Roads, Annual Mean Roadside NO₂ Concentration, 2016 (µg m⁻³)






Figure 5-7 De-seasonalised Trends in NO₂ Concentration, at 8 Long-Running AURN Urban Background Sites, 1992- 2016



All eight long-running urban non-roadside sites in **Figure 5-7** above show a decreasing trend in NO₂; while the magnitude of the year-on year decrease varies (ranging from -0.47 μ g m⁻³ to -1.08 μ g m⁻³ per year), the trend is statistically significant at the 0.001 level for all eight sites.

For the urban traffic sites in **Figure 5-8** below, (for which the dataset is slightly shorter), the pattern of trends is less consistent. Six of the eight sites show a downward trend with varying levels of statistical significance. One site (Camden Kerbside) shows a statistically significant upward trend, and one (London Marylebone Road) shows *no* significant trend.

Figure 5-8 De-seasonalised Trends in NO₂ Concentration at 8 Long-Running AURN Urban Traffic Sites, 1998 - 2016



De-seasonalised NO2 trends for the period 1998 to 2016

Figure 5-9 shows NAEI estimates of total UK annual emission of oxides of nitrogen, in kilotonnes. Total NO_x emissions have decreased substantially over the period shown. While long-running urban background sites show a general decrease in NO₂ concentration as might be expected from the national emissions estimates, the same is not consistently true of urban traffic sites. It is likely that the trend in ambient NO₂ concentration at each individual site depends, at least in part, on the quantity and type of traffic on the adjacent road.

In July 2017, the Government launched its UK Plan for Tackling Roadside Nitrogen Dioxide Concentrations (see section 2.2.5 for more information on this). The national air quality plan is available at <u>https://www.gov.uk/government/publications/air-quality-plan-for-nitrogen-dioxide-no2-in-uk-2017</u>.

Figure 5-9 Estimated Annual UK Emissions of Nitrogen Oxides (kt), 1992 – 2015 Source: NAEI



Nitrogen Oxides (kilotonne)

5.3 PM₁₀ Particulate Matter

5.3.1 PM₁₀: Spatial Distribution

Figure 5-10 shows modelled annual mean urban roadside PM_{10} concentrations in 2016. No roadside locations had an annual mean concentration greater than 40 µg m⁻³. This is consistent with the compliance assessment reported in Section 4.

Figure 5-11 shows modelled annual mean background PM₁₀ concentrations in 2016. Background concentrations were higher in the southern and eastern parts of the country; this is a typical pattern, because these regions receive a larger transboundary contribution of particulate pollution from mainland Europe. The elevated levels of PM₁₀ associated with urban areas – particularly London, and the major cities of the West Midlands and East Midlands - can also be seen. Also visible are the routes of major roads, particularly in the Midlands and east of England.

The concentration bands used in the figures below include the ranges >30.1-30.5 μ g m⁻³, and >30.6-40 μ g m⁻³. The significance of the division at 30.5 μ g m⁻³ is that where the annual mean PM₁₀ concentration exceeds this value, it is likely also that the 24-hour mean has exceeded the daily mean limit value of 50 μ g m⁻³ on more than the permitted 35 occasions. (Note: this threshold is calculated each year on the basis of the measured data. It may therefore change from year to year.)

Road links with annual mean concentrations greater than $30.5 \ \mu g \ m^{-3}$ would be shaded red in **Figure 5-10**. No red shaded road links are visible on the map; in 2016 there were no modelled or measured exceedances of the 24-hour limit value.

Figure 5-10 Urban Major Roads, Annual Mean Roadside PM₁₀ Concentration, 2016 (µg m⁻³) Figure 5-11 Annual Mean Background PM₁₀ Concentration, 2016 (µg m⁻³)



5.3.2 PM₁₀ Changes Over Time

Figure 5-12 shows de-seasonalised trends in ambient PM₁₀ concentration, based on 11 urban background AURN sites, all of which have been operating since at least 1996. The sites are; Belfast Centre, Cardiff Centre, Learnington Spa, Leeds Centre, London Bloomsbury, London North Kensington, Middlesbrough, Newcastle Centre, Nottingham Centre, Southampton Centre and Thurrock. All 11 sites show a downward trend for PM₁₀ over their period of operation, highly statistically significant (at the 0.001 confidence level).

Figure 5-12 De-seasonalised Trends in Ambient PM₁₀, 11 Long-Running Urban Background AURN Sites 1992 - 2016



De-seasonalised PM_{10} trends for the period 1992 to 2016

Figure 5-13 shows de-seasonalised trends in ambient PM₁₀ concentration, based on 11 urban traffic AURN sites. There are few very long-running urban traffic sites: only three began operation before 2008. The sites shown here are the 11 that have been operating since the start of 2009 or earlier. The sites are; Armagh Roadside, Camden Kerbside, Carlisle Roadside, Chepstow A48, Leeds Headingley Kerbside, London Marylebone Road, Sandy Roadside, Stanford-le-Hope Roadside, Stockton-on-Tees Eaglescliffe, Swansea Roadside and York Fishergate.

Although most of these sites show a statistically significant downward trend in PM₁₀ concentration over this period, not all do: Stanford-le-Hope Roadside and Stocktonon-Tees Eaglescliffe show no significant trend, and Swansea Roadside shows a statistically significant increase. As in the case of NO₂, it is possible that trends at urban traffic sites are influenced by changes in the volume and type of traffic on the adjacent road.

Figure 5-13 De-seasonalised Trends in Ambient PM₁₀, 11 Long-Running Urban Traffic AURN Sites 2009 – 2016



De-seasonalised PM10 trends for the period 2009 to 2016

Figure 5-14 shows how the UK's total emissions of PM₁₀ have decreased over the years in which the AURN has been in operation. Total PM₁₀ emissions have steadily decreased over all this period, although in more recent years the rate of decrease has slowed, flattening off after around 2009. However, estimated emissions from road traffic alone have continued to decrease (**Figure 5-15**).

Figure 5-14 Estimated Annual UK Emissions of PM₁₀ (kt), 1992 – 2015 Source: NAEI



PM10 (Particulate Matter < 10μm) (kilotonne)

Figure 5-15 Estimated Annual UK Emissions of PM₁₀ from Road Transport (kt), 2009 – 2015 Source: NAEI



PM10 (Particulate Matter < 10µm) (kilotonne)

(Please note, the colours used in these bar charts are selected automatically and the colour used for road transport here is different from that in the previous graph).

5.4 PM_{2.5} Particulate Matter

5.4.1 PM_{2.5}: Spatial Distribution

Figure 5-16 shows the modelled annual mean urban roadside $PM_{2.5}$ concentrations in 2016. No roadside locations had annual means greater than the target value of 25 μ g m⁻³; even in London, the highest were in the range 15 - 20 μ g m⁻³.

Figure 5-17 shows modelled annual mean background $PM_{2.5}$ concentrations in 2016. Modelled concentrations were in the range 6-10 µg m⁻³ throughout most of England and Wales; concentrations were lower in most parts of Scotland and Northern Ireland. The areas with the highest modelled concentrations for 2016 were London, and the cities of the East and West Midlands; these areas had modelled concentrations greater than 10 µg m⁻³. Also visible are the effects of some major road routes in the middle of the country.

Figure 5-16 Urban Major Roads, Annual Mean Roadside PM_{2.5} Concentration, 2016 (µg m⁻³)

Figure 5-17 Annual Mean Background PM_{2.5} Concentration, 2016 (μg m⁻³)



5.4.2 PM_{2.5}: Changes Over Time

Until 2008, routine monitoring of PM_{2.5} within the AURN was confined to a small number of sites in London. Therefore, in this report, trend analysis for PM_{2.5} concentrates on years 2009 onwards, during which PM_{2.5} monitoring has been widespread.

Figure 5-18 shows trends in PM_{2.5} concentration at 12 long-running urban background AURN sites, 2009-2016. The majority (11 of the 12 sites) show a statistically significant downward trend, however in two of these cases (Newcastle Centre and Southampton Centre) the confidence interval of the trend includes zero so the trend should be treated with caution. Plymouth Centre shows no significant trend.

Figure 5-18 De-seasonalised Trends in Ambient PM_{2.5} Concentration, 12 Long-Running Urban Background AURN Sites 2009-2016



Figure 5-19 shows trends over the same period for PM_{2.5} at 12 urban traffic AURN sites. The majority (10 out of the 12 sites) show decreasing trends, of varying magnitude and level of significance, over this period. However, two do not: of these two (Stockton-on-Tees Eaglescliffe and Swansea Roadside), Swansea Roadside is notable as it has a highly significant *increasing* trend in PM_{2.5} concentration.

Figure 5-19 De-seasonalised Trends in Ambient PM_{2.5} Concentration, 12 Long-Running Urban Traffic AURN Sites 2009-2016



Finally, **Figure 5-20** shows the estimated annual emission of $PM_{2.5}$, from 1992 to 2015. The graph shows that emissions have decreased in a similar manner to emissions of PM_{10} , with a steady decrease from the early 1990s, a clear levelling off, and no further consistent decrease after around 2009.

Figure 5-20 Estimated Annual UK Emissions of PM_{2.5} (kt), 1992 – 2015. Source: NAEI



PM2.5 (Particulate Matter < 2.5um) (kilotonne)

5.5 Benzene

5.5.1 Benzene: Spatial Distribution

Benzene is found in petrol and in vehicle emissions, therefore elevated levels may be expected at roadside locations.

Figure 5-21 shows modelled annual mean benzene concentrations at roadside locations in 2016. **Figure 5-22** shows the modelled annual mean background concentrations of benzene in 2016. Modelled background concentrations were below 0.5 μ g m⁻³ over most of the UK, with marginally higher concentrations for most urban areas. A few small areas, for example in the Midlands and Humberside, had concentrations in excess of 1 μ g m⁻³. However, background concentrations everywhere were well below the limit value of 5 μ g m⁻³ for benzene.

Figure 5-21 Urban Major Roads, Annual Mean Roadside Benzene Concentration, 2016 (µg m⁻³)

Figure 5-22 Annual Mean Background Benzene Concentration, 2016 (µg m⁻³)



5.5.2 Benzene: Changes Over Time

Figure 5-23 shows a smoothed trend plot based on the combined dataset from 15 long-running sites in the Non-Automatic Hydrocarbon Network, which have operated since 2002. These are: Barnsley Gawber, Belfast Centre, Grangemouth, Haringey Roadside, Leamington Spa, Leeds Centre, Liverpool Speke, London Bloomsbury, Manchester Piccadilly, Middlesbrough, Newcastle Centre, Nottingham Centre, Oxford Centre Roadside, Southampton Centre and Stoke on Trent Centre.

The smoothed trend plot for these 15 sites shows a slight increase from 2002 to 2004, followed by a steep decrease between 2004 and 2008. From then on, the graph is much flatter, showing little further fall in ambient concentrations of benzene until 2014 when there is a slight rise. In the past three years, there has been a further slight decrease.

Figure 5-23 Smoothed Trend Plot of Ambient Benzene Concentration, 15 Long-Running Non-Automatic Sites



Figure 5-24 shows the estimated total annual UK emission of benzene (in kilotonnes). The data have been taken from the NAEI (which does not provide a bar chart for benzene). The estimated annual emissions also appear to have decreased over period shown – although more steadily than the average measured ambient concentration.



Figure 5-24 Estimated Annual UK Emissions of Benzene (kt), 2002 – 2015 (data from NAEI)

5.61,3-Butadiene

5.6.1 1,3-Butadiene: Compliance with AQS Objective

The ambient concentration of 1,3-butadiene is not covered by any EU Directives so modelled maps are not routinely produced for this pollutant. However, it is the subject of a UK Air Quality Strategy objective of 2.25 µg m⁻³, as a maximum running annual mean, to have been achieved by 31st December 2003. This objective was met throughout the UK by the due date.

The Automatic Hydrocarbon Network monitors ambient concentrations of 1,3butadiene at its four sites. There is one urban background site (London Eltham), one urban traffic site (London Marylebone Road) and two rural sites: Auchencorth Moss in Lothian, and Chilbolton Observatory in Hampshire. Chilbolton Observatory replaced a previous rural site in Harwell (Oxfordshire) site at the beginning of 2016. Concentrations of this pollutant at all four sites are low: the highest annual mean in 2016 was measured 0.12 μ g m⁻³. Surprisingly, this was measured at one of the two rural sites, Chilbolton Observatory. However, the running annual means at all four sites were well within the Air Quality Strategy objective (2.25 μ g m⁻³) in 2016.

5.6.2 1,3-Butadiene: Changes Over Time

Figure 5-25 shows trends in ambient 1,3-butadiene concentration at the two London sites only, between 2000 and 2016. Concentrations are higher at London Marylebone Road, reflecting its roadside location. Both the London sites show a highly significant downward trend in this pollutant, though in the case of London Eltham it is extremely small (less than 0.01 units per year in magnitude) Auchencorth Moss – not plotted – does not show any significant trend, and Chilbolton Observatory has not been in operation long enough for trends to be evaluated.



Figure 5-25 Trends in Urban 1,3-Butadiene Concentration, 2000-2016

Figure 5-26 shows the total estimated UK annual emission of this compound, in kilotonnes. This appears to have decreased steadily since 2000. The main source of 1,3-butadiene is vehicle emissions, and the use of catalytic converters since the early 1990s has substantially reduced emissions from this source.



Figure 5-26 Estimated Annual UK Emissions of 1,3-Butadiene (kt), 2000 – 2015 (data from NAEI)

5.7 Carbon Monoxide

5.7.1 CO: Spatial Distribution

Ambient concentrations of CO throughout the UK have been well within the limit value for many years, Therefore, since 2010, maps have no longer been routinely produced for CO.

5.7.2 CO: Changes over time

Because concentrations of CO are well within the limit value, relatively few monitoring sites are required. Seven urban AURN sites currently monitor this pollutant, of which six (Belfast Centre, Cardiff Centre, Edinburgh St Leonards, Leeds Centre, London Marylebone Road and London North Kensington) have operated for at least 10 years.

Figure 5-27 shows de-seasonalised trends at these six long-running AURN sites, from 1992 to 2016. All six show a highly significant downward trend over

the period. **Figure 5-28** shows the estimated annual emissions of CO over the same period: a steady, almost linear year-on-year decrease to 2011 is followed by two years in which estimated emissions remained stable, before decreasing further in 2014 and 2015. The decreasing ambient concentrations reflect declining emissions over the last two decades. UK emissions of this pollutant have decreased substantially over recent decades. The NAEI attributes this decrease to *"significant reductions in emissions from road transport, iron and steel production and the domestic sector"*.⁵¹

Figure 5-27 De-seasonalised Trends in CO Concentration, 6 Long-Running AURN Sites 1992-2016



De-seasonalised CO trends for the period 1992 to 2016

Figure 5-28 Estimated Annual UK Emissions of CO (kt), 1992 – 2015 Source: NAEI



Carbon Monoxide (kilotonne)

5.8 Ozone

5.8.1 O₃: Spatial Distribution

Figure 5-29 shows the average number of days per year with ozone concentration > 120 μ g m⁻³, over the **three** years 2014-2016. This was less than six days everywhere apart from a small area of East Anglia. **Figure 5-30** shows the same statistic, for 2016 only (i.e. not averaged over three years). Most of the UK had less than six days above 120 μ g m⁻³ in 2016, with the exception of two areas: a band across the north of England, and a large area of north and mid Wales.

Figure 5-31 shows the AOT40 statistic, averaged over the past **five** complete years, 2012-2016. The AOT40 statistic (expressed in μ g m⁻³.hours) is the sum of the difference between hourly concentrations greater than 80 μ g m⁻³ (= 40 ppb) and 80 μ g m⁻³ over a given period using only the one-hour values measured between 0800 and 2000 Central European Time each day. Highest AOT40 values are seen in a wide band running roughly diagonally across the south of the UK, roughly from the Wash to Cornwall.

Figure 5-32 shows the same statistic, for 2016 only. The highest concentrations occurred in two areas: the north east of England and a large area of central Wales. This pattern is different from that seen in previous recent years' versions of this map, when the model has typically shown highest ozone concentrations across the south east and east of England.

Figure 5-29 Average no. of days with O₃ Concentration > 120 μ g m⁻³ 2014-2016

Figure 5-30 Days with O₃ Concentration > 120 μ g m⁻³, 2016







Figure 5-31 Average AOT40, 2012-2016 (µg m⁻³.hours)

Figure 5-32 Average AOT40, 2016 (µg m⁻³.hours)



5.8.2 O₃: Changes Over Time

Figure 5-33 shows a trend plot of maximum daily 8-hour mean ozone concentrations at 10 long-running rural AURN sites (Aston Hill, Bush Estate, Eskdalemuir, High Muffles, Ladybower, Lough Navar, Lullington Heath, Sibton, Strathvaich and Yarner Wood). Rural sites have been chosen because concentrations of ozone are typically highest in rural areas.

One site (Lullington Heath) shows a highly significant negative trend. One site (Bush Estate) shows a positive trend over this period, though it is significant only at the 10 % (0.1) level and the confidence interval of the slope includes zero. The remaining eight show no statistically significant trend. There is therefore no consistent pattern of upward or downward trends at these rural sites.

Figure 5-33 Trends in Daily Maximum 8-hour Ozone Concentration at 10 Long-Running Rural AURN sites, 1992 - 2016.



Ozone is not emitted in significant quantities directly from any source in the UK (instead, it is formed from reactions involving other pollutants). Ozone is therefore not included in the NAEI, and trends in ozone emissions are not covered by this report.

5.9 Metallic Elements

5.9.1 Metallic Elements: Spatial Distribution

Figure 5-34, **Figure 5-35**, **Figure 5-36** and **Figure 5-37** show modelled annual mean concentrations of lead (Pb), arsenic (As), cadmium (Cd) and nickel (Ni) respectively in 2016. The spatial distribution patterns are discussed below.

Pb: background concentrations were less than 10 ng m⁻³ (0.01 μ g m⁻³) over most of the UK, and well within the limit value of 500 ng m⁻³ (0.5 μ g m⁻³) throughout. There were some small areas (mostly urban, industrial areas) where higher concentrations were modelled: however, the 2016 map shows only the faintest traces of the UK's major roads (which have been evident in previous years' maps).

As: this toxic element is not strictly a metal but is measured by the UK Metals Network. Background concentrations were less than 6.0 ng m⁻³ over the whole UK, and less than 1.9 ng m⁻³ over most of the country. However, concentrations of 1.9 ng m⁻³ and above occurred in some areas – particularly the north-eastern part of England, Yorkshire and Humberside. This pattern reflects the natural sources of airborne arsenic, particularly wind-blown dust. Modelled concentrations were therefore highest in areas where agricultural practices give rise to wind-blown dust (such as parts of eastern England) and where the natural arsenic content of the soil is relatively high (such as parts of Cornwall).

Cd: background concentrations were less than 0.5 ng m⁻³ over almost all of the UK. Some major road routes are visible: this is due to re-suspended road dust. Also there are some small areas with concentrations in the range 0.6 - 5.0 ng m⁻³, relating to specific point sources. Please note that the scale used for Cd concentrations was changed in the 2010 report in this series, reflecting the decrease observed in ambient concentrations over recent years.

Ni: background concentrations of Ni were typically less than 2 ng m⁻³ (well away from urban areas, usually less than 1 ng m⁻³). Some major road routes are visible in the map; like lead, nickel is found in suspended road dust. There are also some small areas with higher concentrations due to industrial activity. Two monitoring sites (in Swansea Urban Area and Sheffield Urban Area) reported an annual mean higher than the target value of 20 ng m⁻³ in 2016 but this is not captured in this background modelling.

Figure 5-34 Annual Mean Background Lead Concentration, Figure 5-35 Annual Mean Background Arsenic 2016 (µg m⁻³)



Concentration, 2016 (ng m⁻³)



Figure 5-36 Annual Mean Background Cadmium Concentration, 2016 (ng m⁻³)



Figure 5-37 Annual Mean Background Nickel Concentration, 2016 (ng m⁻³)



5.9.2 Lead: Changes Over Time

Figure 5-38 shows a time series of annual mean concentration of Pb in the PM₁₀ particulate fraction, as measured from 2004 by urban sites in the UK Metals Network and its predecessors, as described in Section 3.3.2. (Prior to 2004, Pb in the particulate phase was measured by the six sites comprising the former Multi-Element Network. For further information on this, please see earlier reports in this series. However please note that the sampling method used by the Multi-Element Network was not size-selective).

The annual mean of all urban sites in the UK Metals Network is shown: rural sites are not included. In 2016 there were 16 urban sites. The mean for all sites was well below the Air Quality Directive limit value for annual mean Pb, of 500 ng m⁻³.

Figure 5-38 also shows NAEI estimated total annual UK emissions of this metal (plotted as a dotted line, against the right-hand y-axis). Measured ambient concentrations follow the same pattern, generally decreasing (though there is some year-to-year variation) until around 2012 when there appears to be some levelling off of the downward slope, before a further apparent decrease in 2016.





5.9.3 Arsenic: Changes Over Time

Figure 5-39 shows a time series of annual mean concentrations of arsenic (As) in the PM₁₀ fraction, as measured by the urban sites in UK Metals Network and its predecessors, described in Section 3.3.2. (For earlier, non-size selective measurements by the smaller Multi-Element Network, please see previous reports in this series.) The annual mean of all urban sites (of which there were 16 in 2016) is shown by the solid line – rural sites are not included. This parameter is well within the Fourth Daughter Directive target value of 6 ng m⁻³.

Also shown is the UK's estimated total annual emission of As (from the NAEI), in kilotonnes. This is plotted as a dotted line, against the right-hand y-axis. There appears to have been a slight increase in As emissions in 2012 and 2013, which may be reflected in subsequent measured ambient concentrations.





5.9.4 Cadmium: Changes Over Time

Figure 5-40 shows a time series of annual mean concentration of cadmium (Cd) in the PM₁₀ fraction as measured by the UK Metals Network and its predecessors, described in Section 3.3.2. (For earlier, non-size selective measurements from the Multi-Element Network, please see previous reports in this series.) The annual mean of all 16 urban sites is shown by the solid line.





Also shown (plotted as a dotted line, against the right-hand y-axis) is the UK's estimated total annual emission of Cd (in kilotonnes), from the NAEI. The variation in emissions appears to be reflected in the variation in measured ambient concentrations from year to year. In 2011, there was an increase in both the estimated total emission and in the average ambient Cd concentration. The ambient concentration also shows an even larger peak in 2014: although the emission estimates for that year do show some increase, it does not appear to be large enough to be consistent with the peak in ambient Cd.

However, despite the 2014 peak, ambient cadmium concentrations are very low, and well within the Fourth Daughter Directive target value of 5 ng m⁻³ at all sites.

5.9.5 Nickel: Changes Over Time

Figure 5-41 shows a time series of annual mean concentrations of nickel (Ni) in PM₁₀, as measured by urban sites in the UK Metals Network. As with the other metals, information on non-size selective measurements from the older Multi-Element Network can be found in previous reports in this series.



Figure 5-41 Ambient Urban Concentrations of Ni in PM₁₀, and Total Estimated UK Emissions

For the purposes of presenting a national trend, the graph is based on the average annual mean for all 16 urban sites measuring Ni, except one. Pontardawe Tawe Terrace (which began operation in 2011) has been excluded, as it measures ambient nickel concentrations very much higher than the other sites, and if included will dominate the mean for years 2011 onwards. (This site has measured exceedances of the Fourth Daughter Directive target value of 20 ng m⁻³ each year since it started up, in 2011.) Pontardawe Tawe Terrace was the source of the measured exceedance in the Swansea Urban Area, highlighted in **Table 4-7**: the measured annual mean Ni concentration at this site was 47 ng m⁻³. There was also a measured exceedance (annual mean 24 ng m⁻³) at Sheffield Tinsley and a modelled exceedance reported for the South Wales zone.

Figure 5-41 also shows total estimated annual UK emissions of Ni, from the NAEI (as a dotted line, plotted against the right-hand axis). The average measured ambient concentrations appear to generally reflect the year to year variation in

estimated total emissions, though 2014 saw a substantial increase in the measured average.

5.9.6 Mercury: Changes Over Time

At the end of 2013, the UK Metals Network ceased measurement of mercury in PM₁₀ particulate matter at all sites. This decision was taken because the majority of ambient Hg is in the vapour phase. Monitoring of Total Gaseous Mercury (TGM) was also scaled down, continuing at just two of the original sites: Runcorn Weston Point and London Westminster. Runcorn Weston Point is near an industrial installation (a chlor-alkali plant) that used mercury in the past, and measures ambient Hg concentrations an order of magnitude greater than any other sites in the network. London Westminster is an urban background site in central London. Mercury in the vapour phase was also measured during 2016 using the Tekran instrument (see section 3) at two rural sites: Chilbolton Observatory in Hampshire, and Auchencorth Moss in Lothian.

Measurement of TGM therefore continued at four sites: two where it is likely to be highest (Runcorn Weston Point and London Westminster) and two rural background sites, for the purpose of understanding transboundary contribution to ambient Hg concentration. The two original rural sites were Auchencorth Moss, and Harwell in Oxfordshire, until the latter closed at the end of 2015 and was replaced by Chilbolton Observatory.

For information on the measurements of total mercury (TGM plus particulate phase) taken at urban sites in the UK Metals Network and its predecessors from 2004 to 2013, please refer to "*Air Pollution in the UK 2014*" and earlier reports in this series. The present report, and future reports in this series, will focus only on TGM, and on the four sites which have continued monitoring this element through 2016 and beyond: Auchencorth Moss, Chilbolton Observatory, London Westminster and Runcorn Weston Point.

Figure 5-42 shows annual mean concentrations of TGM at the four monitoring sites, from 2004 (when the UK Metals Network began operation) to 2016. It can clearly be seen that the measured annual mean concentrations of Hg at Runcorn Weston Point are an order of magnitude higher than those measured at the two rural sites. However, Hg concentrations at this industrial site appear to have decreased over the past 10 years (though not consistently or steadily). Mercury concentrations at London Westminster do not appear to have decreased: by contrast, they have been slightly higher in recent years than they were a decade ago. Neither of the two rural sites show any consistent pattern of increase or decrease (Harwell is shown in this graph as no mercury data are so far available for Chilbolton Observatory).



Figure 5-42 Measured Urban Concentrations of Total Gaseous Hg, and Total Estimated UK Emissions

5.10 Benzo [a] Pyrene

5.10.1 B[a]P: Spatial Distribution

Figure 5-43 shows the modelled annual mean background concentration of B[a]P. The areas of highest concentration reflect the distribution of industrial sources, and areas where there is widespread domestic use of oil and solid fuels for heating. This has historically included the Belfast area and other urban parts of Northern Ireland: also parts of Yorkshire, Humberside and South Wales.

Recent reports in this series have shown that these areas have reduced in recent years, particularly in Northern Ireland. However, this appears to have reversed slightly in 2016, and once again the UK is reporting areas in Northern Ireland and elsewhere with modelled or measured annual mean B[a]P concentrations in excess of 1 ng m⁻³.

Figure 5-43 Annual mean background B[a]P concentration, 2016 (ng m⁻³)



5.10.2 B[a]P: Changes Over Time

The PAH monitoring network began operation in 1991, comprising a small number of sites, and was increased to over 20 in the late 1990s. However, during the years 2007-2008, the network underwent a further major expansion and re-organisation, including a change of sampling technique.

The newer sampling technique used at most sites from 2008 onwards (the "Digitel" PM_{10} sampler) has been found to give higher results than the older method. The reason for this is likely to be due to a number of factors, predominantly the fact that the new samplers have a shorter collection period. The shorter collection period is likely to decrease the degradation of the PAHs by ozone or other oxidative species⁵².

Because of these changes in the composition of the network, and in particular the techniques used, temporal variation in PAH concentrations has only been analysed from 2008 in this report.

Figure 5-44 shows how the average annual mean B[a]P concentration has changed in the years since 2008. This graph is based on the average of all sites in the PAH

Network: the composition of this network has changed little since 2008 so it is considered appropriate here to use the average of all sites. Following a marked drop in average measured concentrations of B[a]P between 2008 and 2009, average B[a]P concentrations then appear to have remained generally stable until 2014. Further decreases in the average occurred in 2015 and 2016.



Figure 5-44 Ambient Concentrations of Particulate Phase Benzo[a]pyrene, and Total UK Emissions

Figure 5-44 also shows the estimated total annual UK emission of B[a]P (in kg), from the NAEI (shown as a dotted line and plotted against the y-axis on the right). This indicates that emissions have increased since 2008. Emissions of B[a]P at the present time are dominated by domestic combustion of coal, and the NAEI data indicate that it is this source (residential/commercial/institutional) that is increasing.

However, to put this into context, estimated total UK emissions of B[a]P have decreased substantially in recent decades and are an order of magnitude lower than in the early 1990s (**Figure 5-45**). According to the NAEI, most of this reduction is due to decreasing emissions from industry, and the banning of stubble burning in 1993.

Figure 5-45 Estimated Annual UK Emissions of Benzo[a]Pyrene (kg), 1992 – 2015 Source: NAEI



Benzo[a]pyrene (kg)
6 Pollution Events in 2016

6.1 Winter and Spring Particulate Pollution Episodes

Elevated pollution concentrations can occur during winter and spring months when the weather is cold and still. In 2016, there were several periods of elevated particulate pollution recorded by the Automatic Urban and Rural Monitoring Network (AURN) throughout sites across the UK. These episodes occurred in mid-January, mid-March, and again at the end of the year.

6.1.1 Mid-January 2016

Between 19th and 21st January, a spell of cold, still weather occurred across the south of England (Met Office⁵³). This allowed local emissions of nitrogen oxides (NO_X) and particulate matter (PM) to build up in the region. Figure 6-1 shows the episode affected the East Anglia area most severely, but also spread throughout the rest of south and central England and South Wales. The maps in Figure 6-1 are based on the Daily Air Quality Index (DAQI) which has been designed to inform the public about levels of air pollution and provides recommended actions and health advice. The index is numbered 1-10 and divided into four bands, 'Low' (1) to 'Very High' (10), to provide detail about air pollution levels in a simple way⁵⁴.

Figure 6-1 Maps Showing the DAQI between 18th and 22nd January 2016 (Source: UK-AIR)



1 2 3 4 5 6 7 8 9 10 Low Moderate High Very High Using air mass tracking techniques, it is possible to model the paths taken by air masses arriving over the UK at a particular time; these are known as 'back trajectories' and can be shown on a map. Figure 6-2 shows 96-hour back trajectories for 20th January: these show relatively slow-moving air masses circulating over the south and east of the UK.

Figure 6-2 Back Trajectories Showing Air Masses Arriving in the UK on 20th January 2016 (Source: HYSPLIT trajectory model^{55,56} NOAA website⁵⁷) *The authors*

gratefully acknowledge the NOAA Air Resources Laboratory (ARL) for the provision of the HYSPLIT transport and dispersion model and READY website (<u>http://www.ready.noaa.gov</u>) used to prepare this figure.



6.1.2 Mid-March 2016

A particulate pollution episode was also measured by the AURN during mid-March 2016, specifically between the 11th and 13th March. Pollution levels reached 'Moderate' throughout England and much of Wales, with some areas including Yorkshire & Humberside and Greater London, reaching 'High' and 'Very High' levels of pollution (**Figure 6-3**) This pollution episode was caused by low wind speeds and an influx of air from northern Europe. Measurement and analysis of pollution in London (by King's College London, the managers of the London Air Quality Network) showed that this episode was dominated by pollution from Germany and the

Netherlands, and was most likely from a mix of industrial, agricultural and urban sources⁵⁸. The air masses can also be tracked to northern Europe using back trajectories, as shown in **Figure 6-4**. A high-pressure system from the 13th brought dry and settled weather⁵⁹ which also allowed local air pollution to accumulate.





Figure 6-4 Back Trajectories Showing Air Masses Arriving in the UK on 16th March 2016 (Source: NOAA website) The authors gratefully acknowledge the NOAA Air Resources Laboratory (ARL) for the provision of the HYSPLIT transport and dispersion model and READY website (http://www.ready.noaa.gov) used to prepare this figure.



The peak of the mid-March episode was seen on the 12^{th} March; this is evident in **Figure 6-3** (which shows that DAQI values in the 'high' and 'very high' bands were most widespread on this day), and also in **Figure 6-5** and **Figure 6-6**, which show PM₁₀ and PM_{2.5} concentrations, respectively. A slight change in air path and increase in wind speeds brought cleaner air on 13^{th} March⁵⁸ and by 14^{th} March, pollution levels had returned to low. (**Figure 6-6** shows that one site measured an hourly mean PM₁₀ of nearly 180 µg m⁻³ at 3pm on 11^{th} March. This site was Stanford-le-Hope Roadside, in Essex. The reason for this high measurement is not known: it was most likely from a localised source or short-term activity nearby such as roadworks or a bonfire).

Figure 6-5 Daily Mean PM_{10} Concentrations in the UK between 9th and 16th March 2016





Figure 6-6 Hourly Mean PM_{2.5} Concentrations in the UK between 9th and 16th March 2016

6.1.3 Late November and December 2016

Towards the end of 2016, late November and December saw a number of pollution episodes due to cold, settled weather (**Figure 6-7** and **Figure 6-8**).

A prolonged winter smog event affected much of the UK from the end of November until around the 6th December. Northern Ireland was affected by 'Very High' levels of PM on the 25th, 24th and 26th November which coincided with light winds, fog and low overnight temperatures. 'Moderate' PM was also measured widely across northern England and Scotland between these dates. Widespread 'Moderate' PM was then measured across central England on 29th November and reached 'High' and 'Very High' levels across southern England and London on 30th. Again the increases can be linked to the cold, still and foggy weather conditions allowing the build-up of local pollutant emissions. On 1st December, a north-west airflow with a slight increase in wind speed helped reduce pollution levels; however, the increased wind speed was not enough to completely disperse local emissions. The episode continued on 2nd December, but temporarily subsided on the 3rd and 4th due to an easterly wind bringing cleaner air from the North Sea⁶⁰. On 5th and 6th December, light easterly winds imported air from France, which combined with poorly dispersed local emissions, resulted in further widespread NO₂ and particulate pollution⁶⁰.

Figure 6-8 clearly shows the impact of Christmas on NO₂ caused by lower traffic on the roads. This is followed by an increase in PM₁₀ (**Figure 6-7**), NO₂ (**Figure 6-8**) and in PM_{2.5} emissions. This was caused by cold, foggy, calm and settled weather conditions with poor dispersion of local emissions coming from traffic and wood burning⁶¹, which is particularly common between Christmas and New Year.



Figure 6-7 Daily Mean PM_{10} Concentrations in the UK between 22^{nd} November and 31^{st} December 2016

Figure 6-8 Hourly measured NO₂ Concentrations in the UK between 22nd November and 31st December



6.2 Summer Ozone and Particulate Matter Events

Air pollution episodes due to ozone commonly occur in the UK during late spring and summer. In 2016, such ozone pollution episodes occurred in early May, mid-July, and mid-September. Some events were also accompanied by elevated particulate pollution.

6.2.1 Early May 2016

Elevated ozone levels were measured by the AURN between the 5th and 9th May 2016 (**Figure 6-9**). This ozone episode was caused by a combination of strong sunshine which lasted throughout the day, high ambient temperatures, and air arriving in the UK from continental Europe; this air had passed over industrialised areas and therefore contained the 'precursor pollutants' required for the photochemical reactions to produce ozone. The peak of this episode was on the 8th May (**Figure 6-9**). By 10th May, cloudier conditions led to a decrease in the solar radiation required for ozone formation.



Figure 6-9 Ozone Daily Max 8-Hour Running Mean Concentrations from 2nd to 12th May 2016

As well as ozone, this event also saw the accumulation of particulate pollution. Fine and settled conditions between the 6th and 7th May, along with light south-easterly winds, resulted in the import of a mix of industrial, agricultural, urban and traffic related pollution from the continent, to combine with UK emissions⁶². A change in the path of incoming air later on Saturday 7th May resulted in relatively cleaner air from the Mediterranean being imported and an end to the elevated particulate episodes of Friday 6th May and Saturday 7th May⁶².

6.2.2 Mid-late July 2016

Between 19th and 20th July 2016, high levels of sunshine, along with light southerly wind bringing pollution from the continent, allowed ozone to be formed, and to build up. 'Moderate' and 'High' levels of pollution were recorded across the UK, with the worst affected area being Eastern England, on the 19th July. Following the development of a cleaner Atlantic air flow and increasing cloud cover on 21st July, ozone concentrations returned to normal levels⁶³.

6.2.3 Mid-September 2016

The 13th September 2016 saw the UK's highest September temperatures since 1911⁶⁴. Due to strong sunshine at this time, south-easterly air flow importing

emissions from France, and low wind speeds allowing local air pollution to build, 'Moderate' levels of ozone and PM occurred across much of the UK (Figure 6-10).



Figure 6-10 Ozone Daily Max 8-Hour Running Mean Concentrations from 10th to 20th September 2016

6.3 Bonfire Night Particulate Pollution Event

It is common for occasions such as Bonfire Night (5th November) to produce significant short term particulate pollution episodes as a result of bonfires and firework displays. These pollution episodes depend greatly on weather conditions, and whether the celebration falls on a weekday or a weekend. Therefore, they vary significantly from year to year. Wet, windy or unsettled conditions are more likely to result in low concentrations; cold and still weather is more likely to result in elevated levels of PM₁₀ particles and other pollutants as emissions are not effectively dispersed. In recent years, Bonfire Night particulate episodes have been noticeable but small (the exception being 2014, when the UK had its most significant Bonfire Night episode since 2006).

In 2016, 'Moderate' and 'High' levels of PM₁₀ and PM_{2.5} were measured across London during the weekend of Saturday 29th and Sunday 30th October. This coincided with the Hindu festival of Diwali (frequently celebrated with fireworks), which in 2016 was from Sunday 30th October to Thursday 3rd November. Hourly

mean PM₁₀ concentrations of 224 μ g m⁻³ were measured at London Harlington during the evening of 30th October (**Figure 6-11**).

The following week included Halloween on Monday 31st October (also an occasion for fireworks in some areas), and more significantly, Bonfire Night, the following Saturday, 5th November. Low wind speeds over the weekend of 5th – 6th November allowed the build-up of local pollution in some areas from wood burning and fireworks. As well as the local pollution, volatile particulate concentrations and particulate nitrate, which suggest long-range transport, increased over the weekend. The back trajectory models show that air travelled over parts of Germany, Belgium and France before reaching London⁶⁵. Widespread 'Moderate' PM continued on 31st October, with imported air from mainland Europe coinciding with poor dispersion conditions (light winds, fog and low overnight temperatures). **Figure 6-11** shows hourly mean PM₁₀ concentrations averaged over all AURN sites in Greater London, over the period 15th October to 15th November 2016.

Figure 6-11 Hourly Mean PM₁₀ Concentrations in the UK between 15th October and 15th November 2016, for AURN Sites in Greater London



Pollution episodes due to Bonfire Night and other events can be very localised, depending on local variations in the weather over this period, the timing of local fireworks and bonfires, and their proximity to monitoring sites. Other parts of the UK showed different patterns over this period: the East and West Midlands showed several periods of elevated PM₁₀ concentrations between 30th October and 9th November (**Figure 6-12**), while in Wales and the north of England (**Figure 6-13**,

Figure 6-14) PM concentrations were generally lower but some sites recorded a sharp Bonfire Night peak on 5th November. By contrast, other regions such as Scotland (**Figure 6-15**) had no elevated PM concentrations over this period.

Figure 6-12 Hourly Mean PM₁₀ Concentrations in the UK between 15th October and 15th November 2016, for AURN Sites in the Midlands



Figure 6-13 Hourly Mean PM₁₀ Concentrations in the UK between 15th October and 15th November 2016, for AURN Sites in Wales



Figure 6-14 Hourly Mean PM₁₀ Concentrations in the UK between 15th October and 15th November 2016, for AURN Sites in the North of England



Hourly Mean $\rm PM_{10}$ Concentrations, AURN Sites in North of England, 15^{th} Oct - 15^{th} Nov 2016

Figure 6-15 Hourly Mean PM₁₀ Concentrations in the UK between 15th October and 15th November 2016, for AURN Sites in Scotland



6.4 Forecasting Air Pollution Events

As outlined in Section 2.2.4, air quality forecasts are produced by the Met Office using the model AQUM (Air Quality in the Unified Model). AQUM combines predicted meteorological conditions with gridded emissions data to provide forecasts of air pollution concentrations across the UK. Forecasts are produced with a lead time of up to 5 days and are reported using the Daily Air Quality Index (DAQI) to align with associated health advice.

Forecast performance in any particular episode can vary based on a number of factors, including the pollutant type responsible for the episode, the accuracy and complexity of the meteorological conditions modelled by AQUM and the validity of the assumptions made with regard to pollutant emissions.

AQUM uses emissions based on the National Atmospheric Emissions Inventory (NAEI) which presents annual mean emissions at 1km² resolution. Appropriate temporal and vertical profiles (e.g. based on well-established traffic patterns) are applied within AQUM to each emission sector. In the majority of cases this approach provides an appropriate representation of emissions at different times of day and at different times during the year.

7 Where to Find Out More

Defra's web pages relating to air quality can be found at

https://www.gov.uk/government/policies/protecting-and-enhancing-our-urban-andnatural-environment-to-improve-public-health-and-wellbeing/supportingpages/international-european-and-national-standards-for-air-guality

These provide details of what the UK is doing to tackle air pollution, and the science and research programmes in place.

Also, Defra has published a Guide to Air Pollution Information Resources, detailing the types of information that are made available and this can be found at <u>https://uk-air.defra.gov.uk/assets/documents/reports/cat14/1307241318</u> Guide to UK Air Pol <u>lution Information Resources.pdf</u>.

Information on the UK's air quality, now and in the past, is available on UK-AIR, the Defra online air quality resource at http://uk-air.defra.gov.uk/. UK-AIR is the national repository for historical ambient air quality data. It contains measurements from automatic measurement programmes, some dating back to 1972, together with non-automatic sampler measurements dating back to the 1960s. The data archive brings together into one coherent database both data and information from all the UK's measurement networks. New tools recently added to UK-AIR include the UK Ambient Air Quality Interactive Map at http://uk-air.defra.gov.uk/data/gis-mapping that allows you to look at outputs for the national modelling conducted for compliance assessment, based on pollutant, background or roadside and geographical location.

Similar national online air quality resources have also been developed for Scotland, Wales and Northern Ireland:

- The Welsh Air Quality Archive at www.welshairquality.co.uk
- The Scottish Air Quality Archive at <u>www.scottishairquality.co.uk</u>
- The Northern Ireland Archive at <u>www.airqualityni.co.uk</u>

Together, these four national websites provide a comprehensive resource for data and analyses covering all aspects of air quality throughout the UK and all its regions.

The Devolved Administrations each produce their own short annual report, providing more specific information on air quality in their regions. These reports are available from the above websites.

UK-AIR also provides a daily air quality forecast, which is further disseminated via email, RSS feeds and Twitter (see <u>http://uk-air.defra.gov.uk/twitter</u>). Latest forecasts are issued daily, at <u>http://uk-air.defra.gov.uk/forecasting/</u>. Detailed pollutant emission data for the UK are available from the National Atmospheric Emissions Inventory (NAEI) at <u>www.naei.org.uk</u>.

Additional information from the Devolved Administrations of Scotland, Wales and Northern Ireland can be found at:

- The Scottish Government Air Quality web page at <u>http://www.scotland.gov.uk/Topics/Environment/waste-and-pollution/Pollution-1/16215</u>
- The Welsh Government Environment Air Quality web page at <u>http://gov.wales/topics/environmentcountryside/epq/airqualitypollution/airq</u> <u>uality/?lang=en</u>
- The Northern Ireland Department of Agriculture, Environment and Rural Affairs (DAERA) web page at https://www.daera-ni.gov.uk/

References

Section 1

¹ European Parliament and Council of the European Union (2008) '*Council Directive* on ambient air quality and cleaner air for Europe (2008/50/EC)'. [online]. Available at <u>http://eur-lex.europa.eu/LexUriServ/LexUriServ.do?uri=CELEX:32008L0050:EN:NOT</u> (Accessed 01 Aug 2017).

² European Parliament and Council of the European Union (2004) '*Directive* 2004/107/EC of the European Parliament and of the Council of 15 December 2004 relating to arsenic, cadmium, mercury, nickel and polycyclic aromatic hydrocarbons in ambient air'. [online]. Available at <u>http://eur-lex.europa.eu/LexUriServ/LexUriServ.do?uri=CELEX:32004L0107:EN:NOT</u>, (Accessed 01 Aug 2017).

Section 2

³ Air Quality Expert Group (2013) '*Mitigation of United Kingdom PM*_{2.5} *Concentrations*'. Report prepared for Defra. [online]. Available at <u>https://ukair.defra.gov.uk/assets/documents/reports/cat11/1508060903 DEF-</u> <u>PB14161 Mitigation of UK PM25.pdf</u>. (Accessed 01 Aug 2017).

⁴ UK Government *'UK Air Quality Standards Regulations 2010'*. [online]. Available at <u>http://www.legislation.gov.uk/uksi/2010/1001/contents/made</u> (Accessed 01 Aug 2017).

⁵ Scottish Government '*The Air Quality Standards (Scotland) Regulations (2010)*'. [online]. Available at <u>http://www.legislation.gov.uk/ssi/2010/204/made</u> . (Accessed 01 Aug 2017).

⁶ Welsh Government '*The Air Quality Standards (Wales) Regulations 2010*' [online]. Available at <u>http://www.legislation.gov.uk/wsi/2010/1433/contents/made</u> (Accessed 01 Aug 2017).

⁷ Department of Environment, Northern Ireland *'The Air Quality Standards Regulations (Northern Ireland) 2010'.* [online]. Available at <u>http://www.legislation.gov.uk/nisr/2010/188/contents/made</u> (Accessed 01 Aug 2017).

⁸ European Parliament and Council of the European Union (2015) 'Commission Directive (EU) 2015/1480 of 28 August 2015 amending several annexes to Directives 2004/107/EC and 2008/50/EC of the European Parliament and of the Council laying down the rules concerning reference methods, data validation and location of sampling points for the assessment of ambient air quality'. Available at http://eur-lex.europa.eu/legal-content/en/TXT/?uri=CELEX:32015L1480 (Accessed 16 Aug 2017). ⁹ European Parliament and Council of the European Union (2011) *'Commission* Implementing Decision of 12 December 2011 laying down rules for Directives 2004/107/EC and 2008/50/EC of the European Parliament and of the Council as regards the reciprocal exchange of information and reporting on ambient air quality (2011/850/EU)' [online]. Available at <u>http://eur-</u>

lex.europa.eu/LexUriServ/LexUriServ.do?uri=OJ:L:2011:335:0086:0106:EN:PDF (Accessed 01 Aug 2017).

¹⁰ European Parliament and Council of the European Union (1997) 'Council Decision of 27 January 1997 establishing a reciprocal exchange of information and data from networks and individual stations measuring ambient air pollution within the Member States (97/101/EC)' [online]. Available at http://eur-

lex.europa.eu/LexUriServ/LexUriServ.do?uri=OJ:L:1997:035:0014:0022:EN:PDF (Accessed 01 Aug 2017).

¹¹ European Parliament and Council of the European Union (2016) '*Directive (EU)* 2016/2284 of the European Parliament and of the Council of 14 December 2016 on the reduction of national emissions of certain atmospheric pollutants, amending Directive 2003/35/EC and repealing Directive 2001/81/EC' [online]. Available at <u>http://eur-lex.europa.eu/legal-</u>

<u>content/EN/TXT/?uri=uriserv:OJ.L_.2016.344.01.0001.01.ENG&toc=OJ:L:2016:344:</u> <u>TOC</u> . (Accessed 05 Jul 2017).

¹² Department for Environment, Food and Rural Affairs in partnership with the Scottish Executive, Welsh Assembly Government and Department of the Environment Northern Ireland (2007) '*The Air Quality Strategy for England, Scotland, Wales and Northern Ireland (Volume 1)'*, July 2007. [online]. Available at <u>https://www.gov.uk/government/publications/the-air-quality-strategy-for-englandscotland-wales-and-northern-ireland-volume-1</u>. (Accessed 01 Aug 2017).

¹³ Department for Environment, Food and Rural Affairs. *'National Air Quality Objectives and European Directive Limit and Target Values for the Protection of Human Health'*. [online]. Available at <u>https://uk-</u>

air.defra.gov.uk/assets/documents/National air quality objectives.pdf (Accessed 01 Aug 2017).

¹⁴ Department for Environment, Food and Rural Affairs, 2016, '*Air Quality Statistics*' [online]. Available at <u>https://www.gov.uk/government/statistics/air-quality-statistics</u> (Accessed 05 Jul 2017).

¹⁵ Department for Environment, Food and Rural Affairs, 2016 [online]. Available at <u>https://www.gov.uk/government/uploads/system/uploads/attachment_data/file/54775</u> <u>9/ENEI_16_final.pdf</u> (Accessed 01 Aug 2017). ¹⁶ Department of Health, 2012 'Improving outcomes and supporting transparency Part 2: Summary technical specifications of public health indicators, updated November 2013'. Available at

https://www.gov.uk/government/uploads/system/uploads/attachment_data/file/38211 5/PHOF_Part_2_Technical_Specifications_Autumn_2014_refresh_02.12.2014_FINA L.pdf (Accessed 01 Aug 2017).

¹⁷ The Committee on the Medical Effects of Air Pollutants (COMEAP), (2011), *'Review of the UK Air Quality Index'*. ISBN 978-0-85951-699-0 [online]. Available at <u>http://webarchive.nationalarchives.gov.uk/20140505104700/http://comeap.org.uk/im</u> ages/stories/Documents/Reports/comeap%20review%20of%20the%20uk%20air%2 Oquality%20index.pdf . (Accessed 01 Aug 2017).

¹⁸ Department for Environment, Food and Rural Affairs (2014) *'Air Pollution in the UK 2013'* [online]. Available at <u>https://uk-</u>

<u>air.defra.gov.uk/library/annualreport/viewonline?year=2013_issue_1</u> (Accessed 01 Aug 2017)

¹⁹ The Environment (Northern Ireland) Order 2002 (Amendment) Regulations (Northern Ireland) 2010. Available at http://www.legislation.gov.uk/nisr/2010/92/contents/made (01 Aug 2017).

²⁰ *The Air Quality (England) Regulations 2000. (SI 2000 No.928).* Available at <u>http://www.legislation.gov.uk/uksi/2000/928/introduction/made</u> (Accessed 01 Aug 2017).

²¹ *The Air Quality (Scotland) Regulations 2000. (SI 2000 No. 97).* Available at <u>http://www.legislation.gov.uk/ssi/2000/97/contents/made</u> (Accessed 01 Aug 2017).

²² The Air Quality (Wales) Regulations 2000. (SI No. 1940 W. 138). Available at <u>http://www.legislation.gov.uk/wsi/2000/1940/contents/made</u> (Accessed 01 Aug 2017).

²³ *The Air Quality Regulations (Northern Ireland)* 2003 (SI No. 342) Available at <u>http://www.legislation.gov.uk/nisr/2003/342/contents/made</u> (Accessed 01 Aug 2017).

²⁴ The Air Quality (England) (Amendment) Regulations 2002 (SI No. 3043). Available at <u>http://www.legislation.gov.uk/uksi/2002/3043/contents/made</u> (Accessed 01 Aug 2017).

²⁵ The Air Quality (Amendment) (Wales) Regulations 2002 (SI No. 3182 W. 298). Available at <u>http://www.legislation.gov.uk/wsi/2002/3182/contents/made</u> (Accessed 01 Aug 2017).

²⁶ *The Environment (Northern Ireland) Order 2002 (SI No. 3153 (N.I. 7)* Available at <u>http://www.legislation.gov.uk/nisi/2002/3153/contents</u> (Accessed 01 Aug 2017).

²⁷ *The Air Quality (Scotland) Amendment Regulations 2002. (SI No. 297) Available* at <u>http://www.legislation.gov.uk/ssi/2002/297/contents/made</u> (Accessed 01 Aug 2017).

Section 3

²⁸ National Atmospheric Emissions Inventory (2015), *'Overview of air pollutants'* Available at <u>http://naei.beis.gov.uk/overview/ap-overview</u> (Accessed 27 Jul 2017).

²⁹ Department for Environment, Food and Rural Affairs in partnership with the Scottish Executive, Welsh Assembly Government and Department of the Environment Northern Ireland (2007) 'The Air Quality Strategy for England, Scotland, Wales and Northern Ireland' Available at <u>https://www.gov.uk/government/uploads/system/uploads/attachment_data/file/69336/ pb12654-air-quality-strategy-vol1-070712.pdf</u> (Accessed 27 Jul 2017).

³⁰World Health Organization (2005) *'Air quality guidelines. Global update 2005. Particulate matter, ozone, nitrogen dioxide and sulfur dioxide*' (ISBN 92 890 2192 6). Available at <u>http://www.euro.who.int/en/what-we-do/health-topics/environment-and-health/air-quality/publications/pre2009/air-quality-guidelines.-global-update-2005.-particulate-matter,-ozone,-nitrogen-dioxide-and-sulfur-dioxide . (Accessed 27 Jul 2017).</u>

³¹ The Committee on the Medical Effects of Air Pollutants (COMEAP), (2015) *Statement on the Evidence for the Effects of Nitrogen Dioxide on Health'* [online]. Available at <u>https://www.gov.uk/government/publications/nitrogen-dioxide-health-effects-of-exposure</u>. (Accessed 27 Jul 2017).

³² The Committee on the Medical Effects of Air Pollutants (COMEAP), (2011) *'Review of the UK air quality Index'* (ISBN 978-0-85951-699-0). Available at <u>https://www.gov.uk/government/publications/comeap-review-of-the-uk-air-quality-index</u> (Accessed 27 Jul 2017)

³³ The Committee on the Medical Effects of Air Pollutants (COMEAP), (2009) *'Long-Term Exposure to Air Pollution: Effect on Mortality'* (ISBN 978-0-85951-640-2). Available at <u>https://www.gov.uk/government/publications/comeap-long-term-exposure-to-air-pollution-effect-on-mortality</u> (Accessed 27 Jul 2017).

³⁴ The Committee on the Medical Effects of Air Pollutants (COMEAP), (2010) '*The Mortality Effects of Long-Term Exposure to Particulate Air Pollution in the United Kingdom*' (ISBN 978-0-85951-685-3) Available at <u>https://www.gov.uk/government/uploads/system/uploads/attachment_data/file/30464</u> <u>1/COMEAP_mortality_effects_of_long_term_exposure.pdf</u> (Accessed 27 Jul 2017).

³⁵ The Committee on the Medical Effects of Air Pollutants (COMEAP), (2015) '*Quantification of Mortality and Hospital Admissions Associated with Ground-Level Ozone*' (ISBN 978-0-85951-776-8). Available at https://www.gov.uk/government/publications/comeap-quantification-of-mortality-andhospital-admissions-associated-with-ground-level-ozone (Accessed 27 Jul 2017).

³⁶ Defra and the Devolved Administrations' Expert Panel on Air Quality Standards (2009) 'Metals and Metalloids'. Available at <u>http://webarchive.nationalarchives.gov.uk/20130402151656/http://archive.defra.gov.uk/environment/quality/air/airquality/panels/aqs/documents/metals-andmetalloids.pdf</u> (Accessed 27 Jul 2017).

³⁷ World Health Organization (2013) *'Review of Evidence on Health Aspects of Air Pollution – REVIHAAP Project'* Available at <u>http://www.euro.who.int/___data/assets/pdf_file/0004/193108/REVIHAAP-Final-technical-report-final-version.pdf</u>, (Accessed 27 Jul 2017).

³⁸ Carslaw, D., Beevers, S. Westmoreland, E. Williams, M. Tate, J. Murrells, T. Stedman, J. Li, Y., Grice, S., Kent, A. and I. Tsagatakis (2011). *'Trends in NO_x and NO₂ emissions and ambient measurements in the UK. Version: 18th July 2011'.* [online]. Available at <u>https://uk-air.defra.gov.uk/assets/documents/reports/cat05/1108251149</u> 110718 AQ0724 Fin

air.defra.gov.uk/assets/documents/reports/cat05/1108251149_110718_AQ0724_File al_report.pdf (Accessed 27 Jul 2017).

³⁹ CEN 'EN 14625:2005 Ambient air quality – standard method for the measurement of the concentration of ozone by ultraviolet photometry'. 2005

⁴⁰ CEN 'EN 14211:2005 Ambient air quality - Standard method for the measurement of the concentration of nitrogen dioxide and nitrogen monoxide by Chemiluminescence'. 2005

⁴¹ CEN 'EN 14212:2005 Ambient air quality – Standard method for the measurement of the concentration of sulphur dioxide by UV fluorescence'. 2005

⁴² CEN 'EN 14626:2005 Ambient air quality - Standard method for the measurement of the concentration of carbon monoxide by non-dispersive infrared spectroscopy'.
2005

⁴³ CEN 'EN 14907:2005 Ambient air quality - Standard gravimetric measurement method for the determination of the $PM_{2.5}$ mass fraction of suspended particulate matter'. 2005

⁴⁴ CEN 'EN 14902:2005 Reference method for determination of Pb/Cd/As/Ni in ambient air'. 2005

⁴⁵ CEN 'EN 14662-1:2005 Ambient air quality - Standard method for measurement of benzene concentrations - Part 1: Pumped sampling followed by thermal desorption and gas chromatography'. 2005

⁴⁶ European Parliament and Council of the European Union (2008) '*Council Directive* on ambient air quality and cleaner air for Europe (2008/50/EC)'. [online]. Available at <u>http://eur-lex.europa.eu/LexUriServ/LexUriServ.do?uri=CELEX:32008L0050:EN:NOT</u> (Accessed 27 Jul 2017).

⁴⁷ AEA Group (2009) 'QA/QC Procedures for the UK Automatic Urban and Rural Air Quality Monitoring Network (AURN)' Report number AEAT/ENV/R/2837, produced for Defra and the Devolved Administrations. Available at <u>https://uk-</u>

air.defra.gov.uk/assets/documents/reports/cat13/0910081142 AURN QA QC Man ual_Sep_09_FINAL.pdf (Accessed 27 Jul 2017).

⁴⁸ Brookes, D. M. et al. (2015) '*Technical report on UK supplementary assessment under the Air Quality Directive (2008/50/EC), the Air Quality Framework Directive (96/62/EC) and Fourth Daughter Directive (2004/107/EC) for 2014*'. Ricardo Energy & Environment report number R/3459 [online]. Available at <u>https://uk-air.defra.gov.uk/assets/documents/reports/cat09/1611011538_AQ0650_2014_MAA_Q_technical_report.pdf</u> (Accessed 27 Jul 2017).

⁴⁹ *Defra UK-AIR website.* Available at <u>http://uk-air.defra.gov.uk/</u> (Accessed 27 Jul 2017).

⁵⁰ *UK National Atmospheric Emissions Inventory.* Available at <u>http://www.naei.org.uk</u> (Accessed 27 Jul 2017).

Section 5

⁵¹ NAEI *'Pollutant Information: Carbon Monoxide*' [online]. Available at <u>http://naei.defra.gov.uk/overview/pollutants?pollutant_id=4</u>. (Accessed 27 Jul 2017).

⁵² Sarantaridis, D et al (2014) *'Annual Report for 2012 on the UK PAH Monitoring and Analysis Network'* Report produced by NPL for Defra and the Devolved Administrations [online]. Available at <u>https://uk-</u> <u>air.defra.gov.uk/assets/documents/reports/cat05/1402041506_Defra_PAH_Network</u> <u>2012_annual_report_FINAL_v2.pdf</u> (Accessed 27 Jul 2017).

Section 6

⁵³ Met Office (2016a) '*January 2016 weather summary*' [Online]. Available at <u>http://www.metoffice.gov.uk/climate/uk/summaries/2016/january</u> (Accessed 24 Jul 2017).

⁵⁴ Defra (2013) '*Daily Air Quality Index*' [Online]. Available at <u>https://uk-air.defra.gov.uk/air-pollution/daqi</u> (Accessed 24 Jul 2017).

⁵⁵ Stein, A.F., Draxler, R.R, Rolph, G.D., Stunder, B.J.B., Cohen, M.D., and Ngan, F., (2015). *'NOAA's HYSPLIT atmospheric transport and dispersion modeling system'*, Bull. Amer. Meteor. Soc., **96**, pp2059-2077, <u>http://dx.doi.org/10.1175/BAMS-D-14-00110.1</u>

⁵⁶ Rolph, G., Stein, A., and Stunder, B., (2017). *'Real-time Environmental Applications and Display sYstem: READY'*. Environmental Modelling & Software, **95**, pp 210-228.

⁵⁷ Air Resource Laboratory (2016) '*HYSPLiT*' [Online]. Available at <u>http://www.arl.noaa.gov/HYSPLIT_info.php</u> (Accessed 24 Jul 2017).

⁵⁸ London Air (2016a) '*LAQN Pollution Episodes - PM*₁₀, *PM*_{2.5} *Middle March 2016*' [Online]. Available at

http://www.londonair.org.uk/london/asp/PublicEpisodes.asp?region=0&site=&postco de=&la_id=&level=All&bulletindate=19%2F01%2F2016&MapType=Google&zoom=& lat=51.4750&lon=-

0.119824&VenueCode=&bulletin=explanation&episodeID=PM10PM25MiddleMarch2 016&pageID=page1&cm-djitdk-djitdk= (Accessed 24 Jul 2017).

⁵⁹ Met Office (2016b) '*March 2016 weather summary*' [Online]. Available at <u>http://www.metoffice.gov.uk/climate/uk/summaries/2016/march</u> (Accessed 24 Jul 2017).

⁶⁰ London Air (2016b) '*PM*₁₀ and *NO*₂ end *November*' [Online]. Available at <u>https://www.londonair.org.uk/london/asp/publicepisodes.asp?region=0&site=&postco</u><u>de=&la_id=&level=All&bulletindate=27%2F12%2F2016&MapType=Google&zoom=&lat=51.4750&lon=-</u>

<u>0.119824&VenueCode=&bulletin=explanation&episodeID=PM10andNO2endNov&pa</u> <u>geID=page1&cm-djitdk-djitdk=</u> (Accessed 12 Sep 2017).

⁶¹ London Air (2016c) '*Moderate PM*₁₀, *PM*_{2.5}, *NO*₂ *late Dec 2016*' [Online]. Available at

https://www.londonair.org.uk/london/asp/publicepisodes.asp?region=0&site=&postco de=&la_id=&level=All&bulletindate=29%2F11%2F2016&MapType=Google&zoom=& lat=51.4750&lon=-

0.119824&VenueCode=&bulletin=explanation&episodeID=smogLateDecember2016 &pageID=page2&cm-djitdk-djitdk= (Accessed 24 Jul 2017).

⁶² London Air (2016d) '*LAQN Pollution Episodes - Ozone, PM*₁₀ and PM_{2.5} early May 2016' [Online]. Available at

https://www.londonair.org.uk/london/asp/publicepisodes.asp?species=All®ion=0& site=&postcode=&la_id=&level=All&bulletindate=05%2F05%2F2016&MapType=Goo gle&zoom=9&lat=51.4750&lon=-

0.119824&VenueCode=&bulletin=explanation&episodeID=OzoneearlyMay2016&cmdjitdk-djitdk= (Accessed 24 Jul 2017). ⁶³ London Air (2016e) '*LAQN Pollution Episodes - Ozone end July 2016*' [Online]. Available at

https://www.londonair.org.uk/london/asp/publicepisodes.asp?region=0&site=&postco de=&la_id=&level=All&bulletindate=05%2F05%2F2016&MapType=Google&zoom=& lat=51.4750&lon=-

0.119824&VenueCode=&bulletin=explanation&episodeID=OzonelateJuly2016&page ID=page1&cm-djitdk-djitdk= (Accessed 24 Jul 2017).

⁶⁴ Met Office (2016c) 'September 2016 weather summary' [Online]. Available at <u>http://www.metoffice.gov.uk/climate/uk/summaries/2016/september</u> (Accessed 24 Jul 2017).

⁶⁵ London Air (2016f) '*LAQN Pollution Episodes - Moderate PM*₁₀ and PM_{2.5} Oct 2016' [Online]. Available at

http://www.londonair.org.uk/london/asp/publicepisodes.asp?species=All®ion=0&s ite=&postcode=&la_id=&level=All&bulletindate=29%2F10%2F2016&MapType=Goog le&zoom=9&lat=51.4750&lon=-

0.119824&VenueCode=&bulletin=explanation&episodeID=PM10PM25moderateOct1 6&cm-djitdk-djitdk= (Accessed 24 Jul 2017).