



Department
for Environment
Food & Rural Affairs

Air Pollution in the UK 2021

September 2022



We are the Department for Environment, Food and Rural Affairs. We are responsible for improving and protecting the environment, growing the green economy, sustaining thriving rural communities and supporting our world-class food, farming and fishing industries.

We work closely with our 33 agencies and arm's length bodies on our ambition to make our air purer, our water cleaner, our land greener and our food more sustainable. Our mission is to restore and enhance the environment for the next generation, and to leave the environment in a better state than we found it.



© Crown copyright 2022

This information is licensed under the Open Government Licence v3.0. To view this licence, visit www.nationalarchives.gov.uk/doc/open-government-licence/

This publication is available at www.gov.uk/government/publications

Any enquiries regarding this publication should be sent to us at

agevidence@defra.gov.uk

www.gov.uk/defra

Contents

1	Introduction	12
2	Legislative and Policy Framework.....	14
2.1	The Air Quality Standards Regulations	14
2.2	The National Emission Ceilings Regulations 2018.....	16
2.3	The Environmental Permitting Regulations (EPR 2016 & 2018).....	17
2.4	Policies to Improve UK Air Quality	18
2.5	Local Authority Air Quality Management.....	29
3	The Evidence Base	32
3.1	Pollutants of Concern.....	32
3.2	Assessment of Air Quality in the UK	38
3.3	Modelling	49
3.4	Access to Assessment Data	50
4	Assessment of Compliance.....	52
4.1	Definition of Zones	52
4.2	Air Quality Assessment for 2021	54
4.3	Comparison with Previous Years	66
5	Spatial Variation and Changes Over Time	71
5.1	Nitrogen Dioxide	72
5.2	PM ₁₀ Particulate Matter.....	78
5.3	PM _{2.5} Particulate Matter	86
5.4	Ozone	93
5.5	Sulphur Dioxide.....	100
5.6	Carbon Monoxide.....	104
5.7	Benzene.....	106

5.8	1,3-Butadiene.....	110
5.9	Metallic Elements.....	112
5.10	Benzo[a]pyrene.....	125
6	Effects of Covid-19 Restrictions on Air Quality.....	129
6.1	Nitrogen Oxides (NO _x and NO ₂).....	131
6.2	Ozone (O ₃).....	135
6.3	Particulate Matter (PM ₁₀ and PM _{2.5}).....	137
6.4	Other Periods of Interest.....	141
7	Where to Find Out More.....	142
8	References.....	144

Executive Summary

The UK's Air Quality Standards Regulations require reporting of ambient air quality data on an annual basis. Data are reported via the UK-AIR website at <https://uk-air.defra.gov.uk>. This report provides background information on the pollutants covered by these Regulations and the UK's Air Quality Strategy: their sources and effects, the UK's statutory monitoring networks, and the UK's modelling methodology. The report then summarises the UK's ambient air quality assessment for 2021, presenting air quality modelling data and measurements from national air pollution monitoring networks. The pollutants covered in this report are:

- Nitrogen oxides (NO_x) comprising NO and NO₂
- PM₁₀ and PM_{2.5} particles
- Ozone (O₃)
- Sulphur dioxide (SO₂)
- Carbon Monoxide (CO)
- Benzene
- 1,3-Butadiene
- Metals: lead, cadmium, nickel and mercury, and the metalloid arsenic
- Polycyclic aromatic hydrocarbons (PAH).

These data are reported on behalf of Defra (the Department for Environment, Food and Rural Affairs) and the Devolved Administrations of Scotland, Wales and Northern Ireland.

For the purposes of air quality monitoring and assessment of compliance with the Air Quality Standards Regulations, the UK is divided into 43 zones. The 2021 results are detailed in Section 4 of this report and summarised below:

- The UK met the limit value for hourly mean nitrogen dioxide (NO₂) in all 43 zones.
- 33 zones met the limit value for annual mean NO₂, with only 10 zones exceeding.
- The low number of zones exceeding the annual mean limit value is in part attributed to the ongoing Covid-19 restrictions which continued to reduce traffic on many roads in 2021.
- All zones required to meet the critical level for annual NO_x set for protection of vegetation (non-agglomeration zones) did so. This has been the case since 2008.
- All zones met the limit value for daily mean concentration of PM₁₀ particulate matter, without the need for subtraction of the contribution from natural sources.
- All zones met the limit value for annual mean concentration of PM₁₀ particulate matter, without the need for subtraction of the contribution from natural sources.
- All zones met both limit values for annual mean concentration of PM_{2.5} particulate matter. The Stage 1 limit value, which came into force on 1st January 2015, and the Stage 2 limit value came into force in 2020.
- The UK has previously achieved its 2020 national exposure reduction target for PM_{2.5}, based on the Average Exposure Indicator (AEI) statistic. In 2021, the three-

year running mean AEI was $9 \mu\text{g m}^{-3}$; this statistic has therefore remained within the target value.

- All zones met both the target values for ozone; the target value based on the maximum daily eight-hour mean, and the target value based on the AOT40 statistic.
- Four zones were compliant with the long-term objective for ozone, set for the protection of human health. This is based on the maximum daily eight-hour mean.
- 42 zones met the long-term objective for ozone, set for the protection of vegetation. This is based on the AOT40 statistic, explained in Sections 4 and 5 of this report.
- All zones met the limit values for sulphur dioxide, carbon monoxide, benzene and lead, and the target values for arsenic and cadmium.
- Four zones exceeded the target value for nickel.
- Two zones exceeded the target value for benzo[a]pyrene.

A summary of the air quality assessment for 2021, and a comparison with previous years' air quality assessments since 2008 can be found in Section 4 of this report.

Section 6 includes an investigation of the effects of the continuing Covid-19 restrictions on UK air quality during 2021. Traffic flows on many UK roads had been substantially reduced during 2020 and were around 50-60% of pre-pandemic levels at the start of 2021. As restrictions were eased, traffic flows increased, and by May 2021 weekday traffic flows were approaching February 2020 levels. However, traffic flows were lower than normal for a substantial part of the year.

It was reported last year that the pandemic restrictions of 2020 appeared to have substantially increased compliance with the limit values for nitrogen dioxide in 2020 compared to 2019. (This did not mean concentrations of other pollutants were unaffected: however, the UK was already fully compliant with the limit values for other relevant pollutants.) In 2021, measured urban concentrations of NO_2 were typically higher than in 2020 but still lower than their pre-pandemic levels. Ten zones exceeded the annual mean NO_2 limit value in 2021 compared to five in 2020; however, there were still considerably fewer zones exceeding than in pre-pandemic years, when the majority of zones exceeded.

Ozone is also of particular interest in the context of the Covid-19 restrictions. This pollutant is removed from the air by reaction with nitric oxide (NO), which is a component of vehicle emissions. The 2020 data suggested that traffic reductions may have allowed ozone concentrations in some urban areas to become higher than they otherwise would have. However, ambient ozone concentrations are greatly influenced by meteorological conditions in any given year. The year 2021 appears to have been a 'low' year for ozone, as shown by the relatively high level of compliance with the long-term objective for vegetation: just one zone exceeded, the lowest number since 2017.

For more information on air quality in the UK visit the Defra website at www.gov.uk/defra and the UK Air Quality websites at uk-air.defra.gov.uk, scottishairquality.scot/, airquality.gov.wales and airqualityni.co.uk/.

Glossary

Agglomeration Zone. Any urban area with a population greater than 250,000.

Air Quality Directive. The European Union's Directive 2008/50/EC of 21st May 2008, on Ambient Air Quality and Cleaner Air for Europe, which is often referred to as 'the Air Quality Directive'.

Air Quality Standards Regulations. Prior to 31st January 2020, the UK was a Member State of the European Union. As such, the UK was required to incorporate - or 'transpose' - the provisions of EU Directives into their own national law by a specified date. The Air Quality Standards Regulations are the legislation by which the UK fulfilled this requirement.

Air Quality Strategy. The United Kingdom's own National Air Quality Strategy, containing policies for assessment and management of air quality in the UK. This was first published in 1997, as a requirement of The Environment Act 1995.

Air Quality Strategy Objective. The Air Quality Strategy sets objectives for the maximum concentrations of eight pollutants. These are at least as stringent as the limit values of the Air Quality Directive and Air Quality Standards Regulations.

Ambient Air. Outdoor air.

Annual Mean daily maximum 8-hour mean O₃ concentrations. This is an annual mean of the 'maximum daily 8-hour mean' (see below)

Average Exposure Indicator (AEI). The statistic on which the *Air Quality Standards Regulations*' national exposure reduction target is based, for PM_{2.5} between 2010 and 2020. The AEI for the UK is calculated as follows: the arithmetic mean PM_{2.5} concentration at appropriate UK urban background sites is calculated for three consecutive calendar years, and the mean of these values taken as the AEI.

Benzene. A chemical compound that is harmful to human health. As an air pollutant, benzene can be emitted from domestic and industrial combustion processes, and road vehicles. Its chemical formula is C₆ H₆.

Benzo[a]pyrene. One of a group of compounds called *polycyclic aromatic hydrocarbons (PAHs)* that can be air pollutants. The main sources of B[a]P in the UK are domestic coal and wood burning, fires, and industrial processes such as coke production.

Beta Attenuation Monitor (BAM). A type of instrument used for monitoring concentrations of particulate matter, which measures the attenuation of beta rays passing through a paper filter tape on which particulate matter from sampled air has been collected.

1,3-Butadiene. This is an organic compound emitted into the atmosphere mainly from fuel combustion e.g. petrol and diesel vehicles. 1,3-butadiene is also an important chemical in certain industrial processes, particularly the manufacture of synthetic rubber. 1,3-butadiene is known to cause cancer in humans.

Carbon Monoxide (CO) a pollutant gas released in road vehicle exhausts. When breathed in, carbon monoxide affects the blood's ability to carry oxygen around the body.

Clean Air Strategy (CAS): published in 2019, this is the UK Government's framework document setting out policy action to drive down national emissions of five damaging pollutants to achieve statutory emissions reduction commitments, reduce background pollution, and minimise human exposure to harmful concentrations of pollution.

Cleaner Air for Scotland 2 (CAFS2): published in 2021, sets out the framework for air quality policy in Scotland to 2026.

Daily maximum 8-hour mean O₃ concentrations. For a given day the 'daily maximum 8-hour mean' is derived from the highest of the 8-hour averages beginning with the 8-hour period from 17:00 p.m. on the previous day to 01:00 a.m. and ending with the 8-hour period from 16:00 p.m. to 00:00 a.m. The highest of the continuous 8-h averages running through each day is taken for each day.

Digitel™ Sampler. A type of sampler used in the PAH Network: air is drawn through a filter which is subsequently analysed for **polycyclic aromatic hydrocarbons (PAHs)**.

Episode (Air Pollution Episode). An 'air pollution episode' means a period of time (usually a day or several days) when air pollution is high (air quality is poor).

Eutrophication. Increased levels of plant nutrients such as phosphorus and nitrogen, in bodies of water such as lakes or rivers. This can cause an increase in growth of water plants and algae which in turn can affect the water's ability to support other life such as fish.

FDMS. This stands for 'Filter Dynamic Measurement System' and refers to a type of instrument for monitoring concentrations of particulate matter. The FDMS is a modified form of Tapered Element Oscillating Microbalance (TEOM). It measures particulate concentration by measuring the change in oscillating frequency of a tapered element, which holds a filter on which sampled particulate matter is continuously deposited.

Fidas™. A type of instrument which uses an optical technique for monitoring concentrations of particulate matter.

Gravimetric Sampler. A type of instrument used to measure ambient concentrations of **particulate matter**. It works by drawing air through a filter, on which the particulate matter is collected. The filter is subsequently weighed and the ambient concentration of particulate matter calculated.

Fourth Daughter Directive. The European Union's Directive 2004/107/EC, which covers the four metallic elements cadmium, arsenic, nickel and mercury together with **polycyclic aromatic hydrocarbons (PAH)**. (Its name comes from its origin as one of four so-called Daughter Directives set up under an overarching 'framework Directive'.) The provisions of the Fourth Daughter Directive were transposed into UK law by means of the Air Quality Standards Regulations.

Leckel SEQ™. A type of *gravimetric sampler* used for measuring ambient concentrations of PM₁₀ or PM_{2.5}.

Limit value. The *Air Quality Standards Regulations* set 'limit values' for ambient concentrations of pollutants. Limit values are legally binding and must not be exceeded.

Long-Term Objectives. As well as limit values and target values, the *Air Quality Standards Regulations* set 'long-term objectives' for ozone concentration. These are similar to limit values but are not legally mandatory. The UK must take all necessary measures not entailing disproportionate costs to meet the target values and long-term objectives.

Member States. Countries that are part of the European Union.

Microgram per cubic metre ($\mu\text{g m}^{-3}$). Unit often used to express the concentration of a pollutant in air. $1 \mu\text{g} = 1$ millionth of a gram or 1×10^{-6} g.

Micrometre (μm). Unit of length often used for the size of particulate pollutants. $1 \mu\text{m} = 1$ millionth of a metre (1×10^{-6} m) or one thousandth of a millimetre.

Milligram per cubic metre (mg m^{-3}). Unit often used to express the concentration of carbon monoxide in air. $1 \text{mg} = 1$ thousandth of a gram or 1×10^{-3} g.

Nanogram per cubic metre (ng m^{-3}). Unit often used to express concentrations of pollutants such as metallic elements and **PAH**, which are usually found at low concentrations in air. $1 \text{ng} = 1$ billionth of a gram or 1×10^{-9} g.

Net Zero. Net zero emissions are reached when anthropogenic (i.e., human-caused) emissions of greenhouse gases to the atmosphere are balanced by anthropogenic removals over a specified period.

Nickel (Ni) A toxic metallic element found in ambient air as a result of releases from oil and coal combustion, metal processes, manufacturing and other sources.

Nitric oxide (NO). One of the oxides of nitrogen formed in combustion processes. NO is not harmful to human health but combines with oxygen to form nitrogen dioxide.

Nitrogen Dioxide (NO₂) One of the oxides of nitrogen formed in combustion processes. At high concentrations NO₂ is an irritant to the airways. NO₂ can also make people more likely to catch respiratory infections (such as flu), and to react to allergens.

Nitrogen Oxides (NO_x). Compounds formed when nitrogen and oxygen combine. NO_x, which comprises nitric oxide (NO) and nitrogen dioxide (NO₂), is emitted from combustion processes. Main sources include power generation, industrial combustion and road transport.

Non-agglomeration zones. Zones with no single large urban population contained within them.

Ozone (O₃). A pollutant gas which is not emitted directly from any source in significant quantities, but is produced by reactions between other pollutants in the presence of

sunlight. (This is what is known as a '**secondary pollutant**'). Ozone concentrations are greatest in the summer. O₃ can travel long distances and reach high concentrations far away from the original pollutant sources. Ozone is an irritant to the airways of the lungs, throat and eyes: it can also harm vegetation.

Particulate Matter (PM). Small airborne particles. PM may contain many different materials such as soot, wind-blown dust or secondary components, which are formed within the atmosphere as a result of chemical reactions. Some PM is natural and some is human made. Particulate matter can be harmful to human health when inhaled, and research shows a range of health effects associated with PM. In general, the smaller the particle the deeper it can be inhaled into the lung.

Partisol™. A type of **gravimetric sampler** used for measuring ambient concentrations of PM₁₀ or PM_{2.5}.

PM₁₀. Particles which pass through a size-selective inlet with a 50 % efficiency cut-off at 10 µm aerodynamic diameter, as defined in ISO 7708:1995, Clause 6. This size fraction is important in the context of human health, as these particles are small enough to be inhaled into the airways of the lung – described as the 'thoracic convention' in the above ISO standard. PM₁₀ is often described as '*particles of less than 10 micrometres in diameter*' though this is not strictly correct.

PM_{2.5}. Particles which pass through a size-selective inlet with a 50 % efficiency cut-off at 2.5 µm aerodynamic diameter, as defined in ISO 7708:1995, Clause 7.1. This size fraction is important in the context of human health, as these particles are small enough to be inhaled very deep into the lung – described as the 'high risk respirable convention' in the above ISO standard. PM_{2.5} is often described as '*particles of less than 2.5 micrometres in diameter*' though this is not strictly correct.

Primary pollutant. A pollutant which is emitted directly into the atmosphere from a source.

Polycyclic Aromatic Hydrocarbons (PAH). PAHs are a large group of chemical compounds that are toxic and carcinogenic. Once formed, they can remain in the environment for a long time and can be passed up the food chain. The main sources are domestic coal and wood burning, outdoor fires, and some industrial processes. The pollutant **benzo[a]pyrene** is a PAH, and because it is one of the more toxic PAH compounds it is measured as a 'marker' for this group of pollutants.

QA/QC (or QAQC): Quality Assurance and Quality Control.

Secondary pollutant. A pollutant which is formed by chemical reactions from other pollutants in the atmosphere. Ozone, for example, is a secondary pollutant. Particulate matter (PM_{2.5} and PM₁₀) consists of a mix of primary material (directly emitted from sources) and secondary material (formed by reactions in the atmosphere).

Sulphur dioxide (SO₂). An acid gas formed when fuels containing sulphur impurities are burned. SO₂ irritates the airways of the lung.

Target Value. As well as limit values, the **Air Quality Standards Regulations** set target values for some pollutants. These are similar to limit values but are not legally mandatory.

The UK must take all necessary measures not entailing disproportionate costs to meet the target values.

TOMPs. This stands for 'Toxic Organic Micropollutants'. These are compounds that are present in the environment at very low concentrations but are highly toxic and persistent. They include dioxins and dibenzofurans.

1 Introduction

Clean air is vital for people's health and the environment, essential for making sure our cities are welcoming places for people to live and work now and in the future, and for our prosperity. Improving air quality remains a key priority for the UK.

It is therefore important to monitor levels of air pollution. The broad objectives of monitoring air pollution in the UK are:

- To fulfil statutory air quality reporting requirements.
- To provide a sound scientific basis for the development of cost-effective control policies.
- To provide the public with open, reliable and up-to-date information on air pollution, enabling them to take appropriate action to minimise health impacts.
- To evaluate potential impacts on population, ecosystems and our natural environment.

The UK's Air Quality Standards Regulations (UK Government, 2010) , (Scottish Government, 2010), (Welsh Government, 2010) , (Department of Environment Northern Ireland, 2010) require the UK to undertake an air quality assessment and report the findings on an annual basis. The UK has statutory monitoring networks in place to meet the requirements of the above Regulations, with air quality modelling used to supplement the monitored data.

The UK is also required to make the information available to the public. One way in which this is done is by the series of annual 'Air Pollution in the UK' reports. '*Air Pollution in the UK 2021*' continues this series, and has two aims:

- To provide a summary of the UK's 2021 air quality assessment and findings. A separate Compliance Assessment Summary document is also published, based upon Section 4 of this report, and accompanies the UK's 2021 submission. This provides a concise summary aimed at the public.
- To act as a State of the Environment report, making information on the ambient air quality evidence base for the year publicly available. This includes an assessment of trends and spatial distribution, together with information on pollution events during the year.

This report:

- Outlines the air quality legislative and policy framework in the UK (Section 2).

- Describes the evidence base underpinning the UK's air quality assessment: the pollutants of concern, and where and how air pollution is measured and modelled (Section 3).
- Presents an assessment of the UK's compliance in 2021 with the limit values, target values and long-term objectives set out in the Air Quality Standards Regulations (Section 4).
- Compares this with previous recent years (Section 4).
- Explains the spatial distribution of the main pollutants of concern within the UK during 2021 and looks at how ambient concentrations have changed in recent years (Section 5).
- Explains noteworthy pollution events that occurred during 2021 (Section 6). Typically, this section features episodes of high pollution: for the 2021 report, it looks at how the Covid-19 'lockdown' restrictions affected air quality in the UK.
- Explains where to find out more (Section 7).

Further information on air quality in the UK can be found on Defra's online UK Air Information Resource (UK-AIR), at uk-air.defra.gov.uk.

2 Legislative and Policy Framework

The UK air quality framework is currently derived from a mixture of domestic and international legislation and consists of three main strands:

- 1) Legislation regulating concentrations of pollutants in ambient air – the Air Quality Standards Regulations 2010.
- 2) Legislation regulating total national emissions of air pollutants – the National Emission Ceilings Regulations 2002 and the Gothenburg Protocol to the UNECE Convention on Long-range Transboundary Air Pollution; and
- 3) Legislation regulating emissions from specific sources such as UK legislation implementing the Environmental Permitting Regulations and the Clean Air Act. (Note: Northern Ireland does not have Environmental Permitting Regulations but instead regulates industrial emissions via the Northern Ireland Pollution Prevention and Control Regulations, 2013 (Department of Environment Northern Ireland , 2013).)

In February 2022, the UK Government published the Air Quality Common Framework (UK Government, 2022a). This policy paper, which is available online at <https://www.gov.uk/government/publications/air-quality-provisional-common-framework> , explains how the UK Government and the Devolved Administrations propose to work together to develop air quality policy, following the UK's exit from the European Union.

Reducing air pollution requires action to reduce domestic emissions as well as working closely with international partners to reduce transboundary emissions (pollutants blown over from other countries) which, at times, can account for a significant proportion of pollutant concentrations experienced in the UK. For example, a 2013 report prepared by the Air Quality Expert Group on behalf of Defra and the Devolved Administrations estimated that emission sources within the UK only accounted for 50-55% of measured annual average fine particulate matter (PM_{2.5}) concentrations, the remainder being formed or emitted elsewhere (Air Quality Expert Group, 2013).

2.1 The Air Quality Standards Regulations

2.1.1 Background to the Air Quality Standards Regulations

In the UK, concentrations of a range of pollutants in ambient air are regulated by the Air Quality Standards Regulations as follows:

- The Air Quality Standards Regulations 2010 in England (UK Government, 2010), and their December 2016 amendment (UK Government, 2016)

- The Air Quality Standards (Scotland) Regulations 2010 in Scotland (Scottish Government, 2010), and their December 2016 amendment (Scottish Government, 2016)
- The Air Quality Standards (Wales) Regulations 2010 in Wales (Welsh Government, 2010)
- The Air Quality Standards Regulations (Northern Ireland) 2010 (Department of Environment Northern Ireland, 2010) and their December 2016 amendment (DAERA, 2017)
- The Air Quality Standards Regulations (Gibraltar) and their December 2016 amendment (HM Government of Gibraltar, 2016)

These Regulations have their origins in the following European Union legislation:

- Directive 2008/50/EC of 21st May 2008, on Ambient Air Quality and Cleaner Air for Europe (European Parliament and Council of the European Union, 2008). This is referred to in this report as ‘the Air Quality Directive’ and covers the following pollutants: sulphur dioxide, nitrogen oxides, particulate matter (as PM₁₀ and PM_{2.5}), lead, benzene, carbon monoxide and ozone. It revised and consolidated previously existing EU air quality legislation relating to the above pollutants.
- Directive 2004/107/EC of 15th December 2004 (European Parliament and Council of the European, 2004), relating to arsenic, cadmium, mercury, nickel and polycyclic aromatic hydrocarbons in ambient air. This is referred to as ‘the Fourth Daughter Directive’ and covers the four elements cadmium, arsenic, nickel and mercury, together with polycyclic aromatic hydrocarbons (PAH).

2.1.2 Provisions of the Air Quality Standards Regulations

The Air Quality Standards Regulations set ‘limit values’, ‘target values’ and ‘long-term objectives’ for ambient concentrations of pollutants. These are explained below, as well as provisions regarding monitoring, and reporting of data.

Limit values are legally binding and must not be exceeded. They are set for individual pollutants and comprise a concentration value, an averaging period for the concentration value, a number of exceedances allowed (per year) and a date by which this must be achieved. Some pollutants have more than one limit value, for example relating to short-term average concentrations (such as the hourly mean) and long-term average concentrations (such as the annual mean).

Target values and **long-term objectives** are set for some pollutants and are configured in the same way as limit values. These are not legally binding, but the UK must take all necessary measures not entailing disproportionate costs to meet the target values and long-term objectives.

The Air Quality Standards Regulations include detailed provisions on the **monitoring and reporting** of air quality, including:

- The division of the UK into zones for the purposes of compliance reporting.
- The location and number of sampling points.
- The measurement methods to be used.
- Data quality objectives.
- Siting criteria each monitoring station must meet.
- Provision for reporting compliance.
- Provision of information to the public.

The UK has statutory monitoring networks in place to meet the requirements of the above legislation, with air quality modelling used to supplement the monitored data.

2.2 The National Emission Ceilings Regulations 2018

The UK's National Emission Ceilings Regulations (NECR) 2018 (UK Government, 2018) sets emission reduction commitments (ERCs) for the total emissions of oxides of nitrogen (NO_x), oxides of sulphur (SO_x), non-methane volatile organic compounds (NMVOC), ammonia (NH₃) and particulate matter as PM_{2.5} in 2020 and 2030. The 2010 emission ceilings for NO_x, NMVOC, NH₃ and SO_x, in the NECR and Gothenburg Protocol (United Nations Economic Commission for Europe (UNECE), 1999), applied up to the end of 2019 and were then superseded by the 2020 ERCs. The NECR pollutants contribute to acidification and eutrophication of ecosystems whilst also playing a major role in the formation of ground-level ozone. Under the NECR and Gothenburg Protocol, the UK is required to prepare and annually update national emissions inventories for these and a number of other air pollutants.

The Gothenburg Protocol was revised in May 2012 to set emission reduction commitments (ERCs) for 2020 (from the 2005 baseline) for the same four pollutants and PM_{2.5}. The revised National Emission Ceilings Directive (European Parliament and Council of the European Union, 2016) came into force on 31st December 2016. This revised Directive was transposed into UK legislation in February 2018 via the National Emissions Ceilings Regulations 2018, and the new UK legislation came into force on 1st July 2018 (UK Government, 2018). Defra's Clean Air Strategy, published in January 2019, sets out how we will work towards these goals (see Section 2.4.2). In addition, the UK published its National Air Pollution Control Programme (Defra, 2019a) on 1st April 2019 which detailed the potential policies and measures required to meet the 2020 and 2030 emission reduction targets.

The UK has met domestic and international 2020 emission reduction commitments for emissions of NO_x, SO₂, NMVOCs and particulate matter (PM_{2.5}), but UK emissions of ammonia (NH₃) had not fallen by enough to meet the emission reduction commitments for 2020. As permitted under existing regulations, the UK submitted an application to make permitted adjustments to the emissions. This involved preparing an adjusted emissions inventory, on the basis of the non-compliance being a result of our improvements to the methodology for calculating emissions from non-manure digestates (the solid substance produced by anaerobic digestion processes, which can be used as fertiliser). The adjusted inventory has been scrutinised by UNECE experts and accepted by them. The UK's adjusted totals for ammonia are now compliant with National Emission Ceilings Regulations (2018), and the UK is therefore now in compliance with our international commitments, as contained in the amended Gothenburg Protocol.

To fulfil the reporting requirements under the Convention for Long Range Transboundary Air Pollution (CLRTAP) and in the NECR, the UK compiles and reports its air pollutant emissions inventory on an annual basis. The latest emissions data are for the year 2020 and can be found here: <https://naei.beis.gov.uk/data/> .

A revised National Air Pollution Control Programme (NAPCP) is in preparation at the time of writing, with a public consultation having taken place from 25th July 2022 to 4th September 2022. The revised NAPCP will outline UK wide policies and measures which will be considered further in order to reduce emissions in accordance with the National Emission Reduction targets set under the NECR.

2.3 The Environmental Permitting Regulations (EPR 2016 & 2018)

UK's Environmental Permitting Regulations set standards and provisions to reduce the emissions of pollutants from a diverse range of industrial sources - from intensive pig and poultry farms to chemical manufacturing sites and power stations – with the aim of achieving the environmental and human health benefits associated with reduction in pollution.

Under the EPR regulation, industries must use best available techniques (BAT) to reduce their emissions. These techniques, and the emissions limits associated with the use of those techniques, are set out in best available technique reference documents (known as BREFs). BREFs are reviewed regularly, to ensure an ongoing process of improvement to UK's air quality achieved through the continuous reduction of pollution. In addition, the regulation sets emission limits for the emission of pollutants from particular sectors.

The UK is committed to maintaining high environmental standards and has put in place a process for determining future BAT for industrial emissions. The future UK BAT regime will provide a mechanism to address emissions to air, as well as land and water, from our largest industry in an integrated way. The new regime will be based on the principles

followed since the UK originally devised the concept; a detailed, transparent, collaborative, data-led process that builds on existing high levels of environmental protection.

Industrial emissions regulation is a devolved matter but, following EU exit, the UK Government and Devolved Administrations (DA) agreed to create a regime for developing BAT for large industries previously set through an EU process. A joint UK/DA consultation in January 2021 sought views on the design of the regime. A response to the consultation was published in August 2022. Arrangements are different in Northern Ireland, which currently does not have Environmental Permitting Regulations. In Northern Ireland, industrial emissions are regulated via the Northern Ireland Pollution Prevention and Control Regulations, 2013 (Department of Environment Northern Ireland, 2013).

2.4 Policies to Improve UK Air Quality

Domestic, EU and internationally driven environmental legislation introduced over the past seventy years has provided a strong impetus to reduce the levels of harmful air pollutants in the UK. As a result, current concentrations of many recognised pollutants are now at the lowest they have been since measurements began. The UK's 1956 Clean Air Act tackled city smog caused by domestic and industrial coal burning, and significant progress has continued to improve air quality throughout subsequent decades. Between 1970 and 2020 (the most recent year for which data are available), UK estimated emissions of nitrogen oxides have fallen by 76%, UK estimated emissions of PM₁₀ particulate matter have fallen by 80% and UK estimated emissions of PM_{2.5} particulate matter have fallen by 85% (Defra, 2022a).

However, significant challenges remain. Poor air quality remains the greatest environmental risk to public health in the UK. It is known to exacerbate the impact of pre-existing health conditions, such as respiratory and cardio-vascular illnesses, especially for the elderly and infants. The most immediate air quality challenge is tackling nitrogen dioxide concentrations. NO₂ is associated with adverse effects on human health (COMEAP, 2015), (COMEAP, 2018). Estimating the long-term impacts of NO₂ pollution is difficult, because of the challenge of separating its effects from those of other traffic-related pollutants. Although it has been more difficult to estimate the level of impact, there is enough evidence of such health effects to support the need to take action now.

Particulate matter is also of concern. In 2010, the Committee on the Medical Effects of Air Pollutants (COMEAP) produced a report on the mortality effects of long-term exposure to particulate air pollution in the United Kingdom. COMEAP estimated that the long-term impact of particulate pollution in the UK equated to 340,000 years of life lost (COMEAP, 2010).

The UK Environment Act 2021 (UK Government, 2021) received Royal Assent on 9th November 2021. This establishes a duty for the UK Government to set a legally binding target in England to reduce PM_{2.5}, alongside at least one further long-term target on air quality. The long-term target is part of the wider framework for setting legally binding

environmental targets, which also covers biodiversity, water, waste reduction and resource efficiency. The Environment Act requires that all the targets are laid before parliament by 31st October 2022; A public consultation on all the environmental targets including those for PM_{2.5} took place during 2022. The responses are currently being considered by the UK Government, but the targets proposed for air quality are as follows:

- To reduce maximum annual mean PM_{2.5} concentrations in England to 10 µg m⁻³ by 2040.
- To reduce population exposure to PM_{2.5} in England by 35% compared to 2018, by 2040.

The Act also enables English local authorities to take more effective, co-ordinated actions to achieve their air quality objectives and deliver improvements to public health, and tackle emission from domestic burning - a key source of PM_{2.5}. Furthermore, the Act introduces a new power to compel vehicle manufacturers to recall vehicles and non-road mobile machinery if they are found not to meet the environmental standards they were approved to meet. This will enable Government to ensure polluting vehicles are removed from the road and to hold non-compliant manufacturers to account.

In 2021 Welsh Government consulted on a white paper for a Clean Air (Wales) Bill to improve air quality and reduce the impacts of air pollution on public health, biodiversity, the natural environment and the economy. The white paper set out how Welsh Government intends to enable more ambitious air quality targets and put in place a more robust regulatory framework to support them. The white paper can be accessed at <https://gov.wales/white-paper-clean-air-wales-bill> .

2.4.1 The UK Air Quality Strategy

The Environment Act 1995 requires the UK Government and the Devolved Administrations for Scotland and Wales to produce a National Air Quality Strategy, containing standards, objectives and measures for improving ambient air quality; there is equivalent legislation in Northern Ireland. The Air Quality Strategy for England, Scotland, Wales and Northern Ireland was first published in March 1997 and subsequently updated in 2007 (Department for Environment, Food and Rural Affairs in partnership with the Scottish Executive, Welsh Assembly Government and Department of the Environment Northern Ireland, 2007), with a lighter touch update in 2011. The overall objectives of the Strategy are to:

- Map out future ambient air quality policy in the United Kingdom in the medium term.
- Provide best practicable protection to human health by setting health-based objectives for air pollutants.
- Outline measures which are to be taken by local authorities and other persons for the purpose of achieving those objectives.

- Contribute to the protection of the natural environment through objectives for the protection of vegetation and ecosystems.
- Establish a framework to help identify what we all can do to improve air quality.

The Strategy has established objectives for eight key air pollutants, based on the best available medical and scientific understanding of their effects on health, as well as taking into account relevant developments in Europe and advice from the World Health Organisation (Department for Environment, Food and Rural Affairs in partnership with the Scottish Executive, Welsh Assembly Government and Department of the Environment Northern Ireland, 2007). These Air Quality Objectives are at least as stringent as the limit values of the relevant EU Directives and UK Air Quality Standards Regulations – in some cases, more so, e.g. for 8-hour mean ozone, annual mean lead and benzo[a]pyrene. The Strategy also covers an additional pollutant (1,3-butadiene), and includes an additional SO₂ objective, for the 15-minute mean. Scotland has adopted some tighter objectives for particulate matter and benzene.

The Environment Act 2021 requires the Secretary of State to undertake regular five-yearly reviews of the Air Quality Strategy. Defra is currently reviewing the Air Quality Strategy. In a significant change, the forthcoming updated Air Quality Strategy will cover England only, not the Devolved Administrations. This reflects the UK's devolution settlement and the fact that air quality policy is devolved.

2.4.2 The Clean Air Strategy 2019

Clean air is crucial for life, health, the environment and the economy. The UK Government's Clean Air Strategy for England (CAS) (Defra, 2019b) sets out a range of policy actions that will help reduce air pollution, providing healthier air to breathe, enhancing the economy and protecting nature. The CAS begins by outlining our understanding of the problem, and the importance of a robust evidence base, backed by the most up to date science. It focuses on the five damaging pollutants for which the UK has ambitious reduction commitments:

- Fine particulate matter (as PM_{2.5}).
- Ammonia (NH₃).
- Nitrogen oxides (NO_x).
- Sulphur dioxide (SO₂).
- Non-methane volatile organic compounds (NMVOCs).

Ozone (O₃) is a pollutant which is not emitted but formed from chemical reactions in the air. By tackling all sources of pollution, we will continue to reduce the formation of ozone across the whole of the UK.

Next, it deals with **protecting the nation's health**. The health impacts (and their estimated costs to the UK) are summarised: actions are then set out for how public exposure to air pollution can be reduced.

Air pollution is not only a major risk to human health; it also has significant effects on the natural environment. It has damaging impacts on both plants and animal communities. The CAS devotes a chapter to **protecting the environment**.

The CAS highlights the importance of **securing clean growth and driving innovation**. Cleaning our air is about increasing productivity through improved air quality, effective use of resources and moving to a low carbon economy. Improving air quality is linked to tackling climate change, as the sources of 'greenhouse gases' implicated in climate change are in many cases also sources of air pollutants. Government is taking a holistic approach to improving air quality, balancing the need to drive down pollution with other key priorities such as achieving net zero and managing economic burdens on businesses and individuals.

Air pollutants are released in large quantities from sources as diverse as domestic energy generation, industrial energy generation, industrial processes, road, air and rail transport, construction machinery, agriculture, and domestic cleaning and personal care products. The CAS contains subsequent chapters devoted to:

- Action to reduce emissions from **transport** (covering action already taken and action proposed for the future).
- Action to reduce emissions from **the home** (including domestic burning, and the use of products which release NMVOCs).
- Action to reduce emissions from **farming**: this focusses on reduction of emissions of ammonia (which reacts with nitrogen oxides and sulphur dioxide to form secondary particulate matter) and NMVOCs (which contribute to the formation of ozone).
- Action to reduce emissions from **industry** – setting out how we can build upon the existing strong framework and support continuous improvement.

The Strategy also addresses the issue of **leadership at all levels**: international (acknowledging that air pollution is not stopped at national borders, and UK emissions can have an impact across our country, the continent and worldwide); national (including improving the legislative framework); and local (enabling local authorities to take more effective, co-ordinated actions to achieve their air quality objectives). Focusing on the importance of co-operating with all parts of the UK, the document outlines the actions which are already underway throughout Northern Ireland, Scotland and Wales. These actions show how we can address the different sources of air pollution.

Finally, the Strategy summarises **progress** - there are several high-profile agreements to which the UK has signed up to, to improve air quality. Currently, the UK is compliant with ambient air quality limit values set by the Air Quality Standards Regulations 2010 for most pollutants. The annual mean limit value for nitrogen dioxide concentrations is the only statutory air quality limit that the UK is currently failing to meet. The 2017 UK plan for tackling roadside nitrogen dioxide (NO₂) concentrations (Defra, 2017a) and 2018 supplement, sets out action to achieve compliance as swiftly as possible. It has been shown through analysis that the CAS can help meet the ambitious targets to reduce emissions.

In 2015, the Scottish Government published its first air quality strategy separate from the rest of the UK '*Cleaner Air for Scotland – The Road to a Healthier Future*' or CAFS. (Scottish Government, 2015). CAFS brought together into a single framework a number of Government policies impacting on air quality and sets out a series of 40 actions intended to deliver further air improvements. At the end of 2018, an independent review of CAFS was launched, with a remit to review progress to date and identify priorities for additional action. The review report was published in August 2019 and was used as the basis for developing a revised and updated strategy (Scottish Government, 2019). The Scottish Government's new air quality strategy '*Cleaner Air for Scotland 2 – Towards a Better Place for Everyone*' (CAFS2) was published in July 2021 (Scottish Government, 2021) .

CAFS2 is structured around 10 general themes and sets out the air quality policy framework in Scotland to 2026. CAFS was largely concerned with transport emissions and, whilst this continues to be a key part of CAFS2, there is a stronger additional focus on other emissions sources, notably agriculture and domestic fuel combustion, along with a detailed consideration of the human health and wellbeing impacts of poor air quality. Co-ordination with other key Scottish Government plans and policies, such as the National Transport Strategy 2, National Planning Framework 4, the Scottish Nitrogen Balance Sheet, the Climate Change Plan update and the Environment Strategy is a central thread running through CAFS2. The importance of public engagement and behaviour change is also a major theme.

In August 2020, the Welsh Government published its '*Clean Air Plan for Wales: Healthy Air, Healthy Wales*' (Welsh Government, 2020). This Plan sets out a 10-year pathway to achieving cleaner air in Wales. The Plan has been structured around four core themes:

- People: Protecting the health and well-being of current and future generations;
- Environment: Taking action to support our natural environment, ecosystems and biodiversity;
- Prosperity: Working with industry to reduce emissions, supporting a cleaner and more prosperous Wales; and
- Place: Creating sustainable places through better planning, infrastructure and transport.

The themes were designed through the lens of the Well-being of Future Generations (Wales) Act 2015 to enable collaborative and integrated approaches to improving air quality, across a range of policy areas and sectors. Information on this Act can be found online at <https://www.futuregenerations.wales/about-us/future-generations-act/> .

The Department of Agriculture, Environment and Rural Affairs (DAERA) is developing Northern Ireland's first Clean Air Strategy. In autumn 2020, a Discussion Document was issued to public consultation. It invited views on a range of matters relating to air quality and was an opportunity for stakeholders to put ideas to the Department. The consultation closed in spring 2021 and responses were analysed in detail. A synopsis of the responses is available to view at https://www.daera-ni.gov.uk/clean_air_strategy_discussion_document. Preliminary findings have been discussed with the Minister and an inter-departmental working group has been established to further develop proposals and identify policies for cross-departmental consideration. The Draft Clean Air Strategy is to be drafted in autumn 2022. This will be subject to consideration by the Northern Ireland Executive. A further public consultation is planned for the end of December 2022, to seek views on the proposed draft strategy. While work is progressing well at an official level, Northern Ireland currently does not have a functioning Executive, which may impact the planned timeline for this strategy.

2.4.3 National Air Quality Statistics and Indicators

For many years, the UK has reported the following two indicators as National Air Quality Statistics for ambient air:

- **Annual average concentrations of particles and ozone.** These two types of air pollution are believed to have a significant impact on public health.
- **Number of days in the year when air pollution is 'Moderate' or higher.** This may relate to any one of five key air pollutants and is based on the UK's Daily Air Quality Index (see Section 2.4.5 which deals with forecasting). From the 1st January 2012, PM_{2.5} particles replaced carbon monoxide in this suite of pollutants. The thresholds used to define 'Moderate' and higher pollution levels in the air quality index were also revised at the beginning of 2012.

In 2018, new content was added, including the following:

- **Annual mean concentrations of fine particulate matter (PM_{2.5})** at urban roadside and background monitoring sites. The inclusion of PM_{2.5} reflects the increased interest in this size fraction.
- **Annual mean nitrogen dioxide (NO₂) concentrations** at urban roadside, urban background and rural background monitoring sites. The inclusion of NO₂ informs the public and scientific discussion regarding concentrations of this pollutant, particularly at the roadside.

- **Average hours per year in the ‘Moderate’ or higher categories** of the Daily Air Quality Index, for PM₁₀, PM_{2.5}, NO₂ and ozone. This is intended to highlight variation in short-term exposure per year to harmful levels of air pollution.
- **Variation in pollutant concentration by month of the year (for PM_{2.5} and ozone), by day of the week (for NO₂), and by hour of the day - ‘diurnal’ variation – (for PM_{2.5} and NO₂).** These are provided for the most recent year and intended to aid understanding of the nature of variation in pollutant concentrations at different types of site.

The National Air Quality Statistics summary for 2021 was released on 28th April 2022 and is available from the Defra website at <https://www.gov.uk/government/statistics/air-quality-statistics> .

The UK Government’s Public Health Outcomes Framework for England 2016 – 2019 (Department of Health and Social Care, 2016) recognises the burden of ill-health resulting from poor air quality as well as other public health concerns. This Framework sets out 60 health outcome indicators for England, and includes as an indicator:

- The fraction of annual all-cause adult mortality attributable to long-term exposure to current levels of anthropogenic particulate air pollution (measured as fine particulate matter, PM_{2.5}).

This indicator is intended to enable Directors of Public Health to appropriately prioritise action on air quality in their local area. The indicator is calculated for each local authority in England based on modelled concentrations of fine particulate air pollution (PM_{2.5}). Annual estimates of the percentage of mortality attributable to long term exposure to particulate air pollution in England are available from the Public Health Outcomes Framework data tool at <https://fingertips.phe.org.uk/profile/public-health-outcomes-framework>. The most recent estimate for England at the time of writing, which is based on year 2020, is 5.6%.

The Defra document ‘*Air Quality: Public Health Impacts and Local Actions*’ can be found at [https://laqm.defra.gov.uk/documents/air_quality_note_v7a-\(3\).pdf](https://laqm.defra.gov.uk/documents/air_quality_note_v7a-(3).pdf). A toolkit aimed at helping public health professionals appropriately prioritise assessment and action on PM_{2.5} on a local level is available here:

<https://randd.defra.gov.uk/Default.aspx?Menu=Menu&Module=More&Location=None&Completed=0&ProjectID=18580> .

Northern Ireland has a similar Public Health Strategy: ‘*Making Life Better – A Whole System Framework for Public Health 2013-2023*’. This document can be found at <https://www.health-ni.gov.uk/topics/public-health-policy-and-advice/making-life-better-whole-system-strategic-framework-public>, and also includes an air quality indicator.

Wales has a new national indicator under the Well-being of Future Generations (Wales) Act 2015 and the Welsh Public Health Outcomes Framework, which has now been published on StatsWales at <https://statswales.gov.wales/Catalogue/Environment-and->

[Countryside/Air-Quality](#). Guidance has also been published for public health professionals in supporting the collective management of air quality across Wales. *Working together to reduce outdoor air pollution, risks and inequalities* can be found at <https://gov.wales/sites/default/files/publications/2019-06/working-together-to-reduce-outdoor-air-pollution-risks-and-inequalities.pdf> .

The Scottish Government's National Performance Framework (<https://nationalperformance.gov.scot/>) includes 81 National Indicators, many of which relate to environmental and human health.

2.4.4 National Emissions Statistics

The UK reports annual emissions of the following pollutants via an annual National Statistics Release, available at <https://www.gov.uk/government/statistics/emissions-of-air-pollutants>. (This is a large report comprising multiple sections: links to the individual sections for each pollutant are included alongside the main conclusions below).

- Sulphur dioxide (SO₂).
- Oxides of nitrogen (NO_x).
- Non-methane volatile organic compounds (NMVOCs).
- Ammonia (NH₃).
- Particulate matter (as PM₁₀ and PM_{2.5}).

The most recent National Statistics Release covers 1970 to 2020 (the most recent year for which emission statistics are available). The main conclusions are as follows:

- *'Emissions of sulphur dioxide have fallen by 98 per cent since 1970, to 136 thousand tonnes in 2020. Emissions decreased by 13 per cent from 2019 to 2020, dropping to the lowest level in the time series. This was driven by a decline in coal use in power stations, continuing a long-term decrease in emissions from this source. Stricter limits being placed on the sulphur content of liquid fuels has also reduced emissions in the long-term.'* (From Section 2 of 'Sulphur Dioxide' at <https://www.gov.uk/government/statistics/emissions-of-air-pollutants/emissions-of-air-pollutants-in-the-uk-sulphur-dioxide-so2>.)
- *'Emissions of nitrogen oxides have fallen by 76 per cent since 1970, to 702 thousand tonnes in 2020. This trend was driven by a decline in coal use in power stations and modernisation of the road transport fleet. There was a decrease of 13 per cent between 2019 and 2020. This is a larger annual decrease than the long-term trend, since emissions have fallen by an average of 4.6 per cent per year between 1990 and 2020. This is associated with a reduction in vehicle usage during the pandemic.'* (From Section 2 of 'Nitrogen Oxides' at

<https://www.gov.uk/government/statistics/emissions-of-air-pollutants/emissions-of-air-pollutants-in-the-uk-nitrogen-oxides-nox> .)

- *'Emissions of non-methane volatile organic compounds (NMVOCs) have fallen by 67 per cent since 1970, to 785 thousand tonnes in 2020. There was a decrease in emissions of 5 per cent between 2019 and 2020. NMVOC emissions peaked in 1990 and then fell by an average of 5.4 per cent per year between 1990 and 2009, largely due to improvements to emissions standards for road transport and stricter limits applied to industrial processes. Since 2010, annual changes have been much smaller, averaging a decrease of just 1.7 per cent each year. Since 1990, NMVOC emissions have fallen by 72 per cent.'* (From Section 2 of 'Non-methane volatile organic compounds (NMVOCs)' at <https://www.gov.uk/government/statistics/emissions-of-air-pollutants/emissions-of-air-pollutants-in-the-uk-non-methane-volatile-organic-compounds-nmvocs> .)
- *'Emissions of ammonia have fallen by 16 per cent since 1980, to 259 thousand tonnes in 2020. There was a decrease of 3 per cent in emissions of ammonia between 2019 and 2020. Annual emissions of ammonia have remained relatively stable since 2007. Between 2013 to 2017, emissions increased by 7 per cent, then declined by 4 per cent to 2020. Over the longer-term, there was a gradual decrease in annual emissions of ammonia during the 1990s and 2000s. Changes in the trend of emissions of ammonia are largely driven by changes to farming practices and herd sizes.'* (From Section 2 of 'Ammonia' at <https://www.gov.uk/government/statistics/emissions-of-air-pollutants/emissions-of-air-pollutants-in-the-uk-ammonia-nh3> .)
- *'Annual emissions of PM₁₀ have fallen by 80 per cent since 1970, to 137 thousand tonnes in 2020. There was a decrease of 9.1 per cent between 2019 and 2020. Annual emissions of PM_{2.5} have fallen by 85 per cent since 1970, to 80.1 thousand tonnes in 2020. There was a decrease of 7 per cent between 2019 and 2020. Levels of both pollutants generally decreased year-on-year between 1970 and the late-2000s. There are many reasons for this long-term decrease covering most emissions sectors, but the reduction in the burning of coal and improved emission standards for transport and industrial processes are major drivers. Since the late 2000s, annual emissions have fluctuated year-on-year because significant decreases in emissions from some sectors are largely offset by increases in emissions from wood burning in domestic settings and by solid fuel burning by industry (particularly the use of biomass).'* (From Section 1 of 'Particulate Matter (PM₁₀ and PM_{2.5})' at <https://www.gov.uk/government/statistics/emissions-of-air-pollutants/emissions-of-air-pollutants-in-the-uk-particulate-matter-pm10-and-pm25> .)

The report also states that the UK met current emission reduction commitments for sulphur dioxide, oxides of nitrogen, PM_{2.5} and NMVOC in 2020. In the case of ammonia, the UK did not achieve the 2020 emission reduction commitment: however, the UK applied to make permitted adjustments to the emissions. This application was accepted, and the

adjustments applied, so the UK's adjusted totals for ammonia are now compliant with National Emission Ceilings Regulations (2018).

New emission statistics for 2021 will be published in February 2023.

2.4.5 The UK Air Pollution Forecasting System

Daily UK air pollution forecasts are produced for five pollutants; nitrogen dioxide, sulphur dioxide, ozone, PM₁₀ particles and PM_{2.5} particles. The forecasts are communicated using the Daily Air Quality Index (<http://uk-air.defra.gov.uk/air-pollution/daq>) which is a scale of one to ten divided into four bands. This allows the public to see at a glance whether the air pollution is low, moderate, high or very high and to look up any recommended actions to take.

The group of pollutants covered, and the thresholds between the various index bands, were updated by Defra as of 1st January 2012, in the light of recommendations by the Committee on the Medical Effects of Air Pollutants (COMEAP) in their 2011 review of the UK air quality index (COMEAP, 2011). In December 2021, Defra (with support from DHSC and UKHSA) launched a comprehensive review into the way air quality information is communicated to the public. This review is being guided by an independent steering group of multidisciplinary experts. As part of the review process the steering group will make recommendations for any improvements that should be made to the Daily Air Quality Index. Progress on the air quality information system review is being published on the UK-AIR website, at: <https://uk-air.defra.gov.uk/research/aq-system-review>.

Currently, the daily forecast is provided by the Met Office and is available from UK-AIR and from the Scottish, Welsh and Northern Ireland air quality websites (see Section 7), and is further disseminated via e-mail, Twitter and RSS feeds. Anyone may subscribe to the free air pollution bulletins at: <https://uk-air.defra.gov.uk/subscribe>. Latest forecasts are issued daily, at: <https://uk-air.defra.gov.uk/forecasting/>. Defra also provide automated updates on current and forecast air quality via Twitter [@DefraUKAIR](https://twitter.com/DefraUKAIR).

2.4.6 NO₂ Air Quality Plans

In July 2017, the UK Government and Devolved Administrations published the *UK Plan for Tackling Roadside Nitrogen Dioxide Concentrations*, followed by a supplement in October 2018. The Plan and supplement set out how Government will achieve compliance with legal limits for NO₂ in the shortest possible time, supported by a £3.8 billion investment into air quality and cleaner transport. Government has ring-fenced £883 million to help local authorities tackle NO₂ exceedances. This funding supports local authorities deliver their air quality measures to improve the health of their residents and meet legal limits for NO₂. These air quality measures are varied and highly targeted, e.g. traffic management schemes, engineering solutions, grants and loans for vehicle upgrades and encouraging behavioural change. Measures may also include Clean Air Zones; three were implemented in 2021 and are now operational in Bath, Birmingham and Portsmouth.

Further Clean Air Zones are due to be implemented in other cities in 2022 and 2023. The funding also includes a Clean Air Fund, accessible by those local authorities implementing measures to tackle NO₂ exceedances, to help them mitigate the impact of their plans on individuals and businesses. Local authorities have used this funding for grants to individuals and businesses to upgrade their fleets, retrofit buses, Electric Vehicle (EV) charging infrastructure and discounted access to public transport. The UK Air Quality Plan for nitrogen dioxide, together with the supplement published in October 2018, is available at <https://www.gov.uk/government/publications/air-quality-plan-for-nitrogen-dioxide-no2-in-uk-2017> .

The Welsh supplemental plan, which was published in November 2018, can be found at <https://gov.wales/air-quality-plan> .

Scotland has introduced Low Emission Zones for the improvement of air quality in four cities: Glasgow, Aberdeen, Dundee and Edinburgh. These were introduced on 31st May 2022, and local grace periods now apply until enforcement begins. In Glasgow, the LEZ is already in force for buses. For other vehicle types, enforcement will start on 1st June 2023 (a year later for residents within the zone). LEZ enforcement will begin on 30th May 2023 in Dundee, and on 1st June 2024 in Edinburgh and Aberdeen. Eligible households within defined areas may apply for a cash grant to incentivise disposal of a non-LEZ compliant vehicle.

Since 2018/19 significant funding has been made available to local authorities, transport operators and the general public to support LEZ introduction. Other Scottish local authorities with Air Quality Management Areas have completed assessment to determine whether an LEZ would be an appropriate intervention in their areas.

The Scottish Government also provides a total of £3.5 million per year to support local authority air quality work, spends over £1 billion per year on public transport and has committed to investing at least £320 million – or 10% of the total transport budget – on active travel by 2024/25.

2.4.7 Measures to Address Target Value Exceedances of B[a]P and Nickel

The Air Quality Standards Regulations set target values for a number of metallic elements including nickel and for benzo[a]pyrene (B[a]P). The UK exceeded target values for B[a]P and nickel during all years from 2013 to 2021 inclusive, except for nickel in 2017.

These exceedances are reported in September of the following years as part of the UK's annual compliance assessment. For details of exceedances please see earlier 'Air Pollution in the UK' reports in this series which are available at: <https://uk-air.defra.gov.uk/library/annualreport/>.

The UK published reports providing details of the assessment of the exceedances in years 2013 to 2019. These also reported the actions and measures already taken or planned, to help the UK meet the target values. An overview report was provided for each pollutant alongside more detailed information on any exceedances by zone.

The reports explain that we are taking steps to address all the exceedances through existing long-term measures, such as regular coke oven door maintenance and through improvements in our understanding to help target measures appropriately. The nickel overview report details existing and new measures put in place and the continued work with environmental regulators to improve understanding and management of these exceedances.

The reports are available at <https://uk-air.defra.gov.uk/library/bap-nickel-measures> . At the time of writing, the 2019 reports are the most recent in the series.

2.5 Local Authority Air Quality Management

Requirements for local air quality management (LAQM) are set out in Part IV of the Environment Act 1995 (UK Government, 1995) as amended by the Environment Act 2021 (UK Government, 2021), and the Environment (Northern Ireland) Order 2002 (Northern Ireland Government, 2002). Authorities are required to carry out regular 'Review and Assessments' of air quality in their area and take action to improve air quality in those areas where objectives set out in regulation have been shown not to have been achieved, or areas where it is thought there is a risk that they will not be achieved.

Local Authorities in England, Scotland, Wales and Northern Ireland undertake Review and Assessment against the Strategy's objectives prescribed in the Air Quality (England) Regulations 2000 (UK Government, 2000), Air Quality (Scotland) Regulations 2000 (Scottish Government, 2000), Air Quality (Wales) Regulations 2000 (Welsh Government, 2000) and Air Quality (Northern Ireland) Regulations 2003 (Northern Ireland Government, 2003), together with subsequent amendments (UK Government, 2002), (Welsh Government, 2002), (Scottish Government, 2002), (Scottish Government, 2016).

With regards to LAQM statutory reporting requirements, in 2018 authorities in Wales adopted reporting in the form of an Annual Progress Report in line with the streamlined LAQM regime (Welsh Government, 2017). In England and Scotland, reporting in the form of the adopted Annual Status / Progress Reports has continued (Defra, 2016a) (Scottish Government, 2017), whilst London authorities continued working against the revised London specific LLAQM policy guidance (Mayor of London, 2019) through the preparation of Annual Status Reports. Authorities in Northern Ireland commenced Round 8 of the Review and Assessment process in 2021, with appraisal of local air quality via Updating and Screening Assessments in line with the Round based approach to LAQM.

When the Review and Assessment process identifies an exceedance of an Air Quality Strategy objective, the Local Authority must declare an 'Air Quality Management Area'

(AQMA) and develop an Action Plan to tackle problems in the affected areas. Action Plans formally set out the measures the Local Authority proposes to take to work towards meeting the air quality objectives. They may include a variety of measures such as congestion charging, traffic management, planning and financial incentives. Advice for Local Authorities preparing an Action Plan is available from the Defra LAQM web pages at <https://laqm.defra.gov.uk/action-planning/aqap-supporting-guidance.html>.

Information on the UK's AQMAs is summarised in **Table 2-1** below. At the time of writing (August 2022), 256 Local Authorities – 68.5% of those in the UK – have one or more AQMAs. Some AQMAs are for more than one pollutant, and many Local Authorities have more than one AQMA.

Table 2-1 Current UK-wide status of Air Quality Management Areas (AQMAs) and Action Plans (as of August 2022.)

Region	Total LAs	LAs with AQMAs	AQMAs for NO ₂	AQMAs for PM ₁₀	AQMAs for SO ₂
England (outside London)	276	189	509	26	5
London	33	33	34	28	0
Scotland	32	14	25	26	1
Wales	22	11	43	1	0
Northern Ireland	11	9	17	2	0
TOTAL	374	256	628	83	6

Most AQMAs in the UK are in urban areas and have been established to address the contribution to air pollution from traffic emissions of nitrogen dioxide or PM₁₀, or in some cases both. A small number are for SO₂. There are no longer any AQMAs for benzene. The number of AQMAs for PM₁₀ in Scotland is relatively high because of the more stringent objective for PM₁₀ adopted in Scotland.

Where an AQMA is declared, the Local Authority specifies the main sources of pollutants involved – for example road transport, industrial emissions or domestic sources, or a mixture of several. The methodology for counting AQMAs by source has changed since the previous report in this series: the number of AQMAs by source is now split by geographic area rather than pollutant type. This is summarised in **Table 2-2**.

Table 2-2 Current UK Air Quality Management Areas by Source (as of August 2022)

Source	England	Wales	Scotland	Northern Ireland	London
County or Unitary Authority Road	182	21	4	0	1
Domestic Heating	1	0	0	1	0
Strategic Road Network	41	2	0	0	0
Industrial Source	8	1	1	0	0
Mixture of Road Types	79	5	1	1	2
Not Defined	1	0	1	2	0
Railways	1	0	0	0	0
Road Transport (unspecified)	187	15	27	15	27
Transport and Industrial Source	11	0	2	0	4
Transport, Industrial and Domestic Sources	4	0	2	0	1

For up-to-date information on AQMAs throughout the UK, please refer to the interactive map on UK-AIR at <https://uk-air.defra.gov.uk/aqma/maps/> . This interactive map provides information on the location of the AQMA, date declared, the pollutants for which it was declared, and information on the type of pollutant sources.

3 The Evidence Base

A programme of air quality assessment and research is in place in the UK which delivers the evidential needs of Defra and the Devolved Administrations. These needs include assessment of compliance with legislation, as well as the means to assess the effectiveness of air pollution mitigation policies.

This section explains Defra and the Devolved Administrations' evidence base for the annual assessment of compliance with the Air Quality Standards Regulations on ambient air quality. It describes the air pollutants which are of concern and how these are monitored and modelled in the UK.

3.1 Pollutants of Concern

This section summarises the sources, effects (both on human health and the environment) and typical UK concentrations of the pollutants being assessed in relation to the Air Quality Standards Regulations.

The information on sources has largely been summarised from the National Atmospheric Emission Inventory (NAEI) pollutant information pages at <https://naei.beis.gov.uk/> (National Atmospheric Emissions Inventory, 2022) together with Table 1 of the Air Quality Strategy (Defra, 2007). Information on health effects has been summarised from reports produced by the World Health Organization (WHO), the Expert Panel on Air Quality Standards (EPAQS) and the Committee on the Medical Effects of Air Pollutants (COMEAP). The latest estimate is that long-term exposure to the air pollution mixture in the UK has an annual effect equivalent to 29,000 to 43,000 deaths for adults aged 30 and over (UK Health Security Agency, 2022).

3.1.1 Oxides of Nitrogen

There are several oxides of nitrogen. The ones of most interest, for air quality, are nitric oxide (NO) and nitrogen dioxide (NO₂). Together, they are often referred to as NO_x. Nitrogen oxides are emitted from combustion processes. Combustion in industry, passenger cars and other transport are the most important UK sources (National Atmospheric Emissions Inventory, 2022).

NO₂ is a respiratory irritant: short-term exposure to concentrations of NO₂ higher than 200 µg m⁻³ can cause inflammation of the airways and may increase susceptibility to respiratory infections (WHO, 2013). There is a high level of confidence that short-term exposure to NO₂ in outdoor air is associated with all-cause mortality (Orellano, et al., 2020). It has been difficult to identify the direct health effects of NO₂ at ambient concentrations, because it is emitted from the same sources as other pollutants such as particulate matter (PM). However, the WHO's REVIHAAP study (WHO, 2013), COMEAP's 2015 statement, (COMEAP, 2015) and COMEAP's 2018 report on associations of

mortality with NO₂ have reported increasing evidence that NO₂ itself is responsible for health effects. NO is not considered harmful to human health at the concentrations usually found in ambient air but is quickly oxidised to form NO₂.

NO_x can contribute to the formation of other pollutants. In the presence of sunlight, NO_x can react with volatile organic compounds (VOCs) to produce photochemical pollutants including ozone. NO_x also contributes to particulate pollution, via the formation of secondary nitrate particles in the atmosphere.

NO_x can be damaging to the environment. High levels of NO_x deposition can harm plants. It contributes to acidification and eutrophication of terrestrial and aquatic ecosystems, damaging habitats and leading to biodiversity loss.

Peak hourly mean NO₂ concentrations in the UK rarely exceed applicable limit values and objectives, except at some congested urban roadside sites. Prior to 2020, annual mean limit values and objectives were frequently exceeded at roadside sites in the UK, and in many other countries. The extent of these exceedances was substantially reduced in 2020, and in 2021 remained low in comparison with previous years (see sections 4 and 6 for details).

3.1.2 Ozone

Ozone (O₃) is a secondary pollutant produced by the effect of sunlight on NO_x and VOCs from vehicles and industry. O₃ concentrations are therefore typically highest in the summer on hot, sunny, windless days, or days when moderate breezes blow ozone across from continental Europe.

In the upper atmosphere the O₃ layer has a beneficial effect, absorbing harmful ultraviolet radiation from the sun. However, ground level ozone is a pollutant. It irritates the respiratory system and eyes. High levels may exacerbate asthma or trigger asthma attacks in susceptible people and some non-asthmatic individuals may also experience chest discomfort. Evidence is also emerging of links with cardiovascular and metabolic effects and effects due to long-term exposure.

Ozone can cause damage to many plant species leading to loss of yield and quality of crops, damage to forests and impacts on biodiversity. O₃ is also a greenhouse gas implicated in climate change. It can travel long distances, accumulate and reach high concentrations far away from the sources of the pollutants that contributed to its formation. NO_x emitted in cities reduces local O₃ concentrations as NO reacts with O₃ to form NO₂: levels of O₃ are often higher in rural areas than urban areas.

The UK has been compliant with applicable target values since 2009, but most years see long-term objectives exceeded in some areas. Weather conditions during the year determine how widespread such exceedances are.

3.1.3 Particulate Matter: PM₁₀ and PM_{2.5}

PM₁₀ can be 'primary' (emitted directly to the atmosphere) or 'secondary' (formed by the chemical reaction of other pollutants in the air such as SO₂ or NO₂). The main sources of primary PM₁₀ particulate emissions in the UK are: combustion in production processes; industrial; residential and commercial fuel use; as well as agriculture; waste and road transport. In recent years, emissions from residential combustion have increased, both in real terms and as a percentage of the UK total, because of increased use of wood as a domestic fuel. This has offset reductions that have occurred due to decreasing use of coal and other solid fuels. Emissions of particulate matter from road transport include both tailpipe emissions, and tyre and brake wear. Natural sources include wind-blown dust, sea salt, pollens and soil particles.

Like PM₁₀, the finer size fraction PM_{2.5} can be primary or secondary: primary PM_{2.5} has the same main emission sources. Research shows a range of health effects, including respiratory and cardiovascular illness and mortality, associated with PM₁₀. No threshold has been identified below which no adverse health effects occur. In 2016, COMEAP estimated that 722,660 cases of chronic bronchitis could be attributed to anthropogenic particulate pollution, although they considered the evidence insufficient to establish causality (COMEAP, 2016).

PM_{2.5} can penetrate deep into the lungs and research in recent years has strengthened the evidence that both short-term and long-term exposure to PM_{2.5} are linked with a range of health outcomes including (but not restricted to) respiratory and cardiovascular effects.

The UK has been compliant with applicable limit values for PM₁₀ and PM_{2.5} for over a decade. Nonetheless, public health benefits would be expected from further reductions, given that the available evidence has not suggested a threshold for effects.

The environmental effects of particulate pollution are associated with two components of PM: black carbon, which is implicated in climate change, and secondary PM which includes sulphate, nitrate and ammonium. The latter is formed from SO₂, NO_x and NH₃ which are the main drivers for acidification and eutrophication.

3.1.4 Sulphur Dioxide (SO₂)

This acid gas is formed when fuels containing sulphur impurities are burned. The largest UK source of SO₂ is from fuel burning in residential, industrial and commercial settings. Other important sources are manufacturing industry and energy generation. It is a respiratory irritant that can cause constriction of the airways, and people with asthma are considered to be particularly sensitive. Health effects can occur very rapidly, making short-term exposure to peak concentrations important (WHO, 2005), (WHO, 2018), (COMEAP, 2011).

SO₂ deposition is harmful to plants at high concentrations. It contributes to acidification of terrestrial and aquatic ecosystems, damaging habitats and leading to biodiversity loss. SO₂ is also a precursor to the formation of secondary sulphate particles in the atmosphere.

Ambient concentrations of SO₂ in the UK have not exceeded applicable limit values or objectives since 2004.

3.1.5 Carbon Monoxide (CO)

CO is produced when fuels containing carbon are burned with insufficient oxygen to convert all carbon inputs to carbon dioxide (CO₂). Residential fuel use and other stationary combustion are now the largest UK emission sources of CO; road and other transport now account for smaller, but still significant, proportions of emissions (National Atmospheric Emissions Inventory, 2022).

The effects of high levels of CO on human health are well-known. CO is toxic: it affects the ability of the blood to take up oxygen from the lungs and can lead to a range of symptoms, causing death at high concentrations. However, people are more likely to be exposed to dangerous concentrations of CO indoors, due to faulty or poorly ventilated cooking and heating appliances. Cigarette smoke is also a major source of exposure. In the environment, CO can contribute to the formation of ground-level ozone.

The UK has been compliant with all applicable limit values for this pollutant since 1999.

3.1.6 Benzene (C₆H₆)

Benzene (C₆H₆) is an organic chemical compound. Ambient benzene arises from domestic and industrial combustion processes, in addition to road transport. (Defra, 2007).

Benzene is known to cause leukaemia and potentially other cancers in humans (Public Health England, 2019a). Therefore, no safe level can be specified for benzene in ambient air; however the risk increases with increasing exposure. In the environment, benzene can pollute soil and water, leading to exposure via these routes.

Annual mean concentrations of benzene are now low (within limit values and objectives applicable in the UK) due to the introduction of catalytic converters on car exhausts in the 1990s. The UK has been compliant with all applicable limit values for benzene since measurements began in 2003.

3.1.7 Lead (Pb)

Lead (Pb) is a very toxic metallic element. Historically, lead was used as an additive in petrol, and road vehicles were the main source. Leaded petrol was phased out in 1999, resulting in a 98% reduction of pre-1999 UK emissions. Today, the main sources are production processes and transport. However, the contribution from transport comes not

from tailpipe emissions but tyre and brake wear (National Atmospheric Emissions Inventory, 2022). Recent research has found that airborne particulate matter in cities is still 'enriched' with lead, likely due to emissions from historic combustion of leaded petrol (Resongles, et al., 2021).

Lead inhalation can affect red blood cell formation and harm the kidneys, circulatory system, gastrointestinal tract, the joints, reproductive systems, and can cause acute or chronic damage to the central nervous system (CNS). The unborn child and young children are the most sensitive to lead toxicity (Public Health England, 2016). Long-term low-level exposure has been shown to affect intellectual development in young children and the unborn child (EPAQS, 2009).

In the environment, Pb can pollute soil and surface waters. Exposure to contaminated soil and water may then become a health risk. Lead may accumulate in other organisms such as fish and be passed up the food chain. The UK has been compliant with applicable limit values for ambient lead in air for over 20 years.

3.1.8 Nickel (Ni)

Nickel (Ni) is a toxic metallic element found in ambient air as a result of releases from oil and coal combustion, metal processes, manufacturing and other sources. Currently the main UK emission source is the combustion of petroleum coke, solid fuels containing petroleum coke, and heavy fuel oil, in residential and industrial settings (National Atmospheric Emissions Inventory, 2022). A small number of UK zones continue to exceed applicable target values for annual mean Ni, as has been the case for many years, due to local industrial emissions.

Nickel compounds are human carcinogens by inhalation exposure. Nickel can cause irritation to the nose and sinuses and allergic responses and can lead to the loss of the sense of smell. Long-term exposure may lead to respiratory diseases and cancers (WHO, 2018).

As well as ambient air, Ni can pollute soil and water, leading to exposure via these routes.

3.1.9 Arsenic (As)

Arsenic (As) is a toxic metalloid which occurs naturally in the environment. Arsenic is emitted into the atmosphere in the form of particulate matter. Historically the largest source was coal combustion, but this has declined: the largest UK source of arsenic emissions is now the open burning of wood treated with preservatives containing As (National Atmospheric Emissions Inventory, 2022). The UK has been compliant with applicable target values for As for many years. Inhalation of air containing high levels of As can cause lung damage, shortness of breath, chest pain and cough (Public Health England, 2019b). Arsenic compounds may be corrosive and can cause burns to the skin or eyes on contact. Long term inhalation exposure is associated with genotoxic and

carcinogenic effects. Arsenic can also pollute soil and water, leading to exposure via these routes. Arsenic in water or soil can be taken up by plants or fish. Food is the largest source of As exposure for most people in the general population.

3.1.10 Cadmium (Cd)

Cadmium (Cd) is a toxic metallic element. The main emission sources are combustion in the manufacturing industry and production processes. The incineration of municipal solid waste was once a significant source, but improved controls on waste to energy plant in the 1990s have reduced their contribution to 2% of the UK 2020 total (National Atmospheric Emissions Inventory, 2022). The UK has been compliant with applicable target values for Cd for many years.

Acute inhalation exposure to Cd causes effects on the lung such as pulmonary irritation. Chronic exposure via inhalation can lead to lung cancer (WHO, 2018) or cause a build-up of Cd in the kidneys that can lead to kidney disease. In the environment, Cd can pollute soil and water, leading to exposure via these routes.

3.1.11 Mercury (Hg)

Mercury (Hg) is released to the air by human activities. The main current UK sources are coal use in public electricity and heat production and industrial combustion, iron and steel production processes, cremation, and emissions from the disposal of products containing mercury (National Atmospheric Emissions Inventory, 2022).

Acute exposure to high levels of Hg can cause chest pain and shortness of breath and affect the CNS and kidneys. Chronic exposure leads to CNS disorders, kidney damage and stomach problems.

In the environment, Hg can also pollute soil, fresh water and sea water. Exposure to contaminated soil and water may then become a health risk. Mercury may accumulate in other organisms such as fish, and be passed up the food chain.

3.1.12 Polycyclic Aromatic Hydrocarbons (PAH)

Polycyclic aromatic hydrocarbons (PAHs) are a large group of chemical compounds which usually occur as complex mixtures rather than as individual compounds. One particular PAH, **Benzo[a]pyrene (B[a]P)** is used as a 'marker' for this group of compounds. The main sources of B[a]P in the UK are residential, commercial and industrial fuel combustion (National Atmospheric Emissions Inventory, 2022). A small number of UK zones continue to exceed applicable target values for B[a]P, as has been the case for many years.

PAHs are persistent, bio-accumulative, organic compounds with toxic and carcinogenic effects. The International Agency for Research on Cancer (IARC) has classified several PAH, including B[a]P, as causing cancer in humans (Public Health England, 2018). B[a]P

is currently considered the most carcinogenic PAH. PAHs can bio-accumulate and be passed up the food chain.

3.2 Assessment of Air Quality in the UK

The evidence base for the annual assessment of compliance is based on a combination of measurements from the UK national monitoring networks and the results of modelling assessments. The use of models enables air quality to be assessed at locations without monitoring sites and reduces the number of monitoring stations required. It has the added benefit of providing additional information on source apportionment and projections to support the development and implementation of air quality policies.

UK compliance assessment modelling is undertaken using national models known as the Pollution Climate Mapping (PCM) models. The PCM models have been designed to assess compliance with limit values, target values and long-term objectives at locations defined within the Air Quality Standards Regulations. Modelled compliance assessments are undertaken for 11 air pollutants each year. This assessment needs to be completed each year in the relatively short period between the time when the input data (including ratified monitoring data and emission inventories) become available and the reporting deadline at the end of September.

It is important to understand the differences between modelling carried out for compliance assessment purposes, and that carried out for Local Air Quality Management. National air quality modelling for the UK focuses on two components: pollutant concentrations at background locations, on a 1x1 km grid square basis, and roadside pollutant concentrations, at four metres from the kerb of urban major road links¹. By contrast, Local Air Quality Management (LAQM) modelling is different in scope, purpose and methodology from the national modelling and will usually output contour plots showing dispersion away from the source, on a fine resolution grid. The level of detail and resolution of LAQM modelling is therefore much greater in order to focus on local exposure and hotspots and does not necessarily meet the requirements for air quality assessment under the Air Quality Standards Regulations. See **Section 3.5** for more details on the modelling carried out for compliance assessment.

3.2.1 Current UK Air Quality Monitoring

During 2021 there were 529 national air quality monitoring sites across the UK, comprising several networks, each with different objectives, scope and coverage. This section

¹ A road link is a section of road that is greater than 100m in length.

provides a brief description of those used to monitor compliance with the Air Quality Standards Regulations. A summary of the UK national networks is provided in **Table 3-1** (the number of sites shown in this table amounts to considerably more than 529 because some sites belong to more than one network). This table shows the number of sites in operation during part or all of 2021.

Table 3-1 The UK's Air Quality Monitoring Networks in 2021

Network	Pollutants	Number of Sites operating in 2021
Automatic Urban and Rural Network (AURN)	CO, NO _x , NO ₂ , SO ₂ , O ₃ , PM ₁₀ , PM _{2.5} .	157
Automatic London Network (part of AURN)		16
UK Heavy Metals Network	Metals in PM ₁₀ including: As, Cd, Co, Cr, Cu, Fe, Mn, Ni, Pb, Se, V, Zn. Measured deposition including: Al, As, Ba, Be, Cd, Co, Cr, Cs, Cu, Fe, Li, Mn, Mo, Ni, Pb, Rb, Sb, Sc, Se, Sn, Sr, Ti, U, V, W, Zn. Hg deposition Total gaseous mercury	24
Non-Automatic Hydrocarbon	Benzene	35
Automatic Hydrocarbon	Range of volatile organic compounds (VOCs)	5
Polycyclic Aromatic Hydrocarbons (PAH).	27 PAH species including benzo[a]pyrene	34
European Monitoring and Evaluation Programme (EMEP)	Wide range of parameters relating to air quality, precipitation, meteorology and composition of aerosol in PM ₁₀ and PM _{2.5} .	2
Particle Numbers and Concentrations Network	Total particle number, concentration, size distribution, anions, EC/OC, speciation of PM ₁₀ and PM _{2.5} .	4

Network	Pollutants	Number of Sites operating in 2021
Toxic Organic Micropollutants	Range of toxic organics including dioxins and dibenzofurans.	6
UK Eutrophying and Acidifying Pollutants: NO ₂ Net (rural diffusion tubes)	NO ₂ (rural)	24
UK Eutrophying and Acidifying Pollutants: AGANet	HNO ₃ , HONO, SO ₂ , Ca, Cl, Mg, Na, NO ₂ , NO ₃ and SO ₄	27
UK Eutrophying and Acidifying Pollutants: NAMN	NH ₃ and/or NH ₄	74
UK Eutrophying and Acidifying Pollutants: PrecipNet	Major ions in rain water	41
Black Carbon	Black Carbon	14
Upland Waters Monitoring Network	Chemical and biological species in water	10
Rural Mercury Network	Tekran analyser used to measure mercury in PM _{2.5} , reactive mercury and elemental mercury at Auchencorth Moss, and total gaseous mercury at Chilbolton Observatory.	2
UK Urban NO ₂ Network	Diffusion tubes with wind-protection membranes measuring NO ₂ monthly at urban traffic-related sites.	292

3.2.1.1 The Automatic Urban and Rural Network (AURN)

The AURN is currently the largest automatic monitoring network in the UK and forms a large part of the UK's statutory compliance monitoring evidence base. Data from the AURN are available on Defra's online UK Air Information Resource, UK-AIR at <https://uk-air.defra.gov.uk/>. The Automatic London Network (ALN) is a subset of sites in the AURN

which also form part of the wider London Air Quality Network (LAQN). In this report, 'AURN' includes the whole network, i.e. including the ALN subset of sites.

The techniques used for monitoring gaseous pollutants within the AURN are the reference measurement methods defined in the Air Quality Standards Regulations. For particulate matter the AURN uses methods which have demonstrated equivalence to the reference method, but which (unlike the reference method) allow continuous on-line monitoring. Details are provided in **Table 3-2**.

Table 3-2 AURN Measurement Techniques

Pollutant	Method used, including details of CEN Standard Methods
O₃	EN 14625:2012 'Ambient air quality – standard method for the measurement of the concentration of ozone by ultraviolet photometry' (CEN, 2005a)
NO₂/NO_x	EN 14211:2012 'Ambient air quality - Standard method for the measurement of the concentration of nitrogen dioxide and nitrogen monoxide by chemiluminescence' (CEN, 2005b)
SO₂	EN 14212:2012 'Ambient air quality – Standard method for the measurement of the concentration of sulphur dioxide by UV fluorescence' (CEN, 2005c)
CO	EN 14626:2012 'Ambient air quality - Standard method for the measurement of the concentration of carbon monoxide by non-dispersive infrared spectroscopy' (CEN, 2005d)
PM₁₀ and PM_{2.5}	<p>EN 12341:2014 'Ambient air quality - Standard gravimetric measurement method for the determination of the PM₁₀ or PM_{2.5} mass fraction of suspended particulate matter' (BS EN, 2014)</p> <p>In 2021 the AURN used four methods which are equivalent to the reference method for one or both metrics: the Fidas™ 200, an optical technique, the Beta-Attenuation Monitor (BAM), the Filter Dynamic Measurement System (FDMS), and gravimetric samplers (at two sites) that collect daily samples onto a filter for subsequent weighing. Descriptions of these methods are given in the Glossary of this report.</p> <p>The FDMS instruments were of a type no longer supported by the manufacturer and were reaching the end of their working lives. A phased replacement programme began in 2018, in which the AURN's FDMS were all replaced with either a BAM or Fidas™. This was completed in 2021, so at the time of writing there are no longer any FDMS in the AURN.</p>

3.2.1.2 The UK Heavy Metals Network

The UK Heavy Metals Network forms the basis of the UK's compliance monitoring for the Air Quality Standards Regulations, which cover lead, arsenic, cadmium, nickel and mercury.

At the end of 2013 Defra merged the existing Urban and Industrial Network with the Rural Network to form the UK Heavy Metals Network. The merged network monitors a range of elements (not all of which are classified as heavy metals) at urban, industrial and rural sites, using a method equivalent to the CEN standard method (CEN, 2005e). Metals (As, Cd, Co, Cr, Cu, Fe, Mn, Ni, Pb, Se, V and Zn) in PM₁₀ are measured at 23 sites. The network stopped measuring mercury in PM₁₀ in 2014.

Metal deposition (Al, As, Ba, Be, Cd, Co, Cr, Cs, Cu, Fe, Hg, Li, Mn, Mo, Ni, Pb, Rb, Sb, Se, Sn, Sr, Ti, U, V, W, Zn) was measured at the following rural sites: Auchencorth Moss, Chilbolton Observatory, Heigham Holmes and Yarner Wood. The same metals were measured at Lough Navar with the exception of mercury.

The network stopped measuring total gaseous mercury in August 2018.

3.2.1.3 Non-Automatic Hydrocarbon Network

In this network, ambient concentrations of benzene are measured by the CEN standard method (CEN, 2005f). This involves pumping air through an adsorption tube to trap the compound, which is later analysed in a laboratory. This network monitors compliance with the Air Quality Standards Regulations limit value for benzene. All sites in the Non-Automatic Hydrocarbon Network are co-located with AURN sites.

3.2.1.4 Automatic Hydrocarbon Network

The Air Quality Standards Regulations also require measurement and reporting of ozone precursor substances (29 species), which include volatile organic compounds (VOCs). The Air Quality Standards Regulations refer to Annex X (ten) of the Air Quality Directive which provides a list of compounds recommended for measurement.

Ozone precursor measurement is carried out by the Automatic Hydrocarbon Network. Automatic hourly measurements of a range of hydrocarbon species (including all those specified in Annex X of the Air Quality Directive (European Parliament and Council of the European Union, 2008) except formaldehyde and total non-methane hydrocarbons), are made at four sites using automated pumped sampling with *in-situ* gas chromatography. The VOCs monitored include benzene, which is covered by the Air Quality Standards Regulations as a pollutant in its own right.

3.2.1.5 PAH Network

The PAH Network monitors compliance with the Air Quality Standards Regulations, which include a target value of 1 ng m⁻³ for the annual mean concentration of benzo[a]pyrene as

a representative PAH. Samples are collected on filters using the PM₁₀ 'Digitel' sampler. Samples are subsequently analysed in a laboratory for 23 PAH compounds.

3.2.1.6 EMEP

EMEP (European Monitoring and Evaluation Programme) is a programme set up to provide governments with qualified scientific information on air pollutants, under the UNECE Convention on Long-range Transboundary Air Pollution. There are currently two EMEP 'supersites', at Auchencorth Moss in Midlothian (representing the north of the UK) and at Chilbolton Observatory in Hampshire (representing the south). The site at Chilbolton replaced the long running site at Harwell at the start of 2016. A representativeness analyses showed that both sites were similar in their rural background nature. A very wide range of measurements are taken at EMEP sites, supplemented by data from other UK networks which are co-located.

Monitoring includes:

- Hourly meteorological data,
- Soil and vegetation measurements,
- Metallic elements in PM₁₀ and precipitation,
- Deposition of inorganic ions,
- Major ions in PM_{2.5} and PM₁₀, as well as HCl, HNO₂, HNO₃, NH₃ and SO₂,
- Trace gases (ozone, NO_x and SO₂),
- Black carbon, organic carbon (OC) and elemental carbon (EC),
- Ammonia (monthly),
- Daily and hourly PM₁₀ and PM_{2.5} mass,
- Volatile Organic Compounds,
- Carbonyls,
- CH₄ and N₂O fluxes.

3.2.1.7 Particle Numbers and Concentrations Network

The Air Quality Standards Regulations require that the chemical composition of PM_{2.5} is characterised at background locations in the United Kingdom. The Particle Numbers and Concentrations Network sites contribute to this statutory requirement. During 2021, the network consisted of four measurement sites; two rural sites (Auchencorth Moss and

Chilbolton Observatory), and two in London (London Marylebone Road and London Honor Oak Park; the latter site replaced North Kensington in November 2018).

Among the parameters measured are:

- Total particle numbers per cubic centimetre of ambient air,
- Particle numbers in different particle size fractions,
- Major ions (ammonium, nitrate and sulphate) in PM_{2.5} and PM₁.
- Total carbon, organic carbon (OC) and elemental carbon (EC) concentrations in PM_{2.5}.

PM₁₀ speciation was replaced by PM_{2.5} speciation in 2018. PM₁ speciation began at the London sites in 2020.

As well as its statutory function, this network provides data on the chemical composition of particulate matter, primarily for the use of researchers of atmospheric processes, epidemiology and toxicology.

Measurements of elemental carbon (EC) and organic carbon (OC) began at Auchencorth Moss at the start of 2011 and Chilbolton Observatory at the start of 2016. EC and OC measurements were made using a thermal/optical method involving both reflectance and transmission correction methods. Comparing both correction methods aims to provide valuable understanding of the measurement process for EC and OC.

3.2.1.8 TOMPs Network

This research-based network monitors a range of toxic organic micropollutants (compounds that are present in the environment at very low concentrations but are highly toxic and persistent). These include dioxins, dibenzofurans and polychlorinated biphenyls. The TOMPs Network consists of six sites: Auchencorth Moss, Hazelrigg, High Muffles, London Nobel House, Manchester Law Courts and Weybourne.

The purpose of the TOMPs Network is to provide data on these air pollutants, and to support the development of policy to protect the environment and human health. Further information on the TOMPs Network can be found on UK-AIR at <https://uk-air.defra.gov.uk/networks/network-info?view=tomps>. However, this network is not used for compliance monitoring and will not be discussed further in subsequent sections of the report.

3.2.1.9 UK Eutrophying and Acidifying Pollutants Network

The UK Eutrophying and Acidifying Atmospheric Pollutants (UKEAP) network provides information on deposition of eutrophying and acidifying compounds in the UK and

assessment of their potential impacts on ecosystems. The UKEAP network is an 'umbrella' project covering four groups of sites:

- The UKEAP rural NO₂ diffusion tube network (NO₂Net). This measures NO₂ concentrations at 24 locations as required for input to the rural NO_x concentration field in the Pollution Climate Model.
- In 2021 the Acid Gas and Aerosol Network (AGANet) comprised a total of 27 sites. The network measures a range of gases and aerosol components. Samples are collected monthly and are analysed by either inductive coupled plasma optical emission spectrometry (ICP-OES) or ion chromatography.
- The National Ammonia Monitoring Network (NAMN) which characterizes ammonia and ammonium concentrations using both passive samplers (Alpha Samplers) and low volume denuders (Delta Samplers) at 74 locations.
- The Precipitation Network (PrecipNet), measuring major ions in precipitation at 41 rural sites. Eight of these sites form part of the Long-Term Monitoring Network managed by Natural England. The UKEAP network allows estimates of sulphur and nitrogen deposition. Samples are collected fortnightly at all sites and daily at two sites.

3.2.1.10 Black Carbon Network

Black carbon is fine, dark carbonaceous particulate matter produced from the incomplete combustion of materials containing carbon (such as coal, oil, and biomass such as wood). It is of concern due to health effects, and also as a suspected contributor to climate change. In 2021, the Black Carbon Network measured black carbon at 14 sites using the Aethalometer™ automated instrument. The Aethalometer™ measures black carbon directly, using a real-time optical transmission technique. The objectives of the network are as follows:

- To maintain coverage of black carbon measurements across the whole UK;
- To maintain continuity of historic datasets;
- To gather data for epidemiological studies of black carbon and health effects;
- To gather information about black carbon PM sources in the UK;
- To assess PM reductions from air quality management interventions;
- To quantify the contribution of wood burning to black carbon and ambient PM in the UK; and
- To gather data to address future policy considerations including black carbon and climate change.

3.2.1.11 UK Upland Waters Monitoring Network (UK UWMN)

The UK Upland Waters Monitoring Network (UWMN) was set up in 1988 (then called the Acid Waters Monitoring Network) by ENSIS Ltd, at University College London (UCL) under funding from the then Department of the Environment (later DETR and Defra). Its objective was to assess the chemical and biological response of acidified lakes and streams in the UK to the planned reduction in emissions. It was initially designed to provide chemical and biological data on the extent and degree of surface water acidification in the UK uplands and underpin the science linking acid deposition to water quality and aquatic ecosystem health. In recent years it has been adapted to address a wider range of questions, particularly with respect to understanding impacts of nitrogen enrichment, the influence of climate change and land use on upland waters, and interactions between these drivers and recovery from acidification.

The eleven lakes and eleven streams were originally selected to cover a wide deposition gradient and included forest-moorland pairs of sites. Sites were required to be subject to minimal point source pollution and catchment disturbance beyond that caused by traditional upland land use practices such as sheep grazing or forestry. Additional stream sites have recently been added to broaden the acid-sensitivity gradient, while thermistor loggers are now deployed to continuously monitor water temperature. Water chemistry has been monitored monthly in streams and quarterly in lakes ever since the inception of the network to the present. Biological monitoring involves annual assessment of algae (diatoms), higher aquatic plants and macroinvertebrates. Fish monitoring was discontinued in 2015 due to budget cuts. In April 2019, the Centre for Ecology & Hydrology (now UKCEH) took over management of the UWMN from ENSIS Ltd. After a significant funding hiatus from 2016, Defra are again supporting collection and analysis of biological samples in 2021. The UK UWMN also receives funding from the National Environment Research Council (NERC) via the UK Centre for Ecology & Hydrology (UKCEH), Scottish Natural Heritage, the Welsh Government, Natural Resources Wales, Forest Research and Moors for the Future, and also benefits from considerable in-kind support for sampling and survey activity from UCL, Queen Mary University of London, the Scottish Environment Protection Agency (SEPA), the Department of Agriculture, Environment and Rural Affairs (DAERA) in Northern Ireland, and several private volunteers.

3.2.1.12 Rural Mercury Monitoring

The Tekran instrument at Auchencorth Moss measures the mercury composition of PM_{2.5} as well as mercury in its elemental and reactive forms, whereas at Chilbolton Observatory it measures just total gaseous mercury.

3.2.1.13 UK Urban NO₂ Network

The UK Urban NO₂ Network (UUNN) was established in December 2019 with monitoring beginning in January 2020. The objective of the network is to provide additional local roadside NO₂ measurements to enhance the UK's national compliance assessment. Monitoring of NO₂ is undertaken on the UUNN using Palmes-type diffusion tubes with wind

protection membranes. During 2020 monitoring was undertaken at 177 locations across England and Wales: the network expanded further at the start of 2021 and now comprises 292 sites across Great Britain.

3.2.1.14 Air Pollution Impacts on Ecosystem Networks (APIENs)

The following information about UK APIENs is summarised from the APIS website at <https://www.apis.ac.uk/APIENs>. The purpose of UK APIENs is to monitor and report the negative impacts of air pollution (e.g. acidification, eutrophication, ozone damage or changes in biodiversity) on ecosystems that are representative of freshwater, natural and semi-natural habitats and forests in the UK. It was formed in 2018 by integrating UK national air quality and ecosystem monitoring networks and surveys, to meet UK monitoring and reporting obligations under the EU National Emissions Ceilings Directive. Integrated data from APIENs will provide the evidence to determine the state of UK ecosystems, and provide baseline against which any changes and potential recovery can be compared.

3.2.2 Quality Assurance and Quality Control

Air quality monitoring in the UK is subject to rigorous procedures of validation and ratification. The well-established monitoring networks each have a robust and documented Quality Assurance and Quality Control (QA/QC) programme designed to ensure that measurements meet the defined standards of quality with a stated level of confidence. Essentially, each programme serves to ensure that the data obtained are:

- Representative of ambient concentrations existing in the various areas under investigation.
- Sufficiently accurate and precise to meet specified monitoring objectives.
- Comparable and reproducible. Results must be internally consistent and comparable with international or other accepted standards, if these exist.
- Consistent over time. This is particularly important if long-term trend analysis of the data is to be undertaken.
- Representative over the period of measurement; for most purposes, a yearly data capture rate of not less than 90% is usually required for determining compliance with limit values where applicable. An allowance of 5% is made in some cases for down-time due to planned maintenance.
- Consistent with Data Quality Objectives. The uncertainty requirements of the Air Quality Standards Regulations are specified as data quality objectives. In the UK, all air quality data meet the data quality requirements of the Air Quality Standards Regulations in relation to uncertainty.

- Consistent with methodology guidance defined in the Air Quality Standards Regulations for relevant pollutants and measurement techniques. The use of tested and approved analysers that conform to Standard Method (or equivalent) requirements and harmonised on-going QA/QC procedures allows a reliable and consistent quantification of the uncertainties associated with measurements of air pollution.

Most UK networks use a system of regular detailed audits of all monitoring equipment at every site. These audits supplement more regular calibrations and filter changes and test all critical parameters of the measuring equipment including, where appropriate, linearity, converter efficiency (in the case of NO_x analysers) response time, flow rate etc.

Data verification is the process of checking and validating the data. (The term 'ratification' is used in some networks). Data uploaded to the Defra UK Air Information Resource (UK-AIR at <https://uk-air.defra.gov.uk/>) in near real time are provided as provisional data. All these data are then carefully screened and checked via the verification process. The verified data then overwrite the provisional data on the website. It should however be noted that there are occasionally circumstances where data which have been flagged as 'Verified' could be subject to further revision. This may be for example where:

- A QA/QC audit has detected a problem which affects data from earlier verification periods.
- Long-term analysis has detected an anomaly between expected and measured trends which requires further investigation and possible data correction.
- Further research comes to light which indicates that new or tighter QA/QC criteria are required to meet the data quality objectives. This may require review and revision of historical data by applying the new criteria.

Only verified data are included in the UK's assessment of compliance with the Air Quality Standards Regulations.

Further details on the QA/QC procedures appropriate to each network can be obtained from the annual reports of the relevant monitoring networks, and from the report '*Quality Assurance and Quality Control (QA/QC) Procedures for UK Air Quality Monitoring under 2008/50/EC and 2004/107/EC*' available from Defra's UK-AIR website (Defra, 2016b).

3.3 Modelling

3.3.1 Why Do Modelling?

The UK's monitoring programmes are supplemented by air quality modelling. There are several benefits of using modelling to complement the monitoring data gathered across the UK national monitoring networks:

- Coverage of the whole UK rather than specific locations where there is a monitoring site. Whilst our monitoring network is extensive, a monitoring site might not fully represent the wider region in which it is located due to local characteristics such as buildings affecting dispersion, localised or temporary sources.
- Providing information about the sources of pollutants to inform policy development.
- A reduction in the number of fixed continuous monitoring locations required for compliance with the UK Air Quality Standards Regulations – freeing up resources and ensuring value for money.
- Providing a framework within which to assess different air quality scenarios – for example projecting concentrations forward to assess levels in future years, in order to develop policies to continue to improve air quality in the UK.

3.3.2 How the Models Work

The national modelling methodology varies between pollutants. The detailed methodology is explained in a technical report (Pugsley, K. L. et al., 2022) (the latest versions of these can be found in the Library section of Defra's UK-AIR website (Defra, 2022b)).

Defra's air quality national modelling assessment for the UK consists of two components:

- Background concentrations – on a 1x1km resolution, representing ambient air quality concentrations at background locations.
- Roadside concentrations – concentrations at the roadside of urban major road links throughout the UK (i.e. motorways and major A-roads). There are approximately 9,000 of these urban major road links.

Roadside concentrations are not modelled for CO, SO₂, ozone, benzo[a]pyrene and metals as these are deemed not to have significant traffic-related sources.

The models have been designed to assess compliance at locations defined by the Air Quality Standards Regulations (UK Government, 2010) as relevant for air quality assessment.

3.3.3 Background Air Quality

The 1x1 km background maps are made up of several components which are modelled separately and then added together to make the final grid of the UK. These individual components (supplemented by some additional components for certain pollutants) are:

- Large point sources (e.g. power stations, steel works and oil refineries),
- Small point sources (e.g. boilers in town halls, schools or hospitals and crematoria),
- Distant sources (characterised by the rural background concentration),
- Local area sources (e.g. road traffic, domestic and commercial combustion and agriculture).

In order to ensure that these ambient concentrations from area sources are representative of the real-world situation, they are validated against measurements taken from the national networks (including the AURN). After the validation has been completed the large points, small points, distant sources and area source components are added together to provide the final background concentrations.

3.3.4 Roadside Air Quality

Roadside concentrations are determined by using a roadside increment model which estimates the contribution from road traffic sources and adds this to the modelled background concentrations discussed above.

For each of the road links that are modelled, there are emission estimates for each pollutant from the National Atmospheric Emissions Inventory (NAEI, (UK National Atmospheric Emissions Inventory, 2022)) and road traffic counts from the Department for Transport. A measured roadside increment concentration is calculated for road links with a roadside monitoring station by subtracting the link's modelled background concentration (from the 1x1 km modelled maps) from the relevant measured roadside concentration. A roads kernel model (RKM) is used to calculate a modelled roadside increment concentration for each road link by applying the NAEI emissions and road traffic counts (annual average daily traffic flow) in a dispersion model. The RKM is calibrated by comparing the measured roadside increment concentrations at roadside monitoring stations with the modelled roadside increment concentrations for these road links. The application of the RKM ensures that a process-based modelling approach is used to determine the local component of roadside concentrations, including factors influencing dispersion at the roadside e.g. road orientation, width, and additional vehicle induced turbulence.

3.4 Access to Assessment Data

Data from the UK's air quality monitoring networks and annual compliance modelling is available under the Open Government Licence (UK Government, 2022b) from UK-AIR.

Defra has produced a searchable online catalogue of air quality and emissions datasets which allows people to browse the extent of data available and access key metadata. This is available at <https://uk-air.defra.gov.uk/data/data-catalogue>.

Historical monitoring data can be accessed through the data selector tools in UK-AIR, at <https://uk-air.defra.gov.uk/data/>. Modelled data from the Pollution Climate Mapping model are available as .csv files for download from the modelled air quality data pages at <https://uk-air.defra.gov.uk/data/modelling-data> or can be accessed through the Ambient Air Quality Interactive Map at <https://uk-air.defra.gov.uk/data/gis-mapping> - a GIS (geographical information system) based tool which provides enhanced visualisation capability and access to roadside concentration data.

UK-AIR also houses a Compliance Dashboard which displays all the underlying data used in the compliance assessment. The Compliance Dashboard can be found at <https://uk-air.defra.gov.uk/compliance-data> and is made up of three parts:

- **Interactive GIS Compliance map** – a streamlined viewer facilitating summaries of compliance status across different geographies for different pollutant metrics based on modelled background data, modelled roadside data and measurements.
- **Compliance data hub** – a comprehensive data catalogue and extraction tool for underlying data that serves the compliance app. This allows users to acquire the data behind the compliance status either for a specific zone/agglomeration or Local Authority or for the whole of the UK in one process.
- **XML file library** – a catalogue of download links for the machine-readable XML formats of the compliance data.

4 Assessment of Compliance

4.1 Definition of Zones

The UK is divided into 43 zones for air quality assessment. There are 28 agglomeration zones (large urban areas) and 15 non-agglomeration zones. Each zone has an identification code (**Table 4-1**). Zones are shown in **Figure 4-1**.

Table 4-1 UK Zones for Ambient Air Quality Reporting 2021

Zone	Zone code	Zone type
Greater London Urban Area	UK0001	Agglomeration
West Midlands Urban Area	UK0002	Agglomeration
Greater Manchester Urban Area	UK0003	Agglomeration
West Yorkshire Urban Area	UK0004	Agglomeration
Tyneside	UK0005	Agglomeration
Liverpool Urban Area	UK0006	Agglomeration
Sheffield Urban Area	UK0007	Agglomeration
Nottingham Urban Area	UK0008	Agglomeration
Bristol Urban Area	UK0009	Agglomeration
Brighton/Worthing/Littlehampton	UK0010	Agglomeration
Leicester Urban Area	UK0011	Agglomeration
Portsmouth Urban Area	UK0012	Agglomeration
Teesside Urban Area	UK0013	Agglomeration
The Potteries	UK0014	Agglomeration
Bournemouth Urban Area	UK0015	Agglomeration
Reading/Wokingham Urban Area	UK0016	Agglomeration
Coventry/Bedworth	UK0017	Agglomeration
Kingston upon Hull	UK0018	Agglomeration
Southampton Urban Area	UK0019	Agglomeration
Birkenhead Urban Area	UK0020	Agglomeration
Southend Urban Area	UK0021	Agglomeration
Blackpool Urban Area	UK0022	Agglomeration
Preston Urban Area	UK0023	Agglomeration
Glasgow Urban Area	UK0024	Agglomeration
Edinburgh Urban Area	UK0025	Agglomeration
Cardiff Urban Area	UK0026	Agglomeration
Swansea Urban Area	UK0027	Agglomeration
Belfast Metropolitan Urban Area	UK0028	Agglomeration
Eastern	UK0029	Non-agglomeration
South West	UK0030	Non-agglomeration
South East	UK0031	Non-agglomeration
East Midlands	UK0032	Non-agglomeration
North West & Merseyside	UK0033	Non-agglomeration
Yorkshire & Humberside	UK0034	Non-agglomeration
West Midlands	UK0035	Non-agglomeration
North East	UK0036	Non-agglomeration
Central Scotland	UK0037	Non-agglomeration
North East Scotland	UK0038	Non-agglomeration
Highland	UK0039	Non-agglomeration
Scottish Borders	UK0040	Non-agglomeration
South Wales	UK0041	Non-agglomeration
North Wales	UK0042	Non-agglomeration
Northern Ireland	UK0043	Non-agglomeration

Figure 4-1 UK Zones for Ambient Air Quality Reporting 2021



© Crown copyright. All rights reserved Defra, Licence number 100022861 [2021]

4.2 Air Quality Assessment for 2021

The air quality assessment for each pollutant is derived from a combination of measured pollutant concentrations from the Automatic Urban and Rural Network (AURN) and supplementary assessment (that is, modelling using the Pollution Climate Mapping (PCM) model, supplementary NO₂ diffusion tube measurements from the UK Urban Nitrogen Dioxide Network - the UUNN - or objective estimation, as explained in Defra's technical report on UK air quality assessment (Pugsley, K. L. et al., 2022). Where both measurements and supplementary assessment results are available for a zone, the assessment of compliance for each zone is based on the higher concentration of the two.

However, in the case of NO₂, an additional rule has been introduced in 2021. This is used where there is roadside monitoring (an Automatic Urban and Rural Network monitoring site, a UUNN diffusion tube monitoring site, or both) on a major urban road, which is also modelled by the PCM model. This rule determines the order of precedence of these data sources when used in compliance assessment and is described in section 4.2.1 below.

4.2.1 New Approach for Nitrogen Dioxide at Roadside

In previous compliance assessments, including in 2020, the approach taken when assessing NO₂ concentrations at roadside locations where both modelled and measured concentrations were available was to report all concentrations, but to always use the highest concentration to determine the compliance status, whether measured or modelled. This is a conservative approach in which an exceedance is always reported if any of the data indicate one, but it does not consider the quality of the evidence available. The availability of a new source of evidence measurements from the UUNN, which was established in 2020 - prompted a review of the approach for NO₂.

A study led by Defra working closely with members of their independent Air Quality Expert Group (AQEG) compared the quality of modelled NO₂ concentrations from the PCM model to measured concentrations from the UUNN and AURN. This concluded that the AURN provides the most accurate assessment of NO₂ concentrations, followed by the UUNN, and then the PCM model.

The method for determining compliance with the annual mean limit value for NO₂ has therefore been adjusted to reflect this, so that the best available evidence is used. As in previous years, all modelled and measured NO₂ concentrations are reported as part of the assessment, but the order of precedence, for any given major urban road, is as follows:

1. If AURN measurements are available, these have been used to assess compliance in preference to values from the UUNN and/or the PCM model for the same major urban road:
2. If UUNN measurements (but not AURN measurements) are available, the UUNN measurements have been used to assess compliance in preference to values from the PCM model for the same major urban road.

3. If no AURN or UUNN measurements are available, concentrations from the PCM model have been used to assess compliance.

This order of precedence only applies to results for the same major urban road. So, compliance or exceedance of a particular zone for NO₂ could in theory still be determined on the basis of modelling, if the highest concentration for that zone was a modelled value for a location without co-located monitoring.

No change has been made to the method for determining compliance for other pollutants. This means that the most appropriate evidence-based approach is taken for each pollutant.

4.2.2 Compliance Summary

The air quality compliance assessment for each calendar year must be published by Defra by 30th September the following calendar year. The results of the air quality assessment for 2021 are summarised in the tables below. The tables have been completed as follows:

- Where all measurements were within the relevant limit values in 2021, the table shows this as 'OK'.
- In the above cases, where compliance was determined by supplementary assessment only, this is indicated by '(s only)', i.e. 'OK (s only)'.
- Where locations were identified as exceeding a limit value, target value or long-term objective, this is identified as '>LV', '>TV' or '>LTO' as applicable.
- Where a non-compliance was determined by supplementary assessment, this is indicated by '(s only)', as above.
- The abbreviation 'n/a' (not applicable) means that an assessment is not relevant for this zone, such as for the NO_x vegetation critical level in agglomeration zones.
- Zones that complied with the relevant limit values, targets or long-term objectives are shaded blue, while those that did not are shaded red. For ozone, zones that met the relevant target value but not the long-term objective are shaded purple.

There are no longer any zones where margins of tolerance apply.

Sulphur dioxide (SO₂): in 2021, all zones and agglomerations within the UK complied with the limit values for 1-hour mean and 24-hour mean SO₂ concentration, set for protection of human health. All non-agglomeration zones within the UK also complied with the critical levels for annual mean and winter mean SO₂ concentration, set for protection of ecosystems (these are not applicable to built-up areas).

Carbon monoxide (CO), benzene and lead: all zones and agglomerations were compliant with the limit values for these three pollutants in 2021. The 2021 compliance assessment for CO has been based on objective estimation, as explained in Defra's technical report on UK air quality assessment (Pugsley, K. L. et al., 2022). This is underpinned by NAEI emission trends, AURN measurement trends and historical modelling assessments.

Nitrogen dioxide (NO₂): in 2021 not all zones and agglomerations were compliant with the limit values. The results of the air quality assessment for nitrogen dioxide for each zone are summarised in **Table 4-2**.

All zones and agglomerations were compliant with the 1-hour limit value (200 µg m⁻³) in 2021, with none exceeding this limit value on more than the permitted 18 occasions. In recent years only a few zones (typically one or two) have exceeded this limit value, 2021 is the second year in which all zones have been compliant.

Thirty-three zones met the annual mean limit value for NO₂ (40 µg m⁻³) in 2021. The 10 zones that exceeded this limit value were:

- Greater London Urban Area
- West Midlands Urban Area
- Greater Manchester Urban Area
- West Yorkshire Urban Area
- Liverpool Urban Area
- Sheffield Urban Area
- Nottingham Urban Area
- Bristol Urban Area
- Glasgow Urban Area
- South Wales.

The previous year (2020) saw a large reduction in the number of zones exceeding the annual mean limit value: just five zones exceeded in 2020 compared to 33 zones in 2019. This was attributed to the reduced road traffic flows brought about by the Covid-19 pandemic lockdown restrictions. The 2021 data suggest that NO₂ concentrations have increased in 2021 compared to 2020 but remain lower than their pre-pandemic levels.

All non-agglomeration zones within the UK complied with the critical level for annual mean NO_x concentration, set for protection of vegetation, as has been the case for many years.

As part of the 2017 UK plan for tackling roadside nitrogen dioxide concentrations (Defra, 2017b), local authorities with exceedances of the annual mean nitrogen dioxide limit value have been required to develop local plans or studies to consider measures to achieve the statutory limit value within the shortest possible time. These studies or plans may include local scale modelling and/or monitoring data, and in some cases the local data presents different results to the national air quality assessment. This is partly due to local monitoring being sited differently to national monitoring in order to target local pollution hotspots. Where possible, Defra are working to develop and improve the national NO₂ compliance assessment to better reflect local level NO₂ concentrations. This included establishing the UUNN in 2020, to provide more local NO₂ measurement data.

Table 4-2 Results of Air Quality Assessment for Nitrogen Dioxide in 2021

Zone	Zone code	NO ₂ LV for health (1hr mean)	NO ₂ LV for health (annual mean)	NO _x critical level for vegetation (ann. mean)
Greater London Urban Area	UK0001	OK	> LV	n/a
West Midlands Urban Area	UK0002	OK	> LV (s only)	n/a
Greater Manchester Urban Area	UK0003	OK	> LV (s only)	n/a
West Yorkshire Urban Area	UK0004	OK	> LV (s only)	n/a
Tyneside	UK0005	OK	OK	n/a
Liverpool Urban Area	UK0006	OK	> LV (s only)	n/a
Sheffield Urban Area	UK0007	OK	> LV (s only)	n/a
Nottingham Urban Area	UK0008	OK	> LV (s only)	n/a
Bristol Urban Area	UK0009	OK	> LV (s only)	n/a
Brighton/Worthing/Littlehampton	UK0010	OK	OK	n/a
Leicester Urban Area	UK0011	OK	OK	n/a
Portsmouth Urban Area	UK0012	OK	OK	n/a
Teesside Urban Area	UK0013	OK	OK	n/a
The Potteries	UK0014	OK	OK	n/a
Bournemouth Urban Area	UK0015	OK	OK	n/a
Reading/Wokingham Urban Area	UK0016	OK	OK	n/a
Coventry/Bedworth	UK0017	OK	OK	n/a
Kingston upon Hull	UK0018	OK	OK	n/a
Southampton Urban Area	UK0019	OK	OK	n/a
Birkenhead Urban Area	UK0020	OK	OK	n/a
Southend Urban Area	UK0021	OK	OK	n/a
Blackpool Urban Area	UK0022	OK	OK	n/a
Preston Urban Area	UK0023	OK	OK	n/a
Glasgow Urban Area	UK0024	OK	> LV	n/a
Edinburgh Urban Area	UK0025	OK (s only)	OK	n/a
Cardiff Urban Area	UK0026	OK	OK	n/a
Swansea Urban Area	UK0027	OK	OK	n/a
Belfast Urban Area	UK0028	OK	OK	n/a
Eastern	UK0029	OK	OK	OK
South West	UK0030	OK	OK	OK
South East	UK0031	OK	OK	OK
East Midlands	UK0032	OK	OK	OK
North West & Merseyside	UK0033	OK	OK	OK (s only)
Yorkshire & Humberside	UK0034	OK	OK	OK
West Midlands	UK0035	OK	OK	OK (s only)
North East	UK0036	OK	OK	OK (s only)
Central Scotland	UK0037	OK	OK	OK (s only)
North East Scotland	UK0038	OK	OK	OK (s only)
Highland	UK0039	OK	OK	OK (s only)
Scottish Borders	UK0040	OK	OK	OK (s only)
South Wales	UK0041	OK	> LV	OK
North Wales	UK0042	OK	OK	OK
Northern Ireland	UK0043	OK	OK	OK (s only)

LV = limit value, (s only) indicates the compliance or exceedance was determined by supplementary assessment only.

Particulate Matter as PM₁₀: all zones and agglomerations were compliant with the annual mean limit value of 40 µg m⁻³ for PM₁₀. All zones and agglomerations were also compliant with the daily mean limit value of 50 µg m⁻³, which must not be exceeded more than 35 times a year. The results of the air quality assessment for PM₁₀ for each zone, with respect to the daily mean and annual mean limit values, are summarised in **Table 4-3**.

Under the Air Quality Standards Regulations, the UK is required to identify any exceedances of PM₁₀ limit values which are due to natural sources (for example sea salt). Where this is the case, the exceedance does not count as non-compliance. Particulate matter from sea salt is modelled and has been used in the past to determine whether compliance with the limit values has been achieved after contribution from natural sources has been subtracted. However, in 2021 there were no modelled exceedances of either the 24-hr or annual mean limit values, so no subtraction of contribution from natural sources has been carried out (Pugsley, K. L. et al., 2022).

Table 4-3 Results of Air Quality Assessment for PM₁₀ in 2021

Zone	Zone code	PM ₁₀ LV (daily mean)	PM ₁₀ LV (annual mean)
Greater London Urban Area	UK0001	OK	OK
West Midlands Urban Area	UK0002	OK	OK
Greater Manchester Urban Area	UK0003	OK	OK
West Yorkshire Urban Area	UK0004	OK	OK
Tyneside	UK0005	OK	OK
Liverpool Urban Area	UK0006	OK	OK
Sheffield Urban Area	UK0007	OK	OK
Nottingham Urban Area	UK0008	OK	OK
Bristol Urban Area	UK0009	OK	OK
Brighton/Worthing/Littlehampton	UK0010	OK (s only)	OK (s only)
Leicester Urban Area	UK0011	OK	OK
Portsmouth Urban Area	UK0012	OK	OK
Teesside Urban Area	UK0013	OK	OK
The Potteries	UK0014	OK	OK
Bournemouth Urban Area	UK0015	OK (s only)	OK (s only)
Reading/Wokingham Urban Area	UK0016	OK	OK
Coventry/Bedworth	UK0017	OK	OK
Kingston upon Hull	UK0018	OK	OK
Southampton Urban Area	UK0019	OK	OK
Birkenhead Urban Area	UK0020	OK	OK
Southend Urban Area	UK0021	OK	OK
Blackpool Urban Area	UK0022	OK	OK
Preston Urban Area	UK0023	OK	OK
Glasgow Urban Area	UK0024	OK	OK
Edinburgh Urban Area	UK0025	OK	OK
Cardiff Urban Area	UK0026	OK	OK
Swansea Urban Area	UK0027	OK	OK
Belfast Metropolitan Urban Area	UK0028	OK	OK
Eastern	UK0029	OK	OK
South West	UK0030	OK	OK
South East	UK0031	OK	OK
East Midlands	UK0032	OK	OK
North West & Merseyside	UK0033	OK	OK
Yorkshire & Humberside	UK0034	OK	OK
West Midlands	UK0035	OK	OK
North East	UK0036	OK	OK
Central Scotland	UK0037	OK	OK
North East Scotland	UK0038	OK (s only)	OK (s only)
Highland	UK0039	OK	OK
Scottish Borders	UK0040	OK (s only)	OK (s only)
South Wales	UK0041	OK	OK
North Wales	UK0042	OK	OK
Northern Ireland	UK0043	OK	OK

Subtraction of natural source contribution was not carried out for any zones in 2021. LV = limit value, (s only) indicates that the compliance or exceedance was determined by supplementary assessment only.

Particulate Matter as PM_{2.5}: all zones met the Stage 1 limit value (25 µg m⁻³ to be achieved by 1st Jan 2015) which came into force on 1st January 2015, and the Stage 2 limit value (20 µg m⁻³ which was to be achieved by 1st Jan 2020). Both limit values apply to the annual mean, based on the calendar year.

The results of the air quality assessment for PM_{2.5} for each zone are summarised in **Table 4-4**. Subtraction of PM_{2.5} contributions due to natural sources was not necessary for any zone.

Under the Air Quality Standards Regulations, the UK was required to achieve a National Exposure Reduction Target (NERT) for PM_{2.5}, over the period 2010 to 2020. This is based on the Average Exposure Indicator (AEI) statistic. The AEI for the UK is calculated as follows:

- (i) Each year, the annual arithmetic mean PM_{2.5} concentration is calculated for the designated AEI subset of urban background sites².
- (ii) The mean of the most recent three calendar years' values is taken as the AEI.

The AEI for the reference year (2010) was 13 µg m⁻³; based on this, the Air Quality Standards Regulations set an exposure reduction target of 15%, which equated to reducing the AEI to 11 µg m⁻³ by 2020. (The detailed methodology and results of this calculation are presented in Defra's technical report on UK air quality assessment (Pugsley, K. L. et al., 2022).)

The UK achieved the NERT in 2016, well before the 2020 target year, but has continued to report the AEI annually. Most recent annual mean urban background PM_{2.5} concentrations were as follows:

- 2019: 10 µg m⁻³
- 2020: 8 µg m⁻³
- 2021: 8 µg m⁻³

The three-year running mean AEI for 2021 (calculated as the mean of the above annual values, to the nearest integer), is 9 µg m⁻³. Therefore, the UK remained compliant with the NERT in 2021.

² The sites used for calculation of the AEI are all the urban background PM_{2.5} monitoring sites that were in operation in the baseline year. Urban background sites that started monitoring PM_{2.5} later, or were not classified as urban background in the baseline year, are not included. (The exception is where the new site is the relocation of an existing AEI site that has been moved by a short distance, and to a similar environment). This means that the AEI is calculated on a largely consistent group of sites from year to year.

Table 4-4 Results of Air Quality Assessment for PM_{2.5} in 2021

Zone	Zone code	PM _{2.5} Stage 1 limit value (annual mean, for 1 st Jan 2015)	PM _{2.5} Stage 2 limit value (annual mean, for 1 st Jan 2020)
Greater London Urban Area	UK0001	OK	OK
West Midlands Urban Area	UK0002	OK	OK
Greater Manchester Urban Area	UK0003	OK	OK
West Yorkshire Urban Area	UK0004	OK	OK
Tyneside	UK0005	OK	OK
Liverpool Urban Area	UK0006	OK (s only)	OK (s only)
Sheffield Urban Area	UK0007	OK	OK
Nottingham Urban Area	UK0008	OK	OK
Bristol Urban Area	UK0009	OK	OK
Brighton/Worthing/Littlehampton	UK0010	OK	OK
Leicester Urban Area	UK0011	OK	OK
Portsmouth Urban Area	UK0012	OK (s only)	OK (s only)
Teesside Urban Area	UK0013	OK	OK
The Potteries	UK0014	OK	OK
Bournemouth Urban Area	UK0015	OK	OK
Reading/Wokingham Urban Area	UK0016	OK	OK
Coventry/Bedworth	UK0017	OK	OK
Kingston upon Hull	UK0018	OK	OK
Southampton Urban Area	UK0019	OK	OK
Birkenhead Urban Area	UK0020	OK	OK
Southend Urban Area	UK0021	OK	OK
Blackpool Urban Area	UK0022	OK	OK
Preston Urban Area	UK0023	OK	OK
Glasgow Urban Area	UK0024	OK	OK
Edinburgh Urban Area	UK0025	OK	OK
Cardiff Urban Area	UK0026	OK	OK
Swansea Urban Area	UK0027	OK	OK
Belfast Metropolitan Urban Area	UK0028	OK	OK
Eastern	UK0029	OK	OK
South West	UK0030	OK	OK
South East	UK0031	OK	OK
East Midlands	UK0032	OK	OK
North West & Merseyside	UK0033	OK	OK
Yorkshire & Humberside	UK0034	OK	OK
West Midlands	UK0035	OK	OK
North East	UK0036	OK	OK
Central Scotland	UK0037	OK	OK
North East Scotland	UK0038	OK (s only)	OK (s only)
Highland	UK0039	OK	OK
Scottish Borders	UK0040	OK (s only)	OK (s only)
South Wales	UK0041	OK	OK
North Wales	UK0042	OK	OK
Northern Ireland	UK0043	OK	OK

Subtraction of natural source contribution was not carried out for any zones in 2021.

LV = limit value, (s only) indicates the compliance or exceedance was determined by supplementary assessment only.

Ozone: all zones and agglomerations met the target values for health and for protection of vegetation. The results of the air quality assessment for ozone are summarised in **Table 4-5**.

For ozone (O₃), there is a target value based on the maximum daily 8-hour mean. All 43 zones and agglomerations were compliant with this target value. There is also a long-term objective for protection of human health, based on the maximum daily 8-hour mean. Only four of the 43 zones and agglomerations were compliant with the long-term objective (LTO) for health in 2021.

There is also a target value based on the AOT40 statistic. The AOT40 statistic (expressed in $\mu\text{g m}^{-3}\cdot\text{hours}$) is the sum of the difference between hourly concentrations greater than $80 \mu\text{g m}^{-3}$ (= 40 ppb) and $80 \mu\text{g m}^{-3}$ over a given period using only the hourly mean values measured between 08:00 and 20:00 Central European Time each day. All 43 zones and agglomerations met the target value based on the AOT40 statistic. There is also a long-term objective, for protection of vegetation, based on this statistic; one zone exceeded this long-term objective for vegetation in 2021.

The UK met all target values for O₃ in 2021 as it has done for many years. The number of zones exceeding the long-term objective for vegetation (one) was the lowest since 2017.

Ozone concentrations – and hence the number of zones exceeding the LTOs - fluctuate from year to year as ozone is a transboundary pollutant and its formation is influenced by meteorological factors. The year 2021 appears to have been a year of relatively low ozone concentrations, with relatively few exceedances of the LTO for vegetation, and no substantial ozone pollution episodes.

Table 4-5 Results of Air Quality Assessment for Ozone in 2021

Zone	Zone code	O ₃ TV and LTO for health (8hr mean)	O ₃ TV and LTO for vegetation (AOT40)
Greater London Urban Area	UK0001	Met TV, > LTO	OK
West Midlands Urban Area	UK0002	Met TV, > LTO	OK
Greater Manchester Urban Area	UK0003	Met TV, > LTO (s only)	OK
West Yorkshire Urban Area	UK0004	Met TV, > LTO	OK
Tyneside	UK0005	Met TV, > LTO (s only)	OK
Liverpool Urban Area	UK0006	Met TV, > LTO (s only)	OK (s only)
Sheffield Urban Area	UK0007	Met TV, > LTO (s only)	OK
Nottingham Urban Area	UK0008	Met TV, > LTO (s only)	OK
Bristol Urban Area	UK0009	Met TV, > LTO	OK
Brighton/Worthing/Littlehampton	UK0010	Met TV, > LTO	OK
Leicester Urban Area	UK0011	Met TV, > LTO (s only)	OK
Portsmouth Urban Area	UK0012	Met TV, > LTO (s only)	OK
Teesside Urban Area	UK0013	Met TV, > LTO (s only)	OK
The Potteries	UK0014	Met TV, > LTO (s only)	OK
Bournemouth Urban Area	UK0015	Met TV, > LTO	OK
Reading/Wokingham Urban Area	UK0016	Met TV, > LTO (s only)	OK
Coventry/Bedworth	UK0017	Met TV, > LTO	OK
Kingston upon Hull	UK0018	Met TV, > LTO	OK
Southampton Urban Area	UK0019	Met TV, > LTO (s only)	OK
Birkenhead Urban Area	UK0020	Met TV, > LTO	OK
Southend Urban Area	UK0021	Met TV, > LTO	OK
Blackpool Urban Area	UK0022	Met TV, > LTO	OK
Preston Urban Area	UK0023	Met TV, > LTO	OK
Glasgow Urban Area	UK0024	OK	OK
Edinburgh Urban Area	UK0025	OK	OK
Cardiff Urban Area	UK0026	Met TV, > LTO	OK
Swansea Urban Area	UK0027	Met TV, > LTO (s only)	OK
Belfast Metropolitan Urban Area	UK0028	Met TV, > LTO (s only)	OK
Eastern	UK0029	Met TV, > LTO	OK
South West	UK0030	Met TV, > LTO	Met TV, > LTO
South East	UK0031	Met TV, > LTO	OK
East Midlands	UK0032	Met TV, > LTO	OK
North West & Merseyside	UK0033	Met TV, > LTO	OK
Yorkshire & Humberside	UK0034	Met TV, > LTO	OK
West Midlands	UK0035	Met TV, > LTO	OK
North East	UK0036	Met TV, > LTO (s only)	OK
Central Scotland	UK0037	OK	OK
North East Scotland	UK0038	OK (s only)	OK
Highland	UK0039	Met TV, > LTO	OK
Scottish Borders	UK0040	Met TV, > LTO (s only)	OK
South Wales	UK0041	Met TV, > LTO	OK
North Wales	UK0042	Met TV, > LTO	OK
Northern Ireland	UK0043	Met TV, > LTO (s only)	OK

TV = target value, LTO = long-term objective, (s only) indicates that the compliance or exceedance was determined by supplementary assessment only.

In 2021 there was one measured exceedance of the ozone population information threshold of 180 $\mu\text{g m}^{-3}$ (at one site, Rochester Stoke, a rural site in Kent), but no exceedances of the population warning threshold of 240 $\mu\text{g m}^{-3}$. The population information threshold exceedance is detailed in **Table 4-6**. This occurred at 15:00 on 30th March 2021.

Several other AURN sites in the south and east of England also detected an isolated 'spike' in ozone at the same time on the same date, though only at Rochester Stoke was the measured concentration above the population information threshold.

Table 4-6 Measured Exceedances of the Ozone Information Threshold Value in 2021

Site name	Zone code	Number of 1-hour exceedances of information threshold	Maximum 1-hour concentration ($\mu\text{g m}^{-3}$)
Rochester Stoke	UK0031	1	216

Table 4-6 shows the exceedances of the ozone information threshold in the verified dataset.

(Exceedances of the population information threshold were also measured at the Lerwick monitoring site in Shetland, during the period 4th – 6th June: however, these were later confirmed not to be genuine, but due to an instrument malfunction. The spurious data were rejected at the data verification stage.)

The air quality assessments for arsenic (As), cadmium (Cd), nickel (Ni) and benzo[a]pyrene (B[a]P) are summarised in **Table 4-7**. All zones met target values for arsenic and cadmium, but some zones exceeded the target value for nickel, or nickel and benzo[a]pyrene.

Concentrations of Ni exceeded the target value in four zones: Sheffield Urban Area, Yorkshire and Humberside, Swansea Urban Area and South Wales. These exceedances are attributed to emissions from industrial sources.

Concentrations of benzo[a]pyrene were above the target value in two of the same zones; Swansea Urban Area and South Wales. These exceedances are also attributed to emissions from industrial sources.

Table 4-7 Results of Air Quality Assessment for As, Cd, Ni and B[a]P in 2021

Zone	Zone code	As TV	Cd TV	Ni TV	B[a]P TV
Greater London Urban Area	UK0001	OK	OK	OK	OK
West Midlands Urban Area	UK0002	OK	OK	OK	OK
Greater Manchester Urban Area	UK0003	OK (s only)	OK (s only)	OK (s only)	OK
West Yorkshire Urban Area	UK0004	OK (s only)	OK (s only)	OK (s only)	OK
Tyneside	UK0005	OK (s only)	OK (s only)	OK (s only)	OK
Liverpool Urban Area	UK0006	OK (s only)	OK (s only)	OK (s only)	OK (s only)
Sheffield Urban Area	UK0007	OK	OK	> TV (s only)	OK
Nottingham Urban Area	UK0008	OK (s only)	OK (s only)	OK (s only)	OK
Bristol Urban Area	UK0009	OK (s only)	OK (s only)	OK (s only)	OK
Brighton/Worthing/Littlehampton	UK0010	OK (s only)	OK (s only)	OK (s only)	OK (s only)
Leicester Urban Area	UK0011	OK (s only)	OK (s only)	OK (s only)	OK (s only)
Portsmouth Urban Area	UK0012	OK (s only)	OK (s only)	OK (s only)	OK (s only)
Teesside Urban Area	UK0013	OK (s only)	OK (s only)	OK (s only)	OK
The Potteries	UK0014	OK (s only)	OK (s only)	OK (s only)	OK (s only)
Bournemouth Urban Area	UK0015	OK (s only)	OK (s only)	OK (s only)	OK (s only)
Reading/Wokingham Urban Area	UK0016	OK (s only)	OK (s only)	OK (s only)	OK (s only)
Coventry/Bedworth	UK0017	OK (s only)	OK (s only)	OK (s only)	OK (s only)
Kingston upon Hull	UK0018	OK (s only)	OK (s only)	OK (s only)	OK (s only)
Southampton Urban Area	UK0019	OK (s only)	OK (s only)	OK (s only)	OK (s only)
Birkenhead Urban Area	UK0020	OK (s only)	OK (s only)	OK (s only)	OK (s only)
Southend Urban Area	UK0021	OK (s only)	OK (s only)	OK (s only)	OK (s only)
Blackpool Urban Area	UK0022	OK (s only)	OK (s only)	OK (s only)	OK (s only)
Preston Urban Area	UK0023	OK (s only)	OK (s only)	OK (s only)	OK (s only)
Glasgow Urban Area	UK0024	OK (s only)	OK (s only)	OK (s only)	OK
Edinburgh Urban Area	UK0025	OK (s only)	OK (s only)	OK (s only)	OK
Cardiff Urban Area	UK0026	OK (s only)	OK (s only)	OK (s only)	OK
Swansea Urban Area	UK0027	OK	OK	> TV	> TV (s only)
Belfast Urban Area	UK0028	OK	OK	OK	OK
Eastern	UK0029	OK	OK	OK	OK
South West	UK0030	OK	OK	OK	OK
South East	UK0031	OK	OK	OK	OK
East Midlands	UK0032	OK	OK	OK	OK
North West & Merseyside	UK0033	OK (s only)	OK (s only)	OK (s only)	OK
Yorkshire & Humberside	UK0034	OK	OK	> TV (s only)	OK
West Midlands	UK0035	OK	OK	OK	OK (s only)
North East	UK0036	OK (s only)	OK (s only)	OK (s only)	OK
Central Scotland	UK0037	OK	OK	OK	OK
North East Scotland	UK0038	OK (s only)	OK (s only)	OK (s only)	OK (s only)
Highland	UK0039	OK (s only)	OK (s only)	OK (s only)	OK
Scottish Borders	UK0040	OK	OK	OK	OK (s only)
South Wales	UK0041	OK	OK	> TV (s only)	> TV (s only)
North Wales	UK0042	OK (s only)	OK (s only)	OK (s only)	OK (s only)
Northern Ireland	UK0043	OK (s only)	OK (s only)	OK (s only)	OK

TV = target value, (s only) indicates the compliance or exceedance was determined by supplementary assessment only.

4.3 Comparison with Previous Years

This section provides information on non-compliances in previous years from 2008 onwards. (2008 is the year that the Air Quality Directive - which was subsequently transposed into UK legislation by the Air Quality Standards Regulations – came into force.)

For **SO₂**, **PM_{2.5}**, **lead**, **benzene** and **CO**, the UK has been compliant with Air Quality Standards Regulations limit values (apart from the PM_{2.5} Stage 2 indicative limit value) in all years since 2008. For information on compliance with the 1st and 2nd Daughter Directives for all pollutants in earlier years, please see the 2012 or earlier reports in this series, which can be found here: <https://uk-air.defra.gov.uk/library/annualreport/>.

The UK has been compliant with the limit values for both **lead** and **CO** since 2003, and for **benzene** since 2007: these limit values are the same as those contained in the 1st and 2nd Daughter Directives, which the Air Quality Directive (and therefore the Air Quality Standards Regulations) superseded.

For nitrogen dioxide, **Table 4-8** summarises the results of the air quality assessment in years from 2008 to 2021. This table shows the numbers of zones exceeding the limit value (plus any agreed margin of tolerance, in cases where a time extension had been granted). The right-hand column contains notes on the effects of any time extensions, the last of which ended on 1st January 2015.

All non-agglomeration zones within the UK have complied with the critical level for annual mean NO_x concentration, set for protection of vegetation, in years 2008 onwards.

For PM₁₀, **Table 4-9** summarises the results of the air quality assessment in years from 2008 to 2021. There are notes in the right-hand column explaining the effects of the time extensions which were in place up to the end of 2011 for some zones.

For ozone, **Table 4-10** summarises annual exceedances of the target value for human health (based on the maximum daily 8-hour mean), the target value for protection of vegetation (based on the AOT40 statistic), and the two long-term objectives (LTOs) based on these two metrics.

Finally, for the pollutants formerly covered by the Fourth Daughter Directive - arsenic (As), cadmium (Cd), nickel (Ni) and benzo[a]pyrene (B[a]P) - **Table 4-11** summarises the numbers of zones with exceedances of target values in previous years.

Table 4-8 Non-Compliances with Limit Values for Nitrogen Dioxide, 2008-2021

Year	Zones Exceeding NO ₂ LV for health (1hr mean)	Zones Exceeding NO ₂ LV for health (annual mean)	Notes on Time Extensions
2008	3 zones (London, Glasgow, N.E. Scotland)	40 zones	-
2009	2 zones (London, Glasgow)	40 zones	-
2010	3 zones (London, Teesside, Glasgow)	40 zones	-
2011	3 zones (London, Glasgow, South East)	35 zones	<i>A further 5 zones exceeded the annual mean NO₂ LV in 2011 but were covered by time extensions and within the LV+ Margin of Tolerance (MOT), therefore compliant.</i>
2012	2 zones (London, South East)	34 zones	<i>A further 4 zones exceeded the annual mean NO₂ LV in 2012 but were covered by time extensions and within the LV+ MOT, therefore compliant.</i>
2013	1 zone (London)	31 zones	<i>A further 7 zones exceeded the annual mean NO₂ LV in 2013 but were covered by time extensions and within the LV+ MOT, therefore compliant.</i>
2014	2 zones (London, South Wales)	30 zones	<i>A further 8 zones exceeded the annual mean NO₂ LV in 2014 but were covered by time extensions and within the LV+ MOT, therefore compliant.</i>
2015	2 zones (London, South Wales)	37 zones	<i>2015 was the first year with no time extensions for NO₂: this is the reason for the apparent increase in zones exceeding between 2014 and 2015.</i>
2016	2 zones (London, South Wales)	37 zones	<i>No time extensions in place.</i>
2017	2 zones (London, South Wales)	37 zones	<i>No time extensions in place.</i>
2018	2 zones (London, South Wales)	36 zones	<i>No time extensions in place.</i>
2019	1 zone (South Wales)	33 zones	<i>No time extensions in place.</i>
2020	None	5 zones	<i>No time extensions in place.</i>
2021	None	10 zones	<i>No time extensions in place.</i>

Table 4-9 Non-Compliances with the Limit Values for PM₁₀, 2008-2021

Year	PM₁₀ LV (annual mean)	PM₁₀ LV (daily mean)	Notes on Time Extensions and Subtraction of Natural contribution
2008	None	2 zones (1 zone after subtraction of natural contribution)	-
2009	None	3 zones (1 zone after subtraction of natural contribution)	-
2010	None	None (after subtraction of natural contribution)	One zone exceeded the daily mean PM10 limit value more than the permitted 35 times in 2010, after subtraction of natural contribution. This zone was covered by a time extension and was within the LV+MOT so was therefore compliant.
2011	None	None (after subtraction of natural contribution)	One zone exceeded the daily mean PM10 limit value more than the permitted 35 times in 2011, after subtraction of natural contribution. This zone was covered by a time extension and was within the LV+MOT so was therefore compliant.
2012	None	None (after subtraction of natural contribution. No time extension.)	-
2013	None	None (after subtraction of natural contribution. No time extension.)	-
2014	None	None (after subtraction of natural contribution. No time extension.)	-
2015	None	None (after subtraction of natural contribution. No time extension.)	-
2016	None	None	-
2017	None	None	-
2018	None	None	-
2019	None	None	-
2020	None	None	-
2021	None	None	-

Table 4-10 Exceedances of Target Values for Ozone (Health) and Long-Term Objectives, 2008-2021

Year	8-Hour Mean Target Value	AOT40 Target Value	8-Hour Mean LTO	AOT40 LTO
2008	1 zone measured (Eastern)	None	43 zones	41 zones
2009	None	None	39 zones	10 zones
2010	None	None	41 zones	6 zones
2011	None	None	43 zones	3 zones
2012	None	None	41 zones	3 zones
2013	None	None	33 zones	8 zones
2014	None	None	32 zones	3 zones
2015	None	None	43 zones	1 zone
2016	None	None	42 zones	5 zones
2017	None	None	34 zones	None
2018	None	None	43 zones	38 zones
2019	None	None	43 zones	6 zones
2020	None	None	40 zones	16 zones
2021	None	None	39 zones	1 zone

Table 4-11 Zones Exceeding Target Values for As, Cd, Ni and B[a]P, 2008-2021

Year	As	Cd	Ni	B[a]P
2008	None	None	2 (Swansea, South Wales)	6 (Yorks. & Humberside, Teesside, Northern Ireland, Swansea, South Wales, Belfast)
2009	None	None	2 (Swansea, South Wales)	6 (Yorks. & Humberside, Northern Ireland, Teesside, Swansea, North East, South Wales)
2010	None	None	2 (Swansea, South Wales)	8 (Yorks. & Humberside, Northern Ireland, Teesside, Belfast, W Midlands, North East, South Wales, North Wales.)
2011	None	None	2 (Swansea, South Wales)	7 (Yorks. & Humberside, N. Ireland, Teesside, Swansea, Belfast, North East, South Wales)
2012	None	None	2 (Swansea, South Wales)	8 (Yorks. & Humberside, Teesside, Swansea, Belfast, the North East, South Wales, North Wales, Northern Ireland.)
2013	None	None	2 (Swansea, South Wales)	6 (Yorks. & Humberside, Teesside, Swansea, East Midlands, North East, South Wales.)
2014	None	None	3 (Sheffield, Swansea, South Wales)	6 (Yorks. & Humberside, Teesside, Swansea, East Midlands, North East, and South Wales).
2015	None	None	2 (Swansea, South Wales)	5 (Yorks. & Humberside, Teesside, Swansea, the North East and South Wales).
2016	None	None	3 (Sheffield, Swansea, South Wales)	4 (Yorks. & Humberside, Swansea, South Wales and Northern Ireland).
2017	None	None	None	3 (Yorks. & Humberside, Swansea and South Wales)
2018	None	None	4 (Sheffield, Yorks. & Humberside, Swansea and South Wales)	3 (Yorks. & Humberside, Swansea and South Wales)
2019	None	None	4 (Sheffield, Yorks. & Humberside, Swansea and South Wales)	3 (Yorks. & Humberside, Swansea and South Wales)
2020	None	None	4 (Sheffield, Yorks. & Humberside, Swansea and South Wales)	3 (Yorks. & Humberside, Swansea and South Wales)
2021	None	None	4 (Sheffield, Yorks. & Humberside, Swansea and South Wales)	2 (Swansea and South Wales)

5 Spatial Variation and Changes Over Time

This section looks at the spatial distribution of pollutants across the UK, based upon the modelled maps of ambient pollutant concentration discussed in **Section 3.5** of this report, “Modelling”. For some traffic-related pollutants such as NO₂, roadside and background concentrations are discussed separately.

For each pollutant, this section also discusses how ambient concentrations have changed over time, using data from the relevant ambient air quality monitoring networks: the Automatic Urban and Rural Network (AURN), the Automatic Hydrocarbon Network, the Non-Automatic Hydrocarbon Network, the Heavy Metals Network, and the PAH Network.

The AURN has been in operation since 1992. Since that time, it has grown considerably in size, and the proportion of urban traffic (‘roadside’) monitoring sites has increased. Therefore, in most cases, we have based our investigation of trends on data from sub-sets of long-running AURN monitoring sites, rather than the whole network. This is intended to show changes over time, without introducing any spurious effects due to changes in the number and distribution of the sites.

Trend analysis has been carried out using Openair (Carslaw & Ropkins, 2012). Openair provides free, open-source and innovative tools to analyse interpret and understand air pollution data using R; a free and open-source programming language designed for the analysis of data. The Openair package was primarily developed for the analysis of air pollution datasets, and can handle high volumes of data; the AURN, with its long data record, lends itself to this. A further strength of the Openair tools is that they also allow data to be conditioned by one or more variables. For example, plots can be produced that show the inter-relationships between air pollutants and meteorological parameters, or temporal trends. The UK-AIR website provides simplified web access to a customised selection of the Openair tools, including tools that require meteorological measurements.

Openair was developed by King’s College London with the University of Leeds. The Openair project is currently led by Dr David Carslaw, of Ricardo Energy & Environment and the University of York. A range of Openair functions are available on UK-AIR: for more information on the functions and how to use them, please refer to: <https://uk-air.defra.gov.uk/data/openair>.

The Openair ‘TheilSen’ function has been used here for NO₂, PM₁₀, PM_{2.5}, O₃, SO₂ and CO. This uses the Theil-Sen statistical method to quantify trends in concentrations over time. The trend analysis is based on monthly mean pollutant concentrations (at least 75% data capture is required for a valid monthly mean). The ‘TheilSen’ function includes an option to ‘de-seasonalise’ the data (i.e. statistically modify the plotted data to remove the influence of seasonal cycles, thus providing a clearer indication of the overall trend). The ‘de-seasonalise’ option has been used for the AURN pollutants, as indicated in the graph titles. The trend line is shown by a solid red line, with 95% confidence intervals for the trend shown by dotted red lines. The trend is given at the top of the graph, with confidence

intervals shown in square brackets. The trend is given as units (e.g. $\mu\text{g m}^{-3}$) per year, over the period shown. This may be followed by a number of symbols, with + indicating that the trend is statistically significant at the 0.1 level, * indicating that the trend is statistically significant at the 0.05 level, ** indicating significance at the 0.01 level and *** indicating significance at the 0.001 level. It should also be noted that the 'de-seasonalise' option fills in any gaps in the dataset using an interpolation method, so the datasets shown in these trend plots appear uninterrupted, though this is not necessarily the case.

For pollutants measured by the Hydrocarbons, PAH and Heavy Metals networks, time series or smoothed trend plots (not de-seasonalised) have been used to illustrate changes over time.

These changes over time are compared to changes in estimated total UK emissions where appropriate. Estimated UK emissions data are taken from the National Atmospheric Emissions Inventory (NAEI) website at <https://naei.beis.gov.uk/>. (The most recent year for which NAEI emission estimates are available is 2020). The NAEI dataset shows emissions split between various emission source categories, which are different for different pollutants.

Please note that this section only aims to provide a general indication of changes in pollutant concentration over time, based in most cases on averages or groups of long-running sites. Patterns for specific regions or individual sites may be different.

5.1 Nitrogen Dioxide


5.1.1 NO₂: Spatial Distribution in the UK

Figure 5-1 shows the modelled annual mean NO₂ concentrations for 2021, at *urban roadside* locations only. Although not every road link is clearly visible, some road links in urban areas are shaded in the lighter colours, indicating higher concentrations. Some road links are shaded lighter green, indicating that they have modelled annual mean concentrations above the annual mean limit value of $40 \mu\text{g m}^{-3}$. (No road links are shaded yellow, which would mean annual mean concentrations above $60 \mu\text{g m}^{-3}$). Please note that the bands used in the NO₂ maps were updated for the 2020 report, and are therefore different from those used in the reports for 2019 and earlier.

Figure 5-2 shows the modelled annual mean *background* NO₂ concentrations for 2021. All background locations were within the limit value of $40 \mu\text{g m}^{-3}$.


These maps for NO₂, and those for other pollutants in subsequent sections, are intended to provide an illustration of spatial variation in pollutant concentrations across the UK. To see detail for specific areas or individual road links, please use the UK Ambient Air Quality Interactive Map provided by UK-AIR at <https://uk-air.defra.gov.uk/data/gis-mapping/>. (Please note, the online interactive map uses a different colour scheme.)

Figure 5-1 Annual Mean Roadside NO₂ Concentration by Major Urban Roads, 2021 ($\mu\text{g m}^{-3}$)



(An interactive version of this map is available at <https://uk-air.defra.gov.uk/data/gis-mapping/>).

Figure 5-2 Annual Mean Background NO₂ Concentration, 2021 ($\mu\text{g m}^{-3}$)




(An interactive version of this map is available at <https://uk-air.defra.gov.uk/data/gis-mapping/>).

5.1.2 NO₂: Changes Over Time

Figure 5-3 and **Figure 5-4** show how ambient concentrations of NO₂ (as measured by the AURN) have decreased since 1992 (the Network's first year of operation). Time series charts of NO₂ concentration are shown for the following sub-sets of long-running sites:

- Eight urban non-roadside sites operating since 1995 or earlier (**Figure 5-3**); Belfast Centre, Cardiff Centre, Leeds Centre, London Bloomsbury, Middlesbrough, Newcastle Centre, Sheffield Tinsley and Southampton Centre. (These are all urban background, except Middlesbrough which is urban industrial).
- Eight urban traffic sites operating since 1998 or earlier (**Figure 5-4**); Camden Kerbside, Cambridge Roadside, Exeter Roadside, Glasgow Kerbside, Haringey Roadside, London Marylebone Road, Oxford Centre Roadside and Tower Hamlets Roadside.

Figure 5-3 De-seasonalised Trends in NO₂ Concentration, at 8 Long-Running AURN Urban Non-Roadside* Sites, 1992-2021



*Urban Background, except for Middlesbrough which is urban industrial.

All eight long-running urban non-roadside sites in **Figure 5-3** show a decreasing trend in NO₂. While the magnitude of the year-on-year decrease varies (ranging from -1.21 µg m⁻³ to -0.48 µg m⁻³ per year), the trend is statistically highly significant at the 0.001 level for all eight sites, as indicated by the three asterisks (***) on the plots. For the urban traffic sites in **Figure 5-4** below, (for which the dataset is slightly shorter), the pattern of trends has historically been less consistent, as highlighted in previous reports in this series. However,

all eight sites now show a downward trend statistically significant at the 0.001 level. Several of the sites show a dip in NO₂ concentration in 2020, which is likely to be due at least in part to the Covid-19 restrictions.

Figure 5-4 De-seasonalised Trends in NO₂ Concentration at 8 Long-Running AURN Urban Traffic Sites, 1998-2021




Figure 5-5 is taken from Defra’s National Air Quality Statistics web page for NO₂, at <https://www.gov.uk/government/statistics/air-quality-statistics/ntrogen-dioxide> (Defra, 2022c). This shows annual mean NO₂ concentrations averaged over all included sites that had annual data capture greater than or equal to 75% in the relevant year. Roadside (urban traffic), urban background and rural sites are shown by separate, labelled, lines.

As explained at the start of this section, the number of sites in the network has increased substantially over the years. This introduces uncertainty when considering trends for the whole network. Therefore, this graph shows the 95% confidence interval of the annual mean for each site classification, as a shaded area either side of each line. The confidence intervals narrow over time because of an increase in the number of monitoring sites and a reduction in the variation between annual means at monitoring sites for NO₂.

For both urban traffic and urban background sites, there appears to be a sharp decrease in 2020, which is likely to be due at least in part to the reduction in traffic emissions caused by the Covid-19 pandemic restrictions in that year. The ongoing effects of these restrictions are covered in more detail in Section 6.

Figure 5-5 Annual mean concentrations of NO₂ in the UK, by AURN Site Classification, 1990 to 2021. Shaded areas either side of each line show the 95% confidence interval of the mean. (Source: <https://www.gov.uk/government/statistics/air-quality-statistics/nitrogen-dioxide>)





Figure 5-6 shows estimates of total UK annual emission of oxides of nitrogen, in kilotonnes, from 1990 to 2020 (the most recent year for which emission estimates are available). The data are from the National Atmospheric Emissions Inventory (NAEI) website at <https://naei.beis.gov.uk/>. This shows that total NO_x emissions have decreased substantially over this period and are now less than one third of the total emissions in 1990. Emissions from several specific sources, notably public energy and heat production, passenger cars and heavy-duty vehicles, have also shown substantial decreases over the same period.

Figure 5-6 Estimated Annual UK Emissions of Nitrogen Oxides (kt), 1990 – 2020 (Source: NAEI 2022)




5.2 PM₁₀ Particulate Matter

5.2.1 PM₁₀: Spatial Distribution

Figure 5-7 shows modelled annual mean urban roadside PM₁₀ concentrations in 2021. No roadside locations had a modelled annual mean concentration greater than the limit value of 40 µg m⁻³.


Figure 5-8 shows modelled annual mean background PM₁₀ concentrations in 2021. No urban background locations had a modelled annual mean concentration greater than the limit value of 40 µg m⁻³.

Figure 5-7 Annual Mean Roadside PM₁₀ Concentration by Urban Major Roads, 2021 ($\mu\text{g m}^{-3}$)



(An interactive version of this map is available at <https://uk-air.defra.gov.uk/data/gis-mapping/>).


Figure 5-8 Annual Mean Background PM₁₀ Concentration, 2021 ($\mu\text{g m}^{-3}$)



(An interactive version of this map is available at <https://uk-air.defra.gov.uk/data/gis-mapping/>).

Where the annual mean PM_{10} concentration exceeds $31.5 \mu g m^{-3}$, it is likely also that the 24-hour mean has exceeded the daily mean limit value of $50 \mu g m^{-3}$ on more than the permitted 35 occasions. **Figure 5-9** shows roads where the modelled annual mean roadside PM_{10} concentration is less than or equal to this threshold coloured dark blue, and roads where the annual mean exceeds this threshold coloured yellow. All road links are dark blue, indicating that no roads had the modelled annual mean roadside PM_{10} concentration greater than $31.5 \mu g m^{-3}$ in 2021.

Figure 5-9 Compliance with the 24-hour limit value for PM_{10} for major urban roads, 2021 (estimated to be compliant below an annual mean concentration of $31.5 \mu g m^{-3}$)



Similarly, **Figure 5-10** shows areas where the modelled annual mean background PM₁₀ concentration is less than or equal to 31.5 µg m⁻³ in dark blue, and areas where the annual mean exceeds this threshold in yellow. (Note: the value of 31.5 µg m⁻³ is specific to 2021: this threshold is calculated each year on the basis of the measured data. It may therefore change from year to year.) There are no areas coloured yellow, showing that modelled annual mean background PM₁₀ concentration did not exceed 31.5 µg m⁻³ anywhere in the UK, in 2021.


Figure 5-10 Compliance with the 24-hour limit value for PM₁₀ at background locations, 2021 (estimated to be compliant below an annual mean concentration of 31.5 µg m⁻³)



5.2.2 PM₁₀ Changes Over Time

Figure 5-11 shows de-seasonalised trends in ambient PM₁₀ concentration, based on 12 urban non-roadside AURN sites, all of which have been operating since 1997 or earlier. The sites are; Belfast Centre, Cardiff Centre, Leamington Spa, Leeds Centre, London Bloomsbury, London North Kensington, Middlesbrough, Newcastle Centre, Nottingham Centre, Salford Eccles, Southampton Centre and Thurrock. (Middlesbrough is urban industrial, the others are urban background). All 12 sites show a downward trend for PM₁₀ over their period of operation, highly statistically significant (at the 0.001 confidence level).

Figure 5-11 De-seasonalised Trends in Ambient PM₁₀, 12 Long-Running Urban Non-Roadside* AURN Sites 1992 – 2021



*Middlesbrough is urban industrial, the others are urban background.

Figure 5-12 shows de-seasonalised trends in ambient PM₁₀ concentration, based on 12 urban traffic AURN sites. There are few very long-running urban traffic PM₁₀ sites: only three began monitoring PM₁₀ before 2008. The sites shown here have been operating since the start of 2009 or earlier. The sites are; Armagh Roadside, Camden Kerbside, Carlisle Roadside, Chepstow A48, Chesterfield Roadside, Leeds Headingley Kerbside,

London Marylebone Road, Sandy Roadside, Stanford-le-Hope Roadside, Stockton-on-Tees Eaglescliffe, Swansea Roadside and York Fishergate. (Note: Carlisle Roadside has been excluded from the UK's compliance reporting from 2019 onwards, as it was within 25m of a major junction. The site was relocated during 2021, but it is included here for consistency with previous years' reports.)

Ten of these sites show a downward trend which is highly statistically significant (at the 0.001 confidence level). The exceptions are York Fishergate, which shows a downward trend significant at the 0.01 confidence level, and Swansea Roadside which shows a statistically significant increase. As in the case of NO₂, it is likely that trends at urban traffic sites are influenced by changes in the volume and type of traffic on the adjacent road.

Figure 5-12 De-seasonalised Trends in Ambient PM₁₀, 12 Long-Running Urban Traffic AURN Sites 2009 – 2021




Figure 5-13 is taken from Defra's National Air Quality Statistics web page for PM₁₀ and PM_{2.5}, at <https://www.gov.uk/government/statistics/air-quality-statistics/concentrations-of-particulate-matter-pm10-and-pm25>, (Defra, 2022d). This shows annual mean PM₁₀ concentrations averaged over all included AURN sites that had annual data capture

greater than or equal to 75% in the relevant year. Roadside (urban traffic) and urban background sites are shown by separate, labelled, lines.

Shaded areas surrounding the lines show the 95% confidence interval of the annual mean for each site classification. The confidence intervals narrow over time, as the number of monitoring sites has increased: this is particularly the case for urban traffic (roadside) PM₁₀ monitoring sites, which have almost doubled in number since 2008.

Figure 5-13 Annual mean concentrations of PM₁₀ in the UK, by AURN Site Classification, 1992 to 2021. Shaded areas either side of each line show the 95% confidence interval of the mean. (Source: <https://www.gov.uk/government/statistics/air-quality-statistics/concentrations-of-particulate-matter-pm10-and-pm25>)





Figure 5-14 shows NAEI estimates of total UK annual emission of PM₁₀ in kilotonnes, from 1990 to 2020 (the most recent year for which emission estimates are available). Total PM₁₀ emissions have steadily decreased over all this period, although in more recent years the rate of decrease has slowed, flattening off after around 2010 – but with a further decrease in 2020. Emissions from the ‘*combustion in industry, commercial and residential settings*’ sector appears to have increased slightly over the past decade. The NAEI says “*Emissions from residential sector combustion have grown both in real terms and in terms of the contribution to the UK total. This is because of strong growth in the use of wood as a domestic fuel, which has offset reductions that have occurred due to decreasing use of coal and other solid mineral fuels.*”(https://naei.beis.gov.uk/overview/pollutants?pollutant_id=24). By contrast,

estimated emissions from road traffic alone have continued to decrease steadily. 2020 saw a sharp decrease in estimated emissions from road traffic and other traffic, as a result of the Covid-19 pandemic restrictions.

Figure 5-14 Estimated Annual UK Emissions of PM₁₀ (kt), 1990 – 2020 (source: NAEI 2022)




5.3 PM_{2.5} Particulate Matter

5.3.1 PM_{2.5}: Spatial Distribution

Figure 5-15 shows the modelled annual mean urban roadside PM_{2.5} concentrations in 2021. No roadside locations had annual means greater than the Stage 2 limit value of 20 µg m⁻³; this is consistent with the compliance assessment reported in **Section 4**.


Figure 5-16 shows modelled annual mean background PM_{2.5} concentrations for 2021. These were highest in the south and east of the UK, and lower in the north and west, which is a typical pattern. Modelled concentrations ranged from 2 µg m⁻³ or less in northwest Scotland to 9-10 µg m⁻³ over most of southern England. Higher values occurred in cities such as London and Birmingham, but everywhere in the UK was well within the Stage 2 limit value of 20 µg m⁻³, as reported in **Section 4**.

Figure 5-15 Annual Mean Roadside PM_{2.5} Concentration, Urban Major Roads, 2021 ($\mu\text{g m}^{-3}$)



(An interactive version of this map is available at <https://uk-air.defra.gov.uk/data/gis-mapping/>).

Figure 5-16 Annual Mean Background PM_{2.5} Concentration, 2021 ($\mu\text{g m}^{-3}$)



(An interactive version of this map is available at <https://uk-air.defra.gov.uk/data/gis-mapping/>).

5.3.2 PM_{2.5}: Changes Over Time

Until 2008, routine monitoring of PM_{2.5} within the AURN was confined to a small number of sites in London. Therefore, in this report, trend analysis for PM_{2.5} concentrates on years 2009 onwards, during which PM_{2.5} monitoring has been widespread. **Figure 5-17** shows trends in PM_{2.5} concentration at 12 long-running urban background AURN sites, 2009-2021. All 12 sites show a statistically significant downward trend in PM_{2.5} concentration, significant at the 0.001 confidence level.

Figure 5-17 De-seasonalised Trends in Ambient PM_{2.5} Concentration, 12 Long-Running Urban Background AURN Sites 2009-2021




Figure 5-18 shows trends over the same period for PM_{2.5} at 12 urban traffic AURN sites. (As for PM₁₀, Carlisle Roadside has been included for consistency with previous years' trend investigations, despite being excluded from the compliance assessment because it is located on a major junction. It has since been relocated elsewhere in Carlisle.) The majority of the sites (11 out of 12) show decreasing trends, statistically significant at the 0.001 confidence level. This includes Camden Kerbside, which until 2017 showed no significant trend. The exception is Swansea Roadside, which shows an *increasing* trend in

PM_{2.5} concentration, as it also does for PM₁₀, though for PM_{2.5} this is not statistically significant.

Figure 5-18 De-seasonalised Trends in Ambient PM_{2.5} Concentration, 12 Long-Running Urban Traffic AURN Sites 2009-2021





Figure 5-19 is taken from Defra’s National Air Quality Statistics web page for PM₁₀ and PM_{2.5}, at <https://www.gov.uk/government/statistics/air-quality-statistics/concentrations-of-particulate-matter-pm10-and-pm25>, (Defra, 2022d). This shows annual mean PM_{2.5} concentrations averaged over all included AURN sites that had annual data capture greater than or equal to 75% in the relevant year. Roadside (urban traffic) and urban background sites are shown by separate, labelled, lines. This graph shows years from 2009 onwards: although there was some PM_{2.5} monitoring before then, the number of sites was very small.

Shaded areas surrounding the lines show the 95% confidence interval of the annual mean for each site classification. The very wide confidence intervals in 2009 – 2011, especially for roadside sites, reflect the small number of sites measuring PM_{2.5} in these early years.


Figure 5-19 Annual mean concentrations of PM_{2.5} in the UK, by AURN Site Classification, 2009 to 2021. Shaded areas either side of each line show the 95% confidence interval of the mean. (Source: <https://www.gov.uk/government/statistics/air-quality-statistics/concentrations-of-particulate-matter-pm10-and-pm25>)



Finally, **Figure 5-20** shows the estimated annual emission of PM_{2.5}, from 1990 to 2020. The graph shows that emissions have decreased in a similar manner to emissions of PM₁₀, with a steady decrease from the early 1990s, a clear levelling off, and no further consistent decrease after around 2010, until 2020. The largest source category for PM_{2.5} is combustion in industry, residential and commercial premises. Emissions from this source have increased over the past decade. Emissions of PM_{2.5} from residential combustion are estimated to have fallen by 90% between 1970 and 2020, reflecting the phasing out of coal. However, domestic combustion of wood has increased over the long term, and by 42% since 2000; this has now grown to become a substantial source of total PM_{2.5} emissions (rising from 3% in 1990 to 17% of total emissions in 2020). Also, industrial emissions from the combustion of biomass fuels are more than 2.8 times higher than in 2013 (rising from 6% of total PM_{2.5} emissions in 2013, to 19% in 2020). Since the late 2000's, annual emissions of PM_{2.5} have fluctuated year-on-year as the impact of significant decreases in emissions from some sectors has been largely offset by increases in emissions from domestic combustion of wood and from solid fuel burning by industry, particularly the use of biomass.

However, estimated PM_{2.5} emissions from both road transport and other transport showed a marked decrease in 2020, similar to that observed for PM₁₀ and likely reflecting the impact of Covid-19 restrictions.

Figure 5-20 Estimated Annual UK Emissions of PM_{2.5} (kt), 1990 – 2020. (Source: NAEI 2022)



5.4 Ozone

5.4.1 O₃: Spatial Distribution

Figure 5-21 shows the average number of days per year with maximum daily running 8-hour mean ozone concentration $> 120 \mu\text{g m}^{-3}$, over the **three** years 2019-2021. Highest concentrations occurred in a small area on England's south-east coast. Apart from this, elsewhere, concentrations were generally slightly higher in the south and east of the UK, than in more northern and western areas.

This map shows slightly lower values around some major conurbations (such as London, Birmingham, Nottingham, Derby and Cardiff). In some cases, the main routes between them are also just visible. Ozone concentrations tend to be lower in built-up areas, due to the 'scavenging' effect of nitric oxide (NO), which reacts with ozone.

Figure 5-22 shows the number of days per year with maximum daily running 8-hour mean ozone concentration $> 120 \mu\text{g m}^{-3}$, for 2021 only. This value was lower in 2021 than the 2019-2021 average, reflecting generally low ozone concentrations in 2021. Apart from a small area in England's West Country, most of England and Wales had only 1-5 days above this value, while Scotland and most of Northern Ireland had none.

Figure 5-23 shows the AOT40 statistic, averaged over the past **five** complete years, 2016-2021. The AOT40 statistic (expressed in $\mu\text{g m}^{-3}\cdot\text{hours}$) is the sum of the difference between hourly concentrations greater than $80 \mu\text{g m}^{-3}$ (= 40 ppb) and $80 \mu\text{g m}^{-3}$ over a given period using only the one-hour values measured between 0800 and 2000 Central European Time each day. This shows a general pattern of higher AOT40 values to the south and east of the UK (outside of major cities), with lower values to the north and west.

Figure 5-24 shows the same statistic, for 2021 only. This map reflects generally lower ozone concentrations in 2021 compared to the average of the most recent five years. Highest AOT40 values occurred in the south-west of England (Devon and Cornwall) and parts of East Anglia.

Figure 5-21 Average Number of days with Maximum Daily Running 8h Mean O₃ Concentration > 120 µg m⁻³ 2019-2021





Figure 5-22 Days with Maximum Daily Running 8h Mean O₃ Concentration > 120 µg m⁻³, 2021



(An interactive version of this map is available at <https://uk-air.defra.gov.uk/data/gis-mapping/>).

Figure 5-23 Average AOT40, 2017-2021 ($\mu\text{g m}^{-3}\cdot\text{hours}$)





Figure 5-24 Average AOT40, 2021 ($\mu\text{g m}^{-3}\cdot\text{hours}$)




5.4.2 O₃: Changes Over Time

Figure 5-25 shows a trend plot of ozone concentrations at 12 long-running rural AURN sites over the period 1992-2021 (Aston Hill, Bush Estate, Eskdalemuir, High Muffles, Ladybower, Lough Navar, Lullington Heath, Narberth, Rochester Stoke, Sibton, Strathvaich and Yarner Wood). Rural sites have been chosen because concentrations of ozone are typically highest in rural areas.

Seven sites (Bush Estate, Eskdalemuir, High Muffles, Ladybower, Rochester Stoke, Sibton and Yarner Wood) show highly significant positive trends over this period. Two (Aston Hill and Lough Navar) show positive trends of lower significance. One (Lullington Heath) shows no statistically significant trend, and two (Narberth and Strathvaich) show statistically significant negative trends. While there is no consistent pattern, upward trends are present at more than half of these sites. There is evidence that the 'hemispheric background' ozone concentration has increased since the 1950s, and the observed trends may reflect this (Vinzargan, 2004).


Figure 5-25 De-seasonalised Trends in Ozone Concentration at 12 Long-Running Rural AURN sites, 1992 - 2021.



This increasing trend in background ozone concentrations is also illustrated by **Figure 5-26**, taken from Defra's National Air Quality Statistics web page for ozone, at <https://www.gov.uk/government/statistics/air-quality-statistics/concentrations-of-ozone>, (Defra, 2022e).

This shows the annual mean of the daily maximum 8-hour mean, averaged over all sites that had annual data capture greater than or equal to 75%. Separate lines are shown for urban background sites and rural background sites, with the 95% confidence intervals shown as shaded areas either side of each line. Ozone has been monitored in the UK since the 1980s, and this graph shows years from 1987 onwards.

Figure 5-26 Annual mean daily maximum 8-hour mean O₃ concentrations in the UK, 1987 to 2021. (Source: <https://www.gov.uk/government/statistics/air-quality-statistics/concentrations-of-ozone>)



Ozone is not emitted in significant quantities directly from any source in the UK (instead, it is formed from reactions involving other pollutants). Ozone is therefore not included in the NAEI, and trends in ozone emissions are not covered by this report.

5.5 Sulphur Dioxide


5.5.1 SO₂: Spatial Distribution in the UK

Figure 5-27 shows how the modelled 99.73rd percentile³ of hourly mean sulphur dioxide concentration varied across the UK during 2021. This statistic corresponds approximately to the 25th highest hourly mean (in the case of a full year's data); if greater than the hourly mean limit value it indicates that the limit value was exceeded on more than the 24 permitted occasions. There were no areas in which this statistic exceeded the limit value of 350 µg m⁻³.

Figure 5-28 shows the modelled 99.18th percentile of 24-hour means (which corresponds to the 4th highest day in a full year). If greater than the 24-hourly mean limit value of 125 µg m⁻³, this would indicate that there were more than the permitted three exceedances in the year. There were no areas of the UK where this was the case in 2021. The modelled 99.18th percentile is 10 µg m⁻³ or less over most of the UK: it was higher in some small areas due to specific local industrial and other emissions.


³ Where the Directive allows exceedances on a number of occasions (i.e. limit value not to be exceeded more than a specified number of times per year), percentiles are used to illustrate this. These are simply the xth highest hourly mean divided by the number of hours in a year, or yth highest daily mean divided by the days in a year, expressed as a percentage.

Figure 5-27 99.73rd Percentile of 1-hour Mean SO₂ Concentration, 2021 (µg m⁻³)



Also, an interactive map of annual mean SO₂ is available at <https://uk-air.defra.gov.uk/data/gis-mapping/>).

Figure 5-28 99.18th Percentile of 24-hour Mean SO₂ Concentration, 2021 ($\mu\text{g m}^{-3}$)



Also, an interactive map of annual mean SO₂ is available at <https://uk-air.defra.gov.uk/data/gis-mapping/>).

5.5.2 SO₂: Changes Over Time

Figure 5-29 shows how ambient concentrations have changed over the period 1992 to 2021, at the six AURN monitoring stations that have monitored this pollutant for the longest time and remained in operation in 2021. All six stations show a downward trend that is statistically significant at the highly significant 0.001 level.

However, the decrease has not been linear. At most of these sites, the downward trend is steepest for the 1990s and early 2000s: there is a clear flattening-off in more recent years from around 2010. The pattern observed in ambient SO₂ concentrations appears to reflect changes in national emissions.

Figure 5-29 De-seasonalised Trends in SO₂ Concentration, 1992-2021 at 6 Long-running AURN Sites





Figure 5-30 is based on data from the NAEI and shows the UK's estimated annual emissions of sulphur dioxide from 1990 to 2020 (the most recent year for which data are available). The decrease in emissions over time shown here is the continuation of an ongoing trend observed by the NAEI throughout the 1970s and 1980s, partly due to the decline of the UK's heavy industry. The main source of this pollutant is fossil fuel combustion: SO₂ emissions in the UK have decreased substantially since 1990, due to reductions in the use of coal, gas and oil. More stringent legislation restricting the sulphur content of fuel oils and diesel fuel used in road vehicles has also helped to reduce emissions of SO₂.

Around 2009, the graph flattens off, and shows a slight upturn in total SO₂ emissions in 2012. The NAEI pollutant information page for SO₂ (at

https://naei.beis.gov.uk/overview/pollutants?pollutant_id=8) explains this as follows: “As a result of the economic downturn the drive to cut energy costs has resulted in an increase in solid fuel use, particularly in 2012 some coal-sensitive pollutants have seen a significant rise in coal burning emissions.” Following 2012, the downward trend in SO₂ emissions continues.

The UK has met its targets for 2020, set by the National Emission Ceilings Directive and the Gothenburg Protocol. Latest projections suggest that emissions will need to be reduced further to meet 2030 Emission Reduction Commitments.

Figure 5-30 Estimated Annual UK Emissions of SO₂ (kt), 1990 – 2020 (source: NAEI 2022)



5.6 Carbon Monoxide

5.6.1 CO: Spatial Distribution

Ambient concentrations of CO throughout the UK have been well within the limit value for many years, Therefore, since 2010, maps of modelled concentration have no longer been routinely produced for CO.

5.6.2 CO: Changes over time

Because concentrations of CO are well within the limit value, relatively few monitoring sites are required. Seven urban AURN sites currently monitor this pollutant, of which six (Belfast Centre, Cardiff Centre, Edinburgh St Leonards, Leeds Centre, London Marylebone Road and London North Kensington) have operated for at least 10 years.

Figure 5-31 shows de-seasonalised trends at these six long-running AURN sites, from 1992 to 2021. All six show a highly significant downward trend over the period. A small dip in CO concentrations may be visible in the plot of London Marylebone Road in early 2020: this may be due to the Covid restrictions which reduced traffic flow on many roads.

Figure 5-31 De-seasonalised Trends in CO Concentration, 6 Long-Running AURN Sites 1992-2021





Figure 5-32 shows the estimated annual emissions of CO over the same period: a steady, almost linear year-on-year decrease to 2011 is followed by two years in which estimated emissions remained stable, before decreasing further. The decreasing ambient concentrations reflect declining emissions over the last 25 years. The NAEI attributes the decrease in CO emissions to factors including EU-wide emission standards for road vehicles, a decline in industrial use of solid fuels, and a decline in the production of steel and non-ferrous metals (https://naei.beis.gov.uk/overview/pollutants?pollutant_id=4).

Figure 5-32 Estimated Annual UK Emissions of CO (kt), 1990 – 2020 (source: NAEI 2022)




5.7 Benzene

5.7.1 Benzene: Spatial Distribution


Benzene is found in petrol and in vehicle emissions, therefore higher levels may be expected at roadside locations. However, almost all road links had modelled concentrations below $1.0 \mu\text{g m}^{-3}$. **Figure 5-33** shows modelled annual mean benzene concentrations at roadside locations in 2021, and **Figure 5-34** shows the modelled annual mean background concentrations of benzene in 2021. No locations in the UK exceeded the annual mean limit value of $5.0 \mu\text{g m}^{-3}$.

Figure 5-33 Annual Mean Roadside Benzene Concentration for Major Urban Roads, 2021 ($\mu\text{g m}^{-3}$)



(An interactive version of this map is available at <https://uk-air.defra.gov.uk/data/gis-mapping/>).

Figure 5-34 Annual Mean Background Benzene Concentration, 2021 ($\mu\text{g m}^{-3}$)



(An interactive version of this map is available at <https://uk-air.defra.gov.uk/data/gis-mapping/>).

5.7.2 Benzene: Changes Over Time

Figure 5-35 shows a smoothed trend plot of ambient benzene concentration, based on the combined dataset from 14 long-running sites in the Non-Automatic Hydrocarbon Network, which have operated since 2002. These are: Barnsley Gawber, Belfast Centre, Haringey Roadside, Leamington Spa, Leeds Centre, Liverpool Speke, London Bloomsbury, Manchester Piccadilly, Middlesbrough, Newcastle Centre, Nottingham Centre, Oxford Centre Roadside, Southampton Centre and Stoke-on-Trent Centre.

The smoothed trend plot for these 14 sites shows a slight increase from 2002 to 2004, followed by a steep decrease between 2004 and 2008. From then on, the graph is much flatter, showing little further fall in ambient concentrations of benzene until 2012-2014 when there is a slight rise, followed by a further decrease in subsequent years to 2021. Benzene shows seasonal variation, which is illustrated by the graph (which is not de-seasonalised.)

Figure 5-35 Smoothed Trend Plot of Ambient Benzene Concentration, 14 Long-Running Non-Automatic Sites, 2002-2021





Figure 5-36 shows the estimated total annual UK emission of benzene (in kilotonnes), 1990 - 2020. The data are from the NAEI. The largest UK source of benzene is fuel combustion. Like the ambient concentrations, the estimated annual emissions also appear to have decreased over the period 2000 – 2010, but subsequently flattened off. However, the estimated total annual emission for 2020 was the lowest in the time series so far.

Figure 5-36 Estimated Annual UK Emissions of Benzene (kt), 1990 – 2020 (source: NAEI 2022)



5.8 1,3-Butadiene

5.8.1 1,3-Butadiene: Compliance with AQS Objective

The UK Air Quality Strategy objective for 1,3-butadiene is $2.25 \mu\text{g m}^{-3}$, as a maximum running annual mean. This objective was met throughout the UK by the due date of 31st December 2003.


The Automatic Hydrocarbon Network monitors 1,3-butadiene at four sites: London Marylebone Road (urban traffic), London Eltham (urban background), Auchencorth Moss in Lothian (rural background), and Chilbolton Observatory in Hampshire (also rural background). Chilbolton Observatory replaced a previous rural site in Harwell (Oxfordshire) at the beginning of 2016. Measured concentrations of 1,3-butadiene at all three sites were well within the AQS objective in 2021.

5.8.2 1,3-Butadiene: Changes Over Time

Figure 5-37 shows a smoothed trend chart of ambient 1,3-butadiene concentration between 2000 and 2021 at the four automatic sites. This pollutant shows a little seasonal

variation at the urban background site: this can be seen in the chart, which is not de-seasonalised.


Figure 5-37 Time Series Graph of 1,3-Butadiene Concentration, 2000-2021



London Marylebone Road has historically had the highest concentrations of 1,3-butadiene, but these have decreased substantially between 2000 and 2015. London Eltham has also exhibited a decrease, although concentrations have always been lower than at London Marylebone. Chilbolton Observatory, despite its rural location, has typically reported slightly higher concentrations during its five years of operation than London Eltham. All four sites, both urban and rural, are now measuring annual mean concentrations of less than $0.1 \mu\text{g m}^{-3}$.

Figure 5-38 shows the total estimated UK annual emission of this compound, in kilotonnes, between 1990 and 2020. This appears to have decreased steadily since 2000, though flattening off after 2014. The main source of 1,3-butadiene is vehicle emissions, and the use of catalytic converters since the early 1990s has substantially reduced emissions from this source. The estimated total annual emission for 2020 was the lowest in the time series so far.

Figure 5-38 Estimated Annual UK Emissions of 1,3-Butadiene (kt), 1990 – 2020 (source: NAEI 2022)




5.9 Metallic Elements

5.9.1 Metallic Elements: Spatial Distribution

Figure 5-39, Figure 5-40, Figure 5-41 and Figure 5-42 show modelled annual mean background concentrations of lead (Pb), arsenic (As), cadmium (Cd) and nickel (Ni) respectively in 2021. The spatial distribution patterns are discussed below.

Pb: background concentrations were $0.01 \mu\text{g m}^{-3}$ or less over almost all of the UK. (The map shows concentrations in micrograms per cubic metre, as this is the unit used for the Air Quality Standards Regulations limit value.) Some small areas around major cities had concentrations in the $0.02 - 0.05 \mu\text{g m}^{-3}$ range. Also, just visible on the map, with modelled concentrations in the same range, are some major road routes linking the cities. This is not due to vehicle tailpipe emissions (leaded petrol having been phased out from general sale in the UK in 1999), but to re-suspended road dust - tyre and brake wear is now a significant source of Pb emissions in the UK. Modelled concentrations were well within the limit value of $0.5 \mu\text{g m}^{-3}$ throughout the UK.

Figure 5-39 Annual Mean Background Lead Concentration, 2021 ($\mu\text{g m}^{-3}$)




(An interactive version of this map is available at <https://uk-air.defra.gov.uk/data/gis-mapping/>).

As: this toxic element is a metalloid rather than a metal but is nevertheless measured by the Heavy Metals Network. **Figure 5-40** shows that modelled annual mean background concentrations were 1.2 ng m⁻³ or less throughout most of the UK. However, concentrations in the range 2.5 – 3.6 ng m⁻³ occurred in some small areas, particularly the north-eastern part of England, Yorkshire and Humberside. This pattern reflects the natural sources of airborne arsenic, particularly wind-blown dust. Modelled concentrations were therefore highest in areas where agricultural practices give rise to wind-blown dust (such as parts of eastern England) and where the natural arsenic content of the soil is relatively high. Modelled concentrations were well within the limit value of 6 ng m⁻³ throughout the UK.

Cd: background concentrations were less than 0.3 ng m⁻³ over most of the UK, as shown by **Figure 5-41**. Higher concentrations can be seen in some urban areas, which reflects the sources of cadmium which are primarily industrial. Higher concentrations are also just visible along some major road routes: cadmium is also a constituent of suspended road dust. No parts of the UK had modelled concentrations greater than the annual mean limit value of 5 ng m⁻³.


Ni: background concentrations of Ni were typically 2 ng m⁻³ or less (and usually 1 ng m⁻³ or less, away from urban areas). Like As and Cd, Ni is found in suspended road dust, so it is often possible to see the paths of some major road routes, though these do not feature strongly on the 2021 map (**Figure 5-42**). There are also a few small areas with higher concentrations due to industrial activity, including two locations where modelled concentration exceeded the Ni target value of 20 ng m⁻³ in 2021: these are in the Yorkshire area and in South Wales. As reported in section 4, four zones exceeded the target value for Ni in 2021: Sheffield Urban Area, Yorkshire and Humberside, South Wales and Swansea Urban Area.

Figure 5-40 Annual Mean Background Arsenic Concentration, 2021 (ng m⁻³)




(An interactive version of this map is available at <https://uk-air.defra.gov.uk/data/gis-mapping/>).

Figure 5-41 Annual Mean Background Cadmium Concentration, 2021 (ng m⁻³)



(An interactive version of this map is available at <https://uk-air.defra.gov.uk/data/gis-mapping/>).

Figure 5-42 Annual Mean Background Nickel Concentration, 2021 (ng m⁻³)



(An interactive version of this map is available at <https://uk-air.defra.gov.uk/data/gis-mapping/>).

5.9.2 Lead: Changes Over Time

The Heavy Metals Network monitoring stations are very diverse, ranging from remote rural sites to urban industrial locations. The range of measured ambient concentrations reflects this diversity: annual mean concentrations can be an order of magnitude higher at some sites than at others. Consequently, if using a network average concentration to show changes over time, caution is needed. If the arithmetic mean is used, this statistic can be dominated by the sites with highest concentrations. If one of these sites starts or ceases operation, or if its measured concentrations change substantially (e.g. due to changes in local industry), this may cause a discontinuity in the time series.

Therefore, the time series graphs for metals Pb, As, Cd and Ni show the *median* (50th percentile), rather than the arithmetic mean, of the annual mean concentrations at all Heavy Metals Network sites. (This approach, used in 'Air Pollution in the UK' reports for years 2017 onwards, is that used by the network operators, NPL, to investigate trends in ambient concentrations (NPL, 2016). However, please note that 'Air Pollution in the UK' reports for years up to and including 2016 used a different approach; the metals graphs showed the arithmetic mean but included urban sites only.)

Figure 5-43 shows a time series of the median annual mean concentration of Pb in the PM₁₀ particulate fraction, as measured from 2004 by the UK Heavy Metals Network, as described in **Section 3.1**. (Prior to 2004, Pb in the particulate phase was measured by the six sites comprising the former Multi-Element Network. For further information on this, please see earlier reports in this series. However please note that the sampling method used by the Multi-Element Network is not directly comparable with current sampling methods as it was not size-selective).

The median of the annual mean concentrations from all Heavy Metals Network sites, both urban and rural, is shown. (As highlighted above, this is a change from the 2016 and earlier reports in this series, which showed the arithmetic mean for urban sites only). Please also note that for clarity, this graph uses units of ng m⁻³, rather than µg m⁻³ as used in the modelled maps. Ambient concentrations of Pb have decreased substantially, though not consistently, since 2004.

Figure 5-44 shows NAEI estimated total annual UK emissions of this metal from 1990 to 2020. The phasing-out of lead in petrol in the 1990s greatly reduced emissions of Pb from transport, which had previously been the largest UK source. However, Pb is a constituent of dust from tyre and brake wear, so transport remains a significant source, though emissions are now much lower. Other significant sources include production processes and other industrial combustion.

Figure 5-43 Ambient Concentrations of Pb in PM₁₀, 2004-2021





Figure 5-44 Estimated Annual UK Emissions of Pb (kt), 1990 – 2020 (source: NAEI 2022)



5.9.3 Arsenic: Changes Over Time

Figure 5-45 shows a time series of ambient concentration of arsenic (As) in the PM₁₀ fraction, expressed as the median annual mean of all sites in UK Heavy Metals Network, as described in **Section 3.3.2**. (For pre-2004 non-size selective measurements from the Multi-Element Network, please see earlier reports in this series.)

The average used is the median of all sites' annual means, rather than the arithmetic mean, to avoid confounding effects due to changes at sites where concentrations are particularly high. All sites, both urban and rural have been included. Ambient concentrations of As appear to have decreased substantially between 2004 and 2008, remaining relatively stable until 2015 and from then on slightly increasing.

Figure 5-45 Ambient Concentrations of As in PM₁₀, 2004-2021





Figure 5-46 shows the UK's estimated total annual emission of As (from the NAEI), in kilotonnes, from 1990 to 2020. The largest human-made sources of As are production processes, waste (particularly the open burning of treated wood), and metal (iron and steel) production processes.

Changes in measured ambient As concentrations (since 2004) do not appear to reflect estimated total emissions. The reasons for this have not been investigated but it may be that the results from the monitoring sites reflect local rather than national trends. Furthermore – as mentioned in **Section 5.9.1** above – wind-blown dust is a major natural source of airborne arsenic in some areas.

Figure 5-46 Estimated Annual UK Emissions of As (kt), 1990 – 2020 (source: NAEI 2022)



5.9.4 Cadmium: Changes Over Time

Figure 5-47 shows a time series of ambient concentration of cadmium (Cd) in the PM₁₀ fraction as measured by the UK Heavy Metals Network, described in **Section 3.3.2**. (For pre-2004 non-size selective measurements from the Multi-Element Network, please see earlier reports in this series.)

Again, the graph shows the median of all sites’ annual means, rather than the arithmetic mean, to avoid confounding effects due to changes at sites where concentrations are particularly high. All sites – both urban and rural – have been included; there were 24 sites in operation during 2021. Over the network’s years of operation there has been a decrease in ambient Cd concentrations, but it has not been consistent (for example, Cd shows an apparent increase in 2014, as does Pb).

Figure 5-48 shows the UK’s estimated total annual emission of Cd (in kilotonnes), 1990 to 2020, from the NAEI. The main human-made sources of Cd are combustion in manufacturing industry and production processes. Waste incineration was once a large source until control of this source was improved in the 1990s: it now accounts for only 1% of the UK total.

Figure 5-47 Ambient Concentrations of Cd in PM₁₀, 2004 – 2021





Figure 5-48 Estimated Annual UK Emissions of Cd (kt), 1990 – 2020 (source: NAEI 2022)



5.9.5 Nickel: Changes Over Time

Figure 5-49 shows a time series of median annual mean concentrations of nickel (Ni) in PM₁₀, as measured by all sites in the UK Heavy Metals Network. As with the other metals, information on non-size selective measurements from the older Multi-Element Network can be found in earlier reports in this series.

Again, the graph shows the median, rather than the arithmetic mean, of annual mean concentrations at all sites. This avoids confounding effects due to a small number of sites which measure ambient Ni concentrations very much higher than the others. Ambient concentrations also show a general (though not consistent) decrease over the period of operation of the network: the pattern is similar to that for Cd.

Figure 5-50 shows total estimated annual UK emissions of Ni, from the NAEI, from 1990 to 2020. Stationary combustion in industry is the major source. The NAEI data appear to show a general decrease in Ni emissions over the period for which ambient measurements are available (2004 onwards).

Figure 5-49 Ambient Concentrations of Ni in PM₁₀, 2004 – 2021





Figure 5-50 Estimated Annual UK Emissions of Ni (kt), 1990 – 2020 (source: NAEI 2022)



5.9.6 Mercury: Changes Over Time

The Heavy Metals Network ceased measuring mercury (Hg) in PM₁₀ at the end of 2013. Monitoring of Total Gaseous Mercury (TGM) continued at two sites (London Westminster and Runcorn Weston Point) until 2018. For information on previous years’ measurements of mercury carried out by the Heavy Metals Network and its predecessors from 2004 to 2018, please refer to “*Air Pollution in the UK 2018*” and earlier reports in this series.


Mercury deposition (dry deposition and deposition in precipitation) is still carried out at several rural sites (see Section 3). However, ambient concentrations of Hg in air are now only measured at two rural sites: Chilbolton Observatory in Hampshire (which measures TGM), and Auchencorth Moss in Lothian. The latter site measures Hg in PM_{2.5}, Elemental Gaseous Mercury and Reactive Hg in air. These measurements are carried out using the Tekran instrument, as part of the Rural Mercury Network (see **Section 3.3.12**).

Chilbolton Observatory has only been in operation since 2016, which is not considered long enough to evaluate trends. Auchencorth Moss has monitored the above Hg metrics since 2010, but data annual capture has been consistently less than 50%.

Annual mean concentrations of elemental Hg, for Auchencorth Moss only, are available from UK-AIR and have consistently been in the range 1.3 – 1.4 ng m⁻³ since monitoring of this metric began in 2010. There is no clear trend.

Figure 5-51 shows estimated annual UK emissions of Hg, from 1990 to 2020. The main sources are combustion in industry, waste incineration and production processes. Mercury emissions have steadily decreased between 2006 and 2016, though the decrease appears to have flattened off in more recent years. The main sources are industrial, therefore trends in ambient Hg concentrations at the rural sites where monitoring of this element has continued would not necessarily be expected to reflect these emission trends.

Figure 5-51 Estimated Annual UK Emissions of Hg, 1990 - 2020 (source: NAEI 2022)




5.10 Benzo[a]pyrene

5.10.1 B[a]P: Spatial Distribution

Figure 5-52 shows the modelled annual mean background concentration of benzo[a]pyrene (B[a]P). Most of the UK had modelled concentrations of 0.1 ng m^{-3} or less in 2021: areas of higher concentration reflect the distribution of industrial sources, and areas where there is widespread domestic use of oil and solid fuels for heating. There were two very small areas with annual mean B[a]P concentrations in excess of the limit value of 1 ng m^{-3} in 2021: one is in Yorkshire and Humberside and the other in the Swansea Urban Area of South Wales. These reflect industrial sources in those areas.

Figure 5-52 Annual mean background B[a]P concentration, 2021 (ng m⁻³)



(An interactive version of this map is available at <https://uk-air.defra.gov.uk/data/gis-mapping/>).

5.10.2 B[a]P: Changes Over Time

The PAH monitoring network began operation in 1991, comprising a small number of sites, which increased to over 20 in the late 1990s. However, during the years 2007-2008, the network underwent a further major expansion and re-organisation, including a change of sampling technique. The newer sampling technique used at most sites from 2008 onwards (the “Digital” PM₁₀ sampler) was found to give higher results than the older method. The reason for this is likely to be due to a number of factors, predominantly the newer samplers’ shorter collection period. The shorter collection period is likely to decrease the degradation of the PAHs by ozone or other oxidative species (Sarantiridis, 2014).

Because of these changes in the composition of the network, and in particular the techniques used, temporal variation in PAH concentrations has only been analysed from 2008 in this report. **Figure 5-53** shows how the mean B[a]P concentration has changed in the years since 2008. This graph shows a smoothed trend plot, based on combined data from all sites in the PAH Network. (This network takes monthly samples and the graph shows the mean of all sites’ measurements, for each month.) The composition of the PAH network has changed little since 2008, so it is considered appropriate here to use the data from all sites. In recent years, results less than the limit of detection (LoD) have occurred at some sites: these results have been treated as half the LoD, for the purpose of calculating the averages shown here.

B[a]P shows a strong seasonal variation: this is illustrated by this graph (which is not de-seasonalised).

Following a sharp drop in measured concentrations of B[a]P between 2008 and 2009, B[a]P concentrations then appear to have remained generally stable until 2014 when there was a further decrease. Since then, ambient concentrations appear to show a slight overall downward trend, and 2021 was lower than 2020.

Figure 5-54 shows estimated total UK emissions of B[a]P, 1990 to 2020. Emissions have decreased substantially in recent decades compared to the early 1990s, due in part to measures such as the banning of stubble burning.

Emissions of B[a]P at the present time are dominated by domestic combustion of solid fuels, and the NAEI data indicate that this source (described as “residential/commercial/institutional”) is stable, or even slightly increasing.

Figure 5-53 Smoothed Trend Plot of Ambient Concentrations of Particulate Phase B[a]P, 2008-2021





Figure 5-54 Estimated Annual UK Emissions of Benzo[a]pyrene (kg), 1990 – 2020 (source: NAEI)



6 Effects of Covid-19 Restrictions on Air Quality


Section 6 of the 'Air Pollution in the UK' report for 2020 (Defra, 2021) covered the impact of Covid-19 'lockdown' restrictions in 2020 on air quality. During 2021, many restrictions were still in place across the UK. As such, Section 6 of this 2021 report continues the focus on the effect of Covid-19 restrictions on air quality in the UK and the changes observed as restrictions were lifted.

Last year's report also included a discussion of several other air pollution episodes that happened during 2020. However, in 2021 there were relatively few periods when an AURN site recorded pollution concentrations in the Air Quality index 'High' band, and also no periods with widespread or prolonged pollutant concentrations in the 'High' band. The pollution episodes that did occur are discussed briefly in **Section 6.4**.

A timeline of major milestones relating to lockdowns and easing of restrictions for England, Scotland, Northern Ireland and Wales, during 2021 is provided in **Figure 6-1**. In summary, all regions of the UK went into lockdown at the start of 2021 (the lockdown for Wales carried over from December 2020). England and the Devolved Administrations then followed different pathways out of lockdown, with varying dates for easing restrictions. However, by mid-2021, many of the restrictions had been lifted across the UK.

Reports on the effects of the Covid-19 lockdown restrictions on air quality in Wales (Ricardo Energy & Environment, 2020), Scotland (Ricardo Energy & Environment, 2021a) and Northern Ireland (Ricardo Energy & Environment, 2021b) have been produced on behalf of these Devolved Administrations. These reports are available via the air quality websites for Wales and Scotland and Northern Ireland: the Northern Ireland report is also available on the DAERA website.

Figure 6-1 Timeline of milestones relating to Covid-19 restrictions, during 2021




6.1 Nitrogen Oxides (NO_x and NO₂)

In urban areas, road traffic is a large source of ambient nitrogen oxides. In the 2020 report, it was reported that the reductions in road traffic that resulted from the initial lockdown in 2020 had a significant effect on ambient concentrations of nitrogen oxides. This section looks at how traffic levels varied during 2021, and whether this had an impact on ambient concentrations of NO_x (NO + NO₂).

Since 1st March 2020, the Department for Transport (DfT) have published statistics on estimated transport use on roads in Great Britain (i.e. England, Wales and Scotland only). Details on the methodology and full statistics can be found here:

<https://www.gov.uk/government/statistics/transport-use-during-the-coronavirus-covid-19-pandemic>). **Figure 6-2** shows the estimated change in daily traffic volume in Great Britain, as a percentage, from March 2020 to December 2021. Traffic volumes are compared to the first week in February 2020 (which is shown as 100%).

Figure 6-2 Estimated change in the daily volume of motor vehicles on all roads in Great Britain from 1st March 2020 to 31st December 2021, compared to the equivalent day in the first week of February 2020 (before the first lockdown). The colours represent the day of the week.



At the start of 2021, traffic volumes were around 50-60% of pre-pandemic levels. As restrictions began to ease, the traffic flow increased, and by May 2021 weekday traffic flows were close to (though still slightly below) February 2020 levels.

It can be seen from **Figure 6-2** that in early 2021, when restrictions were in place, the volume of traffic at the weekends (relative to levels immediately pre-pandemic) was lower than on weekdays. However, from mid-May onwards traffic volumes at the weekend were higher than pre-pandemic levels, whereas during the week, traffic levels remained just below those in February 2020. This may be due to changes in travelling patterns once restrictions were lifted, with some people who previously commuted to work by car every day, now choosing to continue working from home either full or part-time. The increase in weekend traffic volumes during the summer period could be related to increased leisure travel in the UK when compared to February. In addition, as overseas travel still had some restrictions in 2021, more people may have been vacationing within the UK.

Figure 6-3 and **Figure 6-4** show box plots of annual mean NO and NO₂ concentrations measured at urban traffic-related air quality monitoring sites in the UK from 2016 to 2021. These charts show the distribution of the annual mean dataset. The boxes show the 25th to 75th percentile of each dataset, with the median (50th percentile) as a bar in the middle and the mean shown as a circle. The 'whiskers' extending from the top and bottom of the boxes represent the 5th to 95th percentile of each dataset.

The figures show that NO concentrations measured in 2021 remained low when compared to pre-pandemic years and were similar to concentrations measured in 2020. NO₂ median concentrations increased slightly in 2021 from 2020 levels but were still lower than those in pre-pandemic years. Therefore, roadside concentrations of NO_x were still lower in 2021 compared to before the pandemic. This may in part be due to the general ongoing decreasing trend in NO_x emissions from traffic which has been observed in recent years, as well as the fact that traffic flows were lower than usual over a substantial part of the year.

Figure 6-3 Box plot showing distribution of annual mean NO concentrations measured at urban traffic sites in the UK from 2016 to 2021.





Figure 6-4 Box plot showing NO_2 concentrations measured at urban traffic sites in the UK from 2016 to 2021.



The majority of NO_x emitted from vehicle exhausts is in the form of nitrogen oxide (NO): approximately 83%, based on the National Atmospheric Emission Inventory's primary NO₂ emission factors for road transport (BEIS, 2022). Therefore, we would expect diurnal variations in roadside concentrations of NO to be closely related to changes in traffic patterns. (NO is quickly oxidised to NO₂, so any relationship to traffic variation is likely to be most clearly detectable at roadside sites). **Figure 6-5** shows the diurnal variation in NO measured at a range of road traffic monitoring sites across the UK. There is a distinct diurnal pattern in the NO concentrations observed across all sites shown, with low concentrations overnight followed by a short strong peak during the morning rush hour and a broader peak late afternoon/early evening. This pattern is evident in the 2019, 2020 and 2021 datasets; however, the magnitudes of the concentrations in 2020 and 2021 are lower than in 2019. This is consistent with traffic emissions still being the main source of NO in urban areas close to the roadside, and with NO emissions from this source not having returned to pre-pandemic concentrations.

Figure 6-5 Diurnal Plots of Average hourly mean NO concentrations measured at selected urban traffic sites, for 2019, 2020 and 2021. The shaded regions represent the 95% confidence intervals in the mean.





Figure 6-6 shows average (arithmetic mean) NO concentration for each day of the week measured at urban traffic AURN sites in the UK. Overall, NO concentrations in 2021 were similar to those in 2020, and much lower than those in 2019: this was true for each day of the week.

Weekday average NO concentrations were similar for 2020 and 2021 (although there is some variation on specific days). However, Saturday and Sunday NO concentrations measured at urban traffic sites in the AURN were slightly higher at the weekend in 2021 than in 2020. This is consistent with **Figure 6-2** above, which showed that weekend traffic volumes were above those immediately before the pandemic, from May 2021 onwards.

Figure 6-6 Average NO concentration measured at urban traffic sites the UK, per day of the week, for 2019, 2020 and 2021. The shaded boxes represent the 95% confidence intervals in the mean.



6.2 Ozone (O₃)

Ozone is a secondary pollutant formed from reactions between NO_x and volatile organic compounds (VOCs) in the presence of sunlight. Ozone concentrations typically peak during spring and summer months, when meteorological conditions are favourable for

ozone formation. Long-range transport of pollution from continental Europe can also contribute to elevated ozone concentrations the UK. Once formed, ozone can remain in the atmosphere for a few days, as such high ozone concentrations can occur over wide areas.

Ozone can be removed from the atmosphere by reaction with NO, therefore ozone concentrations are typically highest in rural areas, where there is limited NO, and lower in urban areas (where emissions of NO are usually higher). Therefore, if traffic emissions of NO decrease, urban concentrations of ozone may become higher than they otherwise would be, due to a reduction in NO available to 'scavenge' ozone from the air.


As reported in 'Air Pollution in the UK 2020' (Defra, 2021), air quality measurements provided some evidence of this in 2020. In 'Air Pollution in the UK 2020' the impact of lockdown on ozone at the three urban traffic AURN sites in the UK which measure O₃ were assessed by comparing measured concentrations with the modelled counterfactual or "business as usual" (BAU) scenario, in the period following the spring 2020 lockdown. At all three sites, there was an increase in ozone during the lockdown period in spring 2020, compared to BAU. The observed increase may have been, in part, due to the reduction in road traffic and resulting decrease in NO_x emissions (Defra, 2021).

For this report, a different approach has been used. For the three urban traffic AURN sites which monitor ozone, the annual mean of maximum daily 8-hour mean concentrations has been calculated. This annual average of maximum daily 8-hour means has been compared for the three urban traffic sites, for years 2016 to 2021.

The results are shown in **Figure 6-7**. This shows that the annual mean of the daily maximum 8-hour means was relatively high in 2020 for all three sites, as reported in the 2020 report in this series. During the start of the lockdown in 2020, the weather was settled with sunshine and temperatures above average, conditions which are favourable for ozone formation. So, relatively high concentrations of this pollutant would have been expected at the time: however, it is possible that the reduction in NO_x emissions from traffic may have allowed roadside ozone concentrations to become higher than they otherwise might.

In 2021, ozone levels were lower than in the previous year for all three urban traffic sites. Ozone is strongly influenced by meteorological conditions and long-range transport, and this can mask any local changes due to lockdowns and changes in traffic emissions. As reported in **Section 4** of this report, concentrations of ozone in 2021 throughout the UK appear to have been relatively low compared with 2020 and previous recent years, with no major ozone episodes. The lower 2021 average daily maximum 8-hour means at these sites probably reflect generally lower ozone concentrations in 2021 compared to 2020, which in turn reflect meteorological conditions in 2021 being less conducive to ozone formation.

Figure 6-7 Annual average of the daily 8-hour maximum ozone concentrations from 2016 to 2021, for urban traffic AURN sites in the UK. Error bars represent the standard deviation of the means.



6.3 Particulate Matter (PM₁₀ and PM_{2.5})

Like ozone, PM can be transported long distances in the atmosphere, and a large contribution to PM₁₀ and PM_{2.5} in the UK is from regional and continental pollution, rather than local emission sources. Winds from the north and west typically transport clean air to the UK, whilst winds from the south and east can bring polluted airmasses and high PM concentrations. Long-range transport makes a large contribution to PM pollution in the UK, and varying meteorological conditions can have a significant impact on the measured ambient PM. As such, high PM episodes can be widespread, and it can be challenging to assess the impact of lockdown measures on local PM concentrations.

Figure 6-8 and **Figure 6-9** show the annual mean of PM₁₀ and PM_{2.5} concentrations, respectively, measured at urban traffic AURN sites, from 2016 to 2021. Only those sites with over 75% data capture for PM₁₀ or PM_{2.5} in all six years from 2016 to 2021 have been included in the analysis here. These are the same sites that were assessed for changes in PM₁₀ and PM_{2.5} in ‘Air Pollution in the UK 2020’ report: however, last year the focus was on the comparison during the first lockdown period in 2020 with mean concentrations measured during the same period over the previous five years. This year, the annual means for each year have been considered and compared.

At most of these sites, the annual mean PM₁₀ concentrations (**Figure 6-8**) were lower in 2020 than in 2019, but not necessarily the lowest in all years since 2016 (except at Camden Kerbside, Chesterfield Roadside, Glasgow High Street and London Marylebone Road). Nor were the 2021 annual mean PM₁₀ concentrations consistently higher than those for 2020.

A similar pattern can be seen for PM_{2.5} in **Figure 6-9**: in most cases the annual mean concentration for 2020 is lower than that for 2019 but is not consistently the lowest since 2016. And the 2021 annual mean PM_{2.5} concentrations are higher than the 2020 values at some sites, but not all.

While these results do not contradict the findings of the analysis in the 2020 report, the lack of a consistent pattern does underline the challenges associated with assessing the impact of traffic restrictions on PM concentrations, given the confounding effects of meteorology and the fact that the contribution from traffic – relative to other sources – is likely to vary from site to site.

Figure 6-8 Annual average of the daily mean PM₁₀ concentrations for each year from 2016 to 2021, for urban traffic sites in the UK. Error bars represent the standard deviation of the means.





Figure 6-9 Annual average of the daily mean PM_{2.5} concentrations for each year from 2016 to 2021, for urban traffic sites in the UK. Error bars represent the standard deviation of the means.



As shown in **Figure 6-1** at the beginning of this chapter, restrictions on travel were in place during the first few months of 2021. However, peaks in both PM₁₀ and PM_{2.5} were observed during springtime as shown in **Figure 6-10** and **Figure 6-11**, which was likely due to the impact of increased emissions from agriculture in both the UK and on the European continent during those months. Long-range transport of particulate matter can mask any potential decreases which may have occurred due to the lockdown restrictions and reduced local emissions. The impact of meteorology on PM concentrations during the initial lockdown period in 2020 was explored in detail in Section 6 of the 'Air Pollution in the UK' report for 2020.

Figure 6-10 Monthly averaged PM₁₀ concentrations measured at urban traffic sites across the UK for 2019, 2020 and 2021. The shaded boxes represent the 95% confidence intervals in the mean.





Figure 6-11 Monthly averaged PM_{2.5} concentrations measured at urban traffic sites across the UK for 2019, 2020 and 2021. The shaded boxes represent the 95% confidence intervals in the mean.



6.4 Other Periods of Interest

There were relatively few periods of elevated air pollution during 2021. Those resulting in measurements in the Daily Air Quality Index 'High' band are summarised below.

6.4.1 Ozone

In particular it was a relatively low year for ozone, with no major spring or summer ozone episodes. Ozone (as explained above) is a secondary, transboundary pollutant – formed in the atmosphere from reactions involving other pollutants – and its formation and ambient concentrations are heavily influenced by meteorological factors. As highlighted in Section 4 of this report, in 2021 there was only one measured exceedance of the ozone population information threshold of $180 \mu\text{g m}^{-3}$ as an hourly mean (at Rochester Stoke, a rural site in Kent), and no exceedances of the population warning threshold of $240 \mu\text{g m}^{-3}$. There were no occasions in 2021 when any site reported an ozone measurement in the 'High' band of the Daily Air Quality Index, i.e. a value greater than $160 \mu\text{g m}^{-3}$ as running 8-hour mean.

6.4.2 Particulate Matter

There were only eight days in 2021 when any AURN site recorded a daily mean PM_{10} concentration in the 'High' band (greater than $75 \mu\text{g m}^{-3}$). These were as follows:

- 13th October at Aberdeen Erroll Park, due to nearby demolition work.
- 31st March at Ealing Horn Lane, possibly also due to local activities although it occurred during a period of unusually warm weather when many AURN sites recorded air pollution in the 'Moderate' band.
- 2nd February, 28th March, 6th September, 26th October, 30th December and 31st December at Port Talbot Margam: this is an urban industrial site and these high values are likely to have been due to local industrial activity.

For $\text{PM}_{2.5}$, there were only three days when any AURN site recorded daily mean concentration in the 'High' band (greater than $53 \mu\text{g m}^{-3}$). These were as follows:

- 1st January at Bristol St Paul's.
- 3rd March at two sites, Chesterfield Roadside and Barnsley Road. This was a day when the Met Office reported light winds, low cloud and fog, and also forecast the possibility of Saharan dust arriving in the UK. Many AURN sites measured $\text{PM}_{2.5}$ concentrations in the 'Moderate' band on 2nd and 3rd March, but only these two sites went into the 'High' band.
- 5th November at Swansea Roadside, attributed to Bonfire Night celebrations.

The period around Bonfire Night (5th November) remained in the Low band throughout the UK except in two zones – the South East, where 'Moderate' levels were measured, and South Wales, which as stated above recorded a 'High' daily mean for $\text{PM}_{2.5}$.

7 Where to Find Out More

Defra's web pages relating to air quality can be found at <https://www.gov.uk/government/policies/protecting-and-enhancing-our-urban-and-natural-environment-to-improve-public-health-and-wellbeing/supporting-pages/international-european-and-national-standards-for-air-quality>. These provide details of what the UK is doing to tackle air pollution, and the science and research programmes in place.

Also, Defra has published a Guide to Air Pollution Information Resources, detailing the types of information that are made available and this can be found at https://uk-air.defra.gov.uk/assets/documents/reports/cat14/1307241318_Guide_to_UK_Air_Pollution_Information_Resources.pdf.

Information on the UK's air quality, now and in the past, is available on UK-AIR, the Defra online air quality resource at <https://uk-air.defra.gov.uk/>. UK-AIR is the national repository for historic ambient air quality data. It contains measurements from automatic measurement programmes, some dating back to 1972, together with non-automatic sampler measurements dating back to the 1960s. The data archive brings together into one coherent database both data and information from all the UK's measurement networks. Tools available on UK-AIR include the UK Ambient Air Quality Interactive Map at <https://uk-air.defra.gov.uk/data/gis-mapping> that allows you to look at outputs for the national modelling conducted for this compliance assessment, based on pollutant, background or roadside and geographical location.

Similar national online air quality resources have also been developed for Scotland, Wales and Northern Ireland:

- The Welsh Air Quality Archive at <https://airquality.gov.wales/>
- The Scottish Air Quality Archive at <https://www.scottishairquality.scot/>
- The Northern Ireland Archive at <https://www.airqualityni.co.uk/>

Together, these four national websites provide a comprehensive resource for data and analyses covering all aspects of air quality throughout the UK and all its regions.

The Devolved Administrations each produce their own short annual report, providing more specific information on air quality in their parts of the UK. These reports are available from the above websites.

UK-AIR also provides a daily air quality forecast, which is further disseminated via e-mail, RSS feeds and Twitter (see <https://uk-air.defra.gov.uk/twitter>). Latest forecasts are issued daily, at <https://uk-air.defra.gov.uk/forecasting/>.

Detailed pollutant emission data for the UK are available from the National Atmospheric Emissions Inventory (NAEI) at <https://naei.beis.gov.uk/>.

The Clean Air Hub, at <https://www.cleanairday.org.uk/pages/category/clean-air-hub>, brings together information on air pollution, how it affects our health, and the actions we can take

both to protect ourselves from it, and to help tackle it. There is also information on the annual Clean Air Day. The Clean Air Hub is coordinated by Global Action Plan: more information about Global Action Plan can be found at <https://www.globalactionplan.org.uk/>.

Additional information from the Devolved Administrations of Scotland, Wales and Northern Ireland can be found at:

- The Scottish Government Air Quality web page at <https://www.scotland.gov.uk/Topics/Environment/waste-and-pollution/Pollution-1/16215>
- The Welsh Government Environment and Climate Change web pages at <https://gov.wales/environment-climate-change> .
- The Northern Ireland Department of Agriculture, Environment and Rural Affairs (DAERA) web page at <https://www.daera-ni.gov.uk/topics/protect-environment> .

8 References

Air Quality Expert Group, 2013. *Mitigation of United Kingdom PM2.5 Concentrations*. [Online]

Available at: https://uk-air.defra.gov.uk/assets/documents/reports/cat11/1508060903_DEF-PB14161_Mitigation_of_UK_PM25.pdf

[Accessed 21 July 2022].

BEIS, 2022. *Emission Factors for Transport*. [Online]

Available at: <https://naei.beis.gov.uk/data/ef-transport>

BS EN, 2014. *BS EN 12341:2014: Ambient air quality - Standard gravimetric measurement method for the determination of the PM10 or PM2.5 mass fraction of suspended particulate matter*. s.l.:CEN.

Carshaw, D. C. & Ropkins, K., 2012. Openair – an R package for air quality data analysis. *Environmental Modelling & Software*, Volume 27, pp. 52-61.

CEN, 2005a. *EN 14625:2005 Ambient air quality – standard method for the measurement of the concentration of ozone by ultraviolet photometry*, s.l.: BSI.

CEN, 2005b. *EN 14211:2005 Ambient air quality - Standard method for the measurement of the concentration of nitrogen dioxide and nitrogen monoxide by Chemiluminescence*, s.l.: BSI.

CEN, 2005c. *EN 14212:2005 Ambient air quality – Standard method for the measurement of the concentration of sulphur dioxide by UV fluorescence*, s.l.: BSI.

CEN, 2005d. *EN 14626:2005 Ambient air quality - Standard method for the measurement of the concentration of carbon monoxide by non-dispersive infrared spectroscopy*, s.l.: BSI.

CEN, 2005e. *EN 14902:2005 Reference method for determination of Pb/Cd/As/Ni in ambient air*, s.l.: BSI.

CEN, 2005f. *EN 14662-1:2005 Ambient air quality - Standard method for measurement of benzene concentrations - Part 1: Pumped sampling followed by thermal desorption and gas chromatography*, s.l.: BSI.

COMEAP, 2010. *The Mortality Effects of Long-Term Exposure to Particulate Air Pollution in the United Kingdom*. [Online]

Available at:

https://www.gov.uk/government/uploads/system/uploads/attachment_data/file/304641/COMEAP_mortality_effects_of_long_term_exposure.pdf

[Accessed 21 July 2022].

COMEAP, 2011. *The Review of the UK Air Quality Index*. [Online]

Available at:

<https://webarchive.nationalarchives.gov.uk/20140505104700/http://comeap.org.uk/images/stories/Documents/Reports/comeap%20review%20of%20the%20uk%20air%20quality%20index.pdf>

[Accessed 21 July 2022].

COMEAP, 2015. *Statement on the Evidence for the Effects of Nitrogen Dioxide on Health*. [Online]

Available at: <https://www.gov.uk/government/publications/nitrogen-dioxide-health-effects-of-exposure>

[Accessed 21 July 2022].

COMEAP, 2016. *Long term Exposure to Air Pollution and Chronic Bronchitis*. [Online]

Available at:

https://assets.publishing.service.gov.uk/government/uploads/system/uploads/attachment_data/file/541745/COMEAP_chronic_bronchitis_report_2016_rev_07-16.pdf#:~:text=COMEAP%20has%20therefore%20opted%20to%20assess%20the%20non-lethal,the%20symptoms%20of%20ch

[Accessed 21 July 2022].

COMEAP, 2018. *Associations of long-term average concentrations of nitrogen dioxide with mortality*. [Online]

Available at:

https://assets.publishing.service.gov.uk/government/uploads/system/uploads/attachment_data/file/734799/COMEAP_NO2_Report.pdf

[Accessed 21 July 2022].

DAERA, 2017. *The Air Quality Standards (Amendment) Regulations (Northern Ireland) 2017*. [Online]

Available at: <https://www.legislation.gov.uk/nisr/2017/2/contents/made>

[Accessed 21 July 2022].

Defra, 2007. *The Air Quality Strategy for England, Scotland, Wales and Northern Ireland (Volume 1)*. [Online]

Available at: <https://www.gov.uk/government/publications/the-air-quality-strategy-for-england-scotland-wales-and-northern-ireland-volume-1>

[Accessed 21 July 2022].

Defra, 2016a. *Part IV of the Environment Act 1995 Local Air Quality Management Policy Guidance (PG16)*. [Online]

Available at: <https://laqm.defra.gov.uk/documents/LAQM-PG16-April-16-v1.pdf>

[Accessed 21 July 2022].

Defra, 2016b. *Quality Assurance and Quality Control (QA/QC) Procedures for UK Air Quality Monitoring under 2008/50/EC and 2004/107/EC*. [Online]

Available at: [https://uk-](https://uk-air.defra.gov.uk/assets/documents/reports/cat09/1902040953_All_Networks_QAQC_Docu)

[air.defra.gov.uk/assets/documents/reports/cat09/1902040953_All_Networks_QAQC_Docu](https://uk-air.defra.gov.uk/assets/documents/reports/cat09/1902040953_All_Networks_QAQC_Docu)

ment 2012 Issue2.pdf

[Accessed 21 July 2022].

Defra, 2017a. *Air quality plan for nitrogen dioxide (NO₂) in UK*. [Online]

Available at: <https://www.gov.uk/government/publications/air-quality-plan-for-nitrogen-dioxide-no2-in-uk-2017>

[Accessed 21 July 2022].

Defra, 2017b. *UK plan for tackling roadside nitrogen dioxide concentrations*. [Online]

Available at: <https://www.gov.uk/government/publications/air-quality-plan-for-nitrogen-dioxide-no2-in-uk-2017>

[Accessed 21 July 2022].

Defra, 2019a. *Air Quality: National Air Pollution Control Programme*. [Online]

Available at:

https://assets.publishing.service.gov.uk/government/uploads/system/uploads/attachment_data/file/791025/air-quality-napcp-march2019.pdf

[Accessed 21 July 2022].

Defra, 2019b. *Clean Air Strategy 2019*. [Online]

Available at: https://consult.defra.gov.uk/environmental-quality/clean-air-strategy-consultation/user_uploads/clean-air-strategy-2018-consultation.pdf

[Accessed 21 July 2022].

Defra, 2021. *Air Pollution in the UK' series of reports*. [Online]

Available at: <https://uk-air.defra.gov.uk/library/annualreport/index>

[Accessed 05 August 2022].

Defra, 2022a. *National Statistics*. [Online]

Available at: <https://www.gov.uk/government/statistics/emissions-of-air-pollutants/emissions-of-air-pollutants-in-the-uk-summary>

[Accessed 21 July 2022].

Defra, 2022b. *UK-Air Website*. [Online]

Available at: <https://uk-air.defra.gov.uk/>

[Accessed 21 July 2022].

Defra, 2022c. *National statistics - Concentrations of Nitrogen Dioxide*. [Online]

Available at: <https://www.gov.uk/government/statistics/air-quality-statistics/nitrogen-dioxide>

[Accessed 20 September 2022].

Defra, 2022d. *National Statistics - Concentrations of particulate matter (PM₁₀ and PM_{2.5})*. [Online]

Available at: <https://www.gov.uk/government/statistics/air-quality-statistics/concentrations-of-particulate-matter-pm10-and-pm25>

[Accessed 20 September 2022].

Defra, 2022e. *National Statistics - Concentrations of Ozone*. [Online]
Available at: <https://www.gov.uk/government/statistics/air-quality-statistics/concentrations-of-ozone>
[Accessed 20 September 2022].

Department for Environment, Food and Rural Affairs in partnership with the Scottish Executive, Welsh Assembly Government and Department of the Environment Northern Ireland, 2007. *The Air Quality Strategy for England, Scotland, Wales and Northern Ireland*. [Online]
Available at: <https://www.gov.uk/government/publications/the-air-quality-strategy-for-england-scotland-wales-and-northern-ireland-volume-1>
[Accessed 22 July 2022].

Department of Environment Northern Ireland, 2013. *The Pollution Prevention and Control (Industrial Emissions) Regulations (Northern Ireland) 2013*. [Online]
Available at: <https://www.legislation.gov.uk/nisr/2013/160>
[Accessed 1 September 2022].

Department of Environment Northern Ireland, 2010. *The Air Quality Standards Regulations (Northern Ireland) 2010*. [Online]
Available at: <https://www.legislation.gov.uk/nisr/2010/188/contents/made>
[Accessed 21 July 2022].

Department of Health and Social Care, 2016. *Improving outcomes and supporting transparency Part 2: Summary technical specifications of public health indicators, updated August 2016*. [Online]
Available at: <https://www.gov.uk/government/publications/public-health-outcomes-framework-2016-to-2019>
[Accessed 21 July 2022].

EPAQS, 2009. *Defra and the Devolved Administrations' Expert Panel on Air Quality Standards: Metals and Metalloids*. [Online]
Available at:
<https://webarchive.nationalarchives.gov.uk/20130402151656/http://archive.defra.gov.uk/environment/quality/air/airquality/panels/aqs/documents/metals-and-metalloids.pdf>
[Accessed 21 July 2022].

European Parliament and Council of the European Union, 2008. *Council Directive on ambient air quality and cleaner air for Europe (2008/50/EC)*. [Online]
Available at: <https://eur-lex.europa.eu/LexUriServ/LexUriServ.do?uri=CELEX:32008L0050:EN:NOT>
[Accessed 21 July 2022].

European Parliament and Council of the European Union, 2016. *Directive (EU) 2016/2284 of the European Parliament and of the Council of 14 December 2016 on the reduction of national emissions of certain atmospheric pollutants, amending Directive 2003/35/EC and repealing Directive 2001/81/EC*. [Online]

Available at: https://eur-lex.europa.eu/legal-content/EN/TXT/?uri=uriserv:OJ.L_.2016.344.01.0001.01.ENG&toc=OJ:L:2016:344:TOC [Accessed 21 July 2022].

European Parliament and Council of the European, 2004. *Directive 2004/107/EC of the European Parliament and of the Council of 15 December 2004 relating to arsenic, cadmium, mercury, nickel and polycyclic aromatic hydrocarbons in ambient air.* [Online] Available at: <https://eur-lex.europa.eu/LexUriServ/LexUriServ.do?uri=CELEX:32004L0107:EN:NOT> [Accessed 21 July 2022].

HM Government of Gibraltar, 2016. *Environment (Air Quality Standards) (Amendment) Regulations 2016 (Legal Notice No. 2016/242).* [Online] Available at: <https://www.gibraltarlaws.gov.gi/legislations/environment-air-quality-standards-amendment-regulations-2016-2922> [Accessed 21 July 2022].

Mayor of London, 2019. *The London Local Air Quality Management Framework.* [Online] Available at: <https://www.london.gov.uk/what-we-do/environment/pollution-and-air-quality/working-london-boroughs> [Accessed 21 July 2022].

National Atmospheric Emissions Inventory, 2022. *Overview of air pollutants.* [Online] Available at: <https://naei.beis.gov.uk/overview/ap-overview> [Accessed 21 July 2022].

Northern Ireland Government, 2002. *The Environment (Northern Ireland) Order 2002 (SI No. 3153 (N.I. 7)).* [Online] Available at: <https://www.legislation.gov.uk/nisi/2002/3153/contents> [Accessed 21 July 2022].

Northern Ireland Government, 2003. *The Air Quality Regulations (Northern Ireland) 2003 (SI No. 342).* [Online] Available at: <https://www.legislation.gov.uk/nisr/2003/342/contents/made> [Accessed 21 July 2022].

NPL, 2016. *Annual Report for 2015 on the UK Heavy Metals Network (NPL Report ENV9).* [Online] Available at: https://uk-air.defra.gov.uk/assets/documents/reports/cat13/1611011539_NPL_Heavy_Metals_Annual_Report_FINAL_28072016.pdf [Accessed 21 July 2022].

Orellano, P. et al., 2020. Short-term exposure to particulate matter (PM10 and PM2.5), nitrogen dioxide (NO2), and ozone (O3) and all-cause and cause-specific mortality: Systematic review and meta-analysis. *Environ. Int.*, 142:105876(<https://doi.org/10.1016/j.envint.2020.105876>).

Public Health England, 2016. *Lead: General Information*. [Online]
Available at: https://assets.publishing.service.gov.uk/government/uploads/system/uploads/attachment_data/file/562435/lead_general_information.pdf
[Accessed 21 July 2022].

Public Health England, 2018. *Polycyclic aromatic hydrocarbons (Benzo[a]pyrene): general information*. [Online]
Available at: <https://www.gov.uk/government/publications/benzoapyrene-properties-incident-management-and-toxicology/polycyclic-aromatic-hydrocarbons-benzoapyrene-general-information>
[Accessed 21 July 2022].

Public Health England, 2019a. *Benzene: General Information*. [Online]
Available at: <https://www.gov.uk/government/publications/benzene-general-information-incident-management-and-toxicology/benzene-general-information>
[Accessed 21 July 2022].

Public Health England, 2019b. *Arsenic: General Information*. [Online]
Available at: <https://www.gov.uk/government/publications/arsenic-properties-incident-management-and-toxicology/arsenic-general-information>
[Accessed 21 July 2022].

Pugsley, K. L. et al., 2022. *Technical report on UK supplementary modelling assessment under the Air Quality Standards Regulations 2010 for 2020*. [Online]
Available at: https://uk-air.defra.gov.uk/assets/documents/reports/cat09/2203150935_2020_PCM_technical_report.pdf
[Accessed 04 August 2022].

Resongles, E. et al., 2021. Strong evidence for the continued contribution of lead deposited during the 20th century to the atmospheric environment in London of today. *Proceedings of the National Academy of Sciences of the United States of America*, 29 June, 118(26).

Ricardo Energy & Environment, 2020. *Provisional Analysis of Welsh Air Quality Monitoring Data – Impacts of Covid-19 (on behalf of the Welsh Government)*. [Online]
Available at: https://airquality.gov.wales/sites/default/files/documents/2020-08/Analysis_of_Welsh_Air_Quality_Data_Impacts_of_Covid-19_Final_Issue2.pdf
[Accessed 05 September 2022].

Ricardo Energy & Environment, 2021a. *Impact of Lockdown Measures on Scottish Air Quality in 2020 (on behalf of the Scottish Government)*. [Online]
Available at: <https://www.scottishairquality.scot/news/impact-lockdown-measures-scottish-air-quality-2020-report>
[Accessed 05 September 2022].

Ricardo Energy & Environment, 2021b. *Investigation of Effects of Covid-19 Restrictions on Air Quality in Northern Ireland*. [Online]

Available at: <https://www.daera-ni.gov.uk/publications/effects-covid-19-restrictions-air-quality-northern-ireland>

[Accessed 05 September 2022].

Sarantiridis, D., 2014. *Annual Report for 2012 on the UK PAH Monitoring and Analysis Network*. [Online]

Available at: https://uk-air.defra.gov.uk/assets/documents/reports/cat05/1402041506_Defra_PAH_Network_2012_annual_report_FINAL_v2.pdf

[Accessed 21 July 2022].

Scottish Government, 2019. *Cleaner Air for Scotland Strategy - An Independent Review*. [Online]

Available at: <https://www.gov.scot/publications/cleaner-air-scotland-strategy-independent-review/>.

[Accessed 21 July 2022].

Scottish Government, 2000. *The Air Quality (Scotland) Regulations 2000*. (SSI 2000 No. 97). [Online]

Available at: <https://www.legislation.gov.uk/ssi/2000/97/contents/made>

[Accessed 21 July 2022].

Scottish Government, 2002. *The Air Quality (Scotland) Amendment Regulations 2002*. (SSI No. 297). [Online]

Available at: <https://www.legislation.gov.uk/ssi/2002/297/contents/made>

[Accessed 21 July 2022].

Scottish Government, 2010. *The Air Quality Standards (Scotland) Regulations*. [Online]

Available at: <https://www.legislation.gov.uk/ssi/2010/204/made>

[Accessed 21 July 2022].

Scottish Government, 2015. *Cleaner Air for Scotland - The Road to a Healthier Future*. [Online]

Available at: <https://www2.gov.scot/Resource/0048/00488493.pdf>

[Accessed 21 July 2022].

Scottish Government, 2016. *The Air Quality Standards (Scotland) Amendment Regulations 2016*. [Online]

Available at: <https://www.legislation.gov.uk/ssi/2016/376/contents/made>

[Accessed 21 July 2022].

Scottish Government, 2017. *Local Air Quality Management for Scotland: Policy Guidance*. [Online]

Available at: <https://www.gov.scot/Publications/2016/03/9717>

[Accessed 21 July 2022].

Scottish Government, 2021. *Cleaner Air for Scotland 2 - Towards a Better Place for Everyone*. [Online]

Available at: <https://www.gov.scot/publications/cleaner-air-scotland-2-towards-better-place-everyone/>

[Accessed 21 July 2022].

UK Government, 1995. *Environment Act 1995*. [Online]

Available at: <https://www.legislation.gov.uk/ukpga/1995/25/contents>

[Accessed 21 July 2022].

UK Government, 2000. *The Air Quality (England) Regulations 2000*. (SI 2000 No.928). [Online]

Available at: <https://www.legislation.gov.uk/uksi/2000/928/introduction/made>

[Accessed 21 July 2022].

UK Government, 2002. *The Air Quality (England) (Amendment) Regulations 2002* (SI No. 3043). [Online]

Available at: <https://www.legislation.gov.uk/uksi/2002/3043/contents/made>

[Accessed 21 July 2022].

UK Government, 2010. *UK Air Quality Standards Regulations 2010*. [Online]

Available at: <https://www.legislation.gov.uk/uksi/2010/1001/contents/made>

[Accessed 21 July 2022].

UK Government, 2016. *The Air Quality Standards (Amendment) Regulations 2016*. [Online]

Available at: <https://www.legislation.gov.uk/uksi/2016/1184/contents/made>

[Accessed 21 July 2022].

UK Government, 2018. *The National Emission Ceilings Regulations 2018*. [Online]

Available at: <https://www.legislation.gov.uk/uksi/2018/129/made>

[Accessed 21 July 2022].

UK Government, 2021. *Environment Act 2021*. [Online]

Available at: <https://www.legislation.gov.uk/ukpga/2021/30/contents/enacted>

[Accessed 27 July 2022].

UK Government, 2022a. *Air Quality Provisional Common Framework*. [Online]

Available at: <https://www.gov.uk/government/publications/air-quality-provisional-common-framework>

[Accessed 01 September 2022].

UK Government, 2022b. *Open Government Licence*. [Online]

Available at: <https://www.nationalarchives.gov.uk/doc/open-government-licence/version/2/>

[Accessed 21 July 2022].

UK Health Security Agency, 2022. *Chemical Hazards and Poisons Report Issue 28 – June 2022*. [Online]

Available at:

https://assets.publishing.service.gov.uk/government/uploads/system/uploads/attachment_data/file/1083447/CHaPR_AQ_Special_Edition_2206116.pdf

[Accessed 01 September 2022].

UK National Atmospheric Emissions Inventory, 2022. *National Atmospheric Emissions Inventory*. [Online]

Available at: <https://naei.beis.gov.uk/>

[Accessed 21 July 2022].

United Nations Economic Commission for Europe (UNECE), 1999. *Gothenburg Protocol*. [Online]

Available at: <https://unece.org/environment-policy/air/protocol-abate-acidification-eutrophication-and-ground-level-ozone>

[Accessed 01 September 2022].

Vinzargan, R., 2004. A review of surface ozone background levels and trends.

Atmospheric Environment, Volume 38, pp. 3431-3442 .

Welsh Government, 2000. *The Air Quality (Wales) Regulations 2000*. (SI No. 1940 W. 138). [Online]

Available at: <https://www.legislation.gov.uk/wsi/2000/1940/contents/made>

[Accessed 21 July 2022].

Welsh Government, 2002. *The Air Quality (Amendment) (Wales) Regulations 2002* (SI No. 3182 W. 298). [Online]

Available at: <https://www.legislation.gov.uk/wsi/2002/3182/contents/made>

[Accessed 21 July 2022].

Welsh Government, 2010. *The Air Quality Standards (Wales) Regulations*. [Online]

Available at: <https://www.legislation.gov.uk/wsi/2010/1433/contents/made>

[Accessed 21 July 2022].

Welsh Government, 2017. *Local air quality management in Wales*. [Online]

Available at: <https://gov.wales/sites/default/files/publications/2019-04/local-air-quality-management-in-wales.pdf>

[Accessed 21 July 2022].

Welsh Government, 2020. *The Clean Air Plan for Wales – Healthy Air, Healthy Wales*. [Online]

Available at: <https://gov.wales/sites/default/files/publications/2020-08/clean-air-plan-for-wales-healthy-air-healthy-wales.pdf>

[Accessed 21 July 2022].

WHO, 2005. *Air quality guidelines. Global update 2005. Particulate matter, ozone, nitrogen dioxide and sulfur dioxide*. [Online]

Available at: <https://www.euro.who.int/en/what-we-do/health-topics/environment-and-health/air-quality/publications/pre2009/air-quality-guidelines.-global-update-2005.->

particulate-matter,-ozone,-nitrogen-dioxide-and-sulfur-dioxide

[Accessed 21 July 2022].

WHO, 2013. *Review of Evidence on Health Aspects of Air Pollution – REVIHAAP Project*. [Online]

Available at: https://www.euro.who.int/data/assets/pdf_file/0004/193108/REVIHAAP-Final-technical-report-final-version.pdf

[Accessed 21 July 2022].

WHO, 2018. *Ambient (Outdoor) Air Pollution factsheet*. [Online]

Available at: [https://www.who.int/en/news-room/fact-sheets/detail/ambient-\(outdoor\)-air-quality-and-health](https://www.who.int/en/news-room/fact-sheets/detail/ambient-(outdoor)-air-quality-and-health)

[Accessed 21 July 2022].