

# Annexes

<b>ANNEX 1:</b>	<b>Key Categories</b>	<b>579</b>
A 1.1	GENERAL APPROACH USED TO IDENTIFY KEY CATEGORIES	579
A 1.2	QUALITATIVE ANALYSIS USED TO IDENTIFY KEY CATEGORIES	580
A 1.3	QUANTITATIVE APPROACH 1 KCA FOLLOWING IPCC 2006 GUIDELINES	580
A 1.4	QUANTITATIVE APPROACH 2 KCA FOLLOWING IPCC 2006 GUIDELINES	623
A 1.5	KEY CATEGORY ANALYSIS (KCA) RANKING SYSTEM	645
A 1.6	APPROACH USED TO IDENTIFY KP-LULUCF KEY CATEGORIES	647
A 1.7	USING THE UNCERTAINTY ANALYSIS TO PLAN IMPROVEMENTS IN THE PREPARATION OF THE INVENTORY	647
A 1.8	TABLE NIR 3, AS CONTAINED IN THE ANNEX TO DECISION 6/CMP.3	648
<b>ANNEX 2:</b>	<b>Assessment of Uncertainty</b>	<b>651</b>
A 2.1	ESTIMATION OF UNCERTAINTIES USING AN ERROR PROPAGATION APPROACH (APPROACH 1)	651
A 2.1.1	Key Categories	651
A 2.1.2	Tables of uncertainty estimates from the error propagation approach	652
A 2.2	ESTIMATION OF UNCERTAINTY BY SIMULATION (APPROACH 2)	659
A 2.2.1	Overview of the Method	659
A 2.2.2	Methodological details of the Monte Carlo model	659
A 2.2.3	Quality Control Checks on the Monte Carlo Model Output	660
A 2.3	UNCERTAINTY PARAMETERS	661
A 2.3.1	Uncertainty Parameters used	661
A 2.3.2	General Considerations	677
A 2.3.3	Uncertainty in the Trend	677
A 2.4	UNCERTAINTIES IN GWP WEIGHTED EMISSIONS	678
A 2.4.1	Uncertainty in the emissions	678
A 2.4.2	Uncertainty in the Trend	678
A 2.5	SECTORAL UNCERTAINTIES	680
A 2.5.1	Overview of the Method	680

A 2.6	COMPARISON OF UNCERTAINTIES FROM THE ERROR PROPAGATION AND MONTE CARLO ANALYSES	685
<b>ANNEX 3: Other Detailed Methodological Descriptions for Individual Source or Sink Categories, Including for KP-LULUCF Activities. 687</b>		
A 3.1	ENERGY 687	
	A 3.1.1 Emission factors	687
	A 3.1.2 Commonly used references	687
	A 3.1.3 Feedstocks and Non-Energy Use (NEU) of fuels	690
	A 3.1.4 Aviation (MS 7)	698
	A 3.1.5 Gas leakage	701
A 3.2	INDUSTRIAL PROCESSES (CRF SECTOR 2)	702
A 3.3	AGRICULTURE (CRF SECTOR 3)	702
	A 3.3.1 Enteric Fermentation (3A)	703
	A 3.3.2 Manure Management (3B)	704
	A 3.3.3 Agricultural Soils (3D)	708
A 3.4	LAND USE, LAND USE CHANGE AND FORESTRY (CRF SECTOR 4)	717
	A 3.4.1 Carbon stock changes due to afforestation and forest management (4A)	723
	A 3.4.2 Land Use Change and Soils (4B, 4C, 4E)	734
	A 3.4.3 Changes in stocks of carbon in non-forest biomass due to management and land use change (4B2, 4C2, 4E2)	743
	A 3.4.4 Carbon stock changes and biomass burning emissions due to Deforestation (4B, 4C, 4E, 4G)	748
	A 3.4.5 Biomass Burning – Forest and Non-Forest Wildfires (4A, 4B, 4C)	751
	A 3.4.6 Emissions from drainage (organic soils) (4B1, 4C1)	755
	A 3.4.7 Emissions of N <sub>2</sub> O due to disturbance associated with land use conversion and land management changes (4(III))	755
	A 3.4.8 On-site and off-site emissions from peat extraction (4D)	756
	A 3.4.9 Flooded Lands (4D)	759
	A 3.4.10 Harvested Wood Products (4G)	760
	A 3.4.11 Methods for the Overseas Territories (OTs) and Crown Dependencies (CDs)	761

A 3.4.12	Uncertainty analysis of the LULUCF sector	768
A 3.5	WASTE (CRF SECTOR 5)	769
A 3.5.1	Solid Waste Disposal on Land (5A)	769
A 3.5.2	Biological Treatment of Solid Waste (5B)	778
A 3.5.3	Waste Incineration (5C)	780
A 3.5.4	Wastewater Handling (5D)	783
A 3.6	UK CROWN DEPENDENCIES AND OVERSEAS TERRITORIES	787
A 3.6.1	Overview of Data Sources	787
A 3.6.2	Activity and Emissions Data	787
<b>ANNEX 4: National Energy Balance</b>		<b>793</b>
A 4.1	UK ENERGY BALANCE	793
A 4.2	FUELS DATA	798
A 4.2.1	Reallocations of energy data and differences from UK energy statistics	799
<b>ANNEX 5: Additional Information to be Considered as Part of the Annual Inventory Submission and the Supplementary Information Required Under Article 7, paragraph 1, of the Kyoto Protocol Other Useful Reference Information.</b>		<b>815</b>
A 5.1	ANNUAL INVENTORY SUBMISSION	815
A 5.2	SUPPLEMENTARY INFORMATION UNDER ARTICLE 7, PARAGRAPH 1	815
<b>ANNEX 6: Comparison of Emission Estimates Using Atmospheric Observations</b>		<b>817</b>
A 6.1	MODELLING APPROACH USED FOR COMPARISON WITH THE UK GHGI	817
A 6.2	METHANE	818
A 6.3	NITROUS OXIDE	820
A 6.4	HYDROFLUOROCARBONS	822
A 6.4.1	HFC-134a	822
A 6.4.2	HFC-152a	823
A 6.4.3	HFC-125	824
A 6.4.4	HFC-143a	825

A 6.4.5	HFC-23	826
A 6.4.6	HFC-32	828
A 6.4.7	HFC-43-10mee	829
A 6.4.8	HFC-227ea	830
A 6.4.9	HFC-365mfc	831
A 6.4.10	HFC-245fa	832
A 6.5	PERFLUOROCARBONS	833
A 6.5.1	PFC-14	833
A 6.5.2	PFC-116	834
A 6.5.3	PFC-218	835
A 6.5.4	PFC-318	836
A 6.6	SULPHUR HEXAFLUORIDE	837
A 6.7	NITROGEN TRIFLUORIDE	838
A 6.8	CARBON DIOXIDE	838
<b>ANNEX 7: Analysis of EU ETS Data</b>		<b>841</b>
A 7.1	INTRODUCTION	841
A 7.2	BACKGROUND	844
A 7.2.1	EU ETS Data and GHG Inventories	844
A 7.2.2	Scope of the UK EU ETS and Implications for the GHG Inventory	845
A 7.2.3	Limitations of EU ETS Data Integration with GHG Inventory: Autogeneration	847
A 7.3	DATA PROCESSING	848
A 7.4	EU ETS DATA COVERAGE	849
A 7.5	EU ETS DATA USE IN THE UK GHGI	851
A 7.5.1	Activity and Emissions Data	851
A 7.5.2	Implied Emission Factors	856
<b>ANNEX 8: UK Domestic Emissions Reporting Requirements</b>		<b>867</b>
A 8.1	NATIONAL STATISTICS	868
<b>ANNEX 9: End User Emissions</b>		<b>869</b>
A 9.1	INTRODUCTION	869

A 9.2	DEFINITION OF END USERS	869
A 9.3	OVERVIEW OF THE END USER CALCULATIONS	870
A 9.4	EXAMPLE END USER CALCULATION	871
A 9.5	END USER CALCULATION METHODOLOGY FOR THE UK GREENHOUSE GAS INVENTORY	874
A 9.6	DETAILED EMISSIONS ACCORDING TO END USER CATEGORIES	925

### **Table of Tables:**

Table A 1.3.1	Approach 1 Key Category Analysis for the base year based on level of emissions (including LULUCF) .....	582
Table A 1.3.2	Approach 1 Key Category Analysis for the base year based on level of emissions (excluding LULUCF) .....	588
Table A 1.3.3	Approach 1 Key Category Analysis for the latest reported year based on level of emissions (including LULUCF) .....	594
Table A 1.3.4	Approach 1 Key Category Analysis for the latest reported year based on level of emissions (excluding LULUCF) .....	600
Table A 1.3.5	Approach 1 Key Category Analysis based on trend in emissions (from base year to latest reported year, including LULUCF) .....	606
Table A 1.3.6	Approach 1 Key Category Analysis based on trend in emissions (from base year to latest reported year, excluding LULUCF) .....	615
Table A 1.5.1	KCA Ranking.....	645
Table A 1.8.1	Table NIR 3. Summary overview for Key Categories for Land Use, Land-Use Change and Forestry Activities under the Kyoto Protocol .....	648
Table A 2.1.1	Summary of error propagation uncertainty estimates including LULUCF, base year to the latest reported year .....	652
Table A 2.3.1	Uncertainties in the activity data and emission factors for fuels used in the carbon dioxide (CO <sub>2</sub> ) inventory .....	662
Table A 2.3.2	Estimated uncertainties in the activity data and emission factors used in the methane (CH <sub>4</sub> ) inventory .....	670
Table A 2.3.3	Estimated uncertainties in the activity data and emission factors used in the nitrous oxide (N <sub>2</sub> O) inventory .....	673
Table A 2.3.4	Estimated uncertainties in the activity data and emission factors used in the F-gas inventory .....	675
Table A 2.4.1	Summary of Monte Carlo Uncertainty Estimates .....	679
Table A 2.5.1	Sectoral Uncertainty Estimates.....	681

## Contents of Annexes

---

Table A 2.6.1	Comparison of the central estimates and trends in emissions from the error propagation (Approach 1) and Monte Carlo (Approach 2) uncertainty analyses .....	686
Table A 3.1.1	Summary of Emission Sources for UK Fuels Allocated as Non Energy Use in UK Energy Statistics.....	691
Table A 3.1.2	CAA aircraft types assigned to EMEP-EEA Emissions Inventory Guidebook aircraft types.....	698
Table A 3.1.3	Activity data and methane leakage estimates for Gas leakage at Point of Use, including cooking appliances, gas fires and boilers.....	701
Table A 3.3.1	Livestock Population Data by Animal Type ('000 animal places) .....	702
Table A 3.3.2	Methane Emission Factors for Livestock Emissions for 2017 .....	703
Table A 3.3.3	Methane conversion factors for Manure Management Systems in the UK .	704
Table A 3.3.4	Nitrogen Excretion Factors, kg N animal place <sup>-1</sup> year <sup>-1</sup> for livestock in the UK (1990-2017).....	705
Table A 3.3.5	Distribution of Animal Waste Management Systems (%) used for Different Animal types, 2017 .....	706
Table A 3.3.6	Nitrous Oxide Emission Factors for Animal Waste Handling Systems .....	707
Table A 3.3.7	EF for direct N <sub>2</sub> O emissions from managed soils in the UK inventory .....	708
Table A 3.3.8	Areas of UK Crops and quantities of fertiliser applied for 2017 .....	708
Table A 3.3.9	Trends in area grown ('000 ha) and N fertiliser applied (kg/ha) for the major UK crops, 1990-2017.....	710
Table A 3.3.10	Parameter values for crop residue management. ....	712
Table A 3.3.11	N concentrations in above and below ground biomass .....	713
Table A 3.3.12	Mineralised N from soils .....	715
Table A 3.4.1	Main parameters for forest carbon flow model used to estimate carbon uptake by planting of forests of Sitka spruce ( <i>P. Sitchensis</i> ), yield class 12. ....	724
Table A 3.4.2	Afforestation rate of conifers and broadleaves in the United Kingdom since 1500 based on estimates of woodland area by age from the NFI and administrative records. ....	731
Table A 3.4.3	Grouping of MLC land cover types for soil carbon change modelling.....	734
Table A 3.4.4	Grouping of Countryside Survey Broad Habitat types (Jackson, 2000) for soil carbon change modelling.....	735
Table A 3.4.5	Sources of land use change data used to estimate changes in soil carbon in Great Britain for different periods.....	735
Table A 3.4.6	Sources of land use change data used to estimate changes in soil carbon in Northern Ireland for different periods. ....	736

## Contents of Annexes

---

Table A 3.4.7	Soil carbon stock (TgC = MtC) for depths to 1 m in different land types in the UK.....	737
Table A 3.4.8	Weighted average change in equilibrium soil carbon density (t ha <sup>-1</sup> ) to 1 m deep for changes between different land types in England.....	739
Table A 3.4.9	Weighted average change in equilibrium soil carbon density (t ha <sup>-1</sup> ) to 1 m deep for changes between different land types in Scotland.....	739
Table A 3.4.10	Weighted average change in equilibrium soil carbon density (t ha <sup>-1</sup> ) to 1 m deep for changes between different land types in Wales.....	739
Table A 3.4.11	Weighted average change in equilibrium soil carbon density (t ha <sup>-1</sup> ) to 1 m deep for changes between different land types in Northern Ireland.....	739
Table A 3.4.12	Rates of change of soil carbon for land use change transitions.....	740
Table A 3.4.13	Range of times for soil carbon to reach 99% of a new value after a change in land use in England (E), Scotland (S) and Wales (W).....	740
Table A 3.4.14	Aggregation of Agricultural Census crop types for estimating biomass carbon stock changes from Cropland Management.....	744
Table A 3.4.15	Biomass stock factors for UK Cropland types.....	745
Table A 3.4.16	Aggregation of Countryside Survey Broad Habitats for estimating biomass carbon stock changes from Grassland Management.....	745
Table A 3.4.17	Biomass stock factors for UK Grassland types.....	746
Table A 3.4.18	Mean biomass carbon stock densities, tC/ha.....	747
Table A 3.4.19	Weighted average change in equilibrium biomass carbon density (kg m <sup>2</sup> ) for changes between different land types in England.....	747
Table A 3.4.20	Weighted average change in equilibrium biomass carbon density (kg m <sup>2</sup> ) for changes between different land types in Scotland.....	747
Table A 3.4.21	Weighted average change in equilibrium biomass carbon density(kg m <sup>2</sup> ) for changes between different land types in Wales.....	747
Table A 3.4.22	Weighted average change in equilibrium biomass carbon density (kg m <sup>2</sup> ) for changes between different land types in Northern Ireland.....	748
Table A 3.4.23	Countryside Survey data for Forest conversion.....	750
Table A 3.4.24	Corrected Forest conversion rates.....	750
Table A 3.4.25	IRS database property type descriptions by LULUCF sub-category.....	753
Table A 3.4.26	Activity data for peat extraction sites in Northern Ireland.....	756
Table A 3.4.27	Annual peat production, m <sup>3</sup> for England and Scotland (from Annual Minerals Raised Inquiry/Mineral Extraction in Great Britain reports).....	758
Table A 3.4.28	Information sources for estimating LULUCF emissions from the Overseas Territories and Crown Dependencies.....	763

## Contents of Annexes

---

Table A 3.4.29	Assumptions used in applying the Tier 1 methodology to the Overseas Territories and Crown Dependencies .....	764
Table A 3.4.30	IPCC Tier 1 factors used for estimating LULUCF emissions from Overseas Territories and Crown Dependencies .....	767
Table A 3.5.1	Composition of waste sorting residues and mixed municipal waste .....	770
Table A 3.5.2	DOC, DOCf and composition of waste materials .....	771
Table A 3.5.3	Amount of waste landfilled (1945 to 2017) .....	773
Table A 3.5.4	Amount of waste landfilled and methane generated, captured, utilised, flared, oxidised and emitted.....	774
Table A 3.5.5	Parameters used in landfill emission estimates for overseas territories and crown dependencies.....	776
Table A 3.5.6	Activity Data: Inputs in the composting process.....	778
Table A 3.5.7	Activity Data: Inputs in the anaerobic digestion process .....	779
Table A 3.5.8	Activity Data: UK Waste Incineration .....	780
Table A 3.5.9	Emissions Data: UK Waste Incineration .....	781
Table A 3.5.10	UK Domestic and Commercial Waste Water Treatment (5D1) Activity Data .....	783
Table A 3.5.11	UK Domestic and Commercial Waste Water Treatment (5D1) Implied Emission Factors.....	784
Table A 3.5.12	UK Private Waste Water Management System Emission Estimates Parameters (5D1).....	785
Table A 3.5.13	UK Industrial Waste Water Treatment Activity Data (5D2).....	786
Table A 3.6.1	Isle of Man, Guernsey and Jersey – Emissions of Direct GHGs (Mt CO <sub>2</sub> equivalent).....	787
Table A 3.6.2	Isle of Man, Guernsey and Jersey – Combustion activity data.....	788
Table A 3.6.3	Isle of Man, Guernsey and Jersey – Animal numbers.....	788
Table A 3.6.4	Isle of Man, Guernsey and Jersey – Total emissions from Agricultural Soils (kg N <sub>2</sub> O) .....	789
Table A 3.6.5	Cayman Islands, Falklands Islands, and Bermuda - Emissions of Direct GHGs (Mt CO <sub>2</sub> equivalent) .....	789
Table A 3.6.6	Cayman Islands, Falklands Islands, and Bermuda – Combustion activity data .....	790
Table A 3.6.7	Cayman Islands, Falklands Islands, and Bermuda – Animal numbers.....	790
Table A 3.6.8	Cayman Islands, Falklands Islands, and Bermuda – Total emissions from Agricultural Soils (kg N <sub>2</sub> O) .....	791

## Contents of Annexes

---

Table A 3.6.9	Cayman Islands, Falklands Islands, and Bermuda - Amount of synthetic fertilizer applied .....	791
Table A 3.6.10	Gibraltar – Emissions of Direct GHGs (Mt CO <sub>2</sub> equivalent).....	791
Table A 3.6.11	Gibraltar – Combustion activity data .....	792
Table A 4.1.1	UK Energy Balance for 2017 (thousand tonnes of oil equivalent, gross energy basis) .....	794
Table A 4.2.1	Mapping of fuels used in IPCC and the NAEI .....	798
Table A 4.2.2	Fuel reconciliation - coal use in the latest year (Mtonnes).....	800
Table A 4.2.3	Fuel reconciliation – natural gas use in the latest year (Mtherms).....	803
Table A 4.2.4	Fuel reconciliation – Fuel oils use in latest year (Mtonnes).....	805
Table A 4.2.5	Fuel reconciliation – Gas oil use in latest year (Mtonnes) .....	809
Table A 4.2.6	Fuel reconciliation – Use of petroleum gases in the latest year (Mtherms)	812
Table A 7.1.1	Summary of the use of EU ETS data in the UK inventory .....	843
Table A 7.5.1	Refinery Emissions Data Comparison and Revision to OPG Activity .....	853
Table A 7.5.2	EU ETS data for Fuels used at Power Stations and Autogenerators (Emission Factors in kt / Mt for Coal & Fuel Oil, kt / Mth for Gases) .....	856
Table A 7.5.3	Refinery EU ETS Data for Fuel Oil, OPG and Natural Gas (Emission Factors in kt / Mt for Fuel Oil and kt / Mth for OPG and Natural Gas) .....	858
Table A 7.5.4	EU ETS data for fuels used at integrated steelworks & coke ovens (Emission Factors in kt/Mt for solid & liquid fuels, kt/Mth for gases) .....	860
Table A 7.5.5	EU ETS data for Fuels used at Cement Kilns (kt / Mt) .....	861
Table A 7.5.6	Comparison of Cement Sector Carbon Dioxide Emissions* within the UK GHGI and the EU ETS .....	863
Table A 7.5.7	EU ETS data for Fuels used at Lime Kilns (Emission Factors in kt / Mt for Solid Fuels and kt / Mth for Gases) .....	863
Table A 7.5.8	EU ETS emission factor data for production of lime (kt / Mt lime produced) .....	864
Table A 7.5.9	EU ETS data for Coal, Fuel Oil and Natural Gas used by Industrial Combustion Plant (Emission Factors in kt / Mt for Coal & Fuel Oil, kt / Mth for Natural Gas) .....	865
Table A 8.1.1	Summary table of GHG emissions by NC Category, including net emissions/removals from LULUCF (Mt CO <sub>2</sub> eq) – National Statistics coverage (UK only) .....	868
Table A 8.1.2	Summary table of GHG emissions by Gas, including net emissions/removals from LULUCF (Mt CO <sub>2</sub> eq) – National Statistics coverage (UK only) .....	868
Table A 9.4.1	Example of the outputs during an end user calculation.....	873

Table A 9.5.1	Sources reallocated to end users and the fuels used.....	875
Table A 9.5.2	End user category, IPCC sectors, and NAEI source names and activity names used in the emission calculation .....	878
Table A 9.6.1	End user emissions from all National Communication categories, MtCO <sub>2</sub> equivalent.....	926
Table A 9.6.2	End user CO <sub>2</sub> emissions from all National Communication categories, MtCO <sub>2</sub> equivalent.....	926
Table A 9.6.3	End user CH <sub>4</sub> emissions from all National Communication categories, MtCO <sub>2</sub> equivalent.....	927
Table A 9.6.4	End user N <sub>2</sub> O emissions from all National Communication categories, MtCO <sub>2</sub> equivalent.....	928

### Table of Figures:

Figure A 3.1	Data flow diagrams for each land use sub-category, showing cross-linkages between sectors: (i) 4A and 4G, (ii) 4B, (iii) 4C, (iv) 4D, (v) 4E .....	718
Figure A 3.2	The CARBINE-SCA model .....	728
Figure A 3.3	Stratification of environments across the UK with areas 1 to 6 based on the underlying Land Classification (45 classes). .....	737
Figure A 3.4	Decision tree for assessing the effects of cropland management activities on soil carbon stocks. ....	742
Figure A 3.5	Crop type area in the UK 1989-2017 .....	744
Figure A 3.6	Annual area of FIRMS thermal anomalies for GB for 2001 to the current inventory year (thermal anomalies were filtered to exclude those recorded over urban/industrial areas).....	754
Figure A 3.7	Time series of wildfire burnt areas in the UK 1990 to the current inventory year .....	754
Figure A 3.8	The largest uncertainties in the LULUCF inventory, in terms of standard deviation in the output distributions.....	769
Figure A 6.1	Monthly Northern Hemisphere trend in methane estimated from Mace Head observations (blue line). Red line denotes the annual trend. Data under grey shading are not yet ratified. ....	818
Figure A 6.2	Verification of the UK emission inventory estimates for methane in Gg yr <sup>-1</sup> from 1990. GHGI estimates are shown in black. InTEM (MHD+CBW, 3-year) estimates are shown in blue ( $\pm 1\sigma$ ). InTEM (DECC, 12-mth) estimates are shown in orange ( $\pm 1\sigma$ ).....	819
Figure A 6.3	Verification of the UK emission inventory estimates for methane in Gg yr <sup>-1</sup> from 2013. GHGI estimates are shown in black. InTEM (DECC 2-month) estimates are shown in blue ( $\pm 1\sigma$ ).....	820

Figure A 6.4	Monthly Northern Hemisphere trend in nitrous oxide estimated from Mace Head observations (blue line). Red line denotes the annual trend. Data under grey shading are not yet ratified.....	820
Figure A 6.5	Verification of the UK emission inventory estimates for nitrous oxide in Gg yr <sup>-1</sup> from 1990. GHGI estimates are shown in black. InTEM (MHD-only, 3-year) estimates are shown in blue ( $\pm 1\sigma$ ). InTEM (DECC, 12-mth) estimates are shown in orange ( $\pm 1\sigma$ ).....	821
Figure A 6.6	Verification of the UK emission inventory estimates for nitrous oxide in Gg yr <sup>-1</sup> from 2013. GHGI estimates are shown in black. InTEM (DECC 2-month) estimates are shown in blue ( $\pm 1\sigma$ ).....	822
Figure A 6.7	Monthly Northern Hemisphere trend in HFC-134a estimated from Mace Head observations (blue line). Red line denotes the annual trend. Data under grey shading are not yet ratified. ....	822
Figure A 6.8	Verification of the UK emission inventory estimates for HFC-134a in Gg yr <sup>-1</sup> from 1990. GHGI estimates are shown in black. InTEM (MHD-only, 3-year) estimates are shown in blue ( $\pm 1\sigma$ ). InTEM (DECC, 12-mth) estimates are shown in orange ( $\pm 1\sigma$ ).....	823
Figure A 6.9	Monthly Northern Hemisphere trend in HFC-152a estimated from Mace Head observations (blue line). Red line denotes the annual trend. Data under grey shading are not yet ratified. ....	823
Figure A 6.10	Verification of the UK emission inventory estimates for HFC-152a in Gg yr <sup>-1</sup> from 1990. GHGI estimates are shown in black. InTEM (MHD-only, 3-year) estimates are shown in blue ( $\pm 1\sigma$ ). InTEM (DECC, 12-mth) estimates are shown in orange ( $\pm 1\sigma$ ).....	824
Figure A 6.11	Monthly Northern Hemisphere trend in HFC-125 estimated from Mace Head observations (blue line). Red line denotes the annual trend. Data under grey shading are not yet ratified. ....	824
Figure A 6.12	Verification of the UK emission inventory estimates for HFC-125 in Gg yr <sup>-1</sup> from 1990. GHGI estimates are shown in black. InTEM (MHD-only, 3-year) estimates are shown in blue ( $\pm 1\sigma$ ). InTEM (DECC, 12-mth) estimates are shown in orange ( $\pm 1\sigma$ ).....	825
Figure A 6.13	Monthly Northern Hemisphere trend in HFC-143a estimated from Mace Head observations (blue line). Red line denotes the annual trend. Data under grey shading are not yet ratified. ....	825
Figure A 6.14	Verification of the UK emission inventory estimates for HFC-143a in Gg yr <sup>-1</sup> from 1990. GHGI estimates are shown in black. InTEM (MHD-only, 3-year) estimates are shown in blue ( $\pm 1\sigma$ ). InTEM (DECC, 12-mth) estimates are shown in orange ( $\pm 1\sigma$ ).....	826
Figure A 6.15	Monthly Northern Hemisphere trend in HFC-23 estimated from Mace Head observations (blue line). Red line denotes the annual trend. Data under grey shading are not yet ratified. ....	827

Figure A 6.16	Verification of the UK emission inventory estimates for HFC-23 in Gg yr <sup>-1</sup> from 1990. GHGI estimates are shown in black. InTEM (MHD-only, 3-year) estimates are shown in blue ( $\pm 1\sigma$ ). InTEM (DECC, 12-mth) estimates are shown in orange ( $\pm 1\sigma$ ). The second plot is plotted from the year 2000 on an expanded y-axis. ....	827
Figure A 6.17	Monthly Northern Hemisphere trend in HFC-32 estimated from Mace Head observations (blue line). Red line denotes the annual trend. Data under grey shading are not yet ratified. ....	828
Figure A 6.18	Verification of the UK emission inventory estimates for HFC-32 in Gg yr <sup>-1</sup> from 1990. GHGI estimates are shown in black. InTEM (MHD-only, 3-year) estimates are shown in blue ( $\pm 1\sigma$ ). InTEM (DECC, 12-mth) estimates are shown in orange ( $\pm 1\sigma$ ).....	829
Figure A 6.19	Monthly Northern Hemisphere trend in HFC-43-10mee estimated from Mace Head observations (blue line). Red line denotes the annual trend. Data under grey shading are not yet ratified.....	829
Figure A 6.20	Verification of the UK emission inventory estimates for HFC-43-10mee in Gg yr <sup>-1</sup> from 1990. GHGI estimates are shown in black. InTEM (MHD-only, 3-year) estimates are shown in blue ( $\pm 1\sigma$ ). InTEM (DECC, 12-mth) estimates are shown in orange ( $\pm 1\sigma$ ).....	830
Figure A 6.21	Monthly Northern Hemisphere trend in HFC-227ea estimated from Mace Head observations (blue line). Red line denotes the annual trend. Data under grey shading are not yet ratified. ....	830
Figure A 6.22	Verification of the UK emission inventory estimates for HFC-227ea in Gg yr <sup>-1</sup> from 1990. GHGI estimates are shown in black. InTEM (MHD-only, 3-year) estimates are shown in blue ( $\pm 1\sigma$ ). InTEM (DECC, 12-mth) estimates are shown in orange ( $\pm 1\sigma$ ).....	831
Figure A 6.23	Monthly Northern Hemisphere trend in HFC-365mfc estimated from Mace Head observations (blue line). Red line denotes the annual trend. Data under grey shading are not yet ratified.....	831
Figure A 6.24	Verification of the UK emission inventory estimates for HFC-365mfc in Gg yr <sup>-1</sup> from 1990. GHGI estimates are shown in black. InTEM (MHD-only, 3-year) estimates are shown in blue ( $\pm 1\sigma$ ). InTEM (DECC, 12-mth) estimates are shown in orange ( $\pm 1\sigma$ ).....	832
Figure A 6.25	Monthly Northern Hemisphere trend in HFC-245fa estimated from Mace Head observations (blue line). Red line denotes the annual trend. Data under grey shading are not yet ratified. ....	832
Figure A 6.26	Verification of the UK emission inventory estimates for HFC-245fa in Gg yr <sup>-1</sup> from 1990. GHGI estimates are shown in black. InTEM (MHD-only, 3-year) estimates are shown in blue ( $\pm 1\sigma$ ). InTEM (DECC, 12-mth) estimates are shown in orange ( $\pm 1\sigma$ ).....	833

Figure A 6.27	Monthly Northern Hemisphere trend in PFC-14 estimated from Mace Head observations (blue line). Red line denotes the annual trend. Data under grey shading are not yet ratified. ....	833
Figure A 6.28	Verification of the UK emission inventory estimates for PFC-14 in Gg yr <sup>-1</sup> from 1990. GHGI estimates are shown in black. InTEM (MHD-only, 3-year) estimates are shown in blue ( $\pm 1\sigma$ ). InTEM (DECC, 12-mth) estimates are shown in orange ( $\pm 1\sigma$ ).....	834
Figure A 6.29	Monthly Northern Hemisphere trend in PFC-116 estimated from Mace Head observations (blue line). Red line denotes the annual trend. Data under grey shading are not yet ratified. ....	834
Figure A 6.30	Verification of the UK emission inventory estimates for PFC-116 in Gg yr <sup>-1</sup> from 1990. GHGI estimates are shown in black. InTEM (MHD-only, 3-year) estimates are shown in blue ( $\pm 1\sigma$ ). InTEM (DECC, 12-mth) estimates are shown in orange ( $\pm 1\sigma$ ).....	835
Figure A 6.31	Monthly Northern Hemisphere trend in PFC-218 estimated from Mace Head observations (blue line). Red line denotes the annual trend. Data under grey shading are not yet ratified. ....	835
Figure A 6.32	Verification of the UK emission inventory estimates for PFC-218 in Gg yr <sup>-1</sup> from 1990. GHGI estimates are shown in black. InTEM (MHD-only, 3-year) estimates are shown in blue ( $\pm 1\sigma$ ). InTEM (DECC, 12-mth) estimates are shown in orange ( $\pm 1\sigma$ ).....	836
Figure A 6.33	Monthly Northern Hemisphere trend in PFC-318 estimated from Mace Head observations (blue line). Red line denotes the annual trend. Data under grey shading are not yet ratified. ....	836
Figure A 6.34	Verification of the UK emission inventory estimates for PFC-318 in Gg yr <sup>-1</sup> from 1990. GHGI estimates are shown in black. InTEM (MHD-only, 3-year) estimates are shown in blue ( $\pm 1\sigma$ ). InTEM (DECC, 12-mth) estimates are shown in orange ( $\pm 1\sigma$ ).....	837
Figure A 6.35	Monthly Northern Hemisphere trend in SF <sub>6</sub> estimated from Mace Head observations (blue line). Red line denotes the annual trend. Data under grey shading are not yet ratified. ....	837
Figure A 6.36	Verification of the UK emission inventory estimates for SF <sub>6</sub> in Gg yr <sup>-1</sup> from 1990. GHGI estimates are shown in black. InTEM (MHD-only, 3-year) estimates are shown in blue ( $\pm 1\sigma$ ). InTEM (DECC, 12-mth) estimates are shown in orange ( $\pm 1\sigma$ ).....	838
Figure A 6.37	Monthly Northern Hemisphere trend in NF <sub>3</sub> estimated from Mace Head observations (blue line). Red line denotes the annual trend. Data under grey shading are not yet ratified. ....	838
Figure A 6.38	Monthly Northern Hemisphere trend in CO <sub>2</sub> estimated from Mace Head observations (blue line). Red line denotes annual trend. Data under grey shading are not yet ratified. ....	839

Figure A 9.1	Simplified fuel flows for an end user calculation.....	870
Figure A 9.2	Fuel use in the example calculation.....	872
Figure A 9.3	Comparison of 'direct' and end user emissions of carbon according the sectors considered in the end user example .....	874

## ANNEX 1: Key Categories

This annex contains the key category analysis for the latest GHG inventory<sup>1</sup>. It contains:

- A description of the methodology used for identifying key categories
- Information on the level of disaggregation
- Information to fulfil the reporting requirements of Tables 4.2 and 4.3 of Volume 1 of the 2006 IPCC Guidelines, including and excluding land use, land-use change and forestry (LULUCF).

The annex also contains information relevant to the requirements of reporting under the Kyoto Protocol. The table below contains the additional KP information that Annex 1 needs to contain, and the locations of this information in the Annex<sup>2</sup>.

Requirements	Locations of the relevant information in this Annex
Description of methodology used for identifying key categories, <i>including KP-LULUCF</i>	See sections immediately below including “ <i>General approach used to identify Key Categories</i> ” and “ <i>Approach used to identify KP-LULUCF Key Categories</i> ”.
Reference to the key category tables in the CRF	This Annex of the NIR presents detailed tables of information of the data derived from the key category analysis. These data are used to create the key category tables (Table 7) in the CRF.
Reference to the key category tables in the CRF, <i>including in the KP-LULUCF CRF tables</i>	This Annex of the NIR presents detailed tables of information of the data derived from the key category analysis. These data are used to create the key category KP-LULUCF tables (Table NIR 3) in the CRF.
Information on the level of disaggregation	The tables in this Annex contain information on the level of disaggregation used. The level of disaggregation follows IPCC 2006 Guidelines.
Tables 4.2 to 4.4 of Volume 4 the 2006 IPCC guidelines	The data requested in the 2006 Guidelines tables, including and excluding LULUCF, are provided in <b>Table A 1.3.1 to Table A 1.4.6</b> and <b>Table 1.7 to Table 1.10</b> .
<i>Table NIR.3, as contained in the annex to decision 6/CMP.3</i>	A facsimile of Table NIR 3, provided in the CRF, is given in <b>Table A 1.8.1</b> .

### A 1.1 GENERAL APPROACH USED TO IDENTIFY KEY CATEGORIES

In the UK inventory, certain source categories are particularly significant in terms of their contribution to the overall uncertainty of the inventory. These key source categories have been identified so that the resources available for inventory preparation may be prioritised, and the best possible estimates prepared for the most significant source categories.

The UK completes both quantitative and qualitative Key Category Analyses (KCAs).

<sup>1</sup> Following the requirements to report information about uncertainties as set out in FCCC/CP/2013/10/Add.3. Report of the Conference of the Parties on its nineteenth session, held in Warsaw from 11 to 23 November 2013. Addendum Part two: Action taken by the Conference of the Parties at its nineteenth session.

<sup>2</sup> The information in this table has been taken directly from the UNFCCC document “Annotated outline of the National Inventory Report including reporting elements under the Kyoto Protocol”.

The UK has used the method set out in Section 4.3.1 and Section 4.3.2 of the 2006 IPCC Guidelines Volume 1 General Guidance and Reporting (*Approach 1 to identify key categories*, and *Approach 2 to identify key categories* respectively) to quantitatively determine the key source categories.

The method used in the qualitative KCA is described below, and further descriptions of the methods the UK uses to quantitatively determine key categories are given later in this section.

## **A 1.2 QUALITATIVE ANALYSIS USED TO IDENTIFY KEY CATEGORIES**

Following IPCC good practice, a qualitative analysis of the inventory has been made to identify any additional key source categories, which may not have been identified using the quantitative analysis. The approach set out in Section 4.3.3 of the IPCC 2006 Guidelines has been applied, using the four criteria set out in the guidance, to judge whether a category is a key category. The criteria are:

1. (Use of) mitigation techniques and technologies;
2. Emissions growth (increase or decrease);
3. No quantitative assessment of uncertainties performed;
4. Completeness (examine qualitatively potential key categories that are not yet estimated quantitatively by applying the qualitative considerations above).

In addition, additional criteria have also been taken in account

5. High uncertainty (links to point 3 above);
6. Unexpectedly low or high emissions;
7. External recommendation has also been used as an additional criterion to identify key categories.

The results of the qualitative analysis did not identify any categories that were not already identified by the quantitative key category analysis.

## **A 1.3 QUANTITATIVE APPROACH 1 KCA FOLLOWING IPCC 2006 GUIDELINES**

A key category analysis has been completed for both level and trend. This KCA has been created using the 2006 IPCC Guidelines Approach 1 methodology. The factors that make a source a key category are:

- A high contribution to the level of emissions; and
- A high contribution to the trend;

For example, transport fuel (1A3b) is a key category for carbon dioxide because it is a large source of emissions and nitric acid production (2B2) because it shows a significant trend.

The category groupings are largely aligned to those suggested in Tables 4.5 and 4.6 in Volume 1, Chapter 4 of the 2006 IPCC guidelines, although we deviate in a number of cases, in particular:

- Agriculture and LULUCF. In the 2006 guidelines a different nomenclature for categorising agriculture and LULUCF sources and sinks was used compared to the adopted nomenclature, which means that it would be challenging and confusing to retain this categorisation when sources are grouped differently in the adopted nomenclature. The

agriculture categories are analysed at the level of mostly 3 digit IPCC codes, whilst analysis for LULUCF sources and sinks is more aggregated. The UK inventory agency considers that the level of aggregation used in the UK method for the KCA is sufficiently detailed to target inventory improvements (e.g. consideration of the more important livestock types under agriculture) whilst not introducing unnecessary computational difficulties (e.g. use of “miscellaneous” categories to mop up the remainder within a sector). Further, the level of source/sink category aggregations in the KCA are aligned to how individual methods or models are used to derive the UK inventory estimates, and are therefore at an appropriate level of detail for the UK inventory

- Fugitive Emissions. The suggested categories are at a much more granular level (e.g. 1B2a<sub>ii</sub>) than other sectors. We considered that this would lead to an undue diminishing of these sectors, decreasing their likelihood of being considered key, so have adopted a level of aggregation more consistent with other sectors
- Miscellaneous emissions. The suggested approach was to group a large number of small sources into one category. We considered that this would lead to an undue increase in the significance of these sources, increasing their likelihood of being considered key, so have adopted a level of aggregation more consistent with other sectors

The results of the key category analysis with and without LULUCF, for the base year and the latest reported year and for both Approaches 1 and 2 KCA, are summarised by sector and gas in **Section 1.5.1**. The tables indicate whether a key category arises from the level (L1) assessment or the trend (T1) assessment.

The results of the **level assessment** (based on Approach 1) with and without LULUCF for the base year and the latest reported year are shown **Table A 1.3.1** to **Table A 1.3.4**. The key source categories are highlighted by the shaded cells in the table. The source categories (i.e. rows of the table) were sorted in descending order of magnitude based on the results of the “Level Parameter”, and then the cumulative total was included in the final column of the table. The key source categories are those whose contributions add up to 95% of the sum of the level parameters in the final column after this sorting process, which according to the 2006 IPCC guidelines, should account for 90% of the uncertainty in level.

The results of the **trend assessment** (based on Approach 1) with and without LULUCF for the base year to the latest reported year are shown in **Table A 1.3.5** and **Table A 1.3.6**. The key source categories are highlighted by the shaded cells in the table. The trend parameter was calculated using the absolute value of the result; an absolute function is used since Land Use, Land Use Change and Forestry contains negative sources (sinks) and the absolute function is necessary to produce positive uncertainty contributions for these sinks. The source categories (i.e. rows of the table) were sorted in descending order of the “Trend parameter”, and then the cumulative total was included in the final column of the table. The key source categories are those whose contributions add up to 95% of the sum of the trend parameters in the final column after this sorting process, which according to the 2006 IPCC guidelines, should account for 90% of the uncertainty in trend.

Note that the tables in chapter 1 of the NIR summarise the key categories from both the approach 1 and approach 2 key categories analyses and the aggregations used are slightly different for the two approaches. The category "3A" is therefore total emissions from category 3A, whilst categories 3A1 and 3A2 have also been identified as key categories in their own right. Category "2B Chemical industries - CO<sub>2</sub>" is total CO<sub>2</sub> emissions from category 2B.

**Table A 1.3.1 Approach 1 Key Category Analysis for the base year based on level of emissions (including LULUCF)**

IPCC Code	IPCC Category	GHG	Base year emissions (Gg CO <sub>2</sub> e)	Absolute value of Base year emissions (Gg CO <sub>2</sub> e)	Level Assessment	Cumulative Total
1A1	Energy industries: solid fuels	CO <sub>2</sub>	185,492.23	185,492.23	0.2184	0.2184
1A3b	Road transportation: liquid fuels	CO <sub>2</sub>	108,570.16	108,570.16	0.1278	0.3462
1A4	Other sectors: gaseous fuels	CO <sub>2</sub>	70,373.19	70,373.19	0.0829	0.4290
5A	Solid waste disposal	CH <sub>4</sub>	60,432.98	60,432.98	0.0711	0.5002
1A1	Energy industries: liquid fuels	CO <sub>2</sub>	40,883.76	40,883.76	0.0481	0.5483
1A2	Manufacturing industries and construction: solid fuels	CO <sub>2</sub>	38,484.20	38,484.20	0.0453	0.5936
1A2	Manufacturing industries and construction: liquid fuels	CO <sub>2</sub>	29,832.00	29,832.00	0.0351	0.6288
1A2	Manufacturing industries and construction: gaseous fuels	CO <sub>2</sub>	27,330.36	27,330.36	0.0322	0.6609
1B1	Coal mining and handling	CH <sub>4</sub>	21,826.68	21,826.68	0.0257	0.6866
2B3	Adipic acid production	N <sub>2</sub> O	19,934.61	19,934.61	0.0235	0.7101
3A1	Enteric fermentation from Cattle	CH <sub>4</sub>	19,881.28	19,881.28	0.0234	0.7335
1A4	Other sectors: solid fuels	CO <sub>2</sub>	19,868.96	19,868.96	0.0234	0.7569
1A4	Other sectors: liquid fuels	CO <sub>2</sub>	19,607.73	19,607.73	0.0231	0.7800
2B9	Fluorochemical production	HFCs, PFCs, SF <sub>6</sub> and NF <sub>3</sub>	17,784.67	17,784.67	0.0209	0.8009
4A	Forest land	CO <sub>2</sub>	-15,026.26	15,026.26	0.0177	0.8186
4B	Cropland	CO <sub>2</sub>	14,265.93	14,265.93	0.0168	0.8354
3D	Agricultural soils	N <sub>2</sub> O	13,610.39	13,610.39	0.0160	0.8514
1B2	Oil and gas extraction	CH <sub>4</sub>	12,345.07	12,345.07	0.0145	0.8660
1A1	Energy industries: gaseous fuels	CO <sub>2</sub>	9,229.34	9,229.34	0.0109	0.8768
1A3d	Domestic Navigation: liquid fuels	CO <sub>2</sub>	7,611.13	7,611.13	0.0090	0.8858
2A1	Cement production	CO <sub>2</sub>	7,295.26	7,295.26	0.0086	0.8944
4C	Grassland	CO <sub>2</sub>	-7,111.03	7,111.03	0.0084	0.9028
4E	Settlements	CO <sub>2</sub>	7,011.30	7,011.30	0.0083	0.9110
1B2	Oil and gas extraction	CO <sub>2</sub>	5,777.92	5,777.92	0.0068	0.9178

IPCC Code	IPCC Category	GHG	Base year emissions (Gg CO <sub>2</sub> e)	Absolute value of Base year emissions (Gg CO <sub>2</sub> e)	Level Assessment	Cumulative Total
2C1	Iron and steel production	CO <sub>2</sub>	5,591.54	5,591.54	0.0066	0.9244
1A5	Other: liquid fuels	CO <sub>2</sub>	5,293.44	5,293.44	0.0062	0.9306
3A2	Enteric fermentation from Sheep	CH <sub>4</sub>	4,935.90	4,935.90	0.0058	0.9364
3B1	Manure management from Cattle	CH <sub>4</sub>	4,732.53	4,732.53	0.0056	0.9420
2B8	Petrochemical and carbon black production	CO <sub>2</sub>	4,538.91	4,538.91	0.0053	0.9473
5D	Wastewater treatment and discharge	CH <sub>4</sub>	4,218.90	4,218.90	0.0050	0.9523
2B2	Nitric acid production	N <sub>2</sub> O	3,860.26	3,860.26	0.0045	0.9569
3B2	Manure management from Sheep	N <sub>2</sub> O	3,442.70	3,442.70	0.0041	0.9609
2B1	Ammonia production	CO <sub>2</sub>	1,895.00	1,895.00	0.0022	0.9631
1A3a	Domestic aviation: liquid fuels	CO <sub>2</sub>	1,881.31	1,881.31	0.0022	0.9654
1B1	Coal mining and handling solid fuels	CO <sub>2</sub>	1,698.56	1,698.56	0.0020	0.9674
4G	Harvested wood products	CO <sub>2</sub>	-1,639.08	1,639.08	0.0019	0.9693
2A2	Lime production	CO <sub>2</sub>	1,462.05	1,462.05	0.0017	0.9710
1A3c	Railways: liquid fuels	CO <sub>2</sub>	1,455.18	1,455.18	0.0017	0.9727
2C6	Zinc production	CO <sub>2</sub>	1,358.83	1,358.83	0.0016	0.9743
1A3b	Road transportation: liquid fuels	N <sub>2</sub> O	1,312.28	1,312.28	0.0015	0.9759
5C	Incineration and open burning of waste	CO <sub>2</sub>	1,300.71	1,300.71	0.0015	0.9774
1A4	Other sectors: solid fuels	CH <sub>4</sub>	1,265.22	1,265.22	0.0015	0.9789
1A3b	Road transportation: liquid fuels	CH <sub>4</sub>	1,246.70	1,246.70	0.0015	0.9804
1A1	Energy industries: solid fuels	N <sub>2</sub> O	1,023.72	1,023.72	0.0012	0.9816
4B	Cropland	N <sub>2</sub> O	1,019.85	1,019.85	0.0012	0.9828
3G	Liming	CO <sub>2</sub>	1,015.18	1,015.18	0.0012	0.9840
5D	Wastewater treatment and discharge	N <sub>2</sub> O	783.59	783.59	0.0009	0.9849
2G1	Electrical equipment	HFCs, PFCs, SF <sub>6</sub> and NF <sub>3</sub>	783.25	783.25	0.0009	0.9858

IPCC Code	IPCC Category	GHG	Base year emissions (Gg CO <sub>2</sub> e)	Absolute value of Base year emissions (Gg CO <sub>2</sub> e)	Level Assessment	Cumulative Total
2F4	Aerosols	HFCs, PFCs, SF <sub>6</sub> and NF <sub>3</sub>	663.00	663.00	0.0008	0.9866
2A4	Other process uses of carbonates	CO <sub>2</sub>	640.93	640.93	0.0008	0.9873
4E	Settlements	N <sub>2</sub> O	584.68	584.68	0.0007	0.9880
2D	Non-energy products from fuels and solvent use	CO <sub>2</sub>	552.81	552.81	0.0007	0.9887
2G3	N <sub>2</sub> O from product uses	N <sub>2</sub> O	552.57	552.57	0.0007	0.9893
2F1	Refrigeration and air conditioning	HFCs, PFCs, SF <sub>6</sub> and NF <sub>3</sub>	531.30	531.30	0.0006	0.9900
4D	Wetlands	CO <sub>2</sub>	486.95	486.95	0.0006	0.9905
2C3	Aluminium production	CO <sub>2</sub>	450.32	450.32	0.0005	0.9911
2A3	Glass production	CO <sub>2</sub>	405.54	405.54	0.0005	0.9915
4	Indirect N <sub>2</sub> O emissions from LULUCF	N <sub>2</sub> O	401.49	401.49	0.0005	0.9920
2C4	Magnesium production	HFCs, PFCs, SF <sub>6</sub> and NF <sub>3</sub>	387.17	387.17	0.0005	0.9925
1A4	Other sectors: peat	CO <sub>2</sub>	372.48	372.48	0.0004	0.9929
2C3	Aluminium production	HFCs, PFCs, SF <sub>6</sub> and NF <sub>3</sub>	333.43	333.43	0.0004	0.9933
3H	Urea application to land	CO <sub>2</sub>	327.60	327.60	0.0004	0.9937
3A4	Enteric fermentation from Other livestock	CH <sub>4</sub>	292.27	292.27	0.0003	0.9940
3A3	Enteric fermentation from Swine	CH <sub>4</sub>	283.06	283.06	0.0003	0.9944
2G2	SF <sub>6</sub> and PFCs from other product use	HFCs, PFCs, SF <sub>6</sub> and NF <sub>3</sub>	278.70	278.70	0.0003	0.9947

IPCC Code	IPCC Category	GHG	Base year emissions (Gg CO <sub>2</sub> e)	Absolute value of Base year emissions (Gg CO <sub>2</sub> e)	Level Assessment	Cumulative Total
3J	Agriculture activities in OTs and CDs	CH <sub>4</sub>	276.73	276.73	0.0003	0.9950
1A1	Energy industries: other fuels	CO <sub>2</sub>	246.13	246.13	0.0003	0.9953
1A4	Other sectors: solid fuels	N <sub>2</sub> O	241.68	241.68	0.0003	0.9956
2B7	Soda ash production	CO <sub>2</sub>	231.55	231.55	0.0003	0.9959
4A	Forest land	N <sub>2</sub> O	231.22	231.22	0.0003	0.9961
1A3e	Other transportation: liquid fuels	CO <sub>2</sub>	224.74	224.74	0.0003	0.9964
1A1	Energy industries: gaseous fuels	N <sub>2</sub> O	198.91	198.91	0.0002	0.9966
3F	Field burning of agricultural residues	CH <sub>4</sub>	186.57	186.57	0.0002	0.9968
2F2	Foam blowing agents	HFCs, PFCs, SF <sub>6</sub> and NF <sub>3</sub>	184.49	184.49	0.0002	0.9971
2B10	Other Chemical Industry	CH <sub>4</sub>	174.65	174.65	0.0002	0.9973
1A4	Other sectors: gaseous fuels	CH <sub>4</sub>	157.18	157.18	0.0002	0.9975
1A2	Manufacturing industries and construction: solid fuels	N <sub>2</sub> O	148.72	148.72	0.0002	0.9976
1A1	Energy industries: liquid fuels	N <sub>2</sub> O	140.41	140.41	0.0002	0.9978
1A2	Manufacturing industries and construction: liquid fuels	N <sub>2</sub> O	138.62	138.62	0.0002	0.9980
5C	Incineration and open burning of waste	CH <sub>4</sub>	134.83	134.83	0.0002	0.9981
3J	Agriculture activities in OTs and CDs	N <sub>2</sub> O	132.00	132.00	0.0002	0.9983
1A3d	Domestic Navigation: liquid fuels	N <sub>2</sub> O	105.00	105.00	0.0001	0.9984
2B6	Titanium dioxide production	CO <sub>2</sub>	104.63	104.63	0.0001	0.9985
1A4	Other sectors: liquid fuels	N <sub>2</sub> O	103.32	103.32	0.0001	0.9986
1A1	Energy industries: gaseous fuels	CH <sub>4</sub>	92.30	92.30	0.0001	0.9987
1A2	Manufacturing industries and construction: other fuels	CO <sub>2</sub>	76.86	76.86	0.0001	0.9988
1A4	Other sectors: biomass	CH <sub>4</sub>	62.37	62.37	0.0001	0.9989
1A4	Other sectors: liquid fuels	CH <sub>4</sub>	57.79	57.79	0.0001	0.9990

IPCC Code	IPCC Category	GHG	Base year emissions (Gg CO <sub>2e</sub> )	Absolute value of Base year emissions (Gg CO <sub>2e</sub> )	Level Assessment	Cumulative Total
3F	Field burning of agricultural residues	N <sub>2</sub> O	57.66	57.66	0.0001	0.9990
1A5	Other: liquid fuels	N <sub>2</sub> O	56.12	56.12	0.0001	0.9991
1A1	Energy industries: solid fuels	CH <sub>4</sub>	49.32	49.32	0.0001	0.9992
1A2	Manufacturing industries and construction: solid fuels	CH <sub>4</sub>	47.59	47.59	0.0001	0.9992
5C	Incineration and open burning of waste	N <sub>2</sub> O	46.66	46.66	0.0001	0.9993
1A2	Manufacturing industries and construction: liquid fuels	CH <sub>4</sub>	44.43	44.43	0.0001	0.9993
1A1	Energy industries: liquid fuels	CH <sub>4</sub>	43.36	43.36	0.0001	0.9994
2G4	Other product manufacture and use	N <sub>2</sub> O	41.00	41.00	0.0000	0.9994
1B2	Oil and gas extraction	N <sub>2</sub> O	40.75	40.75	0.0000	0.9995
1A4	Other sectors: gaseous fuels	N <sub>2</sub> O	37.47	37.47	0.0000	0.9995
2C1	Iron and steel production	CH <sub>4</sub>	36.90	36.90	0.0000	0.9996
2F6	Other product uses as substitutes for ODS	HFCs, PFCs, SF <sub>6</sub> and NF <sub>3</sub>	36.88	36.88	0.0000	0.9996
2A4	Other process uses of carbonates	CH <sub>4</sub>	31.10	31.10	0.0000	0.9997
2B8	Petrochemical and carbon black production	CH <sub>4</sub>	29.98	29.98	0.0000	0.9997
1A4	Other sectors: peat	CH <sub>4</sub>	26.35	26.35	0.0000	0.9997
1A2	Manufacturing industries and construction: biomass	N <sub>2</sub> O	19.54	19.54	0.0000	0.9997
1A1	Energy industries: other fuels	CH <sub>4</sub>	18.55	18.55	0.0000	0.9998
5B	Biological treatment of solid waste	CH <sub>4</sub>	18.13	18.13	0.0000	0.9998
1A3a	Domestic aviation: liquid fuels	N <sub>2</sub> O	17.80	17.80	0.0000	0.9998
2C1	Iron and steel production	N <sub>2</sub> O	17.74	17.74	0.0000	0.9998
1A2	Manufacturing industries and construction: gaseous fuels	N <sub>2</sub> O	14.54	14.54	0.0000	0.9998
1A3c	Railways: liquid fuels	N <sub>2</sub> O	13.75	13.75	0.0000	0.9999
5B	Biological treatment of solid waste	N <sub>2</sub> O	12.97	12.97	0.0000	0.9999

IPCC Code	IPCC Category	GHG	Base year emissions (Gg CO <sub>2e</sub> )	Absolute value of Base year emissions (Gg CO <sub>2e</sub> )	Level Assessment	Cumulative Total
1A2	Manufacturing industries and construction: biomass	CH <sub>4</sub>	12.29	12.29	0.0000	0.9999
1A2	Manufacturing industries and construction: gaseous fuels	CH <sub>4</sub>	12.20	12.20	0.0000	0.9999
4C	Grassland	N <sub>2</sub> O	10.33	10.33	0.0000	0.9999
4C	Grassland	CH <sub>4</sub>	9.97	9.97	0.0000	0.9999
2E1	Integrated circuit or semiconductor	HFCs, PFCs, SF <sub>6</sub> and NF <sub>3</sub>	9.56	9.56	0.0000	0.9999
1A4	Other sectors: biomass	N <sub>2</sub> O	9.14	9.14	0.0000	0.9999
1A1	Energy industries: other fuels	N <sub>2</sub> O	6.72	6.72	0.0000	1.0000
1A3a	Domestic aviation: liquid fuels	CH <sub>4</sub>	6.37	6.37	0.0000	1.0000
4D	Wetlands	N <sub>2</sub> O	4.13	4.13	0.0000	1.0000
1A3d	Domestic Navigation: liquid fuels	CH <sub>4</sub>	3.66	3.66	0.0000	1.0000
1A5	Other: liquid fuels	CH <sub>4</sub>	3.56	3.56	0.0000	1.0000
4E	Settlements	CH <sub>4</sub>	3.00	3.00	0.0000	1.0000
4A	Forest land	CH <sub>4</sub>	2.89	2.89	0.0000	1.0000
1A3e	Other transportation: liquid fuels	N <sub>2</sub> O	2.79	2.79	0.0000	1.0000
1A3c	Railways: liquid fuels	CH <sub>4</sub>	2.46	2.46	0.0000	1.0000
2B8	Petrochemical and carbon black production	N <sub>2</sub> O	2.10	2.10	0.0000	1.0000
1A4	Other sectors: peat	N <sub>2</sub> O	1.47	1.47	0.0000	1.0000
2F3	Fire protection	HFCs, PFCs, SF <sub>6</sub> and NF <sub>3</sub>	1.41	1.41	0.0000	1.0000
1A1	Energy industries: biomass	CH <sub>4</sub>	0.47	0.47	0.0000	1.0000
1A2	Manufacturing industries and construction: other fuels	N <sub>2</sub> O	0.38	0.38	0.0000	1.0000
2B1	Ammonia production	N <sub>2</sub> O	0.31	0.31	0.0000	1.0000
1A3e	Other transportation: liquid fuels	CH <sub>4</sub>	0.29	0.29	0.0000	1.0000
2B1	Ammonia production	CH <sub>4</sub>	0.26	0.26	0.0000	1.0000
1A1	Energy industries: biomass	N <sub>2</sub> O	0.25	0.25	0.0000	1.0000

IPCC Code	IPCC Category	GHG	Base year emissions (Gg CO <sub>2</sub> e)	Absolute value of Base year emissions (Gg CO <sub>2</sub> e)	Level Assessment	Cumulative Total
1A2	Manufacturing industries and construction: other fuels	CH <sub>4</sub>	0.16	0.16	0.0000	1.0000
1B1	Coal mining and handling biomass	CH <sub>4</sub>	0.10	0.10	0.0000	1.0000
4B	Cropland	CH <sub>4</sub>	0.09	0.09	0.0000	1.0000
1B1	Coal mining and handling solid fuels	N <sub>2</sub> O	0.09	0.09	0.0000	1.0000
1B1	Coal mining and handling solid fuels	CH <sub>4</sub>	0.08	0.08	0.0000	1.0000
<b>Total</b>			<b>801,848.54</b>	<b>849,401.28</b>	<b>1.0000</b>	

**Table A 1.3.2 Approach 1 Key Category Analysis for the base year based on level of emissions (excluding LULUCF)**

IPCC Code	IPCC Category	GHG	Base year emissions (Gg CO <sub>2</sub> e)	Absolute value of Base year emissions (Gg CO <sub>2</sub> e)	Level Assessment	Cumulative Total
1A1	Energy industries: solid fuels	CO <sub>2</sub>	185,492.23	185,492.23	0.2314	0.3668
1A3b	Road transportation: liquid fuels	CO <sub>2</sub>	108,570.16	108,570.16	0.1354	0.3668
1A4	Other sectors: gaseous fuels	CO <sub>2</sub>	70,373.19	70,373.19	0.0878	0.4546
5A	Solid waste disposal	CH <sub>4</sub>	60,432.98	60,432.98	0.0754	0.5300
1A1	Energy industries: liquid fuels	CO <sub>2</sub>	40,883.76	40,883.76	0.0510	0.5810
1A2	Manufacturing industries and construction: solid fuels	CO <sub>2</sub>	38,484.20	38,484.20	0.0480	0.6290
1A2	Manufacturing industries and construction: liquid fuels	CO <sub>2</sub>	29,832.00	29,832.00	0.0372	0.6663
1A2	Manufacturing industries and construction: gaseous fuels	CO <sub>2</sub>	27,330.36	27,330.36	0.0341	0.7004
1B1	Coal mining and handling	CH <sub>4</sub>	21,826.68	21,826.68	0.0272	0.7276
2B3	Adipic acid production	N <sub>2</sub> O	19,934.61	19,934.61	0.0249	0.7525
3A1	Enteric fermentation from Cattle	CH <sub>4</sub>	19,881.28	19,881.28	0.0248	0.7773
1A4	Other sectors: solid fuels	CO <sub>2</sub>	19,868.96	19,868.96	0.0248	0.8020
1A4	Other sectors: liquid fuels	CO <sub>2</sub>	19,607.73	19,607.73	0.0245	0.8265
2B9	Fluorochemical production	HFCs, PFCs, SF <sub>6</sub> and NF <sub>3</sub>	17,784.67	17,784.67	0.0222	0.8487

IPCC Code	IPCC Category	GHG	Base year emissions (Gg CO <sub>2</sub> e)	Absolute value of Base year emissions (Gg CO <sub>2</sub> e)	Level Assessment	Cumulative Total
3D	Agricultural soils	N <sub>2</sub> O	13,610.39	13,610.39	0.0170	0.8657
1B2	Oil and gas extraction	CH <sub>4</sub>	12,345.07	12,345.07	0.0154	0.8811
1A1	Energy industries: gaseous fuels	CO <sub>2</sub>	9,229.34	9,229.34	0.0115	0.8926
1A3d	Domestic Navigation: liquid fuels	CO <sub>2</sub>	7,611.13	7,611.13	0.0095	0.9021
2A1	Cement production	CO <sub>2</sub>	7,295.26	7,295.26	0.0091	0.9112
1B2	Oil and gas extraction	CO <sub>2</sub>	5,777.92	5,777.92	0.0072	0.9184
2C1	Iron and steel production	CO <sub>2</sub>	5,591.54	5,591.54	0.0070	0.9254
1A5	Other: liquid fuels	CO <sub>2</sub>	5,293.44	5,293.44	0.0066	0.9320
3A2	Enteric fermentation from Sheep	CH <sub>4</sub>	4,935.90	4,935.90	0.0062	0.9381
3B1	Manure management from Cattle	CH <sub>4</sub>	4,732.53	4,732.53	0.0059	0.9440
2B8	Petrochemical and carbon black production	CO <sub>2</sub>	4,538.91	4,538.91	0.0057	0.9497
5D	Wastewater treatment and discharge	CH <sub>4</sub>	4,218.90	4,218.90	0.0053	0.9550
2B2	Nitric acid production	N <sub>2</sub> O	3,860.26	3,860.26	0.0048	0.9598
3B2	Manure management from Sheep	N <sub>2</sub> O	3,442.70	3,442.70	0.0043	0.9641
2B1	Ammonia production	CO <sub>2</sub>	1,895.00	1,895.00	0.0024	0.9664
1A3a	Domestic aviation: liquid fuels	CO <sub>2</sub>	1,881.31	1,881.31	0.0023	0.9688
1B1	Coal mining and handling solid fuels	CO <sub>2</sub>	1,698.56	1,698.56	0.0021	0.9709
2A2	Lime production	CO <sub>2</sub>	1,462.05	1,462.05	0.0018	0.9727
1A3c	Railways: liquid fuels	CO <sub>2</sub>	1,455.18	1,455.18	0.0018	0.9745
2C6	Zinc production	CO <sub>2</sub>	1,358.83	1,358.83	0.0017	0.9762
1A3b	Road transportation: liquid fuels	N <sub>2</sub> O	1,312.28	1,312.28	0.0016	0.9779
5C	Incineration and open burning of waste	CO <sub>2</sub>	1,300.71	1,300.71	0.0016	0.9795
1A4	Other sectors: solid fuels	CH <sub>4</sub>	1,265.22	1,265.22	0.0016	0.9811
1A3b	Road transportation: liquid fuels	CH <sub>4</sub>	1,246.70	1,246.70	0.0016	0.9826
1A1	Energy industries: solid fuels	N <sub>2</sub> O	1,023.72	1,023.72	0.0013	0.9839
3G	Liming	CO <sub>2</sub>	1,015.18	1,015.18	0.0013	0.9852

IPCC Code	IPCC Category	GHG	Base year emissions (Gg CO <sub>2</sub> e)	Absolute value of Base year emissions (Gg CO <sub>2</sub> e)	Level Assessment	Cumulative Total
5D	Wastewater treatment and discharge	N <sub>2</sub> O	783.59	783.59	0.0010	0.9861
2G1	Electrical equipment	HFCs, PFCs, SF <sub>6</sub> and NF <sub>3</sub>	783.25	783.25	0.0010	0.9871
2F4	Aerosols	HFCs, PFCs, SF <sub>6</sub> and NF <sub>3</sub>	663.00	663.00	0.0008	0.9879
2A4	Other process uses of carbonates	CO <sub>2</sub>	640.93	640.93	0.0008	0.9887
2D	Non-energy products from fuels and solvent use	CO <sub>2</sub>	552.81	552.81	0.0007	0.9894
2G3	N <sub>2</sub> O from product uses	N <sub>2</sub> O	552.57	552.57	0.0007	0.9901
2F1	Refrigeration and air conditioning	HFCs, PFCs, SF <sub>6</sub> and NF <sub>3</sub>	531.30	531.30	0.0007	0.9908
2C3	Aluminium production	CO <sub>2</sub>	450.32	450.32	0.0006	0.9913
2A3	Glass production	CO <sub>2</sub>	405.54	405.54	0.0005	0.9919
2C4	Magnesium production	HFCs, PFCs, SF <sub>6</sub> and NF <sub>3</sub>	387.17	387.17	0.0005	0.9923
1A4	Other sectors: peat	CO <sub>2</sub>	372.48	372.48	0.0005	0.9928
2C3	Aluminium production	HFCs, PFCs, SF <sub>6</sub> and NF <sub>3</sub>	333.43	333.43	0.0004	0.9932
3H	Urea application to land	CO <sub>2</sub>	327.60	327.60	0.0004	0.9936
3A4	Enteric fermentation from Other livestock	CH <sub>4</sub>	292.27	292.27	0.0004	0.9940
3A3	Enteric fermentation from Swine	CH <sub>4</sub>	283.06	283.06	0.0004	0.9943
2G2	SF <sub>6</sub> and PFCs from other product use	HFCs, PFCs, SF <sub>6</sub> and NF <sub>3</sub>	278.70	278.70	0.0003	0.9947

IPCC Code	IPCC Category	GHG	Base year emissions (Gg CO <sub>2</sub> e)	Absolute value of Base year emissions (Gg CO <sub>2</sub> e)	Level Assessment	Cumulative Total
3J	Agriculture activities in OTs and CDs	CH <sub>4</sub>	276.73	276.73	0.0003	0.9950
1A1	Energy industries: other fuels	CO <sub>2</sub>	246.13	246.13	0.0003	0.9953
1A4	Other sectors: solid fuels	N <sub>2</sub> O	241.68	241.68	0.0003	0.9956
2B7	Soda ash production	CO <sub>2</sub>	231.55	231.55	0.0003	0.9959
1A3e	Other transportation: liquid fuels	CO <sub>2</sub>	224.74	224.74	0.0003	0.9962
1A1	Energy industries: gaseous fuels	N <sub>2</sub> O	198.91	198.91	0.0002	0.9965
3F	Field burning of agricultural residues	CH <sub>4</sub>	186.57	186.57	0.0002	0.9967
2F2	Foam blowing agents	HFCs, PFCs, SF <sub>6</sub> and NF <sub>3</sub>	184.49	184.49	0.0002	0.9969
2B10	Other Chemical Industry	CH <sub>4</sub>	174.65	174.65	0.0002	0.9971
1A4	Other sectors: gaseous fuels	CH <sub>4</sub>	157.18	157.18	0.0002	0.9973
1A2	Manufacturing industries and construction: solid fuels	N <sub>2</sub> O	148.72	148.72	0.0002	0.9975
1A1	Energy industries: liquid fuels	N <sub>2</sub> O	140.41	140.41	0.0002	0.9977
1A2	Manufacturing industries and construction: liquid fuels	N <sub>2</sub> O	138.62	138.62	0.0002	0.9979
5C	Incineration and open burning of waste	CH <sub>4</sub>	134.83	134.83	0.0002	0.9980
3J	Agriculture activities in OTs and CDs	N <sub>2</sub> O	132.00	132.00	0.0002	0.9982
1A3d	Domestic Navigation: liquid fuels	N <sub>2</sub> O	105.00	105.00	0.0001	0.9983
2B6	Titanium dioxide production	CO <sub>2</sub>	104.63	104.63	0.0001	0.9985
1A4	Other sectors: liquid fuels	N <sub>2</sub> O	103.32	103.32	0.0001	0.9986
1A1	Energy industries: gaseous fuels	CH <sub>4</sub>	92.30	92.30	0.0001	0.9987
1A2	Manufacturing industries and construction: other fuels	CO <sub>2</sub>	76.86	76.86	0.0001	0.9988
1A4	Other sectors: biomass	CH <sub>4</sub>	62.37	62.37	0.0001	0.9989
1A4	Other sectors: liquid fuels	CH <sub>4</sub>	57.79	57.79	0.0001	0.9990
3F	Field burning of agricultural residues	N <sub>2</sub> O	57.66	57.66	0.0001	0.9990

IPCC Code	IPCC Category	GHG	Base year emissions (Gg CO <sub>2e</sub> )	Absolute value of Base year emissions (Gg CO <sub>2e</sub> )	Level Assessment	Cumulative Total
1A5	Other: liquid fuels	N <sub>2</sub> O	56.12	56.12	0.0001	0.9991
1A1	Energy industries: solid fuels	CH <sub>4</sub>	49.32	49.32	0.0001	0.9992
1A2	Manufacturing industries and construction: solid fuels	CH <sub>4</sub>	47.59	47.59	0.0001	0.9992
5C	Incineration and open burning of waste	N <sub>2</sub> O	46.66	46.66	0.0001	0.9993
1A2	Manufacturing industries and construction: liquid fuels	CH <sub>4</sub>	44.43	44.43	0.0001	0.9993
1A1	Energy industries: liquid fuels	CH <sub>4</sub>	43.36	43.36	0.0001	0.9994
2G4	Other product manufacture and use	N <sub>2</sub> O	41.00	41.00	0.0001	0.9994
1B2	Oil and gas extraction	N <sub>2</sub> O	40.75	40.75	0.0001	0.9995
1A4	Other sectors: gaseous fuels	N <sub>2</sub> O	37.47	37.47	0.0000	0.9995
2C1	Iron and steel production	CH <sub>4</sub>	36.90	36.90	0.0000	0.9996
2F6	Other product uses as substitutes for ODS	HFCs, PFCs, SF <sub>6</sub> and NF <sub>3</sub>	36.88	36.88	0.0000	0.9996
2A4	Other process uses of carbonates	CH <sub>4</sub>	31.10	31.10	0.0000	0.9997
2B8	Petrochemical and carbon black production	CH <sub>4</sub>	29.98	29.98	0.0000	0.9997
1A4	Other sectors: peat	CH <sub>4</sub>	26.35	26.35	0.0000	0.9997
1A2	Manufacturing industries and construction: biomass	N <sub>2</sub> O	19.54	19.54	0.0000	0.9998
1A1	Energy industries: other fuels	CH <sub>4</sub>	18.55	18.55	0.0000	0.9998
5B	Biological treatment of solid waste	CH <sub>4</sub>	18.13	18.13	0.0000	0.9998
1A3a	Domestic aviation: liquid fuels	N <sub>2</sub> O	17.80	17.80	0.0000	0.9998
2C1	Iron and steel production	N <sub>2</sub> O	17.74	17.74	0.0000	0.9999
1A2	Manufacturing industries and construction: gaseous fuels	N <sub>2</sub> O	14.54	14.54	0.0000	0.9999
1A3c	Railways: liquid fuels	N <sub>2</sub> O	13.75	13.75	0.0000	0.9999
5B	Biological treatment of solid waste	N <sub>2</sub> O	12.97	12.97	0.0000	0.9999
1A2	Manufacturing industries and construction: biomass	CH <sub>4</sub>	12.29	12.29	0.0000	0.9999

IPCC Code	IPCC Category	GHG	Base year emissions (Gg CO <sub>2e</sub> )	Absolute value of Base year emissions (Gg CO <sub>2e</sub> )	Level Assessment	Cumulative Total
1A2	Manufacturing industries and construction: gaseous fuels	CH <sub>4</sub>	12.20	12.20	0.0000	0.9999
2E1	Integrated circuit or semiconductor	HFCs, PFCs, SF <sub>6</sub> and NF <sub>3</sub>	9.56	9.56	0.0000	0.9999
1A4	Other sectors: biomass	N <sub>2</sub> O	9.14	9.14	0.0000	1.0000
1A1	Energy industries: other fuels	N <sub>2</sub> O	6.72	6.72	0.0000	1.0000
1A3a	Domestic aviation: liquid fuels	CH <sub>4</sub>	6.37	6.37	0.0000	1.0000
1A3d	Domestic Navigation: liquid fuels	CH <sub>4</sub>	3.66	3.66	0.0000	1.0000
1A5	Other: liquid fuels	CH <sub>4</sub>	3.56	3.56	0.0000	1.0000
1A3e	Other transportation: liquid fuels	N <sub>2</sub> O	2.79	2.79	0.0000	1.0000
1A3c	Railways: liquid fuels	CH <sub>4</sub>	2.46	2.46	0.0000	1.0000
2B8	Petrochemical and carbon black production	N <sub>2</sub> O	2.10	2.10	0.0000	1.0000
1A4	Other sectors: peat	N <sub>2</sub> O	1.47	1.47	0.0000	1.0000
2F3	Fire protection	HFCs, PFCs, SF <sub>6</sub> and NF <sub>3</sub>	1.41	1.41	0.0000	1.0000
1A1	Energy industries: biomass	CH <sub>4</sub>	0.47	0.47	0.0000	1.0000
1A2	Manufacturing industries and construction: other fuels	N <sub>2</sub> O	0.38	0.38	0.0000	1.0000
2B1	Ammonia production	N <sub>2</sub> O	0.31	0.31	0.0000	1.0000
1A3e	Other transportation: liquid fuels	CH <sub>4</sub>	0.29	0.29	0.0000	1.0000
2B1	Ammonia production	CH <sub>4</sub>	0.26	0.26	0.0000	1.0000
1A1	Energy industries: biomass	N <sub>2</sub> O	0.25	0.25	0.0000	1.0000
1A2	Manufacturing industries and construction: other fuels	CH <sub>4</sub>	0.16	0.16	0.0000	1.0000
1B1	Coal mining and handling biomass	CH <sub>4</sub>	0.10	0.10	0.0000	1.0000
1B1	Coal mining and handling solid fuels	N <sub>2</sub> O	0.09	0.09	0.0000	1.0000
1B1	Coal mining and handling solid fuels	CH <sub>4</sub>	0.08	0.08	0.0000	1.0000

IPCC Code	IPCC Category	GHG	Base year emissions (Gg CO <sub>2</sub> e)	Absolute value of Base year emissions (Gg CO <sub>2</sub> e)	Level Assessment	Cumulative Total
<b>Total</b>			<b>801,593.08</b>	<b>801,593.08</b>	<b>1.0000</b>	

**Table A 1.3.3 Approach 1 Key Category Analysis for the latest reported year based on level of emissions (including LULUCF)**

IPCC Code	IPCC Category	GHG	2017 emissions (Gg CO <sub>2</sub> e)	Absolute value of 2017 emissions (Gg CO <sub>2</sub> e)	Level Assessment	Cumulative Total
1A3b	Road transportation: liquid fuels	CO <sub>2</sub>	113,364.64	113,364.64	0.2169	0.2169
1A4	Other sectors: gaseous fuels	CO <sub>2</sub>	73,421.79	73,421.79	0.1405	0.3574
1A1	Energy industries: gaseous fuels	CO <sub>2</sub>	61,476.35	61,476.35	0.1176	0.4750
1A2	Manufacturing industries and construction: gaseous fuels	CO <sub>2</sub>	23,723.43	23,723.43	0.0454	0.5204
1A1	Energy industries: solid fuels	CO <sub>2</sub>	20,363.72	20,363.72	0.0390	0.5594
4A	Forest land	CO <sub>2</sub>	-18,211.77	18,211.77	0.0348	0.5942
3A1	Enteric fermentation from Cattle	CH <sub>4</sub>	16,770.98	16,770.98	0.0321	0.6263
1A1	Energy industries: liquid fuels	CO <sub>2</sub>	15,767.40	15,767.40	0.0302	0.6565
1A2	Manufacturing industries and construction: liquid fuels	CO <sub>2</sub>	14,676.48	14,676.48	0.0281	0.6846
5A	Solid waste disposal	CH <sub>4</sub>	14,261.40	14,261.40	0.0273	0.7119
1A4	Other sectors: liquid fuels	CO <sub>2</sub>	13,284.20	13,284.20	0.0254	0.7373
1A2	Manufacturing industries and construction: solid fuels	CO <sub>2</sub>	12,133.91	12,133.91	0.0232	0.7605
2F1	Refrigeration and air conditioning	HFCs, PFCs, SF <sub>6</sub> and NF <sub>3</sub>	11,653.11	11,653.11	0.0223	0.7828
3D	Agricultural soils	N <sub>2</sub> O	11,466.79	11,466.79	0.0219	0.8047
4B	Cropland	CO <sub>2</sub>	10,971.23	10,971.23	0.0210	0.8257
4C	Grassland	CO <sub>2</sub>	-8,861.80	8,861.80	0.0170	0.8427
4E	Settlements	CO <sub>2</sub>	6,451.34	6,451.34	0.0123	0.8550
1A3d	Domestic Navigation: liquid fuels	CO <sub>2</sub>	5,443.73	5,443.73	0.0104	0.8654
1A1	Energy industries: other fuels	CO <sub>2</sub>	5,021.63	5,021.63	0.0096	0.8750
1B2	Oil and gas extraction	CH <sub>4</sub>	4,935.25	4,935.25	0.0094	0.8845

IPCC Code	IPCC Category	GHG	2017 emissions (Gg CO <sub>2</sub> e)	Absolute value of 2017 emissions (Gg CO <sub>2</sub> e)	Level Assessment	Cumulative Total
2A1	Cement production	CO <sub>2</sub>	4,409.79	4,409.79	0.0084	0.8929
1B2	Oil and gas extraction	CO <sub>2</sub>	4,230.64	4,230.64	0.0081	0.9010
3B1	Manure management from Cattle	CH <sub>4</sub>	4,226.81	4,226.81	0.0081	0.9091
3A2	Enteric fermentation from Sheep	CH <sub>4</sub>	4,043.20	4,043.20	0.0077	0.9168
5D	Wastewater treatment and discharge	CH <sub>4</sub>	3,445.75	3,445.75	0.0066	0.9234
2B8	Petrochemical and carbon black production	CO <sub>2</sub>	2,846.51	2,846.51	0.0054	0.9289
3B2	Manure management from Sheep	N <sub>2</sub> O	2,814.79	2,814.79	0.0054	0.9343
2C1	Iron and steel production	CO <sub>2</sub>	2,501.24	2,501.24	0.0048	0.9391
1A4	Other sectors: solid fuels	CO <sub>2</sub>	2,276.82	2,276.82	0.0044	0.9434
4G	Harvested wood products	CO <sub>2</sub>	-2,015.60	2,015.60	0.0039	0.9473
1A3c	Railways: liquid fuels	CO <sub>2</sub>	1,939.08	1,939.08	0.0037	0.9510
1A3a	Domestic aviation: liquid fuels	CO <sub>2</sub>	1,898.39	1,898.39	0.0036	0.9546
2B1	Ammonia production	CO <sub>2</sub>	1,763.86	1,763.86	0.0034	0.9580
2F4	Aerosols	HFCs, PFCs, SF <sub>6</sub> and NF <sub>3</sub>	1,658.10	1,658.10	0.0032	0.9612
1A5	Other: liquid fuels	CO <sub>2</sub>	1,558.20	1,558.20	0.0030	0.9641
5B	Biological treatment of solid waste	CH <sub>4</sub>	1,193.62	1,193.62	0.0023	0.9664
1A3b	Road transportation: liquid fuels	N <sub>2</sub> O	1,095.65	1,095.65	0.0021	0.9685
2A2	Lime production	CO <sub>2</sub>	1,052.06	1,052.06	0.0020	0.9705
3G	Liming	CO <sub>2</sub>	939.13	939.13	0.0018	0.9723
5B	Biological treatment of solid waste	N <sub>2</sub> O	728.03	728.03	0.0014	0.9737
5D	Wastewater treatment and discharge	N <sub>2</sub> O	725.93	725.93	0.0014	0.9751
2G3	N <sub>2</sub> O from product uses	N <sub>2</sub> O	645.41	645.41	0.0012	0.9763
1A2	Manufacturing industries and construction: other fuels	CO <sub>2</sub>	641.74	641.74	0.0012	0.9776
1A4	Other sectors: biomass	CH <sub>4</sub>	601.47	601.47	0.0012	0.9787

IPCC Code	IPCC Category	GHG	2017 emissions (Gg CO <sub>2</sub> e)	Absolute value of 2017 emissions (Gg CO <sub>2</sub> e)	Level Assessment	Cumulative Total
1A3e	Other transportation: liquid fuels	CO <sub>2</sub>	565.53	565.53	0.0011	0.9798
4E	Settlements	N <sub>2</sub> O	524.99	524.99	0.0010	0.9808
2D	Non-energy products from fuels and solvent use	CO <sub>2</sub>	519.22	519.22	0.0010	0.9818
1B1	Coal mining and handling	CH <sub>4</sub>	483.98	483.98	0.0009	0.9827
4B	Cropland	N <sub>2</sub> O	461.97	461.97	0.0009	0.9836
3A4	Enteric fermentation from Other livestock	CH <sub>4</sub>	457.93	457.93	0.0009	0.9845
2F2	Foam blowing agents	HFCs, PFCs, SF <sub>6</sub> and NF <sub>3</sub>	450.52	450.52	0.0009	0.9854
2A4	Other process uses of carbonates	CO <sub>2</sub>	419.65	419.65	0.0008	0.9862
2A3	Glass production	CO <sub>2</sub>	367.98	367.98	0.0007	0.9869
3H	Urea application to land	CO <sub>2</sub>	343.95	343.95	0.0007	0.9875
4D	Wetlands	CO <sub>2</sub>	336.78	336.78	0.0006	0.9882
2F3	Fire protection	HFCs, PFCs, SF <sub>6</sub> and NF <sub>3</sub>	324.69	324.69	0.0006	0.9888
2G2	SF <sub>6</sub> and PFCs from other product use	HFCs, PFCs, SF <sub>6</sub> and NF <sub>3</sub>	318.00	318.00	0.0006	0.9894
1A1	Energy industries: gaseous fuels	N <sub>2</sub> O	289.24	289.24	0.0006	0.9899
2G1	Electrical equipment	HFCs, PFCs, SF <sub>6</sub> and NF <sub>3</sub>	281.17	281.17	0.0005	0.9905
1B1	Coal mining and handling solid fuels	CO <sub>2</sub>	268.38	268.38	0.0005	0.9910
5C	Incineration and open burning of waste	CO <sub>2</sub>	258.91	258.91	0.0005	0.9915
4	Indirect N <sub>2</sub> O emissions from LULUCF	N <sub>2</sub> O	243.62	243.62	0.0005	0.9920

IPCC Code	IPCC Category	GHG	2017 emissions (Gg CO <sub>2</sub> e)	Absolute value of 2017 emissions (Gg CO <sub>2</sub> e)	Level Assessment	Cumulative Total
3J	Agriculture activities in OTs and CDs	CH <sub>4</sub>	206.19	206.19	0.0004	0.9924
2G4	Other product manufacture and use	N <sub>2</sub> O	198.47	198.47	0.0004	0.9927
3A3	Enteric fermentation from Swine	CH <sub>4</sub>	186.33	186.33	0.0004	0.9931
2B6	Titanium dioxide production	CO <sub>2</sub>	181.02	181.02	0.0003	0.9934
2B9	Fluorochemical production	HFCs, PFCs, SF <sub>6</sub> and NF <sub>3</sub>	175.44	175.44	0.0003	0.9938
1A4	Other sectors: solid fuels	CH <sub>4</sub>	173.62	173.62	0.0003	0.9941
1A1	Energy industries: biomass	N <sub>2</sub> O	171.90	171.90	0.0003	0.9944
1A4	Other sectors: gaseous fuels	CH <sub>4</sub>	161.99	161.99	0.0003	0.9947
4A	Forest land	N <sub>2</sub> O	144.65	144.65	0.0003	0.9950
2B7	Soda ash production	CO <sub>2</sub>	140.18	140.18	0.0003	0.9953
1A1	Energy industries: gaseous fuels	CH <sub>4</sub>	137.70	137.70	0.0003	0.9956
1A1	Energy industries: solid fuels	N <sub>2</sub> O	115.89	115.89	0.0002	0.9958
2C4	Magnesium production	HFCs, PFCs, SF <sub>6</sub> and NF <sub>3</sub>	112.04	112.04	0.0002	0.9960
1A1	Energy industries: other fuels	CH <sub>4</sub>	110.63	110.63	0.0002	0.9962
1A1	Energy industries: biomass	CH <sub>4</sub>	109.01	109.01	0.0002	0.9964
1A2	Manufacturing industries and construction: biomass	N <sub>2</sub> O	102.56	102.56	0.0002	0.9966
1A4	Other sectors: biomass	N <sub>2</sub> O	94.30	94.30	0.0002	0.9968
1A2	Manufacturing industries and construction: liquid fuels	N <sub>2</sub> O	93.64	93.64	0.0002	0.9970
1A1	Energy industries: other fuels	N <sub>2</sub> O	93.36	93.36	0.0002	0.9971
1A3b	Road transportation: liquid fuels	CH <sub>4</sub>	93.16	93.16	0.0002	0.9973
3J	Agriculture activities in OTs and CDs	N <sub>2</sub> O	93.01	93.01	0.0002	0.9975
1A3b	Road transportation: biomass	CO <sub>2</sub>	92.80	92.80	0.0002	0.9977

IPCC Code	IPCC Category	GHG	2017 emissions (Gg CO <sub>2e</sub> )	Absolute value of 2017 emissions (Gg CO <sub>2e</sub> )	Level Assessment	Cumulative Total
1B1	Coal mining and handling liquid fuels	CO <sub>2</sub>	91.16	91.16	0.0002	0.9979
1A4	Other sectors: liquid fuels	N <sub>2</sub> O	82.03	82.03	0.0002	0.9980
1A1	Energy industries: liquid fuels	N <sub>2</sub> O	80.97	80.97	0.0002	0.9982
1A3d	Domestic Navigation: liquid fuels	N <sub>2</sub> O	72.08	72.08	0.0001	0.9983
2C3	Aluminium production	CO <sub>2</sub>	71.62	71.62	0.0001	0.9984
1A2	Manufacturing industries and construction: biomass	CH <sub>4</sub>	64.53	64.53	0.0001	0.9986
2F6	Other product uses as substitutes for ODS	HFCs, PFCs, SF <sub>6</sub> and NF <sub>3</sub>	61.71	61.71	0.0001	0.9987
2B10	Other Chemical Industry	CH <sub>4</sub>	48.97	48.97	0.0001	0.9988
1A3c	Railways: solid fuels	CO <sub>2</sub>	41.15	41.15	0.0001	0.9989
5C	Incineration and open burning of waste	N <sub>2</sub> O	39.63	39.63	0.0001	0.9989
1B2	Oil and gas extraction	N <sub>2</sub> O	39.36	39.36	0.0001	0.9990
1A4	Other sectors: gaseous fuels	N <sub>2</sub> O	38.62	38.62	0.0001	0.9991
2B2	Nitric acid production	N <sub>2</sub> O	37.25	37.25	0.0001	0.9991
4C	Grassland	N <sub>2</sub> O	33.24	33.24	0.0001	0.9992
1A4	Other sectors: liquid fuels	CH <sub>4</sub>	32.75	32.75	0.0001	0.9993
1A2	Manufacturing industries and construction: liquid fuels	CH <sub>4</sub>	32.04	32.04	0.0001	0.9993
1A2	Manufacturing industries and construction: solid fuels	N <sub>2</sub> O	30.91	30.91	0.0001	0.9994
1A4	Other sectors: solid fuels	N <sub>2</sub> O	30.40	30.40	0.0001	0.9995
2E1	Integrated circuit or semiconductor	HFCs, PFCs, SF <sub>6</sub> and NF <sub>3</sub>	21.71	21.71	0.0000	0.9995
2F5	Solvents	HFCs, PFCs, SF <sub>6</sub> and NF <sub>3</sub>	18.67	18.67	0.0000	0.9995
1A3c	Railways: liquid fuels	N <sub>2</sub> O	18.30	18.30	0.0000	0.9996
1A1	Energy industries: liquid fuels	CH <sub>4</sub>	18.08	18.08	0.0000	0.9996

IPCC Code	IPCC Category	GHG	2017 emissions (Gg CO <sub>2e</sub> )	Absolute value of 2017 emissions (Gg CO <sub>2e</sub> )	Level Assessment	Cumulative Total
1A3a	Domestic aviation: liquid fuels	N <sub>2</sub> O	17.96	17.96	0.0000	0.9996
4C	Grassland	CH <sub>4</sub>	17.78	17.78	0.0000	0.9997
1A5	Other: liquid fuels	N <sub>2</sub> O	16.48	16.48	0.0000	0.9997
2C3	Aluminium production	HFCs, PFCs, SF <sub>6</sub> and NF <sub>3</sub>	15.07	15.07	0.0000	0.9997
1A2	Manufacturing industries and construction: gaseous fuels	N <sub>2</sub> O	12.48	12.48	0.0000	0.9998
2B8	Petrochemical and carbon black production	CH <sub>4</sub>	12.16	12.16	0.0000	0.9998
1A2	Manufacturing industries and construction: solid fuels	CH <sub>4</sub>	10.86	10.86	0.0000	0.9998
2C1	Iron and steel production	CH <sub>4</sub>	10.68	10.68	0.0000	0.9998
1A2	Manufacturing industries and construction: gaseous fuels	CH <sub>4</sub>	10.47	10.47	0.0000	0.9998
5C	Incineration and open burning of waste	CH <sub>4</sub>	10.45	10.45	0.0000	0.9999
1A2	Manufacturing industries and construction: other fuels	N <sub>2</sub> O	10.05	10.05	0.0000	0.9999
4E	Settlements	CH <sub>4</sub>	8.25	8.25	0.0000	0.9999
1A3e	Other transportation: liquid fuels	N <sub>2</sub> O	7.00	7.00	0.0000	0.9999
1A3d	Domestic Navigation: liquid fuels	CH <sub>4</sub>	6.28	6.28	0.0000	0.9999
2C1	Iron and steel production	N <sub>2</sub> O	6.19	6.19	0.0000	0.9999
1A2	Manufacturing industries and construction: other fuels	CH <sub>4</sub>	6.12	6.12	0.0000	0.9999
2A4	Other process uses of carbonates	CH <sub>4</sub>	5.89	5.89	0.0000	1.0000
1A1	Energy industries: solid fuels	CH <sub>4</sub>	5.69	5.69	0.0000	1.0000
1A4	Other sectors: peat	CO <sub>2</sub>	4.83	4.83	0.0000	1.0000
1B1	Coal mining and handling biomass	CH <sub>4</sub>	3.76	3.76	0.0000	1.0000
4A	Forest land	CH <sub>4</sub>	2.31	2.31	0.0000	1.0000
1A3c	Railways: liquid fuels	CH <sub>4</sub>	1.92	1.92	0.0000	1.0000
1A3a	Domestic aviation: liquid fuels	CH <sub>4</sub>	1.55	1.55	0.0000	1.0000

IPCC Code	IPCC Category	GHG	2017 emissions (Gg CO <sub>2</sub> e)	Absolute value of 2017 emissions (Gg CO <sub>2</sub> e)	Level Assessment	Cumulative Total
2B8	Petrochemical and carbon black production	N <sub>2</sub> O	1.51	1.51	0.0000	1.0000
1A5	Other: liquid fuels	CH <sub>4</sub>	1.03	1.03	0.0000	1.0000
1A3c	Railways: solid fuels	CH <sub>4</sub>	0.94	0.94	0.0000	1.0000
1A3e	Other transportation: liquid fuels	CH <sub>4</sub>	0.35	0.35	0.0000	1.0000
1A4	Other sectors: peat	CH <sub>4</sub>	0.34	0.34	0.0000	1.0000
2B1	Ammonia production	N <sub>2</sub> O	0.33	0.33	0.0000	1.0000
4D	Wetlands	N <sub>2</sub> O	0.30	0.30	0.0000	1.0000
2B1	Ammonia production	CH <sub>4</sub>	0.28	0.28	0.0000	1.0000
1A3c	Railways: solid fuels	N <sub>2</sub> O	0.09	0.09	0.0000	1.0000
4B	Cropland	CH <sub>4</sub>	0.02	0.02	0.0000	1.0000
1A4	Other sectors: peat	N <sub>2</sub> O	0.02	0.02	0.0000	1.0000
1B1	Coal mining and handling solid fuels	N <sub>2</sub> O	0.02	0.02	0.0000	1.0000
1B1	Coal mining and handling solid fuels	CH <sub>4</sub>	0.01	0.01	0.0000	1.0000
<b>Total</b>			<b>464,453.44</b>	<b>522,631.79</b>	<b>1.0000</b>	

**Table A 1.3.4 Approach 1 Key Category Analysis for the latest reported year based on level of emissions (excluding LULUCF)**

IPCC Code	IPCC Category	GHG	2017 emissions (Gg CO <sub>2</sub> e)	Absolute value of 2017 emissions (Gg CO <sub>2</sub> e)	Level Assessment	Cumulative Total
1A3b	Road transportation: liquid fuels	CO <sub>2</sub>	113,364.64	113,364.64	0.2390	0.2390
1A4	Other sectors: gaseous fuels	CO <sub>2</sub>	73,421.79	73,421.79	0.1548	0.3938
1A1	Energy industries: gaseous fuels	CO <sub>2</sub>	61,476.35	61,476.35	0.1296	0.5234
1A2	Manufacturing industries and construction: gaseous fuels	CO <sub>2</sub>	23,723.43	23,723.43	0.0500	0.5734
1A1	Energy industries: solid fuels	CO <sub>2</sub>	20,363.72	20,363.72	0.0429	0.6163
3A1	Enteric fermentation from Cattle	CH <sub>4</sub>	16,770.98	16,770.98	0.0354	0.6517
1A1	Energy industries: liquid fuels	CO <sub>2</sub>	15,767.40	15,767.40	0.0332	0.6849
1A2	Manufacturing industries and construction: liquid fuels	CO <sub>2</sub>	14,676.48	14,676.48	0.0309	0.7159

IPCC Code	IPCC Category	GHG	2017 emissions (Gg CO <sub>2</sub> e)	Absolute value of 2017 emissions (Gg CO <sub>2</sub> e)	Level Assessment	Cumulative Total
5A	Solid waste disposal	CH <sub>4</sub>	14,261.40	14,261.40	0.0301	0.7459
1A4	Other sectors: liquid fuels	CO <sub>2</sub>	13,284.20	13,284.20	0.0280	0.7739
1A2	Manufacturing industries and construction: solid fuels	CO <sub>2</sub>	12,133.91	12,133.91	0.0256	0.7995
2F1	Refrigeration and air conditioning	HFCs, PFCs, SF <sub>6</sub> and NF <sub>3</sub>	11,653.11	11,653.11	0.0246	0.8241
3D	Agricultural soils	N <sub>2</sub> O	11,466.79	11,466.79	0.0242	0.8483
1A3d	Domestic Navigation: liquid fuels	CO <sub>2</sub>	5,443.73	5,443.73	0.0115	0.8597
1A1	Energy industries: other fuels	CO <sub>2</sub>	5,021.63	5,021.63	0.0106	0.8703
1B2	Oil and gas extraction	CH <sub>4</sub>	4,935.25	4,935.25	0.0104	0.8807
2A1	Cement production	CO <sub>2</sub>	4,409.79	4,409.79	0.0093	0.8900
1B2	Oil and gas extraction	CO <sub>2</sub>	4,230.64	4,230.64	0.0089	0.8989
3B1	Manure management from Cattle	CH <sub>4</sub>	4,226.81	4,226.81	0.0089	0.9078
3A2	Enteric fermentation from Sheep	CH <sub>4</sub>	4,043.20	4,043.20	0.0085	0.9164
5D	Wastewater treatment and discharge	CH <sub>4</sub>	3,445.75	3,445.75	0.0073	0.9236
2B8	Petrochemical and carbon black production	CO <sub>2</sub>	2,846.51	2,846.51	0.0060	0.9296
3B2	Manure management from Sheep	N <sub>2</sub> O	2,814.79	2,814.79	0.0059	0.9356
2C1	Iron and steel production	CO <sub>2</sub>	2,501.24	2,501.24	0.0053	0.9408
1A4	Other sectors: solid fuels	CO <sub>2</sub>	2,276.82	2,276.82	0.0048	0.9456
1A3c	Railways: liquid fuels	CO <sub>2</sub>	1,939.08	1,939.08	0.0041	0.9497
1A3a	Domestic aviation: liquid fuels	CO <sub>2</sub>	1,898.39	1,898.39	0.0040	0.9537
2B1	Ammonia production	CO <sub>2</sub>	1,763.86	1,763.86	0.0037	0.9574
2F4	Aerosols	HFCs, PFCs, SF <sub>6</sub> and NF <sub>3</sub>	1,658.10	1,658.10	0.0035	0.9609
1A5	Other: liquid fuels	CO <sub>2</sub>	1,558.20	1,558.20	0.0033	0.9642
5B	Biological treatment of solid waste	CH <sub>4</sub>	1,193.62	1,193.62	0.0025	0.9667

IPCC Code	IPCC Category	GHG	2017 emissions (Gg CO <sub>2</sub> e)	Absolute value of 2017 emissions (Gg CO <sub>2</sub> e)	Level Assessment	Cumulative Total
1A3b	Road transportation: liquid fuels	N <sub>2</sub> O	1,095.65	1,095.65	0.0023	0.9691
2A2	Lime production	CO <sub>2</sub>	1,052.06	1,052.06	0.0022	0.9713
3G	Liming	CO <sub>2</sub>	939.13	939.13	0.0020	0.9733
5B	Biological treatment of solid waste	N <sub>2</sub> O	728.03	728.03	0.0015	0.9748
5D	Wastewater treatment and discharge	N <sub>2</sub> O	725.93	725.93	0.0015	0.9763
2G3	N <sub>2</sub> O from product uses	N <sub>2</sub> O	645.41	645.41	0.0014	0.9777
1A2	Manufacturing industries and construction: other fuels	CO <sub>2</sub>	641.74	641.74	0.0014	0.9790
1A4	Other sectors: biomass	CH <sub>4</sub>	601.47	601.47	0.0013	0.9803
1A3e	Other transportation: liquid fuels	CO <sub>2</sub>	565.53	565.53	0.0012	0.9815
2D	Non-energy products from fuels and solvent use	CO <sub>2</sub>	519.22	519.22	0.0011	0.9826
1B1	Coal mining and handling	CH <sub>4</sub>	483.98	483.98	0.0010	0.9836
3A4	Enteric fermentation from Other livestock	CH <sub>4</sub>	457.93	457.93	0.0010	0.9846
2F2	Foam blowing agents	HFCs, PFCs, SF <sub>6</sub> and NF <sub>3</sub>	450.52	450.52	0.0009	0.9855
2A4	Other process uses of carbonates	CO <sub>2</sub>	419.65	419.65	0.0009	0.9864
2A3	Glass production	CO <sub>2</sub>	367.98	367.98	0.0008	0.9872
3H	Urea application to land	CO <sub>2</sub>	343.95	343.95	0.0007	0.9879
2F3	Fire protection	HFCs, PFCs, SF <sub>6</sub> and NF <sub>3</sub>	324.69	324.69	0.0007	0.9886
2G2	SF <sub>6</sub> and PFCs from other product use	HFCs, PFCs, SF <sub>6</sub> and NF <sub>3</sub>	318.00	318.00	0.0007	0.9893
1A1	Energy industries: gaseous fuels	N <sub>2</sub> O	289.24	289.24	0.0006	0.9899

IPCC Code	IPCC Category	GHG	2017 emissions (Gg CO <sub>2</sub> e)	Absolute value of 2017 emissions (Gg CO <sub>2</sub> e)	Level Assessment	Cumulative Total
2G1	Electrical equipment	HFCs, PFCs, SF <sub>6</sub> and NF <sub>3</sub>	281.17	281.17	0.0006	0.9905
1B1	Coal mining and handling solid fuels	CO <sub>2</sub>	268.38	268.38	0.0006	0.9910
5C	Incineration and open burning of waste	CO <sub>2</sub>	258.91	258.91	0.0005	0.9916
3J	Agriculture activities in OTs and CDs	CH <sub>4</sub>	206.19	206.19	0.0004	0.9920
2G4	Other product manufacture and use	N <sub>2</sub> O	198.47	198.47	0.0004	0.9924
3A3	Enteric fermentation from Swine	CH <sub>4</sub>	186.33	186.33	0.0004	0.9928
2B6	Titanium dioxide production	CO <sub>2</sub>	181.02	181.02	0.0004	0.9932
2B9	Fluorochemical production	HFCs, PFCs, SF <sub>6</sub> and NF <sub>3</sub>	175.44	175.44	0.0004	0.9936
1A4	Other sectors: solid fuels	CH <sub>4</sub>	173.62	173.62	0.0004	0.9939
1A1	Energy industries: biomass	N <sub>2</sub> O	171.90	171.90	0.0004	0.9943
1A4	Other sectors: gaseous fuels	CH <sub>4</sub>	161.99	161.99	0.0003	0.9946
2B7	Soda ash production	CO <sub>2</sub>	140.18	140.18	0.0003	0.9949
1A1	Energy industries: gaseous fuels	CH <sub>4</sub>	137.70	137.70	0.0003	0.9952
1A1	Energy industries: solid fuels	N <sub>2</sub> O	115.89	115.89	0.0002	0.9955
2C4	Magnesium production	HFCs, PFCs, SF <sub>6</sub> and NF <sub>3</sub>	112.04	112.04	0.0002	0.9957
1A1	Energy industries: other fuels	CH <sub>4</sub>	110.63	110.63	0.0002	0.9959
1A1	Energy industries: biomass	CH <sub>4</sub>	109.01	109.01	0.0002	0.9962
1A2	Manufacturing industries and construction: biomass	N <sub>2</sub> O	102.56	102.56	0.0002	0.9964
1A4	Other sectors: biomass	N <sub>2</sub> O	94.30	94.30	0.0002	0.9966
1A2	Manufacturing industries and construction: liquid fuels	N <sub>2</sub> O	93.64	93.64	0.0002	0.9968
1A1	Energy industries: other fuels	N <sub>2</sub> O	93.36	93.36	0.0002	0.9970

IPCC Code	IPCC Category	GHG	2017 emissions (Gg CO <sub>2</sub> e)	Absolute value of 2017 emissions (Gg CO <sub>2</sub> e)	Level Assessment	Cumulative Total
1A3b	Road transportation: liquid fuels	CH <sub>4</sub>	93.16	93.16	0.0002	0.9972
3J	Agriculture activities in OTs and CDs	N <sub>2</sub> O	93.01	93.01	0.0002	0.9974
1A3b	Road transportation: biomass	CO <sub>2</sub>	92.80	92.80	0.0002	0.9976
1B1	Coal mining and handling liquid fuels	CO <sub>2</sub>	91.16	91.16	0.0002	0.9978
1A4	Other sectors: liquid fuels	N <sub>2</sub> O	82.03	82.03	0.0002	0.9979
1A1	Energy industries: liquid fuels	N <sub>2</sub> O	80.97	80.97	0.0002	0.9981
1A3d	Domestic Navigation: liquid fuels	N <sub>2</sub> O	72.08	72.08	0.0002	0.9983
2C3	Aluminium production	CO <sub>2</sub>	71.62	71.62	0.0002	0.9984
1A2	Manufacturing industries and construction: biomass	CH <sub>4</sub>	64.53	64.53	0.0001	0.9985
2F6	Other product uses as substitutes for ODS	HFCs, PFCs, SF <sub>6</sub> and NF <sub>3</sub>	61.71	61.71	0.0001	0.9987
2B10	Other Chemical Industry	CH <sub>4</sub>	48.97	48.97	0.0001	0.9988
1A3c	Railways: solid fuels	CO <sub>2</sub>	41.15	41.15	0.0001	0.9989
5C	Incineration and open burning of waste	N <sub>2</sub> O	39.63	39.63	0.0001	0.9989
1B2	Oil and gas extraction	N <sub>2</sub> O	39.36	39.36	0.0001	0.9990
1A4	Other sectors: gaseous fuels	N <sub>2</sub> O	38.62	38.62	0.0001	0.9991
2B2	Nitric acid production	N <sub>2</sub> O	37.25	37.25	0.0001	0.9992
1A4	Other sectors: liquid fuels	CH <sub>4</sub>	32.75	32.75	0.0001	0.9993
1A2	Manufacturing industries and construction: liquid fuels	CH <sub>4</sub>	32.04	32.04	0.0001	0.9993
1A2	Manufacturing industries and construction: solid fuels	N <sub>2</sub> O	30.91	30.91	0.0001	0.9994
1A4	Other sectors: solid fuels	N <sub>2</sub> O	30.40	30.40	0.0001	0.9995
2E1	Integrated circuit or semiconductor	HFCs, PFCs, SF <sub>6</sub> and NF <sub>3</sub>	21.71	21.71	0.0000	0.9995

IPCC Code	IPCC Category	GHG	2017 emissions (Gg CO <sub>2</sub> e)	Absolute value of 2017 emissions (Gg CO <sub>2</sub> e)	Level Assessment	Cumulative Total
2F5	Solvents	HFCs, PFCs, SF <sub>6</sub> and NF <sub>3</sub>	18.67	18.67	0.0000	0.9995
1A3c	Railways: liquid fuels	N <sub>2</sub> O	18.30	18.30	0.0000	0.9996
1A1	Energy industries: liquid fuels	CH <sub>4</sub>	18.08	18.08	0.0000	0.9996
1A3a	Domestic aviation: liquid fuels	N <sub>2</sub> O	17.96	17.96	0.0000	0.9997
1A5	Other: liquid fuels	N <sub>2</sub> O	16.48	16.48	0.0000	0.9997
2C3	Aluminium production	HFCs, PFCs, SF <sub>6</sub> and NF <sub>3</sub>	15.07	15.07	0.0000	0.9997
1A2	Manufacturing industries and construction: gaseous fuels	N <sub>2</sub> O	12.48	12.48	0.0000	0.9997
2B8	Petrochemical and carbon black production	CH <sub>4</sub>	12.16	12.16	0.0000	0.9998
1A2	Manufacturing industries and construction: solid fuels	CH <sub>4</sub>	10.86	10.86	0.0000	0.9998
2C1	Iron and steel production	CH <sub>4</sub>	10.68	10.68	0.0000	0.9998
1A2	Manufacturing industries and construction: gaseous fuels	CH <sub>4</sub>	10.47	10.47	0.0000	0.9998
5C	Incineration and open burning of waste	CH <sub>4</sub>	10.45	10.45	0.0000	0.9999
1A2	Manufacturing industries and construction: other fuels	N <sub>2</sub> O	10.05	10.05	0.0000	0.9999
1A3e	Other transportation: liquid fuels	N <sub>2</sub> O	7.00	7.00	0.0000	0.9999
1A3d	Domestic Navigation: liquid fuels	CH <sub>4</sub>	6.28	6.28	0.0000	0.9999
2C1	Iron and steel production	N <sub>2</sub> O	6.19	6.19	0.0000	0.9999
1A2	Manufacturing industries and construction: other fuels	CH <sub>4</sub>	6.12	6.12	0.0000	0.9999
2A4	Other process uses of carbonates	CH <sub>4</sub>	5.89	5.89	0.0000	1.0000
1A1	Energy industries: solid fuels	CH <sub>4</sub>	5.69	5.69	0.0000	1.0000
1A4	Other sectors: peat	CO <sub>2</sub>	4.83	4.83	0.0000	1.0000
1B1	Coal mining and handling biomass	CH <sub>4</sub>	3.76	3.76	0.0000	1.0000

IPCC Code	IPCC Category	GHG	2017 emissions (Gg CO <sub>2</sub> e)	Absolute value of 2017 emissions (Gg CO <sub>2</sub> e)	Level Assessment	Cumulative Total
1A3c	Railways: liquid fuels	CH <sub>4</sub>	1.92	1.92	0.0000	1.0000
1A3a	Domestic aviation: liquid fuels	CH <sub>4</sub>	1.55	1.55	0.0000	1.0000
2B8	Petrochemical and carbon black production	N <sub>2</sub> O	1.51	1.51	0.0000	1.0000
1A5	Other: liquid fuels	CH <sub>4</sub>	1.03	1.03	0.0000	1.0000
1A3c	Railways: solid fuels	CH <sub>4</sub>	0.94	0.94	0.0000	1.0000
1A3e	Other transportation: liquid fuels	CH <sub>4</sub>	0.35	0.35	0.0000	1.0000
1A4	Other sectors: peat	CH <sub>4</sub>	0.34	0.34	0.0000	1.0000
2B1	Ammonia production	N <sub>2</sub> O	0.33	0.33	0.0000	1.0000
2B1	Ammonia production	CH <sub>4</sub>	0.28	0.28	0.0000	1.0000
1A3c	Railways: solid fuels	N <sub>2</sub> O	0.09	0.09	0.0000	1.0000
1A4	Other sectors: peat	N <sub>2</sub> O	0.02	0.02	0.0000	1.0000
1B1	Coal mining and handling solid fuels	N <sub>2</sub> O	0.02	0.02	0.0000	1.0000
1B1	Coal mining and handling solid fuels	CH <sub>4</sub>	0.01	0.01	0.0000	1.0000
<b>Total</b>			<b>474,346.12</b>	<b>474,346.12</b>	<b>1.0000</b>	

**Table A 1.3.5 Approach 1 Key Category Analysis based on trend in emissions (from base year to latest reported year, including LULUCF)**

IPCC Code	IPCC Category	GHG	Base year emissions (Gg CO <sub>2</sub> e)	2017 emissions (Gg CO <sub>2</sub> e)	Trend Assessment	Contribution to Trend	Cumulative Total
1A1	Energy industries: solid fuels	CO <sub>2</sub>	185,492.23	20,363.72	0.1025	0.2251	0.2251
1A1	Energy industries: gaseous fuels	CO <sub>2</sub>	9,229.34	61,476.35	0.0661	0.1451	0.3702
1A3b	Road transportation: liquid fuels	CO <sub>2</sub>	108,570.16	113,364.64	0.0594	0.1305	0.5007
1A4	Other sectors: gaseous fuels	CO <sub>2</sub>	70,373.19	73,421.79	0.0385	0.0844	0.5851
5A	Solid waste disposal	CH <sub>4</sub>	60,432.98	14,261.40	0.0244	0.0536	0.6387
1B1	Coal mining and handling	CH <sub>4</sub>	21,826.68	483.98	0.0143	0.0314	0.6702
2B3	Adipic acid production	N <sub>2</sub> O	19,934.61	-	0.0136	0.0298	0.7000

IPCC Code	IPCC Category	GHG	Base year emissions (Gg CO <sub>2</sub> e)	2017 emissions (Gg CO <sub>2</sub> e)	Trend Assessment	Contribution to Trend	Cumulative Total
2F1	Refrigeration and air conditioning	HFCs, PFCs, SF <sub>6</sub> and NF <sub>3</sub>	531.30	11,653.11	0.0134	0.0293	0.7293
1A2	Manufacturing industries and construction: solid fuels	CO <sub>2</sub>	38,484.20	12,133.91	0.0120	0.0263	0.7556
2B9	Fluorochemical production	HFCs, PFCs, SF <sub>6</sub> and NF <sub>3</sub>	17,784.67	175.44	0.0119	0.0262	0.7818
1A4	Other sectors: solid fuels	CO <sub>2</sub>	19,868.96	2,276.82	0.0109	0.0239	0.8056
1A1	Energy industries: liquid fuels	CO <sub>2</sub>	40,883.76	15,767.40	0.0093	0.0205	0.8261
1A2	Manufacturing industries and construction: gaseous fuels	CO <sub>2</sub>	27,330.36	23,723.43	0.0093	0.0204	0.8465
3A1	Enteric fermentation from Cattle	CH <sub>4</sub>	19,881.28	16,770.98	0.0062	0.0136	0.8601
1A1	Energy industries: other fuels	CO <sub>2</sub>	246.13	5,021.63	0.0057	0.0126	0.8727
3D	Agricultural soils	N <sub>2</sub> O	13,610.39	11,466.79	0.0042	0.0093	0.8820
4A	Forest land	CO <sub>2</sub>	-15,026.26	-18,211.77	0.0037	0.0081	0.8901
4B	Cropland	CO <sub>2</sub>	14,265.93	10,971.23	0.0032	0.0070	0.8971
1A2	Manufacturing industries and construction: liquid fuels	CO <sub>2</sub>	29,832.00	14,676.48	0.0031	0.0067	0.9038
4E	Settlements	CO <sub>2</sub>	7,011.30	6,451.34	0.0028	0.0062	0.9100
1B2	Oil and gas extraction	CH <sub>4</sub>	12,345.07	4,935.25	0.0026	0.0057	0.9157
2B2	Nitric acid production	N <sub>2</sub> O	3,860.26	37.25	0.0026	0.0057	0.9214
1A4	Other sectors: liquid fuels	CO <sub>2</sub>	19,607.73	13,284.20	0.0023	0.0050	0.9264
1A5	Other: liquid fuels	CO <sub>2</sub>	5,293.44	1,558.20	0.0018	0.0039	0.9303

IPCC Code	IPCC Category	GHG	Base year emissions (Gg CO <sub>2</sub> e)	2017 emissions (Gg CO <sub>2</sub> e)	Trend Assessment	Contribution to Trend	Cumulative Total
3B1	Manure management from Cattle	CH <sub>4</sub>	4,732.53	4,226.81	0.0017	0.0038	0.9341
2F4	Aerosols	HFCs, PFCs, SF <sub>6</sub> and NF <sub>3</sub>	663.00	1,658.10	0.0015	0.0033	0.9374
4C	Grassland	CO <sub>2</sub>	-7,111.03	-8,861.80	0.0015	0.0032	0.9406
3A2	Enteric fermentation from Sheep	CH <sub>4</sub>	4,935.90	4,043.20	0.0014	0.0031	0.9437
5B	Biological treatment of solid waste	CH <sub>4</sub>	18.13	1,193.62	0.0014	0.0031	0.9467
1A3c	Railways: liquid fuels	CO <sub>2</sub>	1,455.18	1,939.08	0.0013	0.0028	0.9496
1A3d	Domestic Navigation: liquid fuels	CO <sub>2</sub>	7,611.13	5,443.73	0.0012	0.0027	0.9522
5D	Wastewater treatment and discharge	CH <sub>4</sub>	4,218.90	3,445.75	0.0012	0.0026	0.9548
1B2	Oil and gas extraction	CO <sub>2</sub>	5,777.92	4,230.64	0.0010	0.0023	0.9571
3B2	Manure management from Sheep	N <sub>2</sub> O	3,442.70	2,814.79	0.0010	0.0021	0.9592
1A3a	Domestic aviation: liquid fuels	CO <sub>2</sub>	1,881.31	1,898.39	0.0010	0.0021	0.9613
2C6	Zinc production	CO <sub>2</sub>	1,358.83	-	0.0009	0.0020	0.9634
2C1	Iron and steel production	CO <sub>2</sub>	5,591.54	2,501.24	0.0009	0.0019	0.9653
5B	Biological treatment of solid waste	N <sub>2</sub> O	12.97	728.03	0.0008	0.0019	0.9671
1B1	Coal mining and handling solid fuels	CO <sub>2</sub>	1,698.56	268.38	0.0008	0.0018	0.9690
2B1	Ammonia production	CO <sub>2</sub>	1,895.00	1,763.86	0.0008	0.0017	0.9707
1A3b	Road transportation: liquid fuels	CH <sub>4</sub>	1,246.70	93.16	0.0007	0.0016	0.9723

IPCC Code	IPCC Category	GHG	Base year emissions (Gg CO <sub>2</sub> e)	2017 emissions (Gg CO <sub>2</sub> e)	Trend Assessment	Contribution to Trend	Cumulative Total
1A2	Manufacturing industries and construction: other fuels	CO <sub>2</sub>	76.86	641.74	0.0007	0.0015	0.9739
1A4	Other sectors: biomass	CH <sub>4</sub>	62.37	601.47	0.0007	0.0015	0.9753
1A4	Other sectors: solid fuels	CH <sub>4</sub>	1,265.22	173.62	0.0007	0.0014	0.9768
5C	Incineration and open burning of waste	CO <sub>2</sub>	1,300.71	258.91	0.0006	0.0013	0.9781
1A1	Energy industries: solid fuels	N <sub>2</sub> O	1,023.72	115.89	0.0006	0.0012	0.9793
1A3e	Other transportation: liquid fuels	CO <sub>2</sub>	224.74	565.53	0.0005	0.0011	0.9804
3G	Liming	CO <sub>2</sub>	1,015.18	939.13	0.0004	0.0009	0.9813
2F2	Foam blowing agents	HFCs, PFCs, SF <sub>6</sub> and NF <sub>3</sub>	184.49	450.52	0.0004	0.0009	0.9822
1A3b	Road transportation: liquid fuels	N <sub>2</sub> O	1,312.28	1,095.65	0.0004	0.0009	0.9831
2G3	N <sub>2</sub> O from product uses	N <sub>2</sub> O	552.57	645.41	0.0004	0.0008	0.9839
2F3	Fire protection	HFCs, PFCs, SF <sub>6</sub> and NF <sub>3</sub>	1.41	324.69	0.0004	0.0008	0.9848
4G	Harvested wood products	CO <sub>2</sub>	-1,639.08	-2,015.60	0.0004	0.0008	0.9856
3A4	Enteric fermentation from Other livestock	CH <sub>4</sub>	292.27	457.93	0.0003	0.0007	0.9863
5D	Wastewater treatment and discharge	N <sub>2</sub> O	783.59	725.93	0.0003	0.0007	0.9870
2B8	Petrochemical and carbon black production	CO <sub>2</sub>	4,538.91	2,846.51	0.0003	0.0006	0.9876
1A4	Other sectors: peat	CO <sub>2</sub>	372.48	4.83	0.0002	0.0005	0.9881

IPCC Code	IPCC Category	GHG	Base year emissions (Gg CO <sub>2</sub> e)	2017 emissions (Gg CO <sub>2</sub> e)	Trend Assessment	Contribution to Trend	Cumulative Total
2A2	Lime production	CO <sub>2</sub>	1,462.05	1,052.06	0.0002	0.0005	0.9887
2D	Non-energy products from fuels and solvent use	CO <sub>2</sub>	552.81	519.22	0.0002	0.0005	0.9892
2C3	Aluminium production	CO <sub>2</sub>	450.32	71.62	0.0002	0.0005	0.9897
4E	Settlements	N <sub>2</sub> O	584.68	524.99	0.0002	0.0005	0.9901
2A1	Cement production	CO <sub>2</sub>	7,295.26	4,409.79	0.0002	0.0005	0.9906
2C3	Aluminium production	HFCs, PFCs, SF <sub>6</sub> and NF <sub>3</sub>	333.43	15.07	0.0002	0.0005	0.9911
2G4	Other product manufacture and use	N <sub>2</sub> O	41.00	198.47	0.0002	0.0005	0.9915
1A1	Energy industries: gaseous fuels	N <sub>2</sub> O	198.91	289.24	0.0002	0.0004	0.9920
2G1	Electrical equipment	HFCs, PFCs, SF <sub>6</sub> and NF <sub>3</sub>	783.25	281.17	0.0002	0.0004	0.9924
1A1	Energy industries: biomass	N <sub>2</sub> O	0.25	171.90	0.0002	0.0004	0.9929
2G2	SF <sub>6</sub> and PFCs from other product use	HFCs, PFCs, SF <sub>6</sub> and NF <sub>3</sub>	278.70	318.00	0.0002	0.0004	0.9933
3H	Urea application to land	CO <sub>2</sub>	327.60	343.95	0.0002	0.0004	0.9937
2A3	Glass production	CO <sub>2</sub>	405.54	367.98	0.0002	0.0003	0.9940
4B	Cropland	N <sub>2</sub> O	1,019.85	461.97	0.0002	0.0003	0.9944
2B6	Titanium dioxide production	CO <sub>2</sub>	104.63	181.02	0.0001	0.0003	0.9947
2C4	Magnesium production	HFCs, PFCs, SF <sub>6</sub> and NF <sub>3</sub>	387.17	112.04	0.0001	0.0003	0.9950

IPCC Code	IPCC Category	GHG	Base year emissions (Gg CO <sub>2</sub> e)	2017 emissions (Gg CO <sub>2</sub> e)	Trend Assessment	Contribution to Trend	Cumulative Total
1A4	Other sectors: solid fuels	N <sub>2</sub> O	241.68	30.40	0.0001	0.0003	0.9952
1A1	Energy industries: biomass	CH <sub>4</sub>	0.47	109.01	0.0001	0.0003	0.9955
3F	Field burning of agricultural residues	CH <sub>4</sub>	186.57	-	0.0001	0.0003	0.9958
1A1	Energy industries: other fuels	CH <sub>4</sub>	18.55	110.63	0.0001	0.0003	0.9961
1A3b	Road transportation: biomass	CO <sub>2</sub>	-	92.80	0.0001	0.0002	0.9963
1A2	Manufacturing industries and construction: biomass	N <sub>2</sub> O	19.54	102.56	0.0001	0.0002	0.9965
1B1	Coal mining and handling liquid fuels	CO <sub>2</sub>	-	91.16	0.0001	0.0002	0.9968
1A1	Energy industries: other fuels	N <sub>2</sub> O	6.72	93.36	0.0001	0.0002	0.9970
1A4	Other sectors: biomass	N <sub>2</sub> O	9.14	94.30	0.0001	0.0002	0.9972
1A1	Energy industries: gaseous fuels	CH <sub>4</sub>	92.30	137.70	0.0001	0.0002	0.9974
1A4	Other sectors: gaseous fuels	CH <sub>4</sub>	157.18	161.99	0.0001	0.0002	0.9976
5C	Incineration and open burning of waste	CH <sub>4</sub>	134.83	10.45	0.0001	0.0002	0.9978
1A2	Manufacturing industries and construction: biomass	CH <sub>4</sub>	12.29	64.53	0.0001	0.0001	0.9980
1A2	Manufacturing industries and construction: solid fuels	N <sub>2</sub> O	148.72	30.91	0.0001	0.0001	0.9981
4D	Wetlands	CO <sub>2</sub>	486.95	336.78	0.0001	0.0001	0.9982
2B10	Other Chemical Industry	CH <sub>4</sub>	174.65	48.97	0.0001	0.0001	0.9984
2A4	Other process uses of carbonates	CO <sub>2</sub>	640.93	419.65	0.0001	0.0001	0.9985

IPCC Code	IPCC Category	GHG	Base year emissions (Gg CO <sub>2</sub> e)	2017 emissions (Gg CO <sub>2</sub> e)	Trend Assessment	Contribution to Trend	Cumulative Total
3J	Agriculture activities in OTs and CDs	CH <sub>4</sub>	276.73	206.19	0.0001	0.0001	0.9986
1A3c	Railways: solid fuels	CO <sub>2</sub>	-	41.15	0.0000	0.0001	0.9987
2F6	Other product uses as substitutes for ODS	HFCs, PFCs, SF <sub>6</sub> and NF <sub>3</sub>	36.88	61.71	0.0000	0.0001	0.9988
3F	Field burning of agricultural residues	N <sub>2</sub> O	57.66	-	0.0000	0.0001	0.9989
4C	Grassland	N <sub>2</sub> O	10.33	33.24	0.0000	0.0001	0.9990
1A1	Energy industries: solid fuels	CH <sub>4</sub>	49.32	5.69	0.0000	0.0001	0.9990
3A3	Enteric fermentation from Swine	CH <sub>4</sub>	283.06	186.33	0.0000	0.0001	0.9991
1A4	Other sectors: liquid fuels	N <sub>2</sub> O	103.32	82.03	0.0000	0.0001	0.9992
2F5	Solvents	HFCs, PFCs, SF <sub>6</sub> and NF <sub>3</sub>	-	18.67	0.0000	0.0000	0.9992
1A4	Other sectors: gaseous fuels	N <sub>2</sub> O	37.47	38.62	0.0000	0.0000	0.9993
1A2	Manufacturing industries and construction: solid fuels	CH <sub>4</sub>	47.59	10.86	0.0000	0.0000	0.9993
3J	Agriculture activities in OTs and CDs	N <sub>2</sub> O	132.00	93.01	0.0000	0.0000	0.9993
2E1	Integrated circuit or semiconductor	HFCs, PFCs, SF <sub>6</sub> and NF <sub>3</sub>	9.56	21.71	0.0000	0.0000	0.9994
1A5	Other: liquid fuels	N <sub>2</sub> O	56.12	16.48	0.0000	0.0000	0.9994
1B2	Oil and gas extraction	N <sub>2</sub> O	40.75	39.36	0.0000	0.0000	0.9995
1A4	Other sectors: peat	CH <sub>4</sub>	26.35	0.34	0.0000	0.0000	0.9995

IPCC Code	IPCC Category	GHG	Base year emissions (Gg CO <sub>2</sub> e)	2017 emissions (Gg CO <sub>2</sub> e)	Trend Assessment	Contribution to Trend	Cumulative Total
1A2	Manufacturing industries and construction: liquid fuels	N <sub>2</sub> O	138.62	93.64	0.0000	0.0000	0.9995
5C	Incineration and open burning of waste	N <sub>2</sub> O	46.66	39.63	0.0000	0.0000	0.9996
2A4	Other process uses of carbonates	CH <sub>4</sub>	31.10	5.89	0.0000	0.0000	0.9996
4C	Grassland	CH <sub>4</sub>	9.97	17.78	0.0000	0.0000	0.9996
1A3d	Domestic Navigation: liquid fuels	N <sub>2</sub> O	105.00	72.08	0.0000	0.0000	0.9997
4	Indirect N <sub>2</sub> O emissions from LULUCF	N <sub>2</sub> O	401.49	243.62	0.0000	0.0000	0.9997
4A	Forest land	N <sub>2</sub> O	231.22	144.65	0.0000	0.0000	0.9997
2C1	Iron and steel production	CH <sub>4</sub>	36.90	10.68	0.0000	0.0000	0.9997
1A3c	Railways: liquid fuels	N <sub>2</sub> O	13.75	18.30	0.0000	0.0000	0.9998
1A2	Manufacturing industries and construction: other fuels	N <sub>2</sub> O	0.38	10.05	0.0000	0.0000	0.9998
1A3a	Domestic aviation: liquid fuels	N <sub>2</sub> O	17.80	17.96	0.0000	0.0000	0.9998
1A1	Energy industries: liquid fuels	CH <sub>4</sub>	43.36	18.08	0.0000	0.0000	0.9998
4E	Settlements	CH <sub>4</sub>	3.00	8.25	0.0000	0.0000	0.9998
1A2	Manufacturing industries and construction: liquid fuels	CH <sub>4</sub>	44.43	32.04	0.0000	0.0000	0.9999
2B7	Soda ash production	CO <sub>2</sub>	231.55	140.18	0.0000	0.0000	0.9999
1A2	Manufacturing industries and construction: other fuels	CH <sub>4</sub>	0.16	6.12	0.0000	0.0000	0.9999
1A3e	Other transportation: liquid fuels	N <sub>2</sub> O	2.79	7.00	0.0000	0.0000	0.9999

IPCC Code	IPCC Category	GHG	Base year emissions (Gg CO <sub>2</sub> e)	2017 emissions (Gg CO <sub>2</sub> e)	Trend Assessment	Contribution to Trend	Cumulative Total
2B8	Petrochemical and carbon black production	CH <sub>4</sub>	29.98	12.16	0.0000	0.0000	0.9999
1A3d	Domestic Navigation: liquid fuels	CH <sub>4</sub>	3.66	6.28	0.0000	0.0000	0.9999
2C1	Iron and steel production	N <sub>2</sub> O	17.74	6.19	0.0000	0.0000	0.9999
1A2	Manufacturing industries and construction: gaseous fuels	N <sub>2</sub> O	14.54	12.48	0.0000	0.0000	1.0000
1B1	Coal mining and handling biomass	CH <sub>4</sub>	0.10	3.76	0.0000	0.0000	1.0000
1A2	Manufacturing industries and construction: gaseous fuels	CH <sub>4</sub>	12.20	10.47	0.0000	0.0000	1.0000
1A3a	Domestic aviation: liquid fuels	CH <sub>4</sub>	6.37	1.55	0.0000	0.0000	1.0000
4D	Wetlands	N <sub>2</sub> O	4.13	0.30	0.0000	0.0000	1.0000
1A5	Other: liquid fuels	CH <sub>4</sub>	3.56	1.03	0.0000	0.0000	1.0000
1A3c	Railways: solid fuels	CH <sub>4</sub>	-	0.94	0.0000	0.0000	1.0000
1A4	Other sectors: peat	N <sub>2</sub> O	1.47	0.02	0.0000	0.0000	1.0000
1A4	Other sectors: liquid fuels	CH <sub>4</sub>	57.79	32.75	0.0000	0.0000	1.0000
4A	Forest land	CH <sub>4</sub>	2.89	2.31	0.0000	0.0000	1.0000
1A3c	Railways: liquid fuels	CH <sub>4</sub>	2.46	1.92	0.0000	0.0000	1.0000
1A1	Energy industries: liquid fuels	N <sub>2</sub> O	140.41	80.97	0.0000	0.0000	1.0000
2B8	Petrochemical and carbon black production	N <sub>2</sub> O	2.10	1.51	0.0000	0.0000	1.0000
1A3e	Other transportation: liquid fuels	CH <sub>4</sub>	0.29	0.35	0.0000	0.0000	1.0000
2B1	Ammonia production	N <sub>2</sub> O	0.31	0.33	0.0000	0.0000	1.0000
2B1	Ammonia production	CH <sub>4</sub>	0.26	0.28	0.0000	0.0000	1.0000

IPCC Code	IPCC Category	GHG	Base year emissions (Gg CO <sub>2</sub> e)	2017 emissions (Gg CO <sub>2</sub> e)	Trend Assessment	Contribution to Trend	Cumulative Total
1A3c	Railways: solid fuels	N <sub>2</sub> O	-	0.09	0.0000	0.0000	1.0000
1B1	Coal mining and handling solid fuels	N <sub>2</sub> O	0.09	0.02	0.0000	0.0000	1.0000
4B	Cropland	CH <sub>4</sub>	0.09	0.02	0.0000	0.0000	1.0000
1B1	Coal mining and handling solid fuels	CH <sub>4</sub>	0.08	0.01	0.0000	0.0000	1.0000
<b>Total</b>			<b>801,848.54</b>	<b>464,453.44</b>	<b>0.4554</b>	<b>1.0000</b>	

**Table A 1.3.6 Approach 1 Key Category Analysis based on trend in emissions (from base year to latest reported year, excluding LULUCF)**

IPCC Code	IPCC Category	GHG	Base year emissions (Gg CO <sub>2</sub> e)	2017 emissions (Gg CO <sub>2</sub> e)	Trend Assessment	Contribution to Trend	Cumulative Total
1A1	Energy industries: solid fuels	CO <sub>2</sub>	185,492.23	20,363.72	0.1025	0.2312	0.2312
1A1	Energy industries: gaseous fuels	CO <sub>2</sub>	9,229.34	61,476.35	0.0661	0.1490	0.3803
1A3b	Road transportation: liquid fuels	CO <sub>2</sub>	108,570.16	113,364.64	0.0594	0.1340	0.5143
1A4	Other sectors: gaseous fuels	CO <sub>2</sub>	70,373.19	73,421.79	0.0385	0.0867	0.6010
5A	Solid waste disposal	CH <sub>4</sub>	60,432.98	14,261.40	0.0244	0.0551	0.6561
1B1	Coal mining and handling	CH <sub>4</sub>	21,826.68	483.98	0.0143	0.0323	0.6884
2B3	Adipic acid production	N <sub>2</sub> O	19,934.61	-	0.0136	0.0307	0.7190
2F1	Refrigeration and air conditioning	HFCs, PFCs, SF <sub>6</sub> and NF <sub>3</sub>	531.30	11,653.11	0.0134	0.0301	0.7492
1A2	Manufacturing industries and construction: solid fuels	CO <sub>2</sub>	38,484.20	12,133.91	0.0120	0.0270	0.7761

IPCC Code	IPCC Category	GHG	Base year emissions (Gg CO <sub>2</sub> e)	2017 emissions (Gg CO <sub>2</sub> e)	Trend Assessment	Contribution to Trend	Cumulative Total
2B9	Fluorochemical production	HFCs, PFCs, SF <sub>6</sub> and NF <sub>3</sub>	17,784.67	175.44	0.0119	0.0269	0.8030
1A4	Other sectors: solid fuels	CO <sub>2</sub>	19,868.96	2,276.82	0.0109	0.0245	0.8275
1A1	Energy industries: liquid fuels	CO <sub>2</sub>	40,883.76	15,767.40	0.0093	0.0210	0.8485
1A2	Manufacturing industries and construction: gaseous fuels	CO <sub>2</sub>	27,330.36	23,723.43	0.0093	0.0210	0.8695
3A1	Enteric fermentation from Cattle	CH <sub>4</sub>	19,881.28	16,770.98	0.0062	0.0140	0.8835
1A1	Energy industries: other fuels	CO <sub>2</sub>	246.13	5,021.63	0.0057	0.0130	0.8964
3D	Agricultural soils	N <sub>2</sub> O	13,610.39	11,466.79	0.0042	0.0095	0.9059
1A2	Manufacturing industries and construction: liquid fuels	CO <sub>2</sub>	29,832.00	14,676.48	0.0031	0.0069	0.9128
1B2	Oil and gas extraction	CH <sub>4</sub>	12,345.07	4,935.25	0.0026	0.0059	0.9187
2B2	Nitric acid production	N <sub>2</sub> O	3,860.26	37.25	0.0026	0.0058	0.9246
1A4	Other sectors: liquid fuels	CO <sub>2</sub>	19,607.73	13,284.20	0.0023	0.0051	0.9297
1A5	Other: liquid fuels	CO <sub>2</sub>	5,293.44	1,558.20	0.0018	0.0040	0.9337
3B1	Manure management from Cattle	CH <sub>4</sub>	4,732.53	4,226.81	0.0017	0.0039	0.9376
2F4	Aerosols	HFCs, PFCs, SF <sub>6</sub> and NF <sub>3</sub>	663.00	1,658.10	0.0015	0.0034	0.9410
3A2	Enteric fermentation from Sheep	CH <sub>4</sub>	4,935.90	4,043.20	0.0014	0.0031	0.9441
5B	Biological treatment of solid waste	CH <sub>4</sub>	18.13	1,193.62	0.0014	0.0031	0.9473

IPCC Code	IPCC Category	GHG	Base year emissions (Gg CO <sub>2</sub> e)	2017 emissions (Gg CO <sub>2</sub> e)	Trend Assessment	Contribution to Trend	Cumulative Total
1A3c	Railways: liquid fuels	CO <sub>2</sub>	1,455.18	1,939.08	0.0013	0.0029	0.9502
1A3d	Domestic Navigation: liquid fuels	CO <sub>2</sub>	7,611.13	5,443.73	0.0012	0.0027	0.9529
5D	Wastewater treatment and discharge	CH <sub>4</sub>	4,218.90	3,445.75	0.0012	0.0027	0.9556
1B2	Oil and gas extraction	CO <sub>2</sub>	5,777.92	4,230.64	0.0010	0.0023	0.9580
3B2	Manure management from Sheep	N <sub>2</sub> O	3,442.70	2,814.79	0.0010	0.0022	0.9601
1A3a	Domestic aviation: liquid fuels	CO <sub>2</sub>	1,881.31	1,898.39	0.0010	0.0021	0.9623
2C6	Zinc production	CO <sub>2</sub>	1,358.83	-	0.0009	0.0021	0.9644
2C1	Iron and steel production	CO <sub>2</sub>	5,591.54	2,501.24	0.0009	0.0020	0.9663
5B	Biological treatment of solid waste	N <sub>2</sub> O	12.97	728.03	0.0008	0.0019	0.9682
1B1	Coal mining and handling solid fuels	CO <sub>2</sub>	1,698.56	268.38	0.0008	0.0019	0.9701
2B1	Ammonia production	CO <sub>2</sub>	1,895.00	1,763.86	0.0008	0.0018	0.9719
1A3b	Road transportation: liquid fuels	CH <sub>4</sub>	1,246.70	93.16	0.0007	0.0017	0.9736
1A2	Manufacturing industries and construction: other fuels	CO <sub>2</sub>	76.86	641.74	0.0007	0.0016	0.9752
1A4	Other sectors: biomass	CH <sub>4</sub>	62.37	601.47	0.0007	0.0015	0.9767
1A4	Other sectors: solid fuels	CH <sub>4</sub>	1,265.22	173.62	0.0007	0.0015	0.9782
5C	Incineration and open burning of waste	CO <sub>2</sub>	1,300.71	258.91	0.0006	0.0013	0.9795
1A1	Energy industries: solid fuels	N <sub>2</sub> O	1,023.72	115.89	0.0006	0.0013	0.9807

IPCC Code	IPCC Category	GHG	Base year emissions (Gg CO <sub>2</sub> e)	2017 emissions (Gg CO <sub>2</sub> e)	Trend Assessment	Contribution to Trend	Cumulative Total
1A3e	Other transportation: liquid fuels	CO <sub>2</sub>	224.74	565.53	0.0005	0.0012	0.9819
3G	Liming	CO <sub>2</sub>	1,015.18	939.13	0.0004	0.0009	0.9828
2F2	Foam blowing agents	HFCs, PFCs, SF <sub>6</sub> and NF <sub>3</sub>	184.49	450.52	0.0004	0.0009	0.9837
1A3b	Road transportation: liquid fuels	N <sub>2</sub> O	1,312.28	1,095.65	0.0004	0.0009	0.9846
2G3	N <sub>2</sub> O from product uses	N <sub>2</sub> O	552.57	645.41	0.0004	0.0009	0.9855
2F3	Fire protection	HFCs, PFCs, SF <sub>6</sub> and NF <sub>3</sub>	1.41	324.69	0.0004	0.0009	0.9863
3A4	Enteric fermentation from Other livestock	CH <sub>4</sub>	292.27	457.93	0.0003	0.0008	0.9871
5D	Wastewater treatment and discharge	N <sub>2</sub> O	783.59	725.93	0.0003	0.0007	0.9878
2B8	Petrochemical and carbon black production	CO <sub>2</sub>	4,538.91	2,846.51	0.0003	0.0006	0.9884
1A4	Other sectors: peat	CO <sub>2</sub>	372.48	4.83	0.0002	0.0006	0.9890
2A2	Lime production	CO <sub>2</sub>	1,462.05	1,052.06	0.0002	0.0005	0.9895
2D	Non-energy products from fuels and solvent use	CO <sub>2</sub>	552.81	519.22	0.0002	0.0005	0.9900
2C3	Aluminium production	CO <sub>2</sub>	450.32	71.62	0.0002	0.0005	0.9906
2A1	Cement production	CO <sub>2</sub>	7,295.26	4,409.79	0.0002	0.0005	0.9910
2C3	Aluminium production	HFCs, PFCs, SF <sub>6</sub> and NF <sub>3</sub>	333.43	15.07	0.0002	0.0005	0.9915

IPCC Code	IPCC Category	GHG	Base year emissions (Gg CO <sub>2</sub> e)	2017 emissions (Gg CO <sub>2</sub> e)	Trend Assessment	Contribution to Trend	Cumulative Total
2G4	Other product manufacture and use	N <sub>2</sub> O	41.00	198.47	0.0002	0.0005	0.9920
1A1	Energy industries: gaseous fuels	N <sub>2</sub> O	198.91	289.24	0.0002	0.0005	0.9924
2G1	Electrical equipment	HFCs, PFCs, SF <sub>6</sub> and NF <sub>3</sub>	783.25	281.17	0.0002	0.0005	0.9929
1A1	Energy industries: biomass	N <sub>2</sub> O	0.25	171.90	0.0002	0.0005	0.9934
2G2	SF <sub>6</sub> and PFCs from other product use	HFCs, PFCs, SF <sub>6</sub> and NF <sub>3</sub>	278.70	318.00	0.0002	0.0004	0.9938
3H	Urea application to land	CO <sub>2</sub>	327.60	343.95	0.0002	0.0004	0.9942
2A3	Glass production	CO <sub>2</sub>	405.54	367.98	0.0002	0.0004	0.9945
2B6	Titanium dioxide production	CO <sub>2</sub>	104.63	181.02	0.0001	0.0003	0.9949
2C4	Magnesium production	HFCs, PFCs, SF <sub>6</sub> and NF <sub>3</sub>	387.17	112.04	0.0001	0.0003	0.9951
1A4	Other sectors: solid fuels	N <sub>2</sub> O	241.68	30.40	0.0001	0.0003	0.9954
1A1	Energy industries: biomass	CH <sub>4</sub>	0.47	109.01	0.0001	0.0003	0.9957
3F	Field burning of agricultural residues	CH <sub>4</sub>	186.57	-	0.0001	0.0003	0.9960
1A1	Energy industries: other fuels	CH <sub>4</sub>	18.55	110.63	0.0001	0.0003	0.9963
1A3b	Road transportation: biomass	CO <sub>2</sub>	-	92.80	0.0001	0.0002	0.9965
1A2	Manufacturing industries and construction: biomass	N <sub>2</sub> O	19.54	102.56	0.0001	0.0002	0.9968

IPCC Code	IPCC Category	GHG	Base year emissions (Gg CO <sub>2</sub> e)	2017 emissions (Gg CO <sub>2</sub> e)	Trend Assessment	Contribution to Trend	Cumulative Total
1B1	Coal mining and handling liquid fuels	CO <sub>2</sub>	-	91.16	0.0001	0.0002	0.9970
1A1	Energy industries: other fuels	N <sub>2</sub> O	6.72	93.36	0.0001	0.0002	0.9972
1A4	Other sectors: biomass	N <sub>2</sub> O	9.14	94.30	0.0001	0.0002	0.9975
1A1	Energy industries: gaseous fuels	CH <sub>4</sub>	92.30	137.70	0.0001	0.0002	0.9977
1A4	Other sectors: gaseous fuels	CH <sub>4</sub>	157.18	161.99	0.0001	0.0002	0.9979
5C	Incineration and open burning of waste	CH <sub>4</sub>	134.83	10.45	0.0001	0.0002	0.9981
1A2	Manufacturing industries and construction: biomass	CH <sub>4</sub>	12.29	64.53	0.0001	0.0002	0.9982
1A2	Manufacturing industries and construction: solid fuels	N <sub>2</sub> O	148.72	30.91	0.0001	0.0001	0.9984
2B10	Other Chemical Industry	CH <sub>4</sub>	174.65	48.97	0.0001	0.0001	0.9985
2A4	Other process uses of carbonates	CO <sub>2</sub>	640.93	419.65	0.0001	0.0001	0.9986
3J	Agriculture activities in OTs and CDs	CH <sub>4</sub>	276.73	206.19	0.0001	0.0001	0.9988
1A3c	Railways: solid fuels	CO <sub>2</sub>	-	41.15	0.0000	0.0001	0.9989
2F6	Other product uses as substitutes for ODS	HFCs, PFCs, SF <sub>6</sub> and NF <sub>3</sub>	36.88	61.71	0.0000	0.0001	0.9990
3F	Field burning of agricultural residues	N <sub>2</sub> O	57.66	-	0.0000	0.0001	0.9991
1A1	Energy industries: solid fuels	CH <sub>4</sub>	49.32	5.69	0.0000	0.0001	0.9991
3A3	Enteric fermentation from Swine	CH <sub>4</sub>	283.06	186.33	0.0000	0.0001	0.9992

IPCC Code	IPCC Category	GHG	Base year emissions (Gg CO <sub>2</sub> e)	2017 emissions (Gg CO <sub>2</sub> e)	Trend Assessment	Contribution to Trend	Cumulative Total
1A4	Other sectors: liquid fuels	N <sub>2</sub> O	103.32	82.03	0.0000	0.0001	0.9993
2F5	Solvents	HFCs, PFCs, SF <sub>6</sub> and NF <sub>3</sub>	-	18.67	0.0000	0.0000	0.9993
1A4	Other sectors: gaseous fuels	N <sub>2</sub> O	37.47	38.62	0.0000	0.0000	0.9993
1A2	Manufacturing industries and construction: solid fuels	CH <sub>4</sub>	47.59	10.86	0.0000	0.0000	0.9994
3J	Agriculture activities in OTs and CDs	N <sub>2</sub> O	132.00	93.01	0.0000	0.0000	0.9994
2E1	Integrated circuit or semiconductor	HFCs, PFCs, SF <sub>6</sub> and NF <sub>3</sub>	9.56	21.71	0.0000	0.0000	0.9995
1A5	Other: liquid fuels	N <sub>2</sub> O	56.12	16.48	0.0000	0.0000	0.9995
1B2	Oil and gas extraction	N <sub>2</sub> O	40.75	39.36	0.0000	0.0000	0.9996
1A4	Other sectors: peat	CH <sub>4</sub>	26.35	0.34	0.0000	0.0000	0.9996
1A2	Manufacturing industries and construction: liquid fuels	N <sub>2</sub> O	138.62	93.64	0.0000	0.0000	0.9996
5C	Incineration and open burning of waste	N <sub>2</sub> O	46.66	39.63	0.0000	0.0000	0.9997
2A4	Other process uses of carbonates	CH <sub>4</sub>	31.10	5.89	0.0000	0.0000	0.9997
1A3d	Domestic Navigation: liquid fuels	N <sub>2</sub> O	105.00	72.08	0.0000	0.0000	0.9997
2C1	Iron and steel production	CH <sub>4</sub>	36.90	10.68	0.0000	0.0000	0.9998
1A3c	Railways: liquid fuels	N <sub>2</sub> O	13.75	18.30	0.0000	0.0000	0.9998

IPCC Code	IPCC Category	GHG	Base year emissions (Gg CO <sub>2</sub> e)	2017 emissions (Gg CO <sub>2</sub> e)	Trend Assessment	Contribution to Trend	Cumulative Total
1A2	Manufacturing industries and construction: other fuels	N <sub>2</sub> O	0.38	10.05	0.0000	0.0000	0.9998
1A3a	Domestic aviation: liquid fuels	N <sub>2</sub> O	17.80	17.96	0.0000	0.0000	0.9998
1A1	Energy industries: liquid fuels	CH <sub>4</sub>	43.36	18.08	0.0000	0.0000	0.9999
1A2	Manufacturing industries and construction: liquid fuels	CH <sub>4</sub>	44.43	32.04	0.0000	0.0000	0.9999
2B7	Soda ash production	CO <sub>2</sub>	231.55	140.18	0.0000	0.0000	0.9999
1A2	Manufacturing industries and construction: other fuels	CH <sub>4</sub>	0.16	6.12	0.0000	0.0000	0.9999
1A3e	Other transportation: liquid fuels	N <sub>2</sub> O	2.79	7.00	0.0000	0.0000	0.9999
2B8	Petrochemical and carbon black production	CH <sub>4</sub>	29.98	12.16	0.0000	0.0000	0.9999
1A3d	Domestic Navigation: liquid fuels	CH <sub>4</sub>	3.66	6.28	0.0000	0.0000	0.9999
2C1	Iron and steel production	N <sub>2</sub> O	17.74	6.19	0.0000	0.0000	1.0000
1A2	Manufacturing industries and construction: gaseous fuels	N <sub>2</sub> O	14.54	12.48	0.0000	0.0000	1.0000
1B1	Coal mining and handling biomass	CH <sub>4</sub>	0.10	3.76	0.0000	0.0000	1.0000
1A2	Manufacturing industries and construction: gaseous fuels	CH <sub>4</sub>	12.20	10.47	0.0000	0.0000	1.0000
1A3a	Domestic aviation: liquid fuels	CH <sub>4</sub>	6.37	1.55	0.0000	0.0000	1.0000
1A5	Other: liquid fuels	CH <sub>4</sub>	3.56	1.03	0.0000	0.0000	1.0000
1A3c	Railways: solid fuels	CH <sub>4</sub>	-	0.94	0.0000	0.0000	1.0000

IPCC Code	IPCC Category	GHG	Base year emissions (Gg CO <sub>2</sub> e)	2017 emissions (Gg CO <sub>2</sub> e)	Trend Assessment	Contribution to Trend	Cumulative Total
1A4	Other sectors: peat	N <sub>2</sub> O	1.47	0.02	0.0000	0.0000	1.0000
1A4	Other sectors: liquid fuels	CH <sub>4</sub>	57.79	32.75	0.0000	0.0000	1.0000
1A3c	Railways: liquid fuels	CH <sub>4</sub>	2.46	1.92	0.0000	0.0000	1.0000
1A1	Energy industries: liquid fuels	N <sub>2</sub> O	140.41	80.97	0.0000	0.0000	1.0000
2B8	Petrochemical and carbon black production	N <sub>2</sub> O	2.10	1.51	0.0000	0.0000	1.0000
1A3e	Other transportation: liquid fuels	CH <sub>4</sub>	0.29	0.35	0.0000	0.0000	1.0000
2B1	Ammonia production	N <sub>2</sub> O	0.31	0.33	0.0000	0.0000	1.0000
2B1	Ammonia production	CH <sub>4</sub>	0.26	0.28	0.0000	0.0000	1.0000
1A3c	Railways: solid fuels	N <sub>2</sub> O	-	0.09	0.0000	0.0000	1.0000
1B1	Coal mining and handling solid fuels	N <sub>2</sub> O	0.09	0.02	0.0000	0.0000	1.0000
1B1	Coal mining and handling solid fuels	CH <sub>4</sub>	0.08	0.01	0.0000	0.0000	1.0000
<b>Total</b>			<b>801,593.08</b>	<b>474,346.12</b>	<b>0.4434</b>	<b>1.0000</b>	

## A 1.4 GUIDELINES

Following the 2006 IPCC Guidelines, the UK has also completed an Approach 2 KCA for both level and trend, which takes into account uncertainties, using the Approach 1 method for uncertainty estimates. This analysis has been performed using the data shown in **Table A 1.4.1** to **Table A 1.4.4** using the same categorisation and the same estimates of uncertainty.

The results of the level assessment (based on Approach 2) with and without LULUCF for the base year and the latest reported year are shown in **Table A 1.4.1** to **Table A 1.4.4**. The key source categories are highlighted by the shaded cells in the table. The source categories (i.e. rows of the table) were sorted in descending order of magnitude based on the results of the “Level Parameter”, and then the cumulative total was included in the final column of the table. The key source categories are those whose contributions add up to 90% of the sum of the level parameter in the final column after this sorting process, which accounts for 90% of the uncertainty in level.

The results of the trend assessment (based on Approach 2) with and without LULUCF for the base year to the latest reported year to the latest reported year, are shown in **Table A 1.4.5** to **Table A 1.4.6**

The key source categories are highlighted by the shaded cells in the table. The trend parameter was calculated using the absolute value of the result; an absolute function is used since Land Use, Land Use Change and Forestry contains negative sources (sinks) and the absolute function is necessary to produce positive uncertainty contributions for these sinks. The source categories (i.e. rows of the table) were sorted in descending order of magnitude based on the results of the trend parameter, and then the cumulative total was included in the final column of the table. The key source categories are those whose contributions add up to 90% of the sum of the level parameter in the final column after this sorting process, which accounts for 90% of the uncertainty in trend.

Any methodological improvements to the uncertainty analysis are discussed in **Annex 2**.

**Table A 1.4.1 Approach 2 Level Assessment for Base year (including LULUCF) with Key Categories Shaded in Grey**

IPCC Code	IPCC Category	Gas	Base year emissions (Gg CO <sub>2e</sub> )	Absolute value of Base year emissions (Gg CO <sub>2e</sub> )	Level Assessment	Cumulative Total
5A	5A Solid Waste Disposal	CH <sub>4</sub>	60432.98	60432.98	0.2397	0.2397
2B3	2B3 Adipic Acid Production	N <sub>2</sub> O	19934.61	19934.61	0.1634	0.4031
1A	1A Coal	CO <sub>2</sub>	243845.39	243845.39	0.0617	0.4648
4B	4B Cropland	CO <sub>2</sub>	14265.93	14265.93	0.0526	0.5174
1A	1A (Stationary) Oil	CO <sub>2</sub>	95612.77	95612.77	0.0509	0.5683
4A	4A Forest Land	CO <sub>2</sub>	-15026.26	15026.26	0.0431	0.6114
1B1	1B1 Coal Mining	CH <sub>4</sub>	21826.68	21826.68	0.0360	0.6474
5C	5C Waste Incineration	CO <sub>2</sub>	1300.71	1300.71	0.0323	0.6796
4C	4C Grassland	CO <sub>2</sub>	-7111.03	7111.03	0.0291	0.7088
4E	4E Settlements	CO <sub>2</sub>	7011.30	7011.30	0.0287	0.7375
3A	3A Enteric Fermentation	CH <sub>4</sub>	25392.50	25392.50	0.0286	0.7661
1A	1A Natural Gas	CO <sub>2</sub>	106932.90	106932.90	0.0170	0.7831
1B2	1B2 Natural Gas Transmission	CH <sub>4</sub>	10168.33	10168.33	0.0169	0.8000
5D	5D Wastewater Handling	N <sub>2</sub> O	783.59	783.59	0.0159	0.8159
2B	2B Chemical industry	HFCs	17670.77	17670.77	0.0145	0.8304
1A3b	1A3b Gasoline/ LPG	CO <sub>2</sub>	75562.72	75562.72	0.0138	0.8442
3D	3D Agricultural Soils	N <sub>2</sub> O	13610.39	13610.39	0.0124	0.8566
1A1 & 1A2 & 1A4 & 1A5	1A1 & 1A2 & 1A4 & 1A5 Other Combustion	N <sub>2</sub> O	2141.02	2141.02	0.0115	0.8681

IPCC Code	IPCC Category	Gas	Base year emissions (Gg CO <sub>2e</sub> )	Absolute value of Base year emissions (Gg CO <sub>2e</sub> )	Level Assessment	Cumulative Total
1A3d	1A3d Marine fuel	CO <sub>2</sub>	7611.13	7611.13	0.0111	0.8792
2B	2B Chemical industries	CO <sub>2</sub>	6770.09	6770.09	0.0097	0.8890
5D	5D Wastewater Handling	CH <sub>4</sub>	4218.90	4218.90	0.0093	0.8983
1A3b	1A3b Gasoline/ LPG	CH <sub>4</sub>	1157.96	1157.96	0.0071	0.9054
2G	2G Other Product Manufacture and Use	N <sub>2</sub> O	593.57	593.57	0.0069	0.9123
1A3b	1A3b DERV	CO <sub>2</sub>	33005.80	33005.80	0.0060	0.9183
4G	4G Other Activities	CO <sub>2</sub>	-1639.08	1639.08	0.0060	0.9244
1A3b	1A3b Gasoline/ LPG	N <sub>2</sub> O	989.40	989.40	0.0060	0.9304
1A1 & 1A2 & 1A4 & 1A5	1A1 & 1A2 & 1A4 & 1A5 Other Combustion	CH <sub>4</sub>	1893.15	1893.15	0.0055	0.9359
4	4 Indirect LULUCF Emissions	N <sub>2</sub> O	401.49	401.49	0.0054	0.9414
4D	4D Wetland	CO <sub>2</sub>	486.95	486.95	0.0040	0.9454
1A3b	1A3b DERV	N <sub>2</sub> O	322.88	322.88	0.0034	0.9488
1B2	1B2 Offshore Oil& Gas	CH <sub>4</sub>	2176.74	2176.74	0.0033	0.9521
1B2	1B2 Oil & Natural Gas	CO <sub>2</sub>	5777.92	5777.92	0.0033	0.9555
3B	3B Manure Management	CH <sub>4</sub>	4732.53	4732.53	0.0032	0.9587
2C	2C Metal Industries	CO <sub>2</sub>	7400.69	7400.69	0.0032	0.9619
2B2	2B2 Nitric Acid Production	N <sub>2</sub> O	3860.26	3860.26	0.0032	0.9650
1A3a	1A3a Aviation Fuel	CO <sub>2</sub>	1881.31	1881.31	0.0031	0.9681
4B	4B Cropland	N <sub>2</sub> O	1019.85	1019.85	0.0029	0.9710
3B	3B Manure Management	N <sub>2</sub> O	3442.70	3442.70	0.0027	0.9737
1A3	1A3 Other diesel	CO <sub>2</sub>	1679.92	1679.92	0.0020	0.9757
2A1	2A1 Cement Production	CO <sub>2</sub>	7295.26	7295.26	0.0019	0.9776
4A	4A Forest land	N <sub>2</sub> O	231.22	231.22	0.0018	0.9794
3G	3G Liming	CO <sub>2</sub>	1015.18	1015.18	0.0017	0.9812
2D	2D Non Energy Products from Fuels and Solvent Use	CO <sub>2</sub>	552.81	552.81	0.0017	0.9829
3J	3J OT & CD Agriculture	CH <sub>4</sub>	276.73	276.73	0.0016	0.9845
2F	2F Product Uses as Substitutes for ODS	HFCs	1416.64	1416.64	0.0014	0.9859

IPCC Code	IPCC Category	Gas	Base year emissions (Gg CO <sub>2e</sub> )	Absolute value of Base year emissions (Gg CO <sub>2e</sub> )	Level Assessment	Cumulative Total
3H	3H Urea application to agriculture	CO <sub>2</sub>	327.60	327.60	0.0013	0.9872
1A3d	1A3d Marine fuel	N <sub>2</sub> O	105.00	105.00	0.0010	0.9882
1B1	1B1 Solid Fuel Transformation	CO <sub>2</sub>	1698.56	1698.56	0.0010	0.9892
1A4	1A4 Peat	CO <sub>2</sub>	372.48	372.48	0.0010	0.9902
1A3b	1A3b DERV	CH <sub>4</sub>	88.74	88.74	0.0009	0.9911
5C	5C Waste Incineration	N <sub>2</sub> O	46.66	46.66	0.0009	0.9920
3J	3J OT & CD Agriculture	N <sub>2</sub> O	132.00	132.00	0.0008	0.9928
4E	4E Settlements	N <sub>2</sub> O	584.68	584.68	0.0007	0.9935
2A2	2A2 Lime Production	CO <sub>2</sub>	1462.05	1462.05	0.0006	0.9941
2G	2G Other Product Manufacture and Use	PFCs	149.16	149.16	0.0006	0.9946
2C	2C Metal Industries	PFCs	333.43	333.43	0.0005	0.9952
5C	5C Waste Incineration	CH <sub>4</sub>	134.83	134.83	0.0005	0.9957
2G	2G Other Product Manufacture and Use	SF <sub>6</sub>	912.79	912.79	0.0005	0.9962
3F	3F Field Burning	CH <sub>4</sub>	186.57	186.57	0.0004	0.9965
1B2	1B2 Oil & Natural Gas	N <sub>2</sub> O	40.75	40.75	0.0004	0.9969
2B	2B Chemical Industry	CH <sub>4</sub>	204.89	204.89	0.0003	0.9972
1A	1A Other (waste)	CO <sub>2</sub>	247.13	247.13	0.0003	0.9975
2A	2A Minerals industry	CH <sub>4</sub>	31.10	31.10	0.0003	0.9978
2C	2C Metal Industries	SF <sub>6</sub>	387.17	387.17	0.0002	0.9980
2A4	2A4 Other process uses of carbonates	CO <sub>2</sub>	640.93	640.93	0.0002	0.9982
1A3	1A3 Other diesel	N <sub>2</sub> O	16.53	16.53	0.0002	0.9984
2C	2C Iron & Steel	N <sub>2</sub> O	17.74	17.74	0.0002	0.9985
1A4	1A4 Petroleum Coke	CO <sub>2</sub>	81.64	81.64	0.0002	0.9987
2A3	2A3 Glass production	CO <sub>2</sub>	405.54	405.54	0.0002	0.9989
1A3a	1A3a Aviation Fuel	N <sub>2</sub> O	17.80	17.80	0.0002	0.9990
5B	5B Biological treatment of solid waste	CH <sub>4</sub>	18.13	18.13	0.0002	0.9992
2C	2C Iron & Steel Production	CH <sub>4</sub>	36.90	36.90	0.0001	0.9993
3F	3F Field Burning	N <sub>2</sub> O	57.66	57.66	0.0001	0.9994

IPCC Code	IPCC Category	Gas	Base year emissions (Gg CO <sub>2e</sub> )	Absolute value of Base year emissions (Gg CO <sub>2e</sub> )	Level Assessment	Cumulative Total
5B	5B Biological treatment of solid waste	N <sub>2</sub> O	12.97	12.97	0.0001	0.9995
4C	4C Grassland	N <sub>2</sub> O	10.33	10.33	0.0001	0.9996
2B	2B Chemical industry	PFCs	113.90	113.90	0.0001	0.9997
4C	4C Grassland	CH <sub>4</sub>	9.97	9.97	0.0000	0.9998
1A3d	1A3d Marine fuel	CH <sub>4</sub>	3.66	3.66	0.0000	0.9998
4D	4D Grassland	N <sub>2</sub> O	4.13	4.13	0.0000	0.9999
2E	2E Electronics Industry	HFCs	8.73	8.73	0.0000	0.9999
1A3a	1A3a Aviation Fuel	CH <sub>4</sub>	6.37	6.37	0.0000	0.9999
1A3	1A3 Other diesel	CH <sub>4</sub>	2.75	2.75	0.0000	0.9999
2B8	2B8 Petrochemical and Carbon Black Production	N <sub>2</sub> O	2.10	2.10	0.0000	1.0000
4E	4E Settlements	CH <sub>4</sub>	3.00	3.00	0.0000	1.0000
4A	4A Forest Land	CH <sub>4</sub>	2.89	2.89	0.0000	1.0000
2E	2E Electronics Industry	NF <sub>3</sub>	0.83	0.83	0.0000	1.0000
2B1	2B1 Ammonia Production	N <sub>2</sub> O	0.31	0.31	0.0000	1.0000
2F	2F Product Uses as Substitutes for ODS	PFCs	0.44	0.44	0.0000	1.0000
1B1	1B1 Fugitive Emissions from Solid Fuels	N <sub>2</sub> O	0.09	0.09	0.0000	1.0000
1B1	1B1 Solid Fuel Transformation	CH <sub>4</sub>	0.18	0.18	0.0000	1.0000
4B	4B Cropland	CH <sub>4</sub>	0.09	0.09	0.0000	1.0000
1A3c	1A3c Coal	CO <sub>2</sub>	0.00	0.00	0.0000	1.0000
4F	4F Other Land	CO <sub>2</sub>	0.00	0.00	0.0000	1.0000
1A3c	1A3c Coal	CH <sub>4</sub>	0.00	0.00	0.0000	1.0000
2D	2D Non-energy Products from Fuels and Solvent Use	CH <sub>4</sub>	0.00	0.00	0.0000	1.0000
1A3c	1A3c Coal	N <sub>2</sub> O	0.00	0.00	0.0000	1.0000
2D	2D Non-energy Products from Fuels and Solvent Use	N <sub>2</sub> O	0.00	0.00	0.0000	1.0000
2C	2C Metal Industries	HFCs	0.00	0.00	0.0000	1.0000

**Table A 1.4.2 Approach 2 Level Assessment for the latest reported year (including LULUCF) with Key Categories Shaded in Grey**

IPCC Code	IPCC Category	Gas	2017 emissions (Gg CO <sub>2</sub> e)	Absolute value of 2017 emissions (Gg CO <sub>2</sub> e)	Level Assessment	Cumulative Total
5A	5A Solid Waste Disposal	CH <sub>4</sub>	14261.40	14261.40	0.1151	0.1151
4A	4A Forest Land	CO <sub>2</sub>	-18211.77	18211.77	0.1063	0.2214
4B	4B Cropland	CO <sub>2</sub>	10971.23	10971.23	0.0824	0.3038
4C	4C Grassland	CO <sub>2</sub>	-8861.80	8861.80	0.0739	0.3777
4E	4E Settlements	CO <sub>2</sub>	6451.34	6451.34	0.0538	0.4315
1A	1A Natural Gas	CO <sub>2</sub>	158621.57	158621.57	0.0513	0.4828
3A	3A Enteric Fermentation	CH <sub>4</sub>	21458.44	21458.44	0.0492	0.5319
1A	1A (Stationary) Oil	CO <sub>2</sub>	45156.95	45156.95	0.0489	0.5808
5D	5D Wastewater Handling	N <sub>2</sub> O	725.93	725.93	0.0300	0.6108
1A3b	1A3b DERV	CO <sub>2</sub>	77521.13	77521.13	0.0289	0.6397
2F	2F Product Uses as Substitutes for ODS	HFCs	14166.81	14166.81	0.0276	0.6674
1A3b	1A3b DERV	N <sub>2</sub> O	1000.43	1000.43	0.0217	0.6890
3D	3D Agricultural Soils	N <sub>2</sub> O	11466.79	11466.79	0.0213	0.7104
5B	5B Biological treatment of solid waste	CH <sub>4</sub>	1193.62	1193.62	0.0207	0.7311
2G	2G Other Product Manufacture and Use	N <sub>2</sub> O	843.87	843.87	0.0199	0.7510
1A	1A Coal	CO <sub>2</sub>	34774.45	34774.45	0.0179	0.7689
1A3d	1A3d Marine fuel	CO <sub>2</sub>	5443.73	5443.73	0.0162	0.7850
5D	5D Wastewater Handling	CH <sub>4</sub>	3445.75	3445.75	0.0155	0.8005
4G	4G Other Activities	CO <sub>2</sub>	-2015.60	2015.60	0.0151	0.8156
2B	2B Chemical industries	CO <sub>2</sub>	4931.56	4931.56	0.0144	0.8301
1A1 & 1A2 & 1A4 & 1A5	1A1 & 1A2 & 1A4 & 1A5 Other Combustion	N <sub>2</sub> O	1262.84	1262.84	0.0138	0.8439
1A3b	1A3b Gasoline/ LPG	CO <sub>2</sub>	35843.09	35843.09	0.0133	0.8572
5C	5C Waste Incineration	CO <sub>2</sub>	258.91	258.91	0.0131	0.8703
1B2	1B2 Natural Gas Transmission	CH <sub>4</sub>	3764.68	3764.68	0.0127	0.8830
1A	1A Other (waste)	CO <sub>2</sub>	5441.36	5441.36	0.0126	0.8956

IPCC Code	IPCC Category	Gas	2017 emissions (Gg CO <sub>2</sub> e)	Absolute value of 2017 emissions (Gg CO <sub>2</sub> e)	Level Assessment	Cumulative Total
5B	5B Biological treatment of solid waste	N <sub>2</sub> O	728.03	728.03	0.0115	0.9071
1A1 & 1A2 & 1A4 & 1A5	1A1 & 1A2 & 1A4 & 1A5 Other Combustion	CH <sub>4</sub>	1476.32	1476.32	0.0088	0.9158
4	4 Indirect LULUCF Emissions	N <sub>2</sub> O	243.62	243.62	0.0067	0.9225
1A3	1A3 Other diesel	CO <sub>2</sub>	2597.40	2597.40	0.0063	0.9289
1A3a	1A3a Aviation Fuel	CO <sub>2</sub>	1898.39	1898.39	0.0063	0.9352
3B	3B Manure Management	CH <sub>4</sub>	4226.81	4226.81	0.0059	0.9411
4D	4D Wetland	CO <sub>2</sub>	336.78	336.78	0.0056	0.9467
1B2	1B2 Oil & Natural Gas	CO <sub>2</sub>	4230.64	4230.64	0.0050	0.9516
3B	3B Manure Management	N <sub>2</sub> O	2814.79	2814.79	0.0045	0.9561
1B2	1B2 Offshore Oil& Gas	CH <sub>4</sub>	1170.57	1170.57	0.0037	0.9598
2D	2D Non Energy Products from Fuels and Solvent Use	CO <sub>2</sub>	519.22	519.22	0.0033	0.9631
3G	3G Liming	CO <sub>2</sub>	939.13	939.13	0.0033	0.9664
3H	3H Urea application to agriculture	CO <sub>2</sub>	343.95	343.95	0.0029	0.9692
4B	4B Cropland	N <sub>2</sub> O	461.97	461.97	0.0027	0.9719
3J	3J OT & CD Agriculture	CH <sub>4</sub>	206.19	206.19	0.0024	0.9744
2A1	2A1 Cement Production	CO <sub>2</sub>	4409.79	4409.79	0.0023	0.9767
4A	4A Forest land	N <sub>2</sub> O	144.65	144.65	0.0023	0.9790
2C	2C Metal Industries	CO <sub>2</sub>	2572.86	2572.86	0.0022	0.9812
1B1	1B1 Coal Mining	CH <sub>4</sub>	483.98	483.98	0.0016	0.9828
5C	5C Waste Incineration	N <sub>2</sub> O	39.63	39.63	0.0015	0.9844
1A4	1A4 Petroleum Coke	CO <sub>2</sub>	351.76	351.76	0.0015	0.9858
2G	2G Other Product Manufacture and Use	PFCs	183.51	183.51	0.0014	0.9873
1A3d	1A3d Marine fuel	N <sub>2</sub> O	72.08	72.08	0.0014	0.9887
4E	4E Settlements	N <sub>2</sub> O	524.99	524.99	0.0013	0.9900
1A3b	1A3b Gasoline/ LPG	N <sub>2</sub> O	95.21	95.21	0.0012	0.9912
3J	3J OT & CD Agriculture	N <sub>2</sub> O	93.01	93.01	0.0011	0.9923
1A3b	1A3b Gasoline/ LPG	CH <sub>4</sub>	85.31	85.31	0.0011	0.9933

IPCC Code	IPCC Category	Gas	2017 emissions (Gg CO <sub>2</sub> e)	Absolute value of 2017 emissions (Gg CO <sub>2</sub> e)	Level Assessment	Cumulative Total
2A2	2A2 Lime Production	CO <sub>2</sub>	1052.06	1052.06	0.0009	0.9942
1B2	1B2 Oil & Natural Gas	N <sub>2</sub> O	39.36	39.36	0.0007	0.9949
4C	4C Grassland	N <sub>2</sub> O	33.24	33.24	0.0006	0.9955
1A3	1A3 Other diesel	N <sub>2</sub> O	25.30	25.30	0.0006	0.9961
2G	2G Other Product Manufacture and Use	SF <sub>6</sub>	415.66	415.66	0.0004	0.9965
1B1	1B1 Solid Fuel Transformation	CO <sub>2</sub>	359.54	359.54	0.0004	0.9969
1A3a	1A3a Aviation Fuel	N <sub>2</sub> O	17.96	17.96	0.0003	0.9973
2A3	2A3 Glass production	CO <sub>2</sub>	367.98	367.98	0.0003	0.9976
2B	2B Chemical industry	PFCs	172.90	172.90	0.0003	0.9979
2A4	2A4 Other process uses of carbonates	CO <sub>2</sub>	419.65	419.65	0.0003	0.9981
2B	2B Chemical Industry	CH <sub>4</sub>	61.41	61.41	0.0002	0.9983
1A3b	1A3b DERV	CH <sub>4</sub>	7.85	7.85	0.0002	0.9985
2E	2E Electronics Industry	HFCs	21.18	21.18	0.0002	0.9987
4C	4C Grassland	CH <sub>4</sub>	17.78	17.78	0.0002	0.9988
1A3c	1A3c Coal	CO <sub>2</sub>	41.15	41.15	0.0001	0.9990
1A3d	1A3d Marine fuel	CH <sub>4</sub>	6.28	6.28	0.0001	0.9991
2C	2C Metal Industries	SF <sub>6</sub>	109.75	109.75	0.0001	0.9992
2C	2C Iron & Steel	N <sub>2</sub> O	6.19	6.19	0.0001	0.9993
2A	2A Minerals industry	CH <sub>4</sub>	5.89	5.89	0.0001	0.9994
2C	2C Iron & Steel Production	CH <sub>4</sub>	10.68	10.68	0.0001	0.9995
5C	5C Waste Incineration	CH <sub>4</sub>	10.45	10.45	0.0001	0.9996
4E	4E Settlements	CH <sub>4</sub>	8.25	8.25	0.0001	0.9997
2B2	2B2 Nitric Acid Production	N <sub>2</sub> O	37.25	37.25	0.0001	0.9997
2C	2C Metal Industries	PFCs	15.07	15.07	0.0001	0.9998
1A3	1A3 Other diesel	CH <sub>4</sub>	2.27	2.27	0.0000	0.9998
1B1	1B1 Solid Fuel Transformation	CH <sub>4</sub>	3.78	3.78	0.0000	0.9999
1A4	1A4 Peat	CO <sub>2</sub>	4.83	4.83	0.0000	0.9999
2B8	2B8 Petrochemical and Carbon Black Production	N <sub>2</sub> O	1.51	1.51	0.0000	0.9999
4A	4A Forest Land	CH <sub>4</sub>	2.31	2.31	0.0000	0.9999

IPCC Code	IPCC Category	Gas	2017 emissions (Gg CO <sub>2</sub> e)	Absolute value of 2017 emissions (Gg CO <sub>2</sub> e)	Level Assessment	Cumulative Total
1A3c	1A3c Coal	CH <sub>4</sub>	0.94	0.94	0.0000	1.0000
1A3a	1A3a Aviation Fuel	CH <sub>4</sub>	1.55	1.55	0.0000	1.0000
4D	4D Grassland	N <sub>2</sub> O	0.30	0.30	0.0000	1.0000
2C	2C Metal Industries	HFCs	2.29	2.29	0.0000	1.0000
2B	2B Chemical industry	HFCs	2.55	2.55	0.0000	1.0000
2E	2E Electronics Industry	NF <sub>3</sub>	0.53	0.53	0.0000	1.0000
2B1	2B1 Ammonia Production	N <sub>2</sub> O	0.33	0.33	0.0000	1.0000
1A3c	1A3c Coal	N <sub>2</sub> O	0.09	0.09	0.0000	1.0000
1B1	1B1 Fugitive Emissions from Solid Fuels	N <sub>2</sub> O	0.02	0.02	0.0000	1.0000
4B	4B Cropland	CH <sub>4</sub>	0.02	0.02	0.0000	1.0000
4F	4F Other Land	CO <sub>2</sub>	0.00	0.00	0.0000	1.0000
2D	2D Non-energy Products from Fuels and Solvent Use	CH <sub>4</sub>	0.00	0.00	0.0000	1.0000
3F	3F Field Burning	CH <sub>4</sub>	0.00	0.00	0.0000	1.0000
2B3	2B3 Adipic Acid Production	N <sub>2</sub> O	0.00	0.00	0.0000	1.0000
2D	2D Non-energy Products from Fuels and Solvent Use	N <sub>2</sub> O	0.00	0.00	0.0000	1.0000
3F	3F Field Burning	N <sub>2</sub> O	0.00	0.00	0.0000	1.0000
2F	2F Product Uses as Substitutes for ODS	PFCs	0.00	0.00	0.0000	1.0000

**Table A 1.4.3 Approach 2 Level Assessment for Base year (not including LULUCF) with Key Categories Shaded in Grey**

IPCC Code	IPCC Category	Gas	Base year emissions (Gg CO <sub>2</sub> e)	Absolute value of Base year emissions (Gg CO <sub>2</sub> e)	Level Assessment	Cumulative Total
5A	5A Solid Waste Disposal	CH <sub>4</sub>	60432.98	60432.98	0.2904	0.2904
2B3	2B3 Adipic Acid Production	N <sub>2</sub> O	19934.61	19934.61	0.1980	0.4885
1A	1A Coal	CO <sub>2</sub>	243845.39	243845.39	0.0748	0.5632
1A	1A (Stationary) Oil	CO <sub>2</sub>	95612.77	95612.77	0.0616	0.6249
1B1	1B1 Coal Mining	CH <sub>4</sub>	21826.68	21826.68	0.0436	0.6684
5C	5C Waste Incineration	CO <sub>2</sub>	1300.71	1300.71	0.0391	0.7075
3A	3A Enteric Fermentation	CH <sub>4</sub>	25392.50	25392.50	0.0346	0.7422

IPCC Code	IPCC Category	Gas	Base year emissions (Gg CO <sub>2</sub> e)	Absolute value of Base emissions year (Gg CO <sub>2</sub> e)	Level Assessment	Cumulative Total
1A	1A Natural Gas	CO <sub>2</sub>	106932.90	106932.90	0.0206	0.7628
1B2	1B2 Natural Gas Transmission	CH <sub>4</sub>	10168.33	10168.33	0.0204	0.7832
5D	5D Wastewater Handling	N <sub>2</sub> O	783.59	783.59	0.0193	0.8025
2B	2B Chemical industry	HFCs	17670.77	17670.77	0.0176	0.8201
1A3b	1A3b Gasoline/ LPG	CO <sub>2</sub>	75562.72	75562.72	0.0167	0.8368
3D	3D Agricultural Soils	N <sub>2</sub> O	13610.39	13610.39	0.0151	0.8518
1A1 & 1A2 & 1A4 & 1A5	1A1 & 1A2 & 1A4 & 1A5 Other Combustion	N <sub>2</sub> O	2141.02	2141.02	0.0140	0.8658
1A3d	1A3d Marine fuel	CO <sub>2</sub>	7611.13	7611.13	0.0135	0.8793
2B	2B Chemical industries	CO <sub>2</sub>	6770.09	6770.09	0.0118	0.8911
5D	5D Wastewater Handling	CH <sub>4</sub>	4218.90	4218.90	0.0113	0.9023
1A3b	1A3b Gasoline/ LPG	CH <sub>4</sub>	1157.96	1157.96	0.0086	0.9110
2G	2G Other Product Manufacture and Use	N <sub>2</sub> O	593.57	593.57	0.0083	0.9193
1A3b	1A3b DERV	CO <sub>2</sub>	33005.80	33005.80	0.0073	0.9266
1A3b	1A3b Gasoline/ LPG	N <sub>2</sub> O	989.40	989.40	0.0073	0.9340
1A1 & 1A2 & 1A4 & 1A5	1A1 & 1A2 & 1A4 & 1A5 Other Combustion	CH <sub>4</sub>	1893.15	1893.15	0.0067	0.9406
1A3b	1A3b DERV	N <sub>2</sub> O	322.88	322.88	0.0042	0.9448
1B2	1B2 Offshore Oil& Gas	CH <sub>4</sub>	2176.74	2176.74	0.0040	0.9489
1B2	1B2 Oil & Natural Gas	CO <sub>2</sub>	5777.92	5777.92	0.0040	0.9529
3B	3B Manure Management	CH <sub>4</sub>	4732.53	4732.53	0.0039	0.9568
2C	2C Metal Industries	CO <sub>2</sub>	7400.69	7400.69	0.0038	0.9607
2B2	2B2 Nitric Acid Production	N <sub>2</sub> O	3860.26	3860.26	0.0038	0.9645
1A3a	1A3a Aviation Fuel	CO <sub>2</sub>	1881.31	1881.31	0.0037	0.9682
3B	3B Manure Management	N <sub>2</sub> O	3442.70	3442.70	0.0033	0.9715
1A3	1A3 Other diesel	CO <sub>2</sub>	1679.92	1679.92	0.0024	0.9739
2A1	2A1 Cement Production	CO <sub>2</sub>	7295.26	7295.26	0.0023	0.9762
3G	3G Liming	CO <sub>2</sub>	1015.18	1015.18	0.0021	0.9783

IPCC Code	IPCC Category	Gas	Base year emissions (Gg CO <sub>2</sub> e)	Absolute value of Base emissions year (Gg CO <sub>2</sub> e)	Level Assessment	Cumulative Total
2D	2D Non Energy Products from Fuels and Solvent Use	CO <sub>2</sub>	552.81	552.81	0.0021	0.9804
3J	3J OT & CD Agriculture	CH <sub>4</sub>	276.73	276.73	0.0019	0.9824
2F	2F Product Uses as Substitutes for ODS	HFCs	1416.64	1416.64	0.0016	0.9840
3H	3H Urea application to agriculture	CO <sub>2</sub>	327.60	327.60	0.0016	0.9856
1A3d	1A3d Marine fuel	N <sub>2</sub> O	105.00	105.00	0.0012	0.9868
1B1	1B1 Solid Fuel Transformation	CO <sub>2</sub>	1698.56	1698.56	0.0012	0.9880
1A4	1A4 Peat	CO <sub>2</sub>	372.48	372.48	0.0012	0.9892
1A3b	1A3b DERV	CH <sub>4</sub>	88.74	88.74	0.0011	0.9903
5C	5C Waste Incineration	N <sub>2</sub> O	46.66	46.66	0.0011	0.9914
3J	3J OT & CD Agriculture	N <sub>2</sub> O	132.00	132.00	0.0009	0.9923
2A2	2A2 Lime Production	CO <sub>2</sub>	1462.05	1462.05	0.0007	0.9931
2G	2G Other Product Manufacture and Use	PFCs	149.16	149.16	0.0007	0.9938
2C	2C Metal Industries	PFCs	333.43	333.43	0.0007	0.9944
5C	5C Waste Incineration	CH <sub>4</sub>	134.83	134.83	0.0006	0.9950
2G	2G Other Product Manufacture and Use	SF <sub>6</sub>	912.79	912.79	0.0006	0.9956
3F	3F Field Burning	CH <sub>4</sub>	186.57	186.57	0.0005	0.9961
1B2	1B2 Oil & Natural Gas	N <sub>2</sub> O	40.75	40.75	0.0004	0.9965
2B	2B Chemical Industry	CH <sub>4</sub>	204.89	204.89	0.0004	0.9969
1A	1A Other (waste)	CO <sub>2</sub>	247.13	247.13	0.0003	0.9972
2A	2A Minerals industry	CH <sub>4</sub>	31.10	31.10	0.0003	0.9975
2C	2C Metal Industries	SF <sub>6</sub>	387.17	387.17	0.0003	0.9978
2A4	2A4 Other process uses of carbonates	CO <sub>2</sub>	640.93	640.93	0.0002	0.9980
1A3	1A3 Other diesel	N <sub>2</sub> O	16.53	16.53	0.0002	0.9983
2C	2C Iron & Steel	N <sub>2</sub> O	17.74	17.74	0.0002	0.9985
1A4	1A4 Petroleum Coke	CO <sub>2</sub>	81.64	81.64	0.0002	0.9987
2A3	2A3 Glass production	CO <sub>2</sub>	405.54	405.54	0.0002	0.9989
1A3a	1A3a Aviation Fuel	N <sub>2</sub> O	17.80	17.80	0.0002	0.9991

IPCC Code	IPCC Category	Gas	Base year emissions (Gg CO <sub>2e</sub> )	Absolute value of Base year emissions (Gg CO <sub>2e</sub> )	Level Assessment	Cumulative Total
5B	5B Biological treatment of solid waste	CH <sub>4</sub>	18.13	18.13	0.0002	0.9993
2C	2C Iron & Steel Production	CH <sub>4</sub>	36.90	36.90	0.0002	0.9994
3F	3F Field Burning	N <sub>2</sub> O	57.66	57.66	0.0001	0.9996
5B	5B Biological treatment of solid waste	N <sub>2</sub> O	12.97	12.97	0.0001	0.9997
2B	2B Chemical industry	PFCs	113.90	113.90	0.0001	0.9998
1A3d	1A3d Marine fuel	CH <sub>4</sub>	3.66	3.66	0.0000	0.9999
2E	2E Electronics Industry	HFCs	8.73	8.73	0.0000	0.9999
1A3a	1A3a Aviation Fuel	CH <sub>4</sub>	6.37	6.37	0.0000	0.9999
1A3	1A3 Other diesel	CH <sub>4</sub>	2.75	2.75	0.0000	1.0000
2B8	2B8 Petrochemical and Carbon Black Production	N <sub>2</sub> O	2.10	2.10	0.0000	1.0000
2E	2E Electronics Industry	NF <sub>3</sub>	0.83	0.83	0.0000	1.0000
2B1	2B1 Ammonia Production	N <sub>2</sub> O	0.31	0.31	0.0000	1.0000
2F	2F Product Uses as Substitutes for ODS	PFCs	0.44	0.44	0.0000	1.0000
1B1	1B1 Fugitive Emissions from Solid Fuels	N <sub>2</sub> O	0.09	0.09	0.0000	1.0000
1B1	1B1 Solid Fuel Transformation	CH <sub>4</sub>	0.18	0.18	0.0000	1.0000
1A3c	1A3c Coal	CO <sub>2</sub>	0.00	0.00	0.0000	1.0000
1A3c	1A3c Coal	CH <sub>4</sub>	0.00	0.00	0.0000	1.0000
2D	2D Non-energy Products from Fuels and Solvent Use	CH <sub>4</sub>	0.00	0.00	0.0000	1.0000
1A3c	1A3c Coal	N <sub>2</sub> O	0.00	0.00	0.0000	1.0000
2D	2D Non-energy Products from Fuels and Solvent Use	N <sub>2</sub> O	0.00	0.00	0.0000	1.0000
2C	2C Metal Industries	HFCs	0.00	0.00	0.0000	1.0000

**Table A 1.4.4 Approach 2 Level Assessment for the latest reported year (not including LULUCF) with Key Categories Shaded in Grey**

IPCC Code	IPCC Category	Gas	2017 emissions (Gg CO <sub>2e</sub> )	Absolute value of 2017 emissions (Gg CO <sub>2e</sub> )	Level Assessment	Cumulative Total
5A	5A Solid Waste Disposal	CH <sub>4</sub>	14261.40	14261.40	0.1773	0.1773

IPCC Code	IPCC Category	Gas	2017 emissions (Gg CO <sub>2</sub> e)	Absolute value of 2017 emissions (Gg CO <sub>2</sub> e)	Level Assessment	Cumulative Total
1A	1A Natural Gas	CO <sub>2</sub>	158621.57	158621.57	0.0790	0.2563
3A	3A Enteric Fermentation	CH <sub>4</sub>	21458.44	21458.44	0.0757	0.3321
1A	1A (Stationary) Oil	CO <sub>2</sub>	45156.95	45156.95	0.0753	0.4074
5D	5D Wastewater Handling	N <sub>2</sub> O	725.93	725.93	0.0463	0.4537
1A3b	1A3b DERV	CO <sub>2</sub>	77521.13	77521.13	0.0445	0.4982
2F	2F Product Uses as Substitutes for ODS	HFCs	14166.81	14166.81	0.0426	0.5408
1A3b	1A3b DERV	N <sub>2</sub> O	1000.43	1000.43	0.0334	0.5742
3D	3D Agricultural Soils	N <sub>2</sub> O	11466.79	11466.79	0.0329	0.6071
5B	5B Biological treatment of solid waste	CH <sub>4</sub>	1193.62	1193.62	0.0319	0.6390
2G	2G Other Product Manufacture and Use	N <sub>2</sub> O	843.87	843.87	0.0307	0.6697
1A	1A Coal	CO <sub>2</sub>	34774.45	34774.45	0.0276	0.6973
1A3d	1A3d Marine fuel	CO <sub>2</sub>	5443.73	5443.73	0.0249	0.7222
5D	5D Wastewater Handling	CH <sub>4</sub>	3445.75	3445.75	0.0238	0.7460
2B	2B Chemical industries	CO <sub>2</sub>	4931.56	4931.56	0.0222	0.7682
1A1 & 1A2 & 1A4 & 1A5	1A1 & 1A2 & 1A4 & 1A5 Other Combustion	N <sub>2</sub> O	1262.84	1262.84	0.0213	0.7895
1A3b	1A3b Gasoline/ LPG	CO <sub>2</sub>	35843.09	35843.09	0.0205	0.8100
5C	5C Waste Incineration	CO <sub>2</sub>	258.91	258.91	0.0201	0.8302
1B2	1B2 Natural Gas Transmission	CH <sub>4</sub>	3764.68	3764.68	0.0196	0.8497
1A	1A Other (waste)	CO <sub>2</sub>	5441.36	5441.36	0.0194	0.8692
5B	5B Biological treatment of solid waste	N <sub>2</sub> O	728.03	728.03	0.0177	0.8869
1A1 & 1A2 & 1A4 & 1A5	1A1 & 1A2 & 1A4 & 1A5 Other Combustion	CH <sub>4</sub>	1476.32	1476.32	0.0135	0.9004
1A3	1A3 Other diesel	CO <sub>2</sub>	2597.40	2597.40	0.0097	0.9102
1A3a	1A3a Aviation Fuel	CO <sub>2</sub>	1898.39	1898.39	0.0097	0.9199
3B	3B Manure Management	CH <sub>4</sub>	4226.81	4226.81	0.0091	0.9290
1B2	1B2 Oil & Natural Gas	CO <sub>2</sub>	4230.64	4230.64	0.0076	0.9366

IPCC Code	IPCC Category	Gas	2017 emissions (Gg CO <sub>2</sub> e)	Absolute value of 2017 emissions (Gg CO <sub>2</sub> e)	Level Assessment	Cumulative Total
3B	3B Manure Management	N <sub>2</sub> O	2814.79	2814.79	0.0069	0.9435
1B2	1B2 Offshore Oil& Gas	CH <sub>4</sub>	1170.57	1170.57	0.0056	0.9491
2D	2D Non Energy Products from Fuels and Solvent Use	CO <sub>2</sub>	519.22	519.22	0.0051	0.9542
3G	3G Liming	CO <sub>2</sub>	939.13	939.13	0.0050	0.9593
3H	3H Urea application to agriculture	CO <sub>2</sub>	343.95	343.95	0.0044	0.9637
3J	3J OT & CD Agriculture	CH <sub>4</sub>	206.19	206.19	0.0037	0.9674
2A1	2A1 Cement Production	CO <sub>2</sub>	4409.79	4409.79	0.0036	0.9710
2C	2C Metal Industries	CO <sub>2</sub>	2572.86	2572.86	0.0035	0.9745
1B1	1B1 Coal Mining	CH <sub>4</sub>	483.98	483.98	0.0025	0.9770
5C	5C Waste Incineration	N <sub>2</sub> O	39.63	39.63	0.0023	0.9793
1A4	1A4 Petroleum Coke	CO <sub>2</sub>	351.76	351.76	0.0023	0.9816
2G	2G Other Product Manufacture and Use	PFCs	183.51	183.51	0.0022	0.9838
1A3d	1A3d Marine fuel	N <sub>2</sub> O	72.08	72.08	0.0021	0.9859
1A3b	1A3b Gasoline/ LPG	N <sub>2</sub> O	95.21	95.21	0.0018	0.9878
3J	3J OT & CD Agriculture	N <sub>2</sub> O	93.01	93.01	0.0017	0.9895
1A3b	1A3b Gasoline/ LPG	CH <sub>4</sub>	85.31	85.31	0.0016	0.9911
2A2	2A2 Lime Production	CO <sub>2</sub>	1052.06	1052.06	0.0014	0.9925
1B2	1B2 Oil & Natural Gas	N <sub>2</sub> O	39.36	39.36	0.0011	0.9935
1A3	1A3 Other diesel	N <sub>2</sub> O	25.30	25.30	0.0009	0.9944
2G	2G Other Product Manufacture and Use	SF <sub>6</sub>	415.66	415.66	0.0007	0.9950
1B1	1B1 Solid Fuel Transformation	CO <sub>2</sub>	359.54	359.54	0.0007	0.9957
1A3a	1A3a Aviation Fuel	N <sub>2</sub> O	17.96	17.96	0.0005	0.9962
2A3	2A3 Glass production	CO <sub>2</sub>	367.98	367.98	0.0005	0.9967
2B	2B Chemical industry	PFCs	172.90	172.90	0.0004	0.9971
2A4	2A4 Other process uses of carbonates	CO <sub>2</sub>	419.65	419.65	0.0004	0.9975
2B	2B Chemical Industry	CH <sub>4</sub>	61.41	61.41	0.0003	0.9978
1A3b	1A3b DERV	CH <sub>4</sub>	7.85	7.85	0.0003	0.9981
2E	2E Electronics Industry	HFCs	21.18	21.18	0.0003	0.9983

IPCC Code	IPCC Category	Gas	2017 emissions (Gg CO <sub>2</sub> e)	Absolute value of 2017 emissions (Gg CO <sub>2</sub> e)	Level Assessment	Cumulative Total
1A3c	1A3c Coal	CO <sub>2</sub>	41.15	41.15	0.0002	0.9986
1A3d	1A3d Marine fuel	CH <sub>4</sub>	6.28	6.28	0.0002	0.9988
2C	2C Metal Industries	SF <sub>6</sub>	109.75	109.75	0.0002	0.9990
2C	2C Iron & Steel	N <sub>2</sub> O	6.19	6.19	0.0002	0.9991
2A	2A Minerals industry	CH <sub>4</sub>	5.89	5.89	0.0002	0.9993
2C	2C Iron & Steel Production	CH <sub>4</sub>	10.68	10.68	0.0001	0.9994
5C	5C Waste Incineration	CH <sub>4</sub>	10.45	10.45	0.0001	0.9995
2B2	2B2 Nitric Acid Production	N <sub>2</sub> O	37.25	37.25	0.0001	0.9996
2C	2C Metal Industries	PFCs	15.07	15.07	0.0001	0.9997
1A3	1A3 Other diesel	CH <sub>4</sub>	2.27	2.27	0.0001	0.9998
1B1	1B1 Solid Fuel Transformation	CH <sub>4</sub>	3.78	3.78	0.0000	0.9998
1A4	1A4 Peat	CO <sub>2</sub>	4.83	4.83	0.0000	0.9999
2B8	2B8 Petrochemical and Carbon Black Production	N <sub>2</sub> O	1.51	1.51	0.0000	0.9999
1A3c	1A3c Coal	CH <sub>4</sub>	0.94	0.94	0.0000	1.0000
1A3a	1A3a Aviation Fuel	CH <sub>4</sub>	1.55	1.55	0.0000	1.0000
2C	2C Metal Industries	HFCs	2.29	2.29	0.0000	1.0000
2B	2B Chemical industry	HFCs	2.55	2.55	0.0000	1.0000
2E	2E Electronics Industry	NF <sub>3</sub>	0.53	0.53	0.0000	1.0000
2B1	2B1 Ammonia Production	N <sub>2</sub> O	0.33	0.33	0.0000	1.0000
1A3c	1A3c Coal	N <sub>2</sub> O	0.09	0.09	0.0000	1.0000
1B1	1B1 Fugitive Emissions from Solid Fuels	N <sub>2</sub> O	0.02	0.02	0.0000	1.0000
2D	2D Non-energy Products from Fuels and Solvent Use	CH <sub>4</sub>	0.00	0.00	0.0000	1.0000
3F	3F Field Burning	CH <sub>4</sub>	0.00	0.00	0.0000	1.0000
2B3	2B3 Adipic Acid Production	N <sub>2</sub> O	0.00	0.00	0.0000	1.0000
2D	2D Non-energy Products from Fuels and Solvent Use	N <sub>2</sub> O	0.00	0.00	0.0000	1.0000
3F	3F Field Burning	N <sub>2</sub> O	0.00	0.00	0.0000	1.0000
2F	2F Product Uses as Substitutes for ODS	PFCs	0.00	0.00	0.0000	1.0000

**Table A 1.4.5 Approach 2 Assessment for Trend (including LULUCF) with Key Categories Shaded in Grey**

IPCC Code	IPCC Category	Gas	Base year emissions (Gg CO <sub>2</sub> e)	2017 emissions (Gg CO <sub>2</sub> e)	Trend Assessment with Uncertainty	% Contribution to Trend Uncertainty	Cumulative Total
2B3	2B3 Adipic Acid Production	N <sub>2</sub> O	19934.61	0.00	0.0136	23.0%	0.2302
5A	5A Solid Waste Disposal	CH <sub>4</sub>	60432.98	14261.40	0.0118	20.0%	0.4303
1A	1A Coal	CO <sub>2</sub>	243845.39	34774.45	0.0039	6.6%	0.4958
1B1	1B1 Coal Mining	CH <sub>4</sub>	21826.68	483.98	0.0029	4.9%	0.5445
1A	1A Natural Gas	CO <sub>2</sub>	106932.90	158621.57	0.0022	3.7%	0.5819
2F	2F Product Uses as Substitutes for ODS	HFCs	1416.64	14166.81	0.0018	3.1%	0.6130
5C	5C Waste Incineration	CO <sub>2</sub>	1300.71	258.91	0.0018	3.0%	0.6428
1A3b	1A3b DERV	CO <sub>2</sub>	33005.80	77521.13	0.0015	2.6%	0.6689
5B	5B Biological treatment of solid waste	CH <sub>4</sub>	18.13	1193.62	0.0014	2.5%	0.6934
4B	4B Cropland	CO <sub>2</sub>	14265.93	10971.23	0.0014	2.4%	0.7177
4E	4E Settlements	CO <sub>2</sub>	7011.30	6451.34	0.0014	2.4%	0.7415
4A	4A Forest Land	CO <sub>2</sub>	-15026.26	-18211.77	0.0013	2.2%	0.7634
1A3b	1A3b DERV	N <sub>2</sub> O	322.88	1000.43	0.0012	2.1%	0.7845
2B	2B Chemical industry	HFCs	17670.77	2.55	0.0012	2.0%	0.8049
3A	3A Enteric Fermentation	CH <sub>4</sub>	25392.50	21458.44	0.0011	1.8%	0.8234
1A	1A Other (waste)	CO <sub>2</sub>	247.13	5441.36	0.0009	1.5%	0.8381
2G	2G Other Product Manufacture and Use	N <sub>2</sub> O	593.57	843.87	0.0008	1.4%	0.8522
5B	5B Biological treatment of solid waste	N <sub>2</sub> O	12.97	728.03	0.0008	1.4%	0.8658
5D	5D Wastewater Handling	N <sub>2</sub> O	783.59	725.93	0.0008	1.3%	0.8792
1A	1A (Stationary) Oil	CO <sub>2</sub>	95612.77	45156.95	0.0008	1.3%	0.8925
4C	4C Grassland	CO <sub>2</sub>	-7111.03	-8861.80	0.0007	1.2%	0.9048
1A3b	1A3b Gasoline/ LPG	CH <sub>4</sub>	1157.96	85.31	0.0005	0.9%	0.9136
1B2	1B2 Natural Gas Transmission	CH <sub>4</sub>	10168.33	3764.68	0.0005	0.9%	0.9222
3D	3D Agricultural Soils	N <sub>2</sub> O	13610.39	11466.79	0.0005	0.8%	0.9301

IPCC Code	IPCC Category	Gas	Base year emissions (Gg CO <sub>2</sub> e)	2017 emissions (Gg CO <sub>2</sub> e)	Trend Assessment with Uncertainty	% Contribution to Trend Uncertainty	Cumulative Total
1A3b	1A3b Gasoline/ LPG	N <sub>2</sub> O	989.40	95.21	0.0004	0.7%	0.9372
5D	5D Wastewater Handling	CH <sub>4</sub>	4218.90	3445.75	0.0003	0.5%	0.9426
1A3	1A3 Other diesel	CO <sub>2</sub>	1679.92	2597.40	0.0003	0.5%	0.9473
2B2	2B2 Nitric Acid Production	N <sub>2</sub> O	3860.26	37.25	0.0003	0.4%	0.9517
1A3d	1A3d Marine fuel	CO <sub>2</sub>	7611.13	5443.73	0.0002	0.4%	0.9554
2B	2B Chemical industries	CO <sub>2</sub>	6770.09	4931.56	0.0002	0.4%	0.9589
1A3b	1A3b Gasoline/ LPG	CO <sub>2</sub>	75562.72	35843.09	0.0002	0.4%	0.9624
1A3a	1A3a Aviation Fuel	CO <sub>2</sub>	1881.31	1898.39	0.0002	0.3%	0.9656
4G	4G Other Activities	CO <sub>2</sub>	-1639.08	-2015.60	0.0002	0.3%	0.9685
1A1 & 1A2 & 1A4 & 1A5	1A1 & 1A2 & 1A4 & 1A5 Other Combustion	CH <sub>4</sub>	1893.15	1476.32	0.0002	0.3%	0.9711
3B	3B Manure Management	CH <sub>4</sub>	4732.53	4226.81	0.0001	0.2%	0.9736
2C	2C Metal Industries	CO <sub>2</sub>	7400.69	2572.86	0.0001	0.2%	0.9754
3B	3B Manure Management	N <sub>2</sub> O	3442.70	2814.79	0.0001	0.2%	0.9770
3H	3H Urea application to agriculture	CO <sub>2</sub>	327.60	343.95	0.0001	0.2%	0.9785
2D	2D Non Energy Products from Fuels and Solvent Use	CO <sub>2</sub>	552.81	519.22	0.0001	0.2%	0.9800
1A4	1A4 Petroleum Coke	CO <sub>2</sub>	81.64	351.76	0.0001	0.2%	0.9815
3G	3G Liming	CO <sub>2</sub>	1015.18	939.13	0.0001	0.1%	0.9830
1A4	1A4 Peat	CO <sub>2</sub>	372.48	4.83	0.0001	0.1%	0.9843
1B2	1B2 Oil & Natural Gas	CO <sub>2</sub>	5777.92	4230.64	0.0001	0.1%	0.9856
1A3b	1A3b DERV	CH <sub>4</sub>	88.74	7.85	0.0001	0.1%	0.9867
4D	4D Wetland	CO <sub>2</sub>	486.95	336.78	0.0001	0.1%	0.9878
2G	2G Other Product Manufacture and Use	PFCs	149.16	183.51	0.0001	0.1%	0.9887
4B	4B Cropland	N <sub>2</sub> O	1019.85	461.97	0.0001	0.1%	0.9896
1B1	1B1 Solid Fuel Transformation	CO <sub>2</sub>	1698.56	359.54	0.0001	0.1%	0.9905
2C	2C Metal Industries	PFCs	333.43	15.07	0.0000	0.1%	0.9912

IPCC Code	IPCC Category	Gas	Base year emissions (Gg CO <sub>2</sub> e)	2017 emissions (Gg CO <sub>2</sub> e)	Trend Assessment with Uncertainty	% Contribution to Trend Uncertainty	Cumulative Total
3J	3J OT & CD Agriculture	CH <sub>4</sub>	276.73	206.19	0.0000	0.1%	0.9918
4C	4C Grassland	N <sub>2</sub> O	10.33	33.24	0.0000	0.1%	0.9925
5C	5C Waste Incineration	CH <sub>4</sub>	134.83	10.45	0.0000	0.1%	0.9931
5C	5C Waste Incineration	N <sub>2</sub> O	46.66	39.63	0.0000	0.1%	0.9937
4E	4E Settlements	N <sub>2</sub> O	584.68	524.99	0.0000	0.1%	0.9942
3F	3F Field Burning	CH <sub>4</sub>	186.57	0.00	0.0000	0.1%	0.9948
1A3	1A3 Other diesel	N <sub>2</sub> O	16.53	25.30	0.0000	0.0%	0.9952
4	4 Indirect LULUCF Emissions	N <sub>2</sub> O	401.49	243.62	0.0000	0.0%	0.9955
1B2	1B2 Offshore Oil & Gas	CH <sub>4</sub>	2176.74	1170.57	0.0000	0.0%	0.9959
1B2	1B2 Oil & Natural Gas	N <sub>2</sub> O	40.75	39.36	0.0000	0.0%	0.9962
1A1 & 1A2 & 1A4 & 1A5	1A1 & 1A2 & 1A4 & 1A5 Other Combustion	N <sub>2</sub> O	2141.02	1262.84	0.0000	0.0%	0.9965
1A3d	1A3d Marine fuel	N <sub>2</sub> O	105.00	72.08	0.0000	0.0%	0.9968
2A	2A Minerals industry	CH <sub>4</sub>	31.10	5.89	0.0000	0.0%	0.9970
3J	3J OT & CD Agriculture	N <sub>2</sub> O	132.00	93.01	0.0000	0.0%	0.9972
2B	2B Chemical Industry	CH <sub>4</sub>	204.89	61.41	0.0000	0.0%	0.9975
2B	2B Chemical industry	PFCs	113.90	172.90	0.0000	0.0%	0.9977
2A2	2A2 Lime Production	CO <sub>2</sub>	1462.05	1052.06	0.0000	0.0%	0.9979
4A	4A Forest land	N <sub>2</sub> O	231.22	144.65	0.0000	0.0%	0.9981
1A3c	1A3c Coal	CO <sub>2</sub>	0.00	41.15	0.0000	0.0%	0.9983
3F	3F Field Burning	N <sub>2</sub> O	57.66	0.00	0.0000	0.0%	0.9984
1A3a	1A3a Aviation Fuel	N <sub>2</sub> O	17.80	17.96	0.0000	0.0%	0.9986
2C	2C Metal Industries	SF <sub>6</sub>	387.17	109.75	0.0000	0.0%	0.9988
2E	2E Electronics Industry	HFCs	8.73	21.18	0.0000	0.0%	0.9989
2G	2G Other Product Manufacture and Use	SF <sub>6</sub>	912.79	415.66	0.0000	0.0%	0.9990
2A3	2A3 Glass production	CO <sub>2</sub>	405.54	367.98	0.0000	0.0%	0.9992
4C	4C Grassland	CH <sub>4</sub>	9.97	17.78	0.0000	0.0%	0.9993
2A1	2A1 Cement Production	CO <sub>2</sub>	7295.26	4409.79	0.0000	0.0%	0.9994
1A3d	1A3d Marine fuel	CH <sub>4</sub>	3.66	6.28	0.0000	0.0%	0.9995

IPCC Code	IPCC Category	Gas	Base year emissions (Gg CO <sub>2</sub> e)	2017 emissions (Gg CO <sub>2</sub> e)	Trend Assessment with Uncertainty	% Contribution to Trend Uncertainty	Cumulative Total
2C	2C Iron & Steel Production	CH <sub>4</sub>	36.90	10.68	0.0000	0.0%	0.9996
2C	2C Iron & Steel	N <sub>2</sub> O	17.74	6.19	0.0000	0.0%	0.9997
4E	4E Settlements	CH <sub>4</sub>	3.00	8.25	0.0000	0.0%	0.9998
4D	4D Grassland	N <sub>2</sub> O	4.13	0.30	0.0000	0.0%	0.9998
1B1	1B1 Solid Fuel Transformation	CH <sub>4</sub>	0.18	3.78	0.0000	0.0%	0.9999
2A4	2A4 Other process uses of carbonates	CO <sub>2</sub>	640.93	419.65	0.0000	0.0%	0.9999
1A3a	1A3a Aviation Fuel	CH <sub>4</sub>	6.37	1.55	0.0000	0.0%	0.9999
1A3c	1A3c Coal	CH <sub>4</sub>	0.00	0.94	0.0000	0.0%	1.0000
1A3	1A3 Other diesel	CH <sub>4</sub>	2.75	2.27	0.0000	0.0%	1.0000
4A	4A Forest Land	CH <sub>4</sub>	2.89	2.31	0.0000	0.0%	1.0000
2B8	2B8 Petrochemical and Carbon Black Production	N <sub>2</sub> O	2.10	1.51	0.0000	0.0%	1.0000
2C	2C Metal Industries	HFCs	0.00	2.29	0.0000	0.0%	1.0000
1A3c	1A3c Coal	N <sub>2</sub> O	0.00	0.09	0.0000	0.0%	1.0000
2B1	2B1 Ammonia Production	N <sub>2</sub> O	0.31	0.33	0.0000	0.0%	1.0000
2F	2F Product Uses as Substitutes for ODS	PFCs	0.44	0.00	0.0000	0.0%	1.0000
1B1	1B1 Fugitive Emissions from Solid Fuels	N <sub>2</sub> O	0.09	0.02	0.0000	0.0%	1.0000
2E	2E Electronics Industry	NF <sub>3</sub>	0.83	0.53	0.0000	0.0%	1.0000
4B	4B Cropland	CH <sub>4</sub>	0.09	0.02	0.0000	0.0%	1.0000
4F	4F Other Land	CO <sub>2</sub>	0.00	0.00	0.0000	0.0%	1.0000
2D	2D Non-energy Products from Fuels and Solvent Use	CH <sub>4</sub>	0.00	0.00	0.0000	0.0%	1.0000
2D	2D Non-energy Products from Fuels and Solvent Use	N <sub>2</sub> O	0.00	0.00	0.0000	0.0%	1.0000

**Table A 1.4.6 Approach 2 Assessment for Trend (not including LULUCF) with Key Categories Shaded in Grey**

IPCC Code	IPCC Category	Gas	Base year emissions (Gg CO <sub>2</sub> e)	2017 emissions (Gg CO <sub>2</sub> e)	Trend Assessment with Uncertainty	% Contribution to Trend Uncertainty	Cumulative Total
2B3	2B3 Adipic Acid Production	N <sub>2</sub> O	19934.61	0.00	0.0136	25.3%	0.2528
5A	5A Solid Waste Disposal	CH <sub>4</sub>	60432.98	14261.40	0.0118	22.0%	0.4724
1A	1A Coal	CO <sub>2</sub>	243845.39	34774.45	0.0039	7.2%	0.5444
1B1	1B1 Coal Mining	CH <sub>4</sub>	21826.68	483.98	0.0029	5.3%	0.5979
1A	1A Natural Gas	CO <sub>2</sub>	106932.90	158621.57	0.0022	4.1%	0.6389
2F	2F Product Uses as Substitutes for ODS	HFCs	1416.64	14166.81	0.0018	3.4%	0.6730
5C	5C Waste Incineration	CO <sub>2</sub>	1300.71	258.91	0.0018	3.3%	0.7058
1A3b	1A3b DERV	CO <sub>2</sub>	33005.80	77521.13	0.0015	2.9%	0.7344
5B	5B Biological treatment of solid waste	CH <sub>4</sub>	18.13	1193.62	0.0014	2.7%	0.7613
1A3b	1A3b DERV	N <sub>2</sub> O	322.88	1000.43	0.0012	2.3%	0.7844
2B	2B Chemical industry	HFCs	17670.77	2.55	0.0012	2.2%	0.8068
3A	3A Enteric Fermentation	CH <sub>4</sub>	25392.50	21458.44	0.0011	2.0%	0.8271
1A	1A Other (waste)	CO <sub>2</sub>	247.13	5441.36	0.0009	1.6%	0.8432
2G	2G Other Product Manufacture and Use	N <sub>2</sub> O	593.57	843.87	0.0008	1.5%	0.8587
5B	5B Biological treatment of solid waste	N <sub>2</sub> O	12.97	728.03	0.0008	1.5%	0.8737
5D	5D Wastewater Handling	N <sub>2</sub> O	783.59	725.93	0.0008	1.5%	0.8885
1A	1A (Stationary) Oil	CO <sub>2</sub>	95612.77	45156.95	0.0008	1.5%	0.9030
1A3b	1A3b Gasoline/ LPG	CH <sub>4</sub>	1157.96	85.31	0.0005	1.0%	0.9126
1B2	1B2 Natural Gas Transmission	CH <sub>4</sub>	10168.33	3764.68	0.0005	0.9%	0.9220
3D	3D Agricultural Soils	N <sub>2</sub> O	13610.39	11466.79	0.0005	0.9%	0.9308
1A3b	1A3b Gasoline/ LPG	N <sub>2</sub> O	989.40	95.21	0.0004	0.8%	0.9385
5D	5D Wastewater Handling	CH <sub>4</sub>	4218.90	3445.75	0.0003	0.6%	0.9444
1A3	1A3 Other diesel	CO <sub>2</sub>	1679.92	2597.40	0.0003	0.5%	0.9496

IPCC Code	IPCC Category	Gas	Base year emissions (Gg CO <sub>2</sub> e)	2017 emissions (Gg CO <sub>2</sub> e)	Trend Assessment with Uncertainty	% Contribution to Trend Uncertainty	Cumulative Total
2B2	2B2 Nitric Acid Production	N <sub>2</sub> O	3860.26	37.25	0.0003	0.5%	0.9544
1A3d	1A3d Marine fuel	CO <sub>2</sub>	7611.13	5443.73	0.0002	0.4%	0.9585
2B	2B Chemical industries	CO <sub>2</sub>	6770.09	4931.56	0.0002	0.4%	0.9624
1A3b	1A3b Gasoline/ LPG	CO <sub>2</sub>	75562.72	35843.09	0.0002	0.4%	0.9662
1A3a	1A3a Aviation Fuel	CO <sub>2</sub>	1881.31	1898.39	0.0002	0.4%	0.9697
1A1 & 1A2 & 1A4 & 1A5	1A1 & 1A2 & 1A4 & 1A5 Other Combustion	CH <sub>4</sub>	1893.15	1476.32	0.0002	0.3%	0.9727
3B	3B Manure Management	CH <sub>4</sub>	4732.53	4226.81	0.0001	0.3%	0.9754
2C	2C Metal Industries	CO <sub>2</sub>	7400.69	2572.86	0.0001	0.2%	0.9774
3B	3B Manure Management	N <sub>2</sub> O	3442.70	2814.79	0.0001	0.2%	0.9791
3H	3H Urea application to agriculture	CO <sub>2</sub>	327.60	343.95	0.0001	0.2%	0.9808
2D	2D Non Energy Products from Fuels and Solvent Use	CO <sub>2</sub>	552.81	519.22	0.0001	0.2%	0.9825
1A4	1A4 Petroleum Coke	CO <sub>2</sub>	81.64	351.76	0.0001	0.2%	0.9841
3G	3G Liming	CO <sub>2</sub>	1015.18	939.13	0.0001	0.2%	0.9857
1A4	1A4 Peat	CO <sub>2</sub>	372.48	4.83	0.0001	0.1%	0.9872
1B2	1B2 Oil & Natural Gas	CO <sub>2</sub>	5777.92	4230.64	0.0001	0.1%	0.9885
1A3b	1A3b DERV	CH <sub>4</sub>	88.74	7.85	0.0001	0.1%	0.9898
2G	2G Other Product Manufacture and Use	PFCs	149.16	183.51	0.0001	0.1%	0.9908
1B1	1B1 Solid Fuel Transformation	CO <sub>2</sub>	1698.56	359.54	0.0001	0.1%	0.9918
2C	2C Metal Industries	PFCs	333.43	15.07	0.0000	0.1%	0.9925
3J	3J OT & CD Agriculture	CH <sub>4</sub>	276.73	206.19	0.0000	0.1%	0.9932
5C	5C Waste Incineration	CH <sub>4</sub>	134.83	10.45	0.0000	0.1%	0.9939
5C	5C Waste Incineration	N <sub>2</sub> O	46.66	39.63	0.0000	0.1%	0.9945
3F	3F Field Burning	CH <sub>4</sub>	186.57	0.00	0.0000	0.1%	0.9951
1A3	1A3 Other diesel	N <sub>2</sub> O	16.53	25.30	0.0000	0.0%	0.9956
1B2	1B2 Offshore Oil& Gas	CH <sub>4</sub>	2176.74	1170.57	0.0000	0.0%	0.9960

IPCC Code	IPCC Category	Gas	Base year emissions (Gg CO <sub>2</sub> e)	2017 emissions (Gg CO <sub>2</sub> e)	Trend Assessment with Uncertainty	% Contribution to Trend Uncertainty	Cumulative Total
1B2	1B2 Oil & Natural Gas	N <sub>2</sub> O	40.75	39.36	0.0000	0.0%	0.9963
1A1 & 1A2 & 1A4 & 1A5	1A1 & 1A2 & 1A4 & 1A5 Other Combustion	N <sub>2</sub> O	2141.02	1262.84	0.0000	0.0%	0.9967
1A3d	1A3d Marine fuel	N <sub>2</sub> O	105.00	72.08	0.0000	0.0%	0.9969
2A	2A Minerals industry	CH <sub>4</sub>	31.10	5.89	0.0000	0.0%	0.9972
3J	3J OT & CD Agriculture	N <sub>2</sub> O	132.00	93.01	0.0000	0.0%	0.9975
2B	2B Chemical Industry	CH <sub>4</sub>	204.89	61.41	0.0000	0.0%	0.9977
2B	2B Chemical industry	PFCs	113.90	172.90	0.0000	0.0%	0.9979
2A2	2A2 Lime Production	CO <sub>2</sub>	1462.05	1052.06	0.0000	0.0%	0.9982
1A3c	1A3c Coal	CO <sub>2</sub>	0.00	41.15	0.0000	0.0%	0.9984
3F	3F Field Burning	N <sub>2</sub> O	57.66	0.00	0.0000	0.0%	0.9985
1A3a	1A3a Aviation Fuel	N <sub>2</sub> O	17.80	17.96	0.0000	0.0%	0.9987
2C	2C Metal Industries	SF <sub>6</sub>	387.17	109.75	0.0000	0.0%	0.9989
2E	2E Electronics Industry	HFCs	8.73	21.18	0.0000	0.0%	0.9991
2G	2G Other Product Manufacture and Use	SF <sub>6</sub>	912.79	415.66	0.0000	0.0%	0.9992
2A3	2A3 Glass production	CO <sub>2</sub>	405.54	367.98	0.0000	0.0%	0.9994
2A1	2A1 Cement Production	CO <sub>2</sub>	7295.26	4409.79	0.0000	0.0%	0.9995
1A3d	1A3d Marine fuel	CH <sub>4</sub>	3.66	6.28	0.0000	0.0%	0.9996
2C	2C Iron & Steel Production	CH <sub>4</sub>	36.90	10.68	0.0000	0.0%	0.9997
2C	2C Iron & Steel	N <sub>2</sub> O	17.74	6.19	0.0000	0.0%	0.9998
1B1	1B1 Solid Fuel Transformation	CH <sub>4</sub>	0.18	3.78	0.0000	0.0%	0.9999
2A4	2A4 Other process uses of carbonates	CO <sub>2</sub>	640.93	419.65	0.0000	0.0%	0.9999
1A3a	1A3a Aviation Fuel	CH <sub>4</sub>	6.37	1.55	0.0000	0.0%	0.9999
1A3c	1A3c Coal	CH <sub>4</sub>	0.00	0.94	0.0000	0.0%	1.0000
1A3	1A3 Other diesel	CH <sub>4</sub>	2.75	2.27	0.0000	0.0%	1.0000
2B8	2B8 Petrochemical and Carbon Black Production	N <sub>2</sub> O	2.10	1.51	0.0000	0.0%	1.0000
2C	2C Metal Industries	HFCs	0.00	2.29	0.0000	0.0%	1.0000

IPCC Code	IPCC Category	Gas	Base year emissions (Gg CO <sub>2</sub> e)	2017 emissions (Gg CO <sub>2</sub> e)	Trend Assessment with Uncertainty	% Contribution to Trend Uncertainty	Cumulative Total
1A3c	1A3c Coal	N <sub>2</sub> O	0.00	0.09	0.0000	0.0%	1.0000
2B1	2B1 Ammonia Production	N <sub>2</sub> O	0.31	0.33	0.0000	0.0%	1.0000
2F	2F Product Uses as Substitutes for ODS	PFCs	0.44	0.00	0.0000	0.0%	1.0000
1B1	1B1 Fugitive Emissions from Solid Fuels	N <sub>2</sub> O	0.09	0.02	0.0000	0.0%	1.0000
2E	2E Electronics Industry	NF <sub>3</sub>	0.83	0.53	0.0000	0.0%	1.0000
2D	2D Non-energy Products from Fuels and Solvent Use	CH <sub>4</sub>	0.00	0.00	0.0000	0.0%	1.0000
2D	2D Non-energy Products from Fuels and Solvent Use	N <sub>2</sub> O	0.00	0.00	0.0000	0.0%	1.0000

## A 1.5 KEY CATEGORY ANALYSIS (KCA) RANKING SYSTEM

The Key Category Analysis (KCA) ranking system is an additional tool that the UK has developed to aid in the prioritisation of improvement work. The KCA ranking system works by allocating a score based on how high categories rank in the base year and most recent year level assessments and the trend assessment for the approach 1 KCA including LULUCF. For example, if CO<sub>2</sub> from road transport liquid fuel use is the 4<sup>th</sup> highest by the base year level assessment, 3<sup>rd</sup> highest by the most recent year level assessment and has the 5<sup>th</sup> highest trend assessment then it's score would be 4+3+5=12. The categories are then ranked from lowest score to highest, with scores that are equal resolved by the most recent year level assessment.

The assessments used in this ranking exercise are only those *including* LULUCF, because if the additional *excluding* LULUCF assessments were also used, the LULUCF sectors would only be included in half of the assessments and would therefore give an unrepresentative weighting.

The results of this ranking are presented in **Table A 1.5.1**.

**Table A 1.5.1 KCA Ranking**

KCA rank	IPCC Code	IPCC Category	Greenhouse Gas
1	1A3b	Road transportation: liquid fuels	CO <sub>2</sub>
2	1A1	Energy industries: solid fuels	CO <sub>2</sub>
3	1A4	Other sectors: gaseous fuels	CO <sub>2</sub>
4	5A	Solid waste disposal	CH <sub>4</sub>
5	1A1	Energy industries: gaseous fuels	CO <sub>2</sub>

KCA rank	IPCC Code	IPCC Category	Greenhouse Gas
6	1A2	Manufacturing industries and construction: gaseous fuels	CO <sub>2</sub>
7	1A1	Energy industries: liquid fuels	CO <sub>2</sub>
8	1A2	Manufacturing industries and construction: solid fuels	CO <sub>2</sub>
9	3A1	Enteric fermentation from Cattle	CH <sub>4</sub>
10	1A2	Manufacturing industries and construction: liquid fuels	CO <sub>2</sub>
11	4A	Forest land	CO <sub>2</sub>
12	1A4	Other sectors: liquid fuels	CO <sub>2</sub>
13	3D	Agricultural soils	N <sub>2</sub> O
14	4B	Cropland	CO <sub>2</sub>
15	1A4	Other sectors: solid fuels	CO <sub>2</sub>
16	1B2	Oil and gas extraction	CH <sub>4</sub>
17	4E	Settlements	CO <sub>2</sub>
18	1B1	Coal mining and handling	CH <sub>4</sub>
19	4C	Grassland	CO <sub>2</sub>
20	1A3d	Domestic Navigation: liquid fuels	CO <sub>2</sub>
21	2F1	Refrigeration and air conditioning	HFCs, PFCs, SF <sub>6</sub> and NF <sub>3</sub>
22	3B1	Manure management from Cattle	CH <sub>4</sub>
23	1B2	Oil and gas extraction	CO <sub>2</sub>
24	3A2	Enteric fermentation from Sheep	CH <sub>4</sub>
25	1A5	Other: liquid fuels	CO <sub>2</sub>
26	5D	Wastewater treatment and discharge	CH <sub>4</sub>
27	2C1	Iron and steel production	CO <sub>2</sub>
28	2B9	Fluorochemical production	HFCs, PFCs, SF <sub>6</sub> and NF <sub>3</sub>
29	3B2	Manure management from Sheep	N <sub>2</sub> O
30	1A3c	Railways: liquid fuels	CO <sub>2</sub>
31	1A1	Energy industries: other fuels	CO <sub>2</sub>
32	1A3a	Domestic aviation: liquid fuels	CO <sub>2</sub>
33	2A1	Cement production	CO <sub>2</sub>
34	2F4	Aerosols	HFCs, PFCs, SF <sub>6</sub> and NF <sub>3</sub>
35	2B8	Petrochemical and carbon black production	CO <sub>2</sub>
36	4G	Harvested wood products	CO <sub>2</sub>
37	2B2	Nitric acid production	N <sub>2</sub> O

KCA rank	IPCC Code	IPCC Category	Greenhouse Gas
38	2B3	Adipic acid production	N <sub>2</sub> O
39	5B	Biological treatment of solid waste	CH <sub>4</sub>

## A 1.6 APPROACH USED TO IDENTIFY KP-LULUCF KEY CATEGORIES

From the 2010 NIR onwards, the NIR contains a list of the Key Categories for Land Use, Land-Use Change and Forestry Activities under the Kyoto Protocol. The description below explains the Key Category analysis for Article 3.3 activities and any elected activities under Article 3.4.

Five categories are considered to be key: Article 3.3 Afforestation and Reforestation (CO<sub>2</sub>), Article 3.3 Deforestation (CO<sub>2</sub>), Article 3.4 Forest Management (CO<sub>2</sub>), Article 3.4 Cropland Management (CO<sub>2</sub>) and Article 3.4 Grazing Land Management (CO<sub>2</sub>). These have been assessed according to the 2006 IPCC good practice guidance for KP (Chapter 2, Section 2.3.6). The numbers have been compared with Table A 1.4.2. The key category analysis for the latest reported year based on level of emissions (including LULUCF) is given in section 11.6.1 of the main NIR.

## A 1.7 USING THE UNCERTAINTY ANALYSIS TO PLAN IMPROVEMENTS IN THE PREPARATION OF THE INVENTORY

The key category analysis is used to prioritise and plan improvements. The approach the UK takes to achieve this is described in **Section 1.2.2.5. Table 1.7 to Table 1.11 in Chapter 1** show the key category summary tables.

**A 1.8 TABLE NIR 3, AS CONTAINED IN THE ANNEX TO DECISION 6/CMP.3**

**Table A 1.8.1** below is Table NIR 3, containing a summary overview for Key Categories for Land Use, Land-Use Change and Forestry Activities under the Kyoto Protocol<sup>3</sup>.

**Table A 1.8.1 Table NIR 3. Summary overview for Key Categories for Land Use, Land-Use Change and Forestry Activities under the Kyoto Protocol**

KEY CATEGORIES OF EMISSIONS AND REMOVALS	GAS	CRITERIA USED FOR KEY CATEGORY IDENTIFICATION			COMMENTS (3)
		Associated category in UNFCCC inventory (1) is key (indicate which category)	Category contribution is greater than the smallest category considered key in the UNFCCC inventory (1), (4) (including LULUCF)	Other (2)	
Specify key categories according to the national level of disaggregation used (1)					
Afforestation and Reforestation	CO <sub>2</sub>	Land converted to forest land	Yes	Associated UNFCCC category (4A) is key	The associated UNFCCC inventory category is a key category for level.
Deforestation	CO <sub>2</sub>	Land converted to cropland, Land converted to grassland, Land converted to settlements	Yes	Associated UNFCCC categories (4B2, 4C2 and 4E2) are key.	The associated UNFCCC inventory categories are key categories for level and trend.

<sup>3</sup> Table NIR 3 can be found in FCCC/KP/CMP/2007/9/Add.2.

KEY CATEGORIES OF EMISSIONS AND REMOVALS	GAS	CRITERIA USED FOR KEY CATEGORY IDENTIFICATION			COMMENTS (3)
		Associated category in UNFCCC inventory (1) is key (indicate which category)	Category contribution is greater than the smallest category considered key in the UNFCCC inventory (1), (4) (including LULUCF)	Other (2)	
Forest Management	CO <sub>2</sub>	Forest land remaining forest land, Land converted to forest land	Yes	Associated UNFCCC category (4A) is key	The associated UNFCCC inventory category is a key category for level and the Forest Management category contribution is greater than the smallest UNFCCC key category.
Cropland Management	CO <sub>2</sub>	Cropland remaining Cropland, Land converted to Cropland	Yes	Associated UNFCCC category (4B) is key.	The associated UNFCCC inventory category is a key category for level and trend and the Cropland Management category contribution is greater than the smallest UNFCCC key category.
Grazing Land Management	CO <sub>2</sub>	Grassland remaining Grassland, Land converted to Grassland	Yes	Associated UNFCCC category (4C) is key.	The associated UNFCCC inventory category is a key category for level and trend and the Grazing Land Management category contribution is greater than the smallest UNFCCC key category.

- (1) See section 5.4 of the IPCC good practice guidance for LULUCF
- (2) This should include qualitative consideration as per Section 5.4.3 of the IPCC Good Practice Guidance for LULUCF or any other criteria
- (3) Describe the criteria identifying the category as key
- (4) If the emissions or removals of the category exceed the emissions of the smallest category identified as key in the UNFCCC inventory (including LULUCF), Parties should indicate YES. If not, Parties should indicate NO



## **ANNEX 2: Assessment of Uncertainty**

Uncertainty estimates are calculated using two methods: Approach 1 (error propagation) and Approach 2 (Monte Carlo simulation). These are not to be confused with Approaches 1 and 2 for Key Category Analysis, of which Approach 2 KCA uses Approach 1 uncertainties to account for uncertainty in determining Key Categories.

The uncertainty assessment estimates uncertainties according to IPCC sector in addition to presenting estimates by direct greenhouse gas. Estimated uncertainty presented for the sector breakdown used in UK Official Statistics are not reported here, since the categories are not consistent with the requirements of the UK's commitments under the UNFCCC and Kyoto Protocol.

Uncertainty parameters for new sources and sources which have been significantly revised are reviewed each year, particularly for sources which have a significant impact on overall uncertainties.

The overall method used to estimate uncertainties is described below, and the work to improve the accuracy of the uncertainty analysis continues. The key category analysis used data from the uncertainty analysis, and the results of the key category analysis are given in **Annex 1**.

### **A 2.1 ESTIMATION OF UNCERTAINTIES USING AN ERROR PROPAGATION APPROACH (APPROACH 1)**

The IPCC 2006 Guidelines defines error propagation and Monte Carlo modelling approaches to estimating uncertainties in national greenhouse gas inventories. The results of the error propagation approach are shown in **Table A 2.1.1**. The uncertainties used in the error propagation approach are not exactly the same as those used in the Monte Carlo Simulation since the error propagation source categorisation is less detailed and has a more simplistic approach to uncertainties. The Approach 1 uncertainties assumes all parameters are normally distributed (which means it doesn't account for the skew, kurtosis or any other non-normal features of the expected distributions), and does not account for variations in uncertainty in the time series unlike the Monte Carlo approach which takes into account these factors. The parameters used for the Approach 1 uncertainties for both the base year and the most recent year are the values given for the most recent year in **Table A 2.3.1** to **Table A 2.3.4**.

#### **A 2.1.1 Key Categories**

Certain source categories are particularly significant in terms of their contribution to the overall uncertainty of the inventory. Key source categories in this respect are identified using Approach 1 uncertainties in the Approach 2 KCA. These have been identified so that the resources available for inventory preparation may be prioritised, and the best possible estimates prepared for the most significant source categories. We have used the method described in Section 4.3.2 of the 2006 IPCC Guidelines Volume 1 General Guidance and Reporting (Approach 2 to identify key categories).

The results of this key category analysis can be found in **Annex 1**.

**A 2.1.2 Tables of uncertainty estimates from the error propagation approach**

**Table A 2.1.1 Summary of error propagation uncertainty estimates including LULUCF, base year to the latest reported year**

IPCC Category	Gas	Base year emissions (Gg CO <sub>2</sub> e)	2017 emissions (Gg CO <sub>2</sub> e)	Activity data uncertainty (%)	Emission factor uncertainty (%)	Combined uncertainty (%)	Contribution to variance by Category in 2017	Type A sensitivity	Type B sensitivity	Uncertainty in trend in national emissions introduced by emission factor / estimation parameter uncertainty	Uncertainty in trend in national emissions introduced by activity data uncertainty	Uncertainty introduced into the trend in total national emissions
1A (Stationary) Oil	CO <sub>2</sub>	95,612.77	45,156.95	5.97%	2.55%	6.5%	0.0000	1.274%	5.632%	0.0325%	0.4754%	0.0023%
1A Coal	CO <sub>2</sub>	243,845.39	34,774.45	1.28%	2.81%	3.1%	0.0000	13.238%	4.337%	0.3721%	0.0783%	0.0014%
1A Natural Gas	CO <sub>2</sub>	106,932.90	158,621.57	1.01%	1.65%	1.9%	0.0000	12.041%	19.782%	0.1989%	0.2838%	0.0012%
1A Other (waste)	CO <sub>2</sub>	247.13	5,441.36	1.08%	13.86%	13.9%	0.0000	0.661%	0.679%	0.0916%	0.0104%	0.0001%
1A3 Other diesel	CO <sub>2</sub>	1,679.92	2,597.40	14.47%	1.94%	14.6%	0.0000	0.203%	0.324%	0.0039%	0.0663%	0.0000%
1A3a Aviation Fuel	CO <sub>2</sub>	1,881.31	1,898.39	19.65%	3.24%	19.9%	0.0000	0.101%	0.237%	0.0033%	0.0658%	0.0000%
1A3b DERV	CO <sub>2</sub>	33,005.80	77,521.13	1.00%	2.00%	2.2%	0.0000	7.281%	9.668%	0.1456%	0.1367%	0.0004%
1A3b Gasoline/ LPG	CO <sub>2</sub>	75,562.72	35,843.09	0.99%	1.99%	2.2%	0.0000	0.987%	4.470%	0.0196%	0.0629%	0.0000%
1A3c Coal	CO <sub>2</sub>	-	41.15	20.00%	6.00%	20.9%	0.0000	0.005%	0.005%	0.0003%	0.0015%	0.0000%
1A3d Marine fuel	CO <sub>2</sub>	7,611.13	5,443.73	17.71%	1.77%	17.8%	0.0000	0.129%	0.679%	0.0023%	0.1701%	0.0003%
1A4 Peat	CO <sub>2</sub>	372.48	4.83	30.00%	10.00%	31.6%	0.0000	0.026%	0.001%	0.0026%	0.0003%	0.0000%
1A4 Petroleum Coke	CO <sub>2</sub>	81.64	351.76	20.00%	15.00%	25.0%	0.0000	0.038%	0.044%	0.0057%	0.0124%	0.0000%
1B1 Solid Fuel Transformation	CO <sub>2</sub>	1,698.56	359.54	5.17%	4.84%	7.1%	0.0000	0.078%	0.045%	0.0038%	0.0033%	0.0000%
1B2 Oil & Natural Gas	CO <sub>2</sub>	5,777.92	4,230.64	4.49%	5.40%	7.0%	0.0000	0.110%	0.528%	0.0060%	0.0335%	0.0000%
2A1 Cement Production	CO <sub>2</sub>	7,295.26	4,409.79	1.00%	3.00%	3.2%	0.0000	0.023%	0.550%	0.0007%	0.0078%	0.0000%

IPCC Category	Gas	Base year emissions (Gg CO <sub>2</sub> e)	2017 emissions (Gg CO <sub>2</sub> e)	Activity data uncertainty (%)	Emission factor uncertainty (%)	Combined uncertainty (%)	Contribution to variance by Category in 2017	Type A sensitivity	Type B sensitivity	Uncertainty in trend in national emissions introduced by emission factor / estimation parameter uncertainty	Uncertainty in trend in national emissions introduced by activity data uncertainty	Uncertainty introduced into the trend in total national emissions
2A2 Lime Production	CO <sub>2</sub>	1,462.05	1,052.06	0.00%	5.00%	5.0%	0.0000	0.026%	0.131%	0.0013%	0.0000%	0.0000%
2A3 Glass production	CO <sub>2</sub>	405.54	367.98	0.00%	5.00%	5.0%	0.0000	0.017%	0.046%	0.0008%	0.0000%	0.0000%
2A4 Other process uses of carbonates	CO <sub>2</sub>	640.93	419.65	2.00%	3.00%	3.6%	0.0000	0.006%	0.052%	0.0002%	0.0015%	0.0000%
2B Chemical industries	CO <sub>2</sub>	6,770.09	4,931.56	17.29%	2.94%	17.5%	0.0000	0.126%	0.615%	0.0037%	0.1504%	0.0002%
2C Metal Industries	CO <sub>2</sub>	7,400.69	2,572.86	1.18%	5.09%	5.2%	0.0000	0.214%	0.321%	0.0109%	0.0054%	0.0000%
2D Non Energy Products from Fuels and Solvent Use	CO <sub>2</sub>	552.81	519.22	26.78%	27.38%	38.3%	0.0000	0.025%	0.065%	0.0068%	0.0245%	0.0000%
3G Liming	CO <sub>2</sub>	1,015.18	939.13	0.00%	20.90%	20.9%	0.0000	0.044%	0.117%	0.0092%	0.0000%	0.0000%
3H Urea application to agriculture	CO <sub>2</sub>	327.60	343.95	0.00%	50.00%	50.0%	0.0000	0.019%	0.043%	0.0096%	0.0000%	0.0000%
4A Forest Land	CO <sub>2</sub>	-15,026.26	-18,211.77	1.00%	35.00%	35.0%	0.0002	1.186%	2.271%	0.4151%	0.0321%	0.0017%
4B Cropland	CO <sub>2</sub>	14,265.93	10,971.23	1.00%	45.00%	45.0%	0.0001	0.338%	1.368%	0.1519%	0.0193%	0.0002%
4C Grassland	CO <sub>2</sub>	-7,111.03	-8,861.80	1.00%	50.00%	50.0%	0.0001	0.592%	1.105%	0.2958%	0.0156%	0.0009%
4D Wetland	CO <sub>2</sub>	486.95	336.78	1.00%	100.00%	100.0%	0.0000	0.007%	0.042%	0.0068%	0.0006%	0.0000%
4E Settlements	CO <sub>2</sub>	7,011.30	6,451.34	1.00%	50.00%	50.0%	0.0000	0.298%	0.805%	0.1490%	0.0114%	0.0002%
4F Other Land	CO <sub>2</sub>	-	-	0.00%	0.00%	0.0%	-	0.000%	0.000%	0.0000%	0.0000%	0.0000%
4G Other Activities	CO <sub>2</sub>	-1,639.08	-2,015.60	1.00%	45.00%	45.0%	0.0000	0.133%	0.251%	0.0598%	0.0036%	0.0000%
5C Waste Incineration	CO <sub>2</sub>	1,300.71	258.91	300.00%	40.00%	302.7%	0.0000	0.062%	0.032%	0.0247%	0.1370%	0.0002%

IPCC Category	Gas	Base year emissions (Gg CO <sub>2</sub> e)	2017 emissions (Gg CO <sub>2</sub> e)	Activity data uncertainty (%)	Emission factor uncertainty (%)	Combined uncertainty (%)	Contribution to variance by Category in 2017	Type A sensitivity	Type B sensitivity	Uncertainty in trend in national emissions introduced by emission factor / estimation parameter uncertainty	Uncertainty in trend in national emissions introduced by activity data uncertainty	Uncertainty introduced into the trend in total national emissions
1A1 & 1A2 & 1A4 & 1A5 Other Combustion	CH <sub>4</sub>	1,893.15	1,476.32	0.71%	35.55%	35.6%	0.0000	0.047%	0.184%	0.0168%	0.0019%	0.0000%
1A3 Other diesel	CH <sub>4</sub>	2.75	2.27	15.00%	130.00%	130.9%	0.0000	0.000%	0.000%	0.0001%	0.0001%	0.0000%
1A3a Aviation Fuel	CH <sub>4</sub>	6.37	1.55	14.18%	55.65%	57.4%	0.0000	0.000%	0.000%	0.0001%	0.0000%	0.0000%
1A3b DERV	CH <sub>4</sub>	88.74	7.85	1.00%	130.00%	130.0%	0.0000	0.005%	0.001%	0.0071%	0.0000%	0.0000%
1A3b Gasoline/ LPG	CH <sub>4</sub>	1,157.96	85.31	1.00%	74.94%	74.9%	0.0000	0.073%	0.011%	0.0547%	0.0002%	0.0000%
1A3c Coal	CH <sub>4</sub>	-	0.94	20.00%	110.00%	111.8%	0.0000	0.000%	0.000%	0.0001%	0.0000%	0.0000%
1A3d Marine fuel	CH <sub>4</sub>	3.66	6.28	19.14%	124.42%	125.9%	0.0000	0.001%	0.001%	0.0006%	0.0002%	0.0000%
1B1 Coal Mining	CH <sub>4</sub>	21,826.68	483.98	2.00%	20.00%	20.1%	0.0000	1.516%	0.060%	0.3032%	0.0017%	0.0009%
1B1 Solid Fuel Transformation	CH <sub>4</sub>	0.18	3.78	0.00%	49.81%	49.8%	0.0000	0.000%	0.000%	0.0002%	0.0000%	0.0000%
1B2 Natural Gas Transmission	CH <sub>4</sub>	10,168.33	3,764.68	3.00%	20.00%	20.2%	0.0000	0.265%	0.470%	0.0530%	0.0199%	0.0000%
1B2 Offshore Oil& Gas	CH <sub>4</sub>	2,176.74	1,170.57	4.54%	18.15%	18.7%	0.0000	0.011%	0.146%	0.0020%	0.0094%	0.0000%
2A Minerals industry	CH <sub>4</sub>	31.10	5.89	0.00%	100.00%	100.0%	0.0000	0.002%	0.001%	0.0015%	0.0000%	0.0000%
2B Chemical Industry	CH <sub>4</sub>	204.89	61.41	0.00%	20.00%	20.0%	0.0000	0.007%	0.008%	0.0014%	0.0000%	0.0000%
2C Iron & Steel Production	CH <sub>4</sub>	36.90	10.68	1.91%	47.83%	47.9%	0.0000	0.001%	0.001%	0.0006%	0.0000%	0.0000%

IPCC Category	Gas	Base year emissions (Gg CO <sub>2</sub> e)	2017 emissions (Gg CO <sub>2</sub> e)	Activity data uncertainty (%)	Emission factor uncertainty (%)	Combined uncertainty (%)	Contribution to variance by Category in 2017	Type A sensitivity	Type B sensitivity	Uncertainty in trend in national emissions introduced by emission factor / estimation parameter uncertainty	Uncertainty in trend in national emissions introduced by activity data uncertainty	Uncertainty introduced into the trend in total national emissions
2D Non-energy Products from Fuels and Solvent Use	CH <sub>4</sub>	-	-	0.00%	0.00%	0.0%	-	0.000%	0.000%	0.0000%	0.0000%	0.0000%
3A Enteric Fermentation	CH <sub>4</sub>	25,392.50	21,458.44	13.73%	0.00%	13.7%	0.0000	0.842%	2.676%	0.0000%	0.5198%	0.0027%
3B Manure Management	CH <sub>4</sub>	4,732.53	4,226.81	0.00%	8.37%	8.4%	0.0000	0.185%	0.527%	0.0155%	0.0000%	0.0000%
3F Field Burning	CH <sub>4</sub>	186.57	-	25.61%	0.00%	25.6%	-	0.013%	0.000%	0.0000%	0.0000%	0.0000%
3J OT & CD Agriculture	CH <sub>4</sub>	276.73	206.19	50.00%	50.00%	70.7%	0.0000	0.006%	0.026%	0.0029%	0.0182%	0.0000%
4A Forest Land	CH <sub>4</sub>	2.89	2.31	1.00%	55.00%	55.0%	0.0000	0.000%	0.000%	0.0000%	0.0000%	0.0000%
4B Cropland	CH <sub>4</sub>	0.09	0.02	1.00%	55.00%	55.0%	0.0000	0.000%	0.000%	0.0000%	0.0000%	0.0000%
4C Grassland	CH <sub>4</sub>	9.97	17.78	1.00%	55.00%	55.0%	0.0000	0.001%	0.002%	0.0008%	0.0000%	0.0000%
4E Settlements	CH <sub>4</sub>	3.00	8.25	1.00%	55.00%	55.0%	0.0000	0.001%	0.001%	0.0004%	0.0000%	0.0000%
5A Solid Waste Disposal	CH <sub>4</sub>	60,432.98	14,261.40	15.00%	46.00%	48.4%	0.0002	2.585%	1.779%	1.1891%	0.3773%	0.0156%
5B Biological treatment of solid waste	CH <sub>4</sub>	18.13	1,193.62	30.00%	99.50%	103.9%	0.0000	0.148%	0.149%	0.1468%	0.0632%	0.0003%
5C Waste Incineration	CH <sub>4</sub>	134.83	10.45	7.40%	44.45%	45.1%	0.0000	0.008%	0.001%	0.0038%	0.0001%	0.0000%
5D Wastewater Handling	CH <sub>4</sub>	4,218.90	3,445.75	10.00%	25.00%	26.9%	0.0000	0.125%	0.430%	0.0312%	0.0608%	0.0000%

IPCC Category	Gas	Base year emissions (Gg CO <sub>2</sub> e)	2017 emissions (Gg CO <sub>2</sub> e)	Activity data uncertainty (%)	Emission factor uncertainty (%)	Combined uncertainty (%)	Contribution to variance by Category in 2017	Type A sensitivity	Type B sensitivity	Uncertainty in trend in national emissions introduced by emission factor / estimation parameter uncertainty	Uncertainty in trend in national emissions introduced by activity data uncertainty	Uncertainty introduced into the trend in total national emissions
1A1 & 1A2 & 1A4 & 1A5 Other Combustion	N <sub>2</sub> O	2,141.02	1,262.84	0.66%	65.65%	65.7%	0.0000	0.003%	0.157%	0.0019%	0.0015%	0.0000%
1A3 Other diesel	N <sub>2</sub> O	16.53	25.30	15.00%	130.00%	130.9%	0.0000	0.002%	0.003%	0.0025%	0.0007%	0.0000%
1A3a Aviation Fuel	N <sub>2</sub> O	17.80	17.96	19.65%	108.08%	109.9%	0.0000	0.001%	0.002%	0.0010%	0.0006%	0.0000%
1A3b DERV	N <sub>2</sub> O	322.88	1,000.43	1.00%	130.00%	130.0%	0.0000	0.101%	0.125%	0.1319%	0.0018%	0.0002%
1A3b Gasoline/ LPG	N <sub>2</sub> O	989.40	95.21	0.99%	74.47%	74.5%	0.0000	0.060%	0.012%	0.0444%	0.0002%	0.0000%
1A3c Coal	N <sub>2</sub> O	-	0.09	20.00%	110.00%	111.8%	0.0000	0.000%	0.000%	0.0000%	0.0000%	0.0000%
1A3d Marine fuel	N <sub>2</sub> O	105.00	72.08	17.56%	114.16%	115.5%	0.0000	0.001%	0.009%	0.0016%	0.0022%	0.0000%
1B1 Fugitive Emissions from Solid Fuels	N <sub>2</sub> O	0.09	0.02	1.00%	118.00%	118.0%	0.0000	0.000%	0.000%	0.0000%	0.0000%	0.0000%
1B2 Oil & Natural Gas	N <sub>2</sub> O	40.75	39.36	4.77%	105.01%	105.1%	0.0000	0.002%	0.005%	0.0021%	0.0003%	0.0000%
2B1 Ammonia Production	N <sub>2</sub> O	0.31	0.33	2.00%	50.00%	50.0%	0.0000	0.000%	0.000%	0.0000%	0.0000%	0.0000%
2B2 Nitric Acid Production	N <sub>2</sub> O	3,860.26	37.25	0.00%	10.00%	10.0%	0.0000	0.274%	0.005%	0.0274%	0.0000%	0.0000%
2B3 Adipic Acid Production	N <sub>2</sub> O	19,934.61	-	2.00%	100.00%	100.0%	-	1.440%	0.000%	1.4397%	0.0000%	0.0207%
2B8 Petrochemical and Carbon Black Production	N <sub>2</sub> O	2.10	1.51	10.00%	100.00%	100.5%	0.0000	0.000%	0.000%	0.0000%	0.0000%	0.0000%
2C Iron & Steel	N <sub>2</sub> O	17.74	6.19	1.00%	118.00%	118.0%	0.0000	0.001%	0.001%	0.0006%	0.0000%	0.0000%

IPCC Category	Gas	Base year emissions (Gg CO <sub>2</sub> e)	2017 emissions (Gg CO <sub>2</sub> e)	Activity data uncertainty (%)	Emission factor uncertainty (%)	Combined uncertainty (%)	Contribution to variance by Category in 2017	Type A sensitivity	Type B sensitivity	Uncertainty in trend in national emissions introduced by emission factor / estimation parameter uncertainty	Uncertainty in trend in national emissions introduced by activity data uncertainty	Uncertainty introduced into the trend in total national emissions
2D Non-energy Products from Fuels and Solvent Use	N <sub>2</sub> O	-	-	0.00%	0.00%	0.0%	-	0.000%	0.000%	0.0000%	0.0000%	0.0000%
2G Other Product Manufacture and Use	N <sub>2</sub> O	593.57	843.87	100.00%	100.00%	141.4%	0.0000	0.062%	0.105%	0.0624%	0.1488%	0.0003%
3B Manure Management	N <sub>2</sub> O	3,442.70	2,814.79	0.00%	9.53%	9.5%	0.0000	0.102%	0.351%	0.0098%	0.0000%	0.0000%
3D Agricultural Soils	N <sub>2</sub> O	13,610.39	11,466.79	0.00%	11.16%	11.2%	0.0000	0.447%	1.430%	0.0499%	0.0000%	0.0000%
3F Field Burning	N <sub>2</sub> O	57.66	-	25.62%	0.00%	25.6%	-	0.004%	0.000%	0.0000%	0.0000%	0.0000%
3J OT & CD Agriculture	N <sub>2</sub> O	132.00	93.01	50.00%	50.00%	70.7%	0.0000	0.002%	0.012%	0.0010%	0.0082%	0.0000%
4 Indirect LULUCF Emissions	N <sub>2</sub> O	401.49	243.62	1.00%	165.00%	165.0%	0.0000	0.001%	0.030%	0.0023%	0.0004%	0.0000%
4A Forest land	N <sub>2</sub> O	231.22	144.65	1.00%	95.00%	95.0%	0.0000	0.001%	0.018%	0.0013%	0.0003%	0.0000%
4B Cropland	N <sub>2</sub> O	1,019.85	461.97	1.00%	35.00%	35.0%	0.0000	0.016%	0.058%	0.0056%	0.0008%	0.0000%
4C Grassland	N <sub>2</sub> O	10.33	33.24	1.00%	115.00%	115.0%	0.0000	0.003%	0.004%	0.0039%	0.0001%	0.0000%
4D Grassland	N <sub>2</sub> O	4.13	0.30	1.00%	100.00%	100.0%	0.0000	0.000%	0.000%	0.0003%	0.0000%	0.0000%
4E Settlements	N <sub>2</sub> O	584.68	524.99	1.00%	15.00%	15.0%	0.0000	0.023%	0.065%	0.0035%	0.0009%	0.0000%
5B Biological treatment of solid waste	N <sub>2</sub> O	12.97	728.03	30.00%	90.00%	94.9%	0.0000	0.090%	0.091%	0.0809%	0.0385%	0.0001%
5C Waste Incineration	N <sub>2</sub> O	46.66	39.63	7.00%	230.00%	230.1%	0.0000	0.002%	0.005%	0.0036%	0.0005%	0.0000%

IPCC Category	Gas	Base year emissions (Gg CO <sub>2</sub> e)	2017 emissions (Gg CO <sub>2</sub> e)	Activity data uncertainty (%)	Emission factor uncertainty (%)	Combined uncertainty (%)	Contribution to variance by Category in 2017	Type A sensitivity	Type B sensitivity	Uncertainty in trend in national emissions introduced by emission factor / estimation parameter uncertainty	Uncertainty in trend in national emissions introduced by activity data uncertainty	Uncertainty introduced into the trend in total national emissions
5D Wastewater Handling	N <sub>2</sub> O	783.59	725.93	10.00%	248.00%	248.2%	0.0000	0.034%	0.091%	0.0841%	0.0128%	0.0001%
2C Metal Industries	SF <sub>6</sub>	387.17	109.75	5.00%	5.00%	7.1%	0.0000	0.014%	0.014%	0.0007%	0.0010%	0.0000%
2G Other Product Manufacture and Use	SF <sub>6</sub>	912.79	415.66	0.00%	6.17%	6.2%	0.0000	0.014%	0.052%	0.0009%	0.0000%	0.0000%
2B Chemical industry	HFCs	17,670.77	2.55	0.00%	10.00%	10.0%	0.0000	1.276%	0.000%	0.1276%	0.0000%	0.0002%
2C Metal Industries	HFCs	-	2.29	5.00%	10.00%	11.2%	0.0000	0.000%	0.000%	0.0000%	0.0000%	0.0000%
2E Electronics Industry	HFCs	8.73	21.18	0.00%	47.15%	47.1%	0.0000	0.002%	0.003%	0.0009%	0.0000%	0.0000%
2F Product Uses as Substitutes for ODS	HFCs	1,416.64	14,166.81	8.23%	8.30%	11.7%	0.0000	1.664%	1.767%	0.1382%	0.2057%	0.0006%
2E Electronics Industry	NF <sub>3</sub>	0.83	0.53	0.00%	47.15%	47.1%	0.0000	0.000%	0.000%	0.0000%	0.0000%	0.0000%
2B Chemical industry	PFCs	113.90	172.90	0.00%	10.00%	10.0%	0.0000	0.013%	0.022%	0.0013%	0.0000%	0.0000%
2C Metal Industries	PFCs	333.43	15.07	0.00%	20.00%	20.0%	0.0000	0.022%	0.002%	0.0044%	0.0000%	0.0000%
2F Product Uses as Substitutes for ODS	PFCs	0.44	-	0.00%	25.00%	25.0%	-	0.0000%	0.0000%	0.0000%	0.0000%	0.0000%
2G Other Product Manufacture and Use	PFCs	149.16	183.51	0.00%	47.15%	47.1%	0.0000	0.0121%	0.0229%	0.0057%	0.0000%	0.0000%

Percentage uncertainty in total inventory:	3.0%
--	------

Trend uncertainty	2.3%
-------------------	------

## **A 2.2 ESTIMATION OF UNCERTAINTY BY SIMULATION (APPROACH 2)**

### **A 2.2.1 Overview of the Method**

Quantitative estimates of the uncertainties in the emissions were calculated using a Monte Carlo simulation. This corresponds to the IPCC Approach 2 method, discussed in the 2006 Guidelines (IPCC, 2006). The background to the implementation of the Monte Carlo simulation is described in detail by Eggleston *et al* (1998), with the estimates reported here revised to reflect changes in the latest inventory and improvements made in the model. This section gives a brief summary of the methodology, assumptions and results of the simulation.

The computational procedure is detailed below.

- A probability distribution function (PDF) was allocated to each unique emission factor and piece of activity data. The PDFs were mostly normal or log-normal, with more specific distributions given to a handful of sources. The parameters of the PDFs were set by analysing the available data on emission factors and activity data, and by expert judgement;
- A calculation was set up to estimate the total emissions of each gas for the years 1990 and the latest reported year;
- Using the software tool @RISK™, each PDF was sampled at least 20,000 times, such that the emission calculations performed produced a converged output distribution;
- The distribution of errors in the parameter values was calculated from the difference between 2.5 and 97.5 percentile values in the distribution, as a percentage of the distribution mean; and,
- The uncertainty in the trend between 1990 and the latest reported year, according to gas, was also estimated. This is expressed as the 95% confidence interval for the percentage reduction in emissions between the latest year and 1990.

### **A 2.2.2 Methodological details of the Monte Carlo model**

#### **A 2.2.2.1 Uncertainty Distributions**

Nearly all of the distributions of emissions from sources in the inventory are modelled using normal or log normal distributions, with more specific distributions given to a handful of sources. The specific distributions include log-logistic, Pearson and Gamma distributions. The primary use of custom distributions is for agriculture; these are fitted distributions that reflect the results of an agriculture-specific Monte Carlo analysis done by Rothamsted Research which accounts for the various factors that influence the modelled agriculture emissions.

Emissions from landfill have been modelled using a custom distribution. Aitchison *et al.* (cited in Eggleston *et al.*, 1998) estimated the uncertainty for landfill emissions using Monte Carlo analysis and found it to be skewed. The distribution histogram was used to generate an empirical distribution of emissions. We examined the distribution and fitted a log normal distribution to Aitchison's data. The emissions are scaled according to the mean estimate of landfill emissions for each year.

There are a couple of other specific distributions for F-gases and waste water which reflect specific distributions we expect for those sources.

**A 2.2.2.2 Correlations**

The Monte Carlo model contains a number of correlations. If A and B are correlated, then if emissions are under or overestimated from A it would be expected to be over or underestimated by a similar amount from B.

The type and implementation of the correlations has been examined as part of a review (Abbott *et al.*, 2007). The sensitivity analysis that we have completed on the Monte Carlo model suggest that the uncertainties are not sensitive to the correlations between emission factors for fuel used, and for LULUCF sources.

**A 2.2.2.2.1 Across years**

In running this simulation, it was necessary to make assumptions about the degree of correlation between sources in 1990 and the latest reported year. If source emission factors are correlated this will have the effect of reducing the trend uncertainty, but will not affect uncertainties on emission totals in 1990 or the latest inventory year. The trend estimated by the Monte Carlo model is particularly sensitive to N<sub>2</sub>O emissions from agricultural soils.

**A 2.2.2.2.2 Between Sources in the same year**

In many cases the same factors, or factors derived on the same basis are used for multiple sources. In these cases, we'd say that the emission factors are correlated. For example, the coal emissions factors for N<sub>2</sub>O used for cement industry use may be the same as coal use in other industrial combustion due to lack of a more specific factor, in this case we may say the two factors are correlated. Omitting these correlations leads to an underestimate of emissions in any given year.

**A 2.2.2.3 Simulation Method**

Following recommendations in the 2006 IPCC Guidelines, the model uses a true Monte Carlo sampling method.

**A 2.2.3 Quality Control Checks on the Monte Carlo Model Output**

A number of quality control checks are completed as part of the uncertainty analysis.

**A 2.2.3.1 Checks against totals of the national emissions**

To ensure the emissions in the Monte Carlo model closely agree with the reported totals in the NIR, the emissions in the model were checked against the national totals both before and after the simulation was run. The central estimates from the model are expected to be similar to the reported emissions totals, but are not expected to match exactly.

**A 2.2.3.2 Inter-comparison between the output of the error propagation and Monte Carlo models**

A formal check to compare the output of the error propagation and Monte Carlo model is completed. The results of this comparison are discussed in **Section A 2.6**.

**A 2.2.3.3 Calculation of uncertainty on the total**

The uncertainty on the 1990 and the most recent year emissions was calculated using two different methods;

i) Using  $\frac{1.96s.d}{\mu}$

ii) Using  $\frac{(97.5 \text{ percentile} - 2.5 \text{ percentile})}{2 \times \mu}$

The first method uses the standard deviation calculated by @RISK and the mean to give an percentage uncertainty, while the second method uses the 95% confidence interval given by the percentiles quoted. When a distribution is completely normally distributed, the two methods should give the same results. However, when a distribution is skewed the two methods diverge, since the variance is dominated by outliers which aren't necessarily accounted for in the 95% confidence interval. The overall uncertainty quoted in **Table A 2.4.1** is calculated using the second method so that uncertainties in sectors that show a skewed distribution (such as agricultural soils and N<sub>2</sub>O) are better represented.

Calculating the uncertainty using both of these methods allows us to check that the Monte Carlo analysis is behaving in the way we would expect, and that convergence of the distributions is being achieved. Comparing the results using both calculations showed that the uncertainties were almost the same for gases where the distributions used were predominantly normal, but higher for N<sub>2</sub>O and the GWP weighted total, as expected.

## **A 2.3 UNCERTAINTY PARAMETERS**

The following sections present the uncertainties in emissions, and the trend in emissions according to gas.

### **A 2.3.1 Uncertainty Parameters used**

**Table A 2.3.1** to **Table A 2.3.4** summarise the uncertainty parameters used for both Approach 1 and 2 uncertainties. For all of these tables the following apply:

- Uncertainties expressed as 0.5\*R/E where R is the difference between 2.5 and 97.5 percentiles and E is the mean;
- Where custom distributions are used for the Approach 2 uncertainties the parameters are not used directly, but the below parameters should still be a reasonable indicator of the uncertainty in the distribution used for Approach 2;
- (r) means revised in comparison to previous NIR; and
- (a) means uncertainty for emission factors and activity cannot be separated, so one uncertainty that represents both is displayed.

**Table A 2.3.1 Uncertainties in the activity data and emission factors for fuels used in the carbon dioxide (CO<sub>2</sub>) inventory**

Category	Fuel	1990		2017		Justification for key sources
		Activity uncertainty (%)	Emission factor uncertainty (%)	Activity uncertainty (%)	Emission factor uncertainty (%)	
1A	Lubricants	50.00%	5.00%	50.00%	5.00%	It's challenging to determine the proportion of lubricant used as a fuel, hence a high activity uncertainty.
1A1	Blast Furnace Gas	1.50%	10.00%	1.00%	10.00%	Overall uncertainty for all coke & steelmaking emissions is quite low but allocation to individual sources is definitely higher - we've assumed 10%
1A1	Coke Oven Coke	1.00%	10.00%	1.00%	10.00%	Overall uncertainty for all coke & steelmaking emissions is quite low but allocation to individual sources is definitely higher - we've assumed 10%
1A1	Coke Oven Gas	1.50%	10.00%	1.00%	10.00%	Overall uncertainty for all coke & steelmaking emissions is quite low but allocation to individual sources is definitely higher - we've assumed 10%
1A1	Colliery Methane	5.00%	5.00%	5.00%	5.00%	(Minor fuel in sector context)
1A1	Gas/Diesel Oil	1.80%	2.10%	1.75%	2.10%	ETS-based data, so low uncertainty.
1A1	Liquefied Petroleum Gas	25.70%	2.10%	2.50%	2.10%	The DUKES data from 2009 onwards were revised considerably in the energy / NEU split for LPG, and we have created a new split for earlier years. Chosen 2.1% EF uncertainty to be consistent with gas oil - the makeup of LPG is well understood and documented
1A1	Motor Gasoline	2.50%	2.10%	2.50%	2.10%	Outside of 1A3, the motor gasoline allocations are probably much more uncertain as they are reliant on the off-road model etc., so chosen 2.5%. Chosen 2.1% EF uncertainty to be consistent with gas oil - the makeup of motor gasoline is well understood and documented
1A1	Municipal Solid Waste	1.00%	15.00%	1.00%	15.00%	MSW quantity is known accurately. Uncertainty is in mass of fossil carbon per tonne of residual MSW. This is based on reasonable waste composition data from peer reviewed sources, adapted from landfill data.

Category	Fuel	1990		2017		Justification for key sources
		Activity uncertainty (%)	Emission factor uncertainty (%)	Activity uncertainty (%)	Emission factor uncertainty (%)	
1A1	Naphtha	50.00%	5.00%	50.00%	5.00%	DUKES are uncertain about where naphtha is used (or not), so a high activity uncertainty has been chosen. EF uncertainty chosen as 5%. The content of naphtha is quite variable - it contains a huge range of hydrocarbons from C5 up to C70+, so the exact carbon content is variable and there are about 5 different grades of naphtha according to UKPIA.
1A1	Natural Gas	2.80%	2.00%	1.00%	2.00%	ETS-based data, so low uncertainties.
1A1	Orimulsion	5.00%	5.00%	5.00%	5.00%	(Minor fuel in sector context)
1A1	Other Bituminous Coal	2.00%	2.00%	2.00%	2.00%	ETS-based data, so low uncertainties.
1A1	Other Kerosene	1.25%	5.00%	1.25%	5.00%	ETS-based data, so low uncertainties.
1A1	Other Oil: Other	11.90%	5.00%	10.00%	5.00%	(Minor fuel in sector context)
1A1	Petroleum Coke	7.80%	10.00%	5.00%	10.00%	ETS-based data, so low uncertainties. 10% chosen for EF uncertainty as there is only a small dataset for the quality of petcoke used in the sector and the CEF could be quite variable depending on the source of the pet coke.
1A1	Refinery Gas	50.00%	20.00%	25.00%	15.00%	Comparisons between EUETS and DUKES are variable over time. Risk that in earlier years the "own use" may have been mis-reported to energy stats. High uncertainty on AD. Also a variable quality fuel, so the EF is also uncertain.
1A1	Residual Fuel Oil	5.50%	2.55%	1.25%	2.55%	ETS-based data, so low uncertainties.
1A1	Scrap Tyres	15.00%	10.00%	15.00%	10.00%	Limited reported use of this fuel; only a small amount of reporting (typically cement kilns) within EUETS and a modest number of fuel quality analyses either through the BCA/MPA (trade body) or the EUETS. Also some variability in the fossil C versus bio-C content of the tyres adds to EF uncertainty.
1A2	Blast Furnace Gas	1.50%	10.00%	1.00%	10.00%	Overall uncertainty for all coke & steelmaking emissions is quite low but allocation to individual sources is definitely higher - we've assumed 10%

Category	Fuel	1990		2017		Justification for key sources
		Activity uncertainty (%)	Emission factor uncertainty (%)	Activity uncertainty (%)	Emission factor uncertainty (%)	
1A2	Coke Oven Coke	3.00%	10.00%	1.00%	10.00%	Overall uncertainty for all coke & steelmaking emissions is quite low but allocation to individual sources is definitely higher - we've assumed 10%
1A2	Coke Oven Gas	3.00%	10.00%	1.00%	10.00%	Overall uncertainty for all coke & steelmaking emissions is quite low but allocation to individual sources is definitely higher - we've assumed 10%
1A2	Colliery Methane	5.00%	5.00%	5.00%	5.00%	(Minor fuel in sector context)
1A2	Gas/Diesel Oil	20.00%	2.00%	20.00%	2.00%	Low EF uncertainty as the composition of gas oil is well understood across the time series. The AD for stationary combustion in industrial sectors is quite uncertain, however. DUKES does not distinguish between mobile and stationary sources, and other AD data sources (e.g. EU ETS) have limited coverage of gas oil use across all of 1A2.
1A2	Liquefied Petroleum Gas	25.70%	2.10%	2.50%	2.10%	The DUKES data from 2009 onwards were revised considerably in the energy / NEU split for LPG, and we have created a new split for earlier years. Chosen 2.1% EF uncertainty to be consistent with gas oil - the makeup of LPG is well understood and documented
1A2	Motor Gasoline	20.00%	2.10%	20.00%	2.10%	Outside of 1A3, the motor gasoline allocations are probably much more uncertain. Chosen 2.1% EF uncertainty to be consistent with gas oil - the makeup of motor gasoline is well understood and documented
1A2	Municipal Solid Waste	5.00%	15.00%	5.00%	15.00%	MSW quantity is known accurately. Uncertainty is in mass of fossil carbon per tonne of residual MSW. This is based on reasonable waste composition data from peer reviewed sources, adapted from landfill data.
1A2	Natural Gas	2.80%	3.00%	1.00%	3.00%	Low EF uncertainty as gas composition is monitored and reported across much of the time series, and the fuel has narrow compositional range. AD are also well understood and low uncertainty. Gas supplier data to DUKES can be checked against periodic data matching (meter point data against industry sector information).

Category	Fuel	1990		2017		Justification for key sources
		Activity uncertainty (%)	Emission factor uncertainty (%)	Activity uncertainty (%)	Emission factor uncertainty (%)	
1A2	Other Bituminous Coal	5.00%	10.00%	5.00%	10.00%	Limited compositional data over time (e.g. EU ETS data for coal is incomplete), so EF uncertainty reflects the range of composition for coal types in 1A2. AD uncertainty is moderate for 1A2, reflecting energy supplier reporting to BEIS.
1A2	Other Kerosene	6.00%	2.00%	6.00%	2.00%	(Minor fuel in sector context)
1A2	Other Oil: Other	5.00%	50.00%	5.00%	3.00%	(Minor fuel in sector context)
1A2	Patent Fuel	10.00%	3.00%	10.00%	3.00%	(Minor fuel in sector context)
1A2	Petroleum Coke	25.00%	15.00%	20.00%	15.00%	EF uncertainty reflects range of petcoke composition that may be used for fuel in 1A2. AD uncertainty is quite high as we have limited data from DUKES and not much AD from EU ETS on petcoke use.
1A2	Refinery Gas	50.00%	15.00%	50.00%	15.00%	(Minor fuel in sector context)
1A2	Residual Fuel Oil	5.50%	2.10%	1.50%	2.10%	Low EF uncertainty as the composition of fuel oil is well understood across the time series. The AD uncertainty is low in recent years as the fuel is not widely used other than by larger operators that report under EUETS. Moderate uncertainty in earlier years, when fewer routine annual AD sources.
1A2	Scrap Tyres	15.00%	10.00%	15.00%	10.00%	(See 1A1 comment – same applies here.)
1A2	non-fuel combustion	50.00%	100.00%	50.00%	100.00%	(Minor emission source in sector context)
1A3	Aviation Gasoline	20.00%	3.30%	20.00%	3.30%	Activity uncertainty is higher than many other similar fuels because of the uncertainty in the international-domestic split
1A3	Other Gas/Diesel Oil	15.00%	2.00%	15.00%	2.00%	(Minor fuel in sector context)
1A3	Jet Gasoline	20.00%	3.30%	20.00%	3.30%	Activity uncertainty is higher than many other similar fuels because of the uncertainty in the international-domestic split
1A3	Other Bituminous Coal	20.00%	6.00%	20.00%	6.00%	(Minor fuel in sector context)

Category	Fuel	1990		2017		Justification for key sources
		Activity uncertainty (%)	Emission factor uncertainty (%)	Activity uncertainty (%)	Emission factor uncertainty (%)	
1A3	liquid biofuels	5.00% (n)	5.00% (n)	5.00% (n)	5.00% (n)	Activity data are not very uncertain, as it's taken from RTFO data. There is a total potential range of 10% variability in the fossil fuel carbon content of FAME (i.e. judging from the contents of the different fatty acid types used to synthesize the FAME, the highest content is around 44.8g/kg, whilst the lowest is 40.2g/kg). In reality, these are the extremes, so a lower overall uncertainty is expected. the other liquid biofuels are consumed in much smaller quantities than FAME.
1A3b	Gas/Diesel Oil	1.80%	2.00%	1.00%	2.00%	Low EF uncertainty as the composition of gas oil is well understood across the time series. Low AD uncertainty as good corroboration between fuel sales data and estimates based on vehicle movement data.
1A3b	Liquefied Petroleum Gas	5.00%	2.00%	5.00%	2.00%	EF uncertainty is consistent with gas oil - the makeup of LPG is well understood and documented. Not a major fuel in the sector but AD are considered moderately uncertain.
1A3b	Motor Gasoline	1.00%	2.00%	1.00%	2.00%	Low EF uncertainty as the composition of petrol is well understood across the time series. Low AD uncertainty as good corroboration between fuel sales data and estimates based on vehicle movement data.
1A3d	Gas/Diesel Oil	20.00%	2.00%	20.00%	2.00%	Activity uncertainty is higher than many other similar fuels because of the uncertainty in the international-domestic split
1A3d	Residual Fuel Oil	20.00%	2.00%	20.00%	2.00%	Activity uncertainty is higher than many other similar fuels because of the uncertainty in the international-domestic split
1A4	Anthracite	1.50%	6.00%	1.00%	6.00%	Low AD uncertainty as tax data helps establish residential use. EF uncertainty reflects variability in anthracite composition.
1A4	Coke Oven Coke	3.00%	10.00%	1.00%	10.00%	(Minor fuel in sector context)
1A4	Gas/Diesel Oil	30.00%	2.00%	30.00%	2.00%	Low EF uncertainty as the composition of gas oil is well understood across the time series. High AD uncertainty as scarce data on use of this fuel, e.g. in mobile machinery, in 1A4.
1A4	Gas Works Gas	5.00%	5.00%	5.00%	5.00%	(Minor fuel in sector context)

Category	Fuel	1990		2017		Justification for key sources
		Activity uncertainty (%)	Emission factor uncertainty (%)	Activity uncertainty (%)	Emission factor uncertainty (%)	
1A4	Liquefied Petroleum Gas	25.70%	2.10%	2.50%	2.10%	The DUKES data from 2009 onwards were revised considerably in the energy / NEU split for LPG, and we have created a new split for earlier years. Chosen 2.1% EF uncertainty to be consistent with gas oil - the makeup of LPG is well understood and documented
1A4	Motor Gasoline	50.00%	2.00%	50.00%	2.00%	Low EF uncertainty as the composition of petrol is well understood across the time series. High AD uncertainty as scarce data on use of this fuel in mobile machinery in 1A4.
1A4	Natural Gas	2.80%	3.00%	2.00%	3.00%	(As for 1A2)
1A4	Other Bituminous Coal	3.00%	10.00%	3.00%	10.00%	Chosen 3% activity uncertainty as we know that there are some limitations on the coal allocation to small-scale users.
1A4	Other Kerosene	3.00%	2.00%	3.00%	2.00%	Low AD uncertainty as tax data helps establish residential use. EF uncertainty reflects narrow range of fuel composition.
1A4	Patent Fuel	3.30%	3.00%	2.00%	3.00%	(Minor fuel in sector context)
1A4	Peat	30.00%	10.00%	30.00%	10.00%	(Minor fuel in sector context)
1A4	Petroleum Coke	20.00%	15.00%	20.00%	15.00%	Limited information on the AD of use in domestic fuels which increases uncertainty. Moderate emission factor uncertainty as there is only a small dataset for the quality of petcoke used in the sector and the CEF could be quite variable depending on the source of the pet coke.
1A4	Residual Fuel Oil	5.50%	2.10%	3.00%	2.10%	(Minor fuel in sector context)
1A5	Gas/Diesel Oil	6.25%	2.05%	6.25%	2.05%	Moderate AD uncertainty as data from very few data suppliers. EF uncertainty reflects narrow range of fuel composition.
1A5	Jet Gasoline	10.00%	3.00%	10.00%	3.00%	Activity Data comes directly from fuel users so should have high confidence.
1B1	Coke Oven Gas	1.50%	10.00%	1.00%	10.00%	(Minor fuel in sector context)
1B1	Other Bituminous Coal	1.50%	6.00%	1.50%	6.00%	EF uncertainty reflects the range of composition of coal types in SSF manufacture. AD uncertainty is quite low, reflecting the small number of operators and high level of AD reporting.
1B1	petroleum coke	20.00%	10.00%	20.00%	10.00%	(Minor fuel in sector context)

Category	Fuel	1990		2017		Justification for key sources
		Activity uncertainty (%)	Emission factor uncertainty (%)	Activity uncertainty (%)	Emission factor uncertainty (%)	
1B2a	non-fuel combustion	5.00%	6.00%	5.00%	6.00%	(Minor source in UK context)
1B2b	non-fuel combustion	3.00%	6.00%	3.00%	6.00%	(Minor source in UK context)
1B2c	non-fuel combustion	5.00%	6.00%	5.00%	6.00%	(Minor source in UK context)
2A1	non-fuel combustion	1.00%	3.00%	1.00%	3.00%	EU ETS-type data collected from BCA for all sites so assume very good quality and complete.
2A2	non-fuel combustion	10.00%	5.00%	(a)	5.00%	High level of reporting in EUETS for recent years and EF reflects small range of data for carbonates used in lime production. AD uncertainty higher in earlier years.
2A3	non-fuel combustion	(a)	5.00%	(a)	5.00%	Mostly based on ETS data. Very small sites outside EU ETS; it's not certain how well EU ETS factor will apply to these non-EU ETS sites.
2A4	non-fuel combustion	2.00%	3.00%	2.00%	3.00%	(Minor source in UK context)
2B	Natural Gas	2.80%	1.25%	1.75%	1.25%	Covers both feedstock and fuel (i.e. total fuel used at the sites), so AD should be very good.
2B	Coke	1.00%	20.00%	1.00%	10.00%	(Minor source in UK context)
2B	petroleum coke	1.00%	10.00%	1.00%	10.00%	(Minor source in UK context)
2B	OPG	(a)	5.00%	(a)	5.00%	Moderate uncertainty in EF reflecting good level of reporting of fuel quality in EUETS but range of variability of process off-gases that are generated and used in the chemical sector.
2B	non-fuel combustion	2.00%	5.00%	2.00%	5.00%	(Minor source in UK context)
2B	coke oven coke	(a)	20.00%	(a)	20.00%	(Minor source in UK context)
2B	refinery gas	30.00%	5.00%	30.00%	5.00%	High uncertainty, as we deviate from DUKES. Low emission factor uncertainty, but not a well-characterised fuel.
2C	Blast Furnace Gas	2.00%	10.00%	2.00%	10.00%	Overall uncertainty in 2C is quite low and uncertainty is more about where the carbon input (from the coking coal) ends up being emitted, and less about the overall amount of carbon emitted.

Category	Fuel	1990		2017		Justification for key sources
		Activity uncertainty (%)	Emission factor uncertainty (%)	Activity uncertainty (%)	Emission factor uncertainty (%)	
2C	non-fuel combustion	2.00%	10.00%	2.00%	10.00%	Overall uncertainty in 2C is quite low and uncertainty is more about where the carbon input (from the coking coal) ends up being emitted, and less about the overall amount of carbon emitted.
2C	Coke	2.00%	10.00%	2.00%	10.00%	Good level of reporting from I&S operators across the time series.
2C	Petroleum Coke	10.00%	7.50%	10.00%	7.50%	(Minor source in UK context)
2C	coke oven coke	2.00%	5.00%	2.00%	5.00%	Activity data has low uncertainty since it's based on ETS/ISSB/DUKES. Emissions are based on regulator data, so low uncertainty.
2D	Lubricants	50.00%	50.00%	50.00%	50.00%	It's challenging to determine the size of the recovered lubricant market, as this is outside the scope of energy statistics, hence a high activity uncertainty. The fraction of lubricant incidentally oxidised is also highly uncertain, so should be reflected in a high EF uncertainty.
2D	Petroleum Coke	20.00%	30.00%	20.00%	30.00%	(Minor source in UK context)
2D	Petroleum Waxes	10.00%	50.00%	10.00%	50.00%	(Minor source in UK context)
2D	non-fuel combustion	25.00%	2.00%	25.00%	2.00%	Some uncertainty as to the proportion of HDVs requiring urea and how much is needed per vehicle. Very low EF uncertainty because carbon content of urea solution known accurately.
3G	non-fuel combustion	(a)	20.90%	(a)	20.90%	Reflects overall uncertainty of AD and EF for carbonate application to soils.
3H	non-fuel combustion	(a)	50.00%	(a)	50.00%	(Minor source in UK context)
4A	non-fuel combustion	1.00%	35.00%	1.00%	35.00%	High uncertainty reflects modelled assumptions and limited AD, and is focussed in the EF parameter.
4B	non-fuel combustion	1.00%	45.00%	1.00%	45.00%	High uncertainty reflects modelled assumptions and limited AD, and is focussed in the EF parameter.
4C	non-fuel combustion	1.00%	50.00%	1.00%	50.00%	High uncertainty reflects modelled assumptions and limited AD, and is focussed in the EF parameter.
4D	non-fuel combustion	1.00%	100.00%	1.00%	100.00%	High uncertainty reflects modelled assumptions and limited AD, and is focussed in the EF parameter.

Category	Fuel	1990		2017		Justification for key sources
		Activity uncertainty (%)	Emission factor uncertainty (%)	Activity uncertainty (%)	Emission factor uncertainty (%)	
4E	non-fuel combustion	1.00%	50.00%	1.00%	50.00%	High uncertainty reflects modelled assumptions and limited AD, and is focussed in the EF parameter.
4F	non-fuel combustion	(a)	50.00%	(a)	50.00%	High uncertainty reflects modelled assumptions and limited AD, and is focussed in the EF parameter.
4G	non-fuel combustion	1.00%	45.00%	1.00%	45.00%	High uncertainty reflects modelled assumptions and limited AD, and is focussed in the EF parameter.
5C	Municipal Solid Waste	300.00%	40.00%	1.00%	15.00%	(Minor source in UK context)
5C	Clinical waste	300.00%	40.00%	5.00%	20.00%	(Minor source in UK context)
5C	Chemical waste	300.00%	40.00%	10.00%	30.00%	(Minor source in UK context)
5C	non-fuel combustion	300.00%	40.00%	300.00%	40.00%	Unauthorised and widely dispersed activity, estimated from indirect data sources so high uncertainty. Significant uncertainty in the composition of material burnt.

**Table A 2.3.2 Estimated uncertainties in the activity data and emission factors used in the methane (CH<sub>4</sub>) inventory**

Category	Fuel	1990		2017		Justification for key sources
		Activity uncertainty (%)	Emission factor uncertainty (%)	Activity uncertainty (%)	Emission factor uncertainty (%)	
1A1		1.50%	50.00%	1.00%	50.00%	Minor source but uncertainty mainly reflects uncertainty in the EF from combustion of biomass.
1A2		1.50%	50.00%	1.00%	50.00%	As above.
1A3	Aviation Gasoline	20.00%	78.50%	20.00%	78.50%	(Minor source in UK context)
1A3	Jet Gasoline	20.00%	78.50%	20.00%	78.50%	(Minor source in UK context)
1A3	Other Gas/Diesel Oil	15.00%	130.00%	15.00%	130.00%	(Minor source in UK context)
1A3	Other Bituminous Coal	20.00%	110.00%	20.00%	110.00%	(Minor source in UK context)

Category	Fuel	1990		2017		Justification for key sources
		Activity uncertainty (%)	Emission factor uncertainty (%)	Activity uncertainty (%)	Emission factor uncertainty (%)	
1A3b	Gas/Diesel Oil	1.80%	130.00%	1.00%	130.00%	Road transport fuel sales well documented, so uncertainty in AD should be low. Uncertainty in EF reflects the variability in EFs for the range of vehicle (car, van, HGV) and road types.
1A3b	Motor Gasoline	1.00%	75.00%	1.00%	75.00%	Road transport dominates consumption of these fuels, so uncertainty in AD should be low. Uncertainty in EF reflects the variability in EFs for petrol cars and road types. Lower uncertainty than diesel vehicles because consumption dominated by only one vehicle type.
1A3b	Liquefied Petroleum Gas	5.00%	130.00%	5.00%	130.00%	(Minor source in UK context)
1A3d	Gas/Diesel Oil	20.00%	130.00%	20.00%	130.00%	Uncertainty in AD due to uncertainty in getting domestic/international split from bottom-up method. Uncertainty in EF should be consistent with other 1A3 gas oil
1A3d	Residual Fuel Oil	20.00%	130.00%	20.00%	130.00%	Uncertainty in AD due to uncertainty in getting domestic/international split from bottom-up method.
1A4		1.50%	50.00%	1.00%	50.00%	Minor source but uncertainty mainly reflects uncertainty in the EF from combustion of biomass.
1A5		7.07%	65.55%	7.07%	65.55%	(Minor source in UK context)
1B1	Coke Oven Gas	1.50%	50.00%	1.00%	50.00%	(Minor source in UK context)
1B1	non-fuel combustion	2.00%	20.00%	2.00%	20.00%	High EF uncertainty reflects the modelled estimates of emissions from coal mines.
1B1	wood	(a)	50.00%	(a)	50.00%	(Minor source in UK context)
1B2a	non-fuel combustion	5.00%	20.00%	5.00%	20.00%	Good level of operator reporting. EF uncertainty reflects limited scope for verifying the factors applied in deriving operator estimates from fugitives especially.
1B2b	non-fuel combustion	3.00%	20.00%	3.00%	20.00%	As above.
1B2c	non-fuel combustion	5.00%	20.00%	5.00%	20.00%	As above.
2A4		(a)	100.00%	(a)	100.00%	(Minor source in UK context)
2B		(a)	20.00%	(a)	20.00%	(Minor source in UK context)
2C	Blast Furnace Gas	2.00%	50.00%	2.00%	50.00%	(Minor source in UK context)

Category	Fuel	1990		2017		Justification for key sources
		Activity uncertainty (%)	Emission factor uncertainty (%)	Activity uncertainty (%)	Emission factor uncertainty (%)	
2C	non-fuel combustion	1.00%	50.00%	1.00%	50.00%	(Minor source in UK context)
2C	coke oven coke	2.00%	50.00%	2.00%	50.00%	Activity data has low uncertainty since it's based on ETS/ISSB/DUKES. Emissions are based on literature factors, so a high EF uncertainty.
2D		50.00%	50.00%	50.00%	50.00%	(Minor source in UK context)
3A	non-fuel combustion	13.73%	(a)	13.73%	(a)	Based on monte carlo analysis for the agriculture model
3B	non-fuel combustion	(a) (r)	8.37% (r)	(a) (r)	8.37% (r)	Based on monte carlo analysis for the agriculture model
3F	non-fuel combustion	25.61%	(a)	25.61%	(a)	Based on monte carlo analysis for the agriculture model
3J	non-fuel combustion	50.00% (n)	50.00% (n)	50.00% (n)	50.00% (n)	Based on monte carlo analysis for the agriculture model
4A	non-fuel combustion	1.00%	55.00%	1.00%	55.00%	(Minor source in UK context)
4B	non-fuel combustion	1.00%	55.00%	1.00%	55.00%	(Minor source in UK context)
4C	non-fuel combustion	1.00%	55.00%	1.00%	55.00%	(Minor source in UK context)
4E	non-fuel combustion	1.00%	55.00%	1.00%	55.00%	(Minor source in UK context)
5A	non-fuel combustion	15.00%	46.00%	15.00%	46.00%	Moderate/high uncertainty in historical waste data, rates of decomposition and generation of methane in the modelled approach. Some extrapolation of data needed for methane utilisation, hence high uncertainty overall, across AD and EF.
5B		30.00%	99.50%	30.00%	99.50%	Scarce data for UK biological treatments. High uncertainty.
5C	Municipal Solid Waste	5.00%	75.00%	1.00%	75.00%	(Minor source in UK context)
5C	non-fuel combustion	5.00%	50.00%	5.00%	50.00%	(Minor source in UK context)
5C	wood	50.00% (n)	50.00% (n)	50.00% (n)	50.00% (n)	(Minor source in UK context)
5D	non-fuel combustion	10.00%	25.00%	10.00%	25.00%	UK industry research and model. Moderate-high uncertainty.

**Table A 2.3.3 Estimated uncertainties in the activity data and emission factors used in the nitrous oxide (N<sub>2</sub>O) inventory**

Category	Fuel	1990		2017		Justification for key sources
		Activity uncertainty (%)	Emission factor uncertainty (%)	Activity uncertainty (%)	Emission factor uncertainty (%)	
1A1		1.50%	100.00%	1.00%	100.00%	
1A2		1.50%	100.00%	1.00%	100.00%	
1A3	Aviation Gasoline	20.00%	110.00%	20.00%	110.00%	
1A3	Jet Gasoline	20.00%	110.00%	20.00%	110.00%	
1A3	Other Gas/Diesel Oil	15.00%	130.00%	15.00%	130.00%	
1A3	Other Bituminous Coal	20.00%	110.00%	20.00%	110.00%	
1A3b	Gas/Diesel Oil	1.80%	130.00%	1.00%	130.00%	Road transport dominates consumption of these fuels, so uncertainty in AD should be low. Uncertainty in EF reflects the variability in Efs for different diesel vehicle types and road types.
1A3b	Motor Gasoline	1.00%	75.00%	1.00%	75.00%	Road transport dominates consumption of these fuels, so uncertainty in AD should be low. Uncertainty in EF reflects the variability in Efs for petrol cars and road types. Lower uncertainty than diesel vehicles because consumption dominated by only one vehicle type.
1A3b	Liquefied Petroleum Gas	5.00%	130.00%	5.00%	130.00%	
1A3d	Gas/Diesel Oil	20.00%	130.00%	20.00%	130.00%	Uncertainty in AD due to uncertainty in getting domestic/international split from bottom-up method. Uncertainty in EF should be consistent with other 1A3 gas oil
1A3d	Residual Fuel Oil	20.00%	130.00%	20.00%	130.00%	Uncertainty in AD due to uncertainty in getting domestic/international split from bottom-up method.
1A4		1.50%	100.00%	1.00%	100.00%	
1A5		7.07%	85.15%	7.07%	85.15%	
1B1		1.50%	118.00%	1.00%	118.00%	
1B2a	non-fuel combustion	5.00%	110.00%	5.00%	110.00%	
1B2b	non-fuel combustion	5.00%	110.00%	5.00%	110.00%	

Category	Fuel	1990		2017		Justification for key sources
		Activity uncertainty (%)	Emission factor uncertainty (%)	Activity uncertainty (%)	Emission factor uncertainty (%)	
1B2c	non-fuel combustion	5.00%	110.00%	5.00%	110.00%	
2B1		2.00%	50.00%	2.00%	50.00%	Strong activity data, so low activity uncertainty. Assume a high uncertainty for the literature factor.
2B2		10.00%	100.00%	(a) (r)	10.00% (r)	Emission estimates for recent years have been based partially (1998-2008) or wholly (2009-2017) on continuous monitoring, and therefore will be subject to low uncertainty. The monitoring systems used at the 2 sites currently in operation are subject to an uncertainty of 5-10%. Uncertainty in earlier years is much higher due to more limited information
2B3		2.00%	100.00%	2.00%	100.00%	
2B8		10.00%	100.00%	10.00%	100.00%	
2C		1.50%	118.00%	1.00%	118.00%	
2D		50.00%	100.00%	50.00%	100.00%	
2G		100.00%	100.00%	100.00%	100.00%	
3B		(a)	68.07%	(a)	9.53% (r)	Based on separate monte carlo analysis for the agriculture model
3D		(a)	53.28%	(a)	11.16% (r)	Based on separate monte carlo analysis for the agriculture model
3F		25.63%	(a)	25.62% (r)	(a)	
3J		50.00% (n)	50.00% (n)	50.00% (n)	50.00% (n)	
4A	non-fuel combustion	1.00%	95.00% (r)	1.00%	95.00% (r)	
4B	non-fuel combustion	1.00%	55.00%	1.00%	35.00%	
4C	non-fuel combustion	1.00%	115.00% (r)	1.00%	115.00% (r)	
4D	non-fuel combustion	1.00%	100.00%	1.00%	100.00%	
4E	non-fuel combustion	1.00%	20.00%	1.00%	15.00%	
4only	non-fuel combustion	1.00%	165.00%	1.00%	165.00%	

Category	Fuel	1990		2017		Justification for key sources
		Activity uncertainty (%)	Emission factor uncertainty (%)	Activity uncertainty (%)	Emission factor uncertainty (%)	
5B		30.00%	90.00%	30.00%	90.00%	
5C		7.00%	230.00%	7.00%	230.00%	
5D		10.00%	248.00%	10.00%	248.00%	

**Table A 2.3.4 Estimated uncertainties in the activity data and emission factors used in the F-gas inventory**

Gas	Category	1990		2017		Justification for key sources
		Activity uncertainty (%)	Emission factor uncertainty (%)	Activity uncertainty (%)	Emission factor uncertainty (%)	
HFCs	2B9	(a)	10.00%	(a)	10.00%	
HFCs	2C4	5.00%	10.00%	5.00%	10.00%	
HFCs	2E1	(a)	47.15%	(a)	47.15%	
HFCs	2F1	10.00%	10.00%	10.00%	10.00%	Good UK data on refrigerant supply is used to tune the model of emissions for this sector, which means that there is a high confidence in the overall estimates of a activity for this sector. Good activity data helps mitigate the uncertainty in emissions, as leakage and disposal is directly linked to refrigerant demand.
HFCs	2F2	(a)	15.00%	(a)	15.00%	
HFCs	2F3	(a)	25.00%	(a)	25.00%	
HFCs	2F4a	5.00%	10.00%	5.00%	10.00%	
HFCs	2F4b	(a)	10.00%	(a)	10.00%	
HFCs	2F5	(a)	25.50%	(a)	25.50%	
HFCs	2F6	(a)	20.00%	(a)	20.00%	
NF <sub>3</sub>	2E1	(a)	47.15%	(a)	47.15%	

Gas	Category	1990		2017		Justification for key sources
		Activity uncertainty (%)	Emission factor uncertainty (%)	Activity uncertainty (%)	Emission factor uncertainty (%)	
PFCs	2B9	(a)	10.00%	(a)	10.00%	
PFCs	2C3	(a)	20.00%	(a)	20.00%	
PFCs	2F1	10.00%	10.00%	10.00%	10.00%	
PFCs	2F3	(a)	25.00%	(a)	25.00%	
PFCs	2G2e	(a)	47.15%	(a)	47.15%	
SF <sub>6</sub>	2C4	5.00%	5.00%	5.00%	5.00%	
SF <sub>6</sub>	2G1	(a)	5.00%	(a)	5.00%	
SF <sub>6</sub>	2G2a	(a)	17.50%	(a)	17.50%	
SF <sub>6</sub>	2G2b	(a)	40.00%	(a)	40.00%	
SF <sub>6</sub>	2G2e	(a)	47.15%	(a)	47.15%	

### A 2.3.2 General Considerations

The uncertainty parameters presented in above are based primarily on expert judgment, but where applicable will account for:

- The uncertainty range presented for data (for example the confidence interval in the 2006 IPCC guidelines for default factors)
- Monte Carlo Analysis of some of the more sophisticated models, most notably for agriculture, LULUCF and F-gases

In some cases, the individual uncertainties for the activity data and the emission factor are difficult to separate, but the uncertainty on the total emission can more easily be estimated. In these cases, the uncertainties are listed in the column marked “uncertainty in emission”.

The analysis of the uncertainties in the nitrous oxide emissions is particularly difficult because emissions sources are diverse, and few data are available to form an assessment of the uncertainties in each source. Emission factor data for the combustion sources are scarce and for some fuels are not available. The uncertainty assumed for agricultural soils (IPCC category 3D) uses a custom distribution. These parameterised functions have been defined and provided by Rothamsted Research as the best possible fit to the expected distribution of uncertainties in 1990 and the most recent year’s emissions, and are normalised in the Approach 2 methodology such that the resultant mean is consistent with the current inventory emissions in 1990 and the most recent year.

Many of the uncertainties in the emissions of HFCs, PFCs, NF<sub>3</sub> and SF<sub>6</sub> (collectively known as F-gases) are based on the recent study to update emissions and projections of F-gases (ICF, 2014). Some sources have been updated since then and the uncertainties for those sources have been revisited accordingly.

We assume that all F-gas emissions are independent between years as the technologies, gases (which have a very wide range of GWPs) used and regulations have changed drastically between the base year and the most recent year. Many HFCs in particular were not in use until the early 90s.

### A 2.3.3 Uncertainty in the Trend

The uncertainty in the trend between 1990 and the most recent year is given in **Section A 2.4.2**. In running this simulation, it was necessary to make assumptions about the degree of correlation between sources and between 1990 and the most recent year. The assumptions were as follows:

- Activity data are uncorrelated;
- Emission factors of some similar fuels are correlated;
- Land Use Change and forestry emissions are correlated (e.g. 1990 4A CO<sub>2</sub> with 4A CO<sub>2</sub> for the most recent year);
- Offshore emissions are not correlated since they are based on separate studies using emission factors appropriate for the time;
- Emission factors covered by the Carbon Factors Review (Baggott et al, 2004) are not correlated;
- Process emissions from blast furnaces, coke ovens and ammonia plants are not correlated;
- Landfill emissions were partly correlated across years in the simulation. It is likely that the emission factors used in the model will be correlated, and also the historical estimates of waste arisings will be correlated since they are estimated by extrapolation

from the year of the study. However, the reduction in emissions is due to flaring and utilisation systems installed since 1990 and this is unlikely to be correlated. As a simple estimate it was assumed that the degree of correlation should reflect the reduction in emissions since 1990;

- Emissions from agricultural soils and manure management are correlated in the base and inventory year;
- The emission factor used for sewage treatment was assumed to be correlated between years, though the protein consumption data used as activity data were assumed not to be correlated between years; and,
- Nitric acid production emission factors were assumed not to be correlated, since the mix of operating plants is very different in the most recent year compared with 1990 – only two of the original eight units are still operating in the latest inventory year, all of which now have differing levels of abatement fitted.

## **A 2.4 UNCERTAINTIES IN GWP WEIGHTED EMISSIONS**

### **A 2.4.1 Uncertainty in the emissions**

The uncertainty in the combined GWP weighted emission is given in **Table A 2.4.1**, along with uncertainties for each of the seven categorised GHGs. This is calculated as half of the 95% confidence range, i.e. the limits between which there is a 95% probability that the actual value of emissions falls. Note that the uncertainty in the GWP is not accounted for.

### **A 2.4.2 Uncertainty in the Trend**

The uncertainty estimates for all gases are summarised in **Table A 2.4.1** under 'Range of likely % change'. This indicates the range between which there is a 95% probability that the actual trend in inventory emissions falls. Note that the uncertainty in the GWP is not accounted for.

**Table A 2.4.1 Summary of Monte Carlo Uncertainty Estimates**

IPCC Source Category	Gas	1990 Emissions	2017 Emissions	95% confidence interval for 1990 emissions		Uncertainty in 1990 emissions as % of emissions in category	95% confidence interval for 2017 emissions		Uncertainty in 2017 emissions as % of emissions in category	% change in emissions between 1990 and 2017	95% confidence interval for the % change in emissions between 1990 and 2017	
				2.5 percentile	97.5 percentile		2.5 percentile	97.5 percentile			2.5 percentile	97.5 percentile
		Gg CO <sub>2</sub> e	Gg CO <sub>2</sub> e	Gg CO <sub>2</sub> e	Gg CO <sub>2</sub> e	%	Gg CO <sub>2</sub> e	Gg CO <sub>2</sub> e	%	%	%	%
TOTAL	CO <sub>2</sub> (net)	599,409	376,757	585,999	612,867	2.2%	365,909	387,601	2.9%	-37%	-39%	-36%
	CH <sub>4</sub>	132,883	51,872	106,504	171,910	24.6%	44,346	61,629	16.7%	-60%	-70%	-50%
	N <sub>2</sub> O	48,427	21,173	36,939	66,435	30.5%	18,540	24,896	15.0%	-55%	-69%	-41%
	HFC	14,405	14,194	12,257	16,535	14.8%	12,890	15,484	9.1%	-1%	-17%	19%
	PFC	1,652	371	1,571	1,732	4.9%	290	465	23.6%	-78%	-83%	-72%
	SF <sub>6</sub>	1,305	526	1,169	1,443	10.5%	464	586	11.6%	-60%	-66%	-53%
	NF <sub>3</sub>	0.4	0.5	0.2	0.6	43.6%	0.3	0.8	47.1%	41%	-35%	151%
	<b>All</b>	<b>798,081</b>	<b>464,893</b>	<b>764,043</b>	<b>842,085</b>	<b>4.9%</b>	<b>450,970</b>	<b>479,682</b>	<b>3.1%</b>	<b>-42%</b>	<b>-45%</b>	<b>-39%</b>

Uncertainty calculated as  $0.5 \cdot R/E$  where R is the difference between 2.5 and 97.5 percentiles and E is the mean calculated in the simulation.

Emissions of CO<sub>2</sub> are net emissions (i.e. sum of emissions and removals).

Emissions in this table are taken from the Monte Carlo model output. The central estimates, according to gas, for 1990 and the latest inventory year are very similar but not identical to the emission estimates in the inventory. The Executive Summary of this NIR and the accompanying CRF tables present the agreed national GHG emissions and removals reported to the UNFCCC.

## **A 2.5 SECTORAL UNCERTAINTIES**

### **A 2.5.1 Overview of the Method**

Sectoral uncertainties were calculated from the same base data used for the “by gas” analysis. The emissions and uncertainties per sector are presented in **Table A 2.5.1**. The estimates are presented in IPCC categories, which is consistent with the reporting format used within this submission to the UNFCCC, but we recommend that these estimates should only be considered as indicative.

**Table A 2.5.1 Sectoral Uncertainty Estimates**

IPCC Source Category	1990 Emissions	2017 Emissions	95% confidence interval for 2017 emissions		Uncertainty in 2017 emissions as % of emissions in category	% change in emissions between 1990 and 2017	95% confidence interval for the % change in emissions between 1990 and 2017	
			2.5 percentile	97.5 percentile			2.5 percentile	97.5 percentile
1A1a	205,372	74,445	72,649	76,245	2.4%	-64%	-65%	-62%
1A1b	17,849	13,585	11,622	15,763	15.2%	-24%	-43%	3%
1A1c	14,184	15,734	15,274	16,256	3.1%	11%	5%	18%
1A2a	21,547	9,328	8,522	10,137	8.7%	-57%	-62%	-51%
1A2b	4,359	734	696	772	5.2%	-83%	-86%	-79%
1A2c	12,117	5,316	5,041	5,595	5.2%	-56%	-60%	-52%
1A2d	4,617	1,417	1,333	1,501	5.9%	-69%	-73%	-65%
1A2e	7,614	4,115	3,905	4,326	5.1%	-46%	-50%	-41%
1A2f	7,138	2,578	2,278	2,898	12.0%	-64%	-72%	-53%
1A2g	38,769	28,063	26,609	29,530	5.2%	-28%	-34%	-21%
1A3a	1,908	1,918	1,539	2,291	19.6%	1%	-24%	33%
1A3b	111,129	114,660	112,665	116,678	1.8%	3%	1%	6%
1A3c	1,471	2,002	1,640	2,366	18.1%	36%	6%	74%
1A3d	7,717	5,522	4,546	6,497	17.7%	-28%	-44%	-10%
1A3e	228	573	467	679	18.5%	151%	94%	223%
1A4a	25,638	19,304	18,663	19,942	3.3%	-25%	-30%	-19%
1A4b	80,234	65,600	62,997	68,281	4.0%	-18%	-23%	-13%

IPCC Source Category	1990 Emissions	2017 Emissions	95% confidence interval for 2017 emissions		Uncertainty in 2017 emissions as % of emissions in category	% change in emissions between 1990 and 2017	95% confidence interval for the % change in emissions between 1990 and 2017	
			2.5 percentile	97.5 percentile			2.5 percentile	97.5 percentile
1A4c	6,309	5,300	3,737	6,858	29.4%	-16%	-48%	48%
1A5b	5,355	1,576	1,455	1,697	7.7%	-71%	-74%	-67%
1B1	23,531	848	756	941	10.9%	-96%	-97%	-96%
1B2	18,166	9,197	6,178	12,242	33.0%	-49%	-65%	-33%
2A1	7,295	4,410	4,269	4,550	3.2%	-40%	-40%	-39%
2A2	1,462	1,052	1,000	1,105	5.0%	-28%	-35%	-20%
2A3	406	368	349	387	5.1%	-9%	-10%	-9%
2A4	672	426	410	442	3.7%	-37%	-40%	-33%
2B1	1,895	1,764	1,727	1,802	2.1%	-7%	-11%	-3%
2B2	3,861	37	17	71	72.6%	-99%	-100%	-97%
2B3	19,974	-	-	-	n/a	-100%	-100%	-100%
2B6	105	181	163	199	10.0%	73%	40%	120%
2B7	232	140	132	148	5.6%	-39%	-45%	-33%
2B8	4,568	2,859	1,997	3,739	30.4%	-37%	-58%	-9%
2B9	14,418	175	158	193	9.8%	-99%	-99%	-99%
2B10	175	49	37	61	24.4%	-72%	-80%	-60%
2C	9,395	2,716	2,582	2,847	4.9%	-71%	-73%	-69%
2D	554	518	344	761	40.3%	-6%	-52%	111%

IPCC Source Category	1990 Emissions	2017 Emissions	95% confidence interval for 2017 emissions		Uncertainty in 2017 emissions as % of emissions in category	% change in emissions between 1990 and 2017	95% confidence interval for the % change in emissions between 1990 and 2017	
			2.5 percentile	97.5 percentile			2.5 percentile	97.5 percentile
2E	5	22	13	33	46.1%	359%	142%	769%
2F	0	14,168	12,865	15,457	9.1%	3964969%	2628871%	4753039%
2G	1,593	1,441	755	2,652	65.8%	-10%	-58%	92%
3A	25,393	21,464	19,220	23,935	11.0%	-15%	-28%	-1%
3B	8,174	7,523	6,675	8,498	12.1%	-8%	-22%	9%
3D	13,617	11,462	9,659	13,555	17.0%	-16%	-32%	4%
3F	244	-	-	-	n/a	-100%	-100%	-100%
3G	1,015	939	740	1,135	21.0%	-7%	-32%	25%
3H	328	344	224	501	40.3%	5%	-41%	87%
3J	410	300	182	422	39.9%	-27%	-60%	33%
4	398	246	82	572	99.5%	-38%	-84%	142%
4A	-14,785	-18,055	-24,360	-11,677	35.1%	22%	14%	30%
4B	15,253	11,408	6,544	16,263	42.6%	-25%	-35%	-15%
4C	-7,105	-8,827	-13,131	-4,435	49.3%	24%	13%	36%
4D	491	337	160	638	70.9%	-31%	-59%	-15%
4E	7,608	6,993	4,755	9,982	37.4%	-8%	-12%	-5%
4F	-	-	-	-	n/a	n/a	n/a	n/a
4G	-1,640	-2,016	-2,930	-1,098	45.4%	23%	12%	34%

IPCC Source Category	1990 Emissions	2017 Emissions	95% confidence interval for 2017 emissions		Uncertainty in 2017 emissions as % of emissions in category	% change in emissions between 1990 and 2017	95% confidence interval for the % change in emissions between 1990 and 2017	
			2.5 percentile	97.5 percentile			2.5 percentile	97.5 percentile
5A	60,302	14,227	8,133	23,049	52.4%	-76%	-88%	-55%
5B	31	1,922	1,188	3,025	47.8%	6079%	2931%	12594%
5C	1,483	308	122	709	95.1%	-79%	-94%	-33%
5D	5,005	4,177	2,867	7,070	50.3%	-17%	-41%	16%
<b>Grand Total</b>	<b>798,082</b>	<b>464,893</b>	<b>450,971</b>	<b>479,682</b>	<b>3.1%</b>	<b>-42%</b>	<b>-45%</b>	<b>-39%</b>

**Note:** Emissions in this table are taken from the Monte Carlo model output. The central estimates, according to gas, for 1990 and the latest inventory year are very similar but not identical to the emission estimates in the inventory. The Executive Summary of this NIR and the accompanying CRF tables present the agreed national GHG emissions and removals reported to the UNFCCC.

## **A 2.6 COMPARISON OF UNCERTAINTIES FROM THE ERROR PROPAGATION AND MONTE CARLO ANALYSES**

Comparing the results of the error propagation approach, and the Monte Carlo estimation of uncertainty by simulation, is a useful quality control check on the behaviour of the Monte Carlo model.

The reason that the error propagation approach is used as a reference is because the approach to the error propagation approach has been defined and checked by the IPCC, and is clearly set out in the IPCC 2000 Good Practice Guidance and the 2006 Guidelines. The UK has implemented the IPCC error propagation approach as set out in this guidance. The implementation of an uncertainty estimation by simulation cannot be prescriptive, and will depend on how the country constructs its model, and the correlations included within that model. Therefore, there is a greater likelihood of errors being introduced in the model used to estimate uncertainty by Monte Carlo simulation.

If all the distributions in the Monte Carlo model were normal, and the assumed correlations were identical, the estimated errors on the trend from the Monte Carlo model should approach those estimated by the error propagation approach if enough iterations are done. The error propagation approach assumes 100% correlation between EFs in the base and inventory year, and no correlation between sources, however in reality the nature and degree of correlation varies by source, and many distributions are not normal but heavily skewed, particularly those with very high uncertainty. These differences interact in various ways, but would be expected broadly to result in higher trend uncertainty, and lower uncertainty on the most recent year's total in the Monte Carlo uncertainty estimates compared to the error propagation approach. This can be seen in **Table A 2.6.1** which shows differences in the trend uncertainty between the error propagation and Monte Carlo approaches. These differences mostly arise from the fact that the error propagation approach only uses normal distributions, cannot account for different uncertainty parameters between the 1990 and the latest inventory year, cannot account for correlations between sources, and automatically assumes a correlation between the emission factor uncertainty in 1990 and the most recent year.

The central estimates of emissions generated by the Monte Carlo model in 1990, and those in the latest inventory year, are very close. We would not expect the central estimates from the two methods to be identical, but with a very large number of iterations we would expect the difference to tend to zero. It should be noted that the Approach 1 uncertainties base year is 1990 for N<sub>2</sub>O, CH<sub>4</sub> and CO<sub>2</sub>, but is 1995 for the F-gases; this differs from the Approach 2 uncertainties which uses 1990 emission for all gases for the starting year.

**Table A 2.6.1 Comparison of the central estimates and trends in emissions from the error propagation (Approach 1) and Monte Carlo (Approach 2) uncertainty analyses**

Method of uncertainty estimation	Central estimate (Gg CO <sub>2</sub> equivalent) <sup>b</sup>		Uncertainty on trend, 95% CI (1990 / base year to 2017) <sup>a</sup>
	Base year	2017	
Error propagation	801,849	464,453	2.3%
Monte Carlo	801,711	464,893	2.8%

**Notes:**

CI Confidence Interval

<sup>a</sup> Calculated as half the difference between 2.5 and 97.5 percentiles, assuming a normal distribution is equal to  $\pm 1.96$  standard deviations on the central estimate.

<sup>b</sup> Net emissions, including emissions and removals from LULUCF

## **ANNEX 3: Other Detailed Methodological Descriptions for Individual Source or Sink Categories, Including for KP-LULUCF Activities.**

This Annex contains background information about methods used to estimate emissions in the UK GHG inventory. This information has not been incorporated in the main body of the report because of the level of detail, and because the methods used to estimate emissions cut across sectors.

This Annex provides background information on the fuels used in the UK GHG inventory, mapping between IPCC and NAEI source categories and detailed description of methods used to estimate GHG emissions, and emission factors used in those methods.

### **A 3.1 ENERGY**

Methods for calculating emissions within the energy sector are detailed in the method statements set out in **Chapter 3**. This Annex details the emission factors used and their source, and elaborates on references commonly used within the Energy sector. The national energy balance (and how it is used) is described in **Annex 4**.

#### **A 3.1.1 Emission factors**

Emission factors used for the 2019 submission for sectors 1A and 1B can be found in the accompanying excel file: 'Energy\_background\_data\_uk\_2019.xlsx'. This can be found as one of the additional documents in on [http://naei.beis.gov.uk/reports/reports?report\\_id=954](http://naei.beis.gov.uk/reports/reports?report_id=954). Note that there can be a delay between the NIR being published on the NAEI website after official submission.

#### **A 3.1.2 Commonly used references**

This section describes data sources that are used across multiple emission sources within the energy sector, and how they are used.

##### **A 3.1.2.1 Baggott et al., 2004 – Carbon factors review**

A review of the carbon factors used in the UK GHG inventory was carried out in 2004. The report detailing this study is available from:

[http://naei.beis.gov.uk/reports/reports?report\\_id=417](http://naei.beis.gov.uk/reports/reports?report_id=417)

This aimed to validate existing emission factors and seek new data for country specific emission factors for the UK. At the time of publication this reference provided new emission factors for:

- coal from power stations;
- fuels used in the cement industry;
- a number of petroleum based fuels;

- natural gas; and
- coke oven and blast furnace gas.

Since then following updates are made to the following emission factors based on new information:

1. Coal emission factors are adjusted based on the annual variations in the GCV of the fuels using methods developed as part of the 2004 analysis (Baggott et al., 2004).

$$EF_y = EF_{ref} / GCV_{ref} * GCV_y$$

Where

$EF_y$  is the emission factor (EF) in year y

$EF_{ref}$  is the EF in the reference year (the year for which data are available)

$GCV_{ref}$  is the GCV in the reference year

$GCV_y$  is the GCV in year y

2. Since the advent of EU ETS in 2005, a number of sources of emissions from coal which had previously been reliant on Baggott et al., 2004 have now been replaced with data from the ETS, where the data set was considered suitable (high proportion of source included, and high proportion of T3 plant specific data). In addition, in 2014 the use of oxidation factors from this report was reviewed, and where suitable background evidence to support the factors used were not available, the IPCC default (of 1, IPCC 2006) has been used.
3. Emission factors for petroleum based fuels (where ETS data are not available) are still largely based on Baggott et al., 2004. These were reviewed in 2014 and compared with the defaults in the 2006 IPCC Guidelines and found to be largely within the range of the 2006 Guidelines. No new data for the UK has been identified and the emission factors from Baggott et al., 2004 are still considered to be relevant country specific emission factors.
4. During 2017-18, a review of the UK's shipping inventory was conducted (Scarborough et al., 2018). This identified new carbon emission factors for marine fuels, which replace the factors identified as part of Baggott et al., 2004.
5. Emission factors for natural gas are updated annually based on analyses from the gas network operators (Personal Communications from network operators, 2015).
6. Emission factors for coke oven gas and blast furnace gas are estimated based on a carbon balance approach (as described in Chapter 3, **MS 4**).
7. The Mineral Products Association provide data for fuels used in the cement industry annually on a confidential basis, and these are validated with EU ETS data (Personal Communication, MPA, 2018).

A new review of carbon emission factors was conducted during 2017, focusing on those factors retained from the 2004 review (Brown et al., 2017). This concluded that the factors that are currently in use are slightly more conservative than more recent values identified, and that there was no new robust evidence upon which we could justify changing the current factors. This report is available here: [http://naei.beis.gov.uk/reports/reports?report\\_id=947](http://naei.beis.gov.uk/reports/reports?report_id=947)

### **A 3.1.2.2 The Pollution Inventory and other regulators' inventories**

The Pollution Inventory (PI) has, since 1998, provided emission data for the six Kyoto gases and other air pollutant for installations regulated by the Environment Agency (EA) in England and Natural Resources Wales (NRW) in Wales. This is part of the UK's process for managing

regulated emissions from industry processes under the IPPC permitting system. The PI does contain earlier data of carbon dioxide emissions at some sites reported from 1994 onwards. The Scottish Pollutant Release Inventory (SPRI) covers processes regulated by the Scottish Environment Protection Agency (SEPA), and contains data from 2002 and 2004 onwards. The Northern Ireland Pollution Inventory (NIPI) covers processes regulated by the Northern Ireland Environment Agency and includes data for 1999 onwards.

These data are subject to some very significant limitations:

- Emissions of each pollutant are reported for each permitted installation as a whole, so emissions data for carbon dioxide, for example, can cover emissions from fuel use as well as from an industrial process. No information is given on what the source of emissions is, so a judgement has to be made about the scope of reporting;
- Permitting arrangements have changed over time, so the reporting of data is not on a consistent basis across the time-series. In general, the tendency has been to reduce the number of permits, so that whereas in the early 1990s there might have been separate permits at an industrial installation covering the boiler plant and the chemical processes, from the late 1990s onwards the tendency would be to issue a single permit to cover both. Therefore, the problems with the scope of emissions data mentioned in the first bullet point are most severe for the second half of the GHGI time series; and,
- Since 1998, process operators need only report emissions of each pollutant if those emissions exceed a reporting threshold. For example, where emissions from an installation are less than 10,000 tonnes of CO<sub>2</sub>, or 10 tonnes of methane, the operator does not need to report any emissions data for that substance in that year. Reporting thresholds are irrelevant for many of the sectors of interest to this study, since emissions would be many times higher than the thresholds, but the reporting thresholds do mean that it is necessary to consider whether the data available in the PI (and in the SPRI & NIPI for later years) will be complete.

Despite these limitations, these data are still a useful source of information for the UK GHG inventory. A considerable amount of effort is put into manually interpreting the individual returns and allocating these to appropriate categories for use in the inventory estimates by the Inventory Agency.

### **A 3.1.2.3 The Environmental and Emissions Monitoring System (EEMS) Reporting System**

Emissions from upstream oil and gas production facilities, including onshore terminals, are estimated based on operator reporting via EEMS, regulated by the BEIS Offshore Inspectorate and developed in conjunction with the trade association Oil & Gas UK (formerly the UK Offshore Operators' Association, UKOOA). The EEMS data provides a detailed inventory of point source emissions estimates, based on operator returns for the years 1995-2017. However, the EEMS data for 1995 to 1997 are not complete, frequently exhibiting duplicate entries with identical submissions by operators across years. Since the 1995 – 1997 data are not considered reliable, the EEMS dataset is only used directly to inform national inventory estimates from 1998 onwards for the following sources:

- gas flaring;
- own gas combustion;
- well testing; and
- oil loading (onshore and offshore).

*[Activity data are not routinely collected via EEMS for sources including: fugitive releases, direct process activities, oil storage or gas venting. The emissions from these sources are reported as annual estimates by operators and used directly within the inventory.]*

These EEMS-derived activity data enable greater analysis of the oil & gas emissions and related emission factors at the installation level, providing a high degree of data transparency and improving the level of detail for performing quality checks by source, by site, by year. For those sources, this has led to an improvement in data transparency and easier query of Implied Emission Factor trends. However, the EEMS activity data are only available back to 1998, and hence the activity data back to 1990 are extrapolated using the oil and gas production time-series that were collected at that time for the purposes of energy data reporting.

#### **A 3.1.2.4 Fynes & Sage (1994)**

Fynes and Sage is a country-specific reference from the mid-1990s and it includes analysis of solid fuels typically used in the UK economy in that period, deriving mass-based emission factors that are used within the UK GHGI. In the 1990s, coal used in the UK economy was predominantly mined in the UK, whereas over the time series of the inventory there has been a decline in the share of coal from UK sources and an increase in coal imports from around the world.

For recent years, for the more significant emission sources, e.g. energy industries and manufacturing industries, the inventory agency uses EFs that are derived from EU ETS data, but for smaller emission sources in the UK that still use solid fuels (such as residential, collieries) the Fynes and Sage data are retained, as there are no EU ETS data for fuels used in these sectors. There is some uncertainty regarding how representative the EFs from Fynes and Sage may be for these smaller combustion sources, but we note that the use of coal-fired technology in sectors such as collieries and residential is predominantly in the UK coal production areas, where local supplies are still available.

#### **A 3.1.3 Feedstocks and Non-Energy Use (NEU) of fuels**

The estimation methods are described within individual sections of the NIR, but are summarised here. The general approach adopted in the UK GHG inventory is to assume that emissions from all non-energy uses of fuels are zero (i.e. the carbon is assumed to be sequestered in products such as plastics and other chemicals), except for cases where emission sources can be identified and emission estimates included in the inventory. There is one exception to this, for petroleum coke where we have no information on any non-emissive uses at all, and so we adopt the conservative approach of assuming that all petroleum coke use is emissive.

The UK Inventory Agency conducts periodic studies into the fate of fuels reported as non-energy use, in order to assess the levels of stored carbon and carbon emitted for different fuels over the time series. These detailed studies are supplemented through annual data gathering and consultation with stakeholders to maintain an accurate representation of the emitted and stored carbon in the inventory.

The assumptions and estimates for individual sources are based on a review conducted in 2013-14 (Ricardo-AEA, 2014b) which included research into UK-specific activities and data sources as well as a review of the National Inventory Reports (NIRs) of other countries.

The sections below outline the emission sources from feedstock and NEU of fuels that are included in the UK GHGI, the source data and estimation methods and a summary of the time series for each of the fuel types where there is a stored carbon component in the UK energy balance. The estimates are all presented in CRF Tables 1.Ab and 1.Ad.

**Table A 3.1.1 Summary of Emission Sources for UK Fuels Allocated as Non Energy Use in UK Energy Statistics**

Fuel	IPCC	Source Category
<i>Liquid Fossil</i>		
Naphtha, Liquid Petroleum Gases (LPG), Refinery Fuel Gas (RFG) / Other Petroleum Gases (OPG), gas oil and Ethane	1A1a	Scrap tyre combustion in power stations (1994 to 2000 only).  Fossil carbon in MSW combustion in energy from waste plant.  <i>Emissions of carbon from chemical feedstock via combustion of products such as synthetic rubbers and plastics.</i>
	1A1b	Other petroleum gas use in refineries (2004, 2006 to 2011, 2013 to 2017 only).  <i>Re-allocated from non-energy use as EU ETS and trade association data indicates that DUKES data on OPG combustion are an under-report.</i>
	1A2f	Waste solvents, waste-derived fuels containing fossil carbon, in cement kilns.  Scrap tyres and waste plastics etc. combusted in cement kilns.
	1A2g	Industrial combustion of waste solvents.  <i>Emissions of carbon from chemical feedstock via combustion of products such as synthetic rubbers and solvents.</i>
	2B8	Energy recovery from process off-gases in the chemical industry.  <i>Large quantities of naphtha, butane, propane, ethane, and other petroleum gases are listed in DUKES as used for non-energy applications and these fuels are known to be used extensively as chemical feedstocks. However, EU ETS and operator data indicate that process off-gases, derived from the chemical feedstocks, are a major fuel for ethylene production processes and other petrochemical sites. Emissions of carbon are reported in 2B8.</i>
	5C	Fossil carbon in chemical waste incineration.  Fossil carbon in MSW incineration.  Fossil carbon in clinical waste incineration.  <i>Emissions of carbon from chemical feedstock via combustion of products such as synthetic rubbers and plastics.</i>
Lubricants	1A1a	Waste oil combustion in power stations.
	1A2f	Waste oil combustion in cement kilns.
	1A2g	Waste oil combustion in unclassified industry (including road-stone coating plant)
	1A3biv	Lubricant combustion in moped engines
	2D1	Lubricant combustion in aircraft, industrial, road vehicle (except moped), marine shipping and agricultural engines.
	5C	Incineration of waste oil.
Bitumen	n/a	<i>No known UK applications that lead to GHG emissions.</i>

Fuel	IPCC	Source Category
Petroleum coke	1A2f 1A2g 1A4b 2A4 2B6 2C1 2C3 2D4	<p>Based on reported energy use data by specific industries within datasets such as EU ETS and also from direct dialogue with industry representatives, the Inventory Agency re-allocates a small proportion of the reported “NEU” allocation from DUKES, and reports emissions within the UK GHG inventory. This re-allocation generates emissions for the mineral processing sector (1A2f) and other industry (1A2g) and for petcoke use in the domestic sector (1A4b).</p> <p>There are also non-combustion, emissive uses of petcoke in the UK through the use of petcoke-derived anodes in the metal processing industries. Emissions from these uses of petcoke are reported in 2C1 (electrode use in electric arc furnaces) and 2C3 (anode use in aluminium manufacture). Petroleum coke is also used in the minerals (2A4) and chemicals industries (2B6) leading to further emissions. The remaining consumption of petroleum coke is also assumed to be emissive, with emissions reported under 2D4.</p> <p><i>Note that DUKES already includes allocations of petcoke use as a fuel in combustion in power stations (1A1a) and refineries (1A1b), which are included in the UK GHG inventory.</i></p>
Other Oil	2D2	Carbon released from use of petroleum waxes. Uses of petroleum waxes includes candles, with carbon emitted during use.
<i>Solid Fossil</i>		
Coking coal (coal oils and tars)	n/a	<i>Unknown quantities of coal tar pitch are used in the manufacture of anodes for industrial processes. In the UK inventory the emissions from the use of these anodes are allocated only against petroleum coke (also used in anode production). This is a small mis-allocation of emissions between the two fuels since the carbon emitted is likely to arise from both petroleum coke and the coal tar pitch, but it is due to lack of detailed data, and does not affect the accuracy of UK inventory emissions.</i>
<i>Gaseous Fossil</i>		
Natural Gas	2B1 2B8	Ammonia and methanol production leading to direct release of CO <sub>2</sub> from natural gas used to provide the energy for steam reforming and from natural gas feedstock to the reformer. Carbon originating in the natural gas feedstock which is converted into methanol is assumed stored, however.

**A 3.1.3.1 Naphtha, Ethane, Gas Oil, Refinery/Other Fuel Gas (RFG/OPG) Propane and Butane (LPG)**

Ethane, LPG (given separately as propane & butane in the energy statistics), gas oil, refinery / other fuel gas (RFG/OPG) and naphtha are all consumed in very significant quantities for non-energy uses, primarily as feedstock in chemical manufacturing. In the UK, several major petrochemical production facilities are supplied with Natural Gas Liquid (NGL) feedstock directly from upstream production pipelines, and then utilise NGL fractions such as ethane, propane and butane in their manufacturing processes. In addition, several integrated refinery / petrochemical complexes in the UK use a proportion of the refinery fuel gas as a feedstock in petrochemical production.

The NEU allocations presented in DUKES reflect the reported disposals of these commodities as feedstocks to chemical and petrochemical companies. There are several sources of GHG emissions from this stock of “NEU” feedstock carbon, although a high proportion of carbon is stored into products and not emitted.

One large emission source known to occur in the UK is the use of carbon-containing process off-gases as a fuel within the chemical facilities. Whilst the exact source of the carbon cannot be

traced directly to a specific feedstock commodity within the UK sectoral approach, the available information from EU ETS and from consultation with operators enables the Inventory Agency to derive estimates of the GHG emissions across the time series from this emission source.

The majority of emissions are from installations manufacturing ethylene, but a number of other chemical sites report additional emissions in the EU ETS that can be attributed to the combustion of process off-gases and residues derived from the chemical feedstock. As a result, the UK inventory emissions in 2B8 now include estimates of emissions from use of process off-gases and residues at 5 ethylene manufacturing installations and 16 other chemical manufacturing installations in the UK (some of these 16 may be using process off-gases from neighbouring ethylene plant). The derivation of a time series of emission estimates from these sources is based as far as possible on reported data by plant operators within trading scheme data and other regulatory reporting mechanisms. For the early part of the time series, data on changes in plant capacity over time is used to derive the best estimates of activity and emissions by extrapolation back from later emission estimates, whilst for later years the completeness and transparency of operator reporting is greater. Therefore, whilst the uncertainty for the emission estimates in the early part of the time series is significantly greater than for those in recent years, the Inventory Agency has made best use of the available data to derive the time series estimates of emissions from "NEU" activity. Consultation with a sector trade association has also confirmed that there are no other sector estimates of this activity, or of production data across the time series, that could be used to further improve the time series (Personal communication: Chemical Industries Association, 2014).

Other emissions included within the UK GHG inventory include emissions from the destruction of chemical products, e.g. when wastes are incinerated or used as fuels. Although emissions from incineration and combustion of wastes are estimated, we cannot relate the carbon in these wastes back to individual feedstock, so it is not possible to generate reliable UK estimates of the proportion of carbon that is ultimately emitted from each individual fuel. Incineration of wastes derived from chemical feedstocks will be reported in 1A1a (in the case of plastics etc. in municipal waste incinerated with energy recovery) and in 6C (in the case of chemical, clinical and municipal wastes incinerated without energy recovery). Waste-derived fuels, including waste solvents, waste plastics and scrap tyres are used as fuels in cement kilns and other industrial plants, and emissions reported in 1A2. Tyres contain a mixture of natural and synthetic rubbers, and so where waste tyres are used as a fuel, the emission estimates take into account that only some of the carbon emitted is derived from fossil fuels.

Some propane / butane mixtures are used as a propellant in aerosols and are emitted as VOC. The UK inventory contains estimates of these VOC emissions, combined with emissions of solvents used in aerosols.

We assume that all gas oil used for non-energy purposes is used as a feedstock material, and consultation with DECC (now BEIS) energy statisticians supports this (Personal communication: Will Spry, DECC Energy Statistics team, 2014). A possible alternative use would be in explosives, but consultation with the Health and Safety Executive, who regulate the UK explosives industry, has confirmed that no UK installations manufacture explosives using gas oil or fuel oil as a feedstock (HSE, 2013).

### **A 3.1.3.2      Lubricants**

Lubricants are listed separately in the UK energy statistics and are used in vehicles and in machinery. The inventory includes estimates of emissions of carbon due to oxidation of lubricants

during use, and also includes estimates of emissions from the combustion of waste lubricants and other oils used as fuel.

UK GHG inventory estimates of the quantities of lubricants burnt as fuels are based on data from Recycling Advisory Unit, 1999; BLF/UKPIA/CORA, 1994; Oakdene Hollins Ltd, 2001 & ERM, 2008, as well as recent research to access information regarding the UK market for waste oils and the impact of European Directives to consolidate industrial emission regulations such as the Waste Incineration Directive (Oil Recycling Association, 2010). Estimates of waste oil combustion are derived for the following source categories:

- 1A1a Power stations;
- 1A2f Cement kilns; and
- 1A2f Other (unclassified) industry.

The estimated emissions for other industry assume that waste oils are used by two sectors: road-stone coating plant and garages. Other sectors may use waste oils as a fuel or as a reductant, but research to date provides no compelling evidence that there is a significant gap in the UK inventory for waste oil use by industrial operators.

The emission trends from power station use of waste lubricants reflect the fact that the Waste Incineration Directive (WID) had a profound impact on the market for waste oil, used as a fuel. It is assumed that no waste oil was burnt in power stations for the years 2006-2008, on the basis that the classification of waste oil as a fuel would have led to users being subject to the requirements of WID. In 2009 a Quality Protocol<sup>4</sup> was introduced that allowed compliant fuel produced from waste oils to be burned as non-waste and this has encouraged a resumption in the consumption of waste oil-derived fuels from 2009 onwards.

Carbon dioxide emission estimates for the oxidation of lubricants within vehicle engines and machinery, and the use of waste oils for energy are all based on a single carbon emission factor derived from analysis of the elemental composition of a series of UK-sourced samples of waste oil (Passant, 2004). The UK inventory adopts the IPCC Tier 1 methodology for lubricant use i.e. assuming that 20% of all lubricants are oxidized during use. This assumption is used for the various sub-categories of lubricant use (including road, rail, marine, off-road and air transport) given in DUKES.

### **A 3.1.3.3 Bitumen**

In the UK, bitumen is used only for applications where the carbon is stored. By far the most important of these is the use of bitumen in road dressings. The inventory does assume that a very small proportion of the carbon in the bitumen itself is emitted as VOC during road-stone coating but does not include any estimates of direct carbon emissions from uses of bitumen. Industry consultation in 2013 (UK Petroleum Industries Association, 2013; Refined Bitumen Association, 2013) has confirmed that there are no emissive applications of bitumen in the UK. Around 85% of bitumen is used in road paving, with the remaining proportion used almost entirely in the manufacture of weather-proofing materials.

### **A 3.1.3.4 Coal Oils and Tars**

Coal-tars and benzole are by-products of coke ovens. Consultation with the operators of coal ovens (Tata, 2013) and also the UK company that refines and processes coal tars and benzole

---

<sup>4</sup> <http://webarchive.nationalarchives.gov.uk/20140328084622/http://www.environment-agency.gov.uk/business/topics/waste/116133.aspx>

(Koppers UK, 2013) has confirmed that all of these materials are collected, refined and processed into a range of products that are not used as fuels. The carbon within coal tars and oils are entirely used within chemical processes. In some cases, the carbon is processed into anodes used in the ferrous and non-ferrous metals industries and then used (in the UK and overseas) within emissive applications. The UK inventory already includes estimates of emissions from UK consumption of carbon anodes within these industries, using methods based on UK metal production statistics.

Based on the evidence from process operators, the Inventory Agency allocates all of the reported coal tars and oils to Non Energy Use, i.e. assuming that all carbon is stored and there are no GHG emissions from this source-activity. The Digest of UK Energy Statistics (BEIS, 2018) also report the use of tars and benzole entirely to Non Energy Use.

Coal-tar pitch is used in the manufacture of electrodes, together with petroleum coke and a proportion of the carbon ultimately emitted, but details of input materials are scarce; emissions of carbon from these sources are included in the inventory attributed to petroleum coke. This may introduce a small mis-allocation of emissions between petroleum coke and coal oils and tars, but does not affect the UK inventory emissions total.

### **A 3.1.3.5 Natural Gas**

Natural gas is used as a chemical feedstock for the manufacture of ammonia and formerly for methanol as well, though production of the latter ceased in 2001. Emissions occur directly as a result of a) combustion of gas used to power the steam reforming process that is required for manufacture of both ammonia and methanol; b) oxidation of gas in the steam reforming, producing CO<sub>2</sub> which in the case of ammonia production is not needed and is instead emitted. The emissions are reported under 2B1 for ammonia and 2B8 for methanol.

Most of the emissions from feedstock use of natural gas in ammonia production are at source, i.e. waste gases containing carbon are emitted directly from the ammonia plant. Up until 2001, some was exported to a neighbouring methanol plant and here converted into methanol, and this CO<sub>2</sub> is treated as stored. Further CO<sub>2</sub> is captured and sold for use elsewhere, for example, in carbonated drinks and this CO<sub>2</sub> is assumed all to be emitted in the UK.

### **A 3.1.3.6 Other Oil (industrial spirit, white spirit, petroleum wax, miscellaneous products)**

White Spirit and Special Boiling Point (SBP) spirits are used exclusively for non-energy applications, and are listed in CRF Table 1.A(d) within the category 'other oil'. They are used as solvents; SBP spirits are used for industrial applications where quick drying times are needed (e.g. adhesives and other coatings) while white spirit is used as a solvent for decorative paint, as a cleaning solvent and for other applications. Estimates of VOC emissions are included in the UK inventory but no estimates are made of direct emissions of carbon from these products, as they are regarded as "not occurring".

The only emissions from this group of petroleum feedstock that are included in the UK GHG inventory are the releases of carbon from petroleum waxes which are reported under 2D2. These are accounted for in the UK inventory under the fuel category "Other Oils" in CRF Table 1Ad.

### **A 3.1.3.7 Petroleum Coke**

The evidence from industrial reporting of fuel use and from periodic surveys of fuel producers that use petroleum coke to produce domestic fuels (including smokeless fuels) indicates that the allocation of petroleum coke to combustion activities in the UK energy balance is an underestimate across all years. Therefore, the Inventory Agency generates revised estimates for all

combustion activities and effectively re-allocates some of the petroleum coke reported in DUKES as non-energy use to energy-related emission sources in the UK inventory.

Within the UK inventory, petroleum coke is included for the following energy and non-energy source categories:

- 1A1a: Power station use of petroleum coke, primarily within blends with coal at a small number of UK facilities;
- 1A1b: Refinery emissions from regeneration of catalysts;
- 1A2f: Cement industry use of petroleum coke as a fuel;
- 1A2g: Other industry use of petroleum coke as a fuel;
- 1A4b: Petroleum coke use within domestic fuels;
- 2A4: Use in brick manufacture (reported combined with other emissions e.g. from use of carbonate minerals in brickmaking);
- 2B6: Use in chemicals manufacturing;
- 2C1: Carbon emissions from electrodes used in electric arc furnaces and ladle arc furnaces and petroleum coke added to furnaces as a carbon source;
- 2C3: Carbon emissions from anode use in primary aluminium production; and
- 2D4: Petroleum coke used for non-energy applications not included elsewhere.

The UK energy balance tables in DUKES contain data on the energy use in power stations (1A1a) and refineries (1A1b), although the former are only available for 2007 onwards, and both sets of data do not always agree with the available activity data from EU ETS. The remaining energy uses in industrial combustion (1A2f, 1A2g) and the domestic sector (1A4b) are not included in DUKES. The UK Inventory Agency therefore makes independent estimates of the consumption of petroleum coke in all of these sectors.

Petroleum coke is burnt in **cement kilns** (1A2f) and has been burnt in some years at a handful of **power stations** (1A1a). A few other **large industrial sites** (1A2g) have also used the fuel. Good estimates of the consumption of petroleum coke by these large sites are available from the operators themselves, from trade associations and from EU ETS data (from 2005 onwards).

Fuel grade petroleum coke is also used as a **domestic fuel** (both smokeless and non-smokeless types, reported in 1A4b). The Inventory Agency uses data supplied by the UK fuel supply industry to estimate petroleum coke consumption for domestic fuels over the period 1990 to 2016; these estimates are broadly consistent with fuel use data published in earlier editions of DUKES for a few years in the late 1990s.

Carbon deposits build up with time on catalysts used in **refinery** processes such as catalytic cracking. These deposits need to be burnt off to regenerate the surface area of the catalyst and ensure continued effectiveness of the catalyst; emissions from this process are reported within EU ETS since 2005, with the time series estimates provided by the trade association (UKPIA, 2018) and the catalyst regeneration is treated in the inventory as use of a fuel (since heat from the process is used) and are reported under 1A1b.

Estimates of carbon released from electrodes and anodes during **metal processes** are estimated based on operator data and reported in 2C1 and 2C3. Petroleum coke content of these electrodes and anodes is estimated based on operator data and literature sources such as Best available techniques REFerence documents (BREF notes). EU ETS data also show that some petroleum coke is added to electric arc furnaces as a carbon source, and the emissions from this use are also reported in 2C1. EU ETS data are also used for emission estimates for brickmaking, which

include a component from petroleum coke. Finally, petroleum coke is used in the manufacture of titanium dioxide, with emission estimates generated from EU ETS and other operator data.

Based on data from DUKES we believe that there is some additional non-energy use of petroleum coke for most years; we assigned this residue to 2D4 and assume that it is all eventually emitted. The total fuel assigned to sector 2 is what we report as 'excluded carbon' in the CRF, table 1A(d). The consumption estimates for industrial users of petcoke as a fuel or in industrial processes are associated with low uncertainty as they are primarily based on operator reported data within the EU ETS or other regulatory reporting mechanisms. Whilst it is conceivable that other sectors may also use petroleum coke as a fuel, there is no evidence from resources such as EU ETS and Climate Change Agreement reporting that this is the case in the UK. The remaining petroleum coke consumption given in DUKES is therefore assumed to be used in various unidentified non-energy uses, all of which are assumed to be emissive. The estimates of petroleum coke used to generate fuels for the domestic sector are associated with higher uncertainty as they are based on periodic consultation with fuel suppliers to that market, and expert judgement of stakeholders.

As well as the total UK supply figure from UK energy statistics, DUKES has data on UK production, imports and exports of petroleum coke, which together provide more information on the nature of the UK consumption of petroleum coke. These data cover three distinct types of petroleum coke – catalyst coke, produced and consumed at refineries only (so no import/export or supply of fuel to other UK sectors), and then two products made in a refinery process known as coking: fuel grade (green) coke and anode-grade coke, with the former being used as a fuel, and the latter being a calcined<sup>5</sup> version of the former, used in various non-energy processes. Consultation with the DECC energy statistics team and the only UK refinery with a coking process (DECC, 2013) has confirmed that the UK produces only anode-grade coke, and exports will also be anode-grade coke, whilst imports will be fuel grade coke for use as a cost-effective fuel source or raw material for production processes under NEU.

Carbon factors for petroleum coke use are derived from industry-specific data (including EU ETS fuel analysis) in the case of cement kilns (MPA, 2018), power stations and other industrial sites (EA, 2018; SEPA, 2018). The petroleum coke factor for refinery consumption is based on trade association analysis conducted as part of the 2004 Carbon Factors Review (UKPIA, 2004) while the factor for domestic consumption is based on compositional analysis of samples of petroleum coke sold as domestic fuels (Loader et al, 2008).

These factors do show quite a large variation from sector to sector: this is probably primarily a reflection of the different requirements of fuels for different sectors (higher quality, higher carbon for some, less so for others). The highest carbon factor is for 'petroleum coke' burnt in sector 1A1b, but this fuel is actually of a different nature from the fuel burnt as petroleum coke in sectors 1A1a, 1A2f and 1A4b. In the case of 1A1b, the fuel is a build-up of carbon on catalysts used in various refinery process units, while in the other three cases, the petroleum coke is a solid by-product of a totally different refinery process (coking) which has different characteristics.

### **A 3.1.3.8 Carbon Storage Fractions: Import-Export balance for Carbon-containing Materials**

The analysis within the UK energy statistics or GHG inventory compilation system cannot accurately account for the variable (over time) import-export balance of carbon-containing

---

<sup>5</sup> Calcined petroleum coke is a processed petroleum coke that has a very high carbon content; the resulting fuel is somewhat similar to coke oven coke

materials in the UK economy. For example, where the Inventory Agency accounts for the carbon emissions from scrap tyres burned in cement kilns, power stations, incinerators and so on within the inventory estimates or from the incineration of plastics or synthetic fibres, there is no way of tracing the quantity that is derived from imported tyres/plastics/fibres.

The reported estimate of the fate of the reported NEU of fuels from the UK energy balance is based on an assumed “closed system”, whereby we account for all emissions from carbon-containing products and fuel types that are allocated as NEU as if they are derived from the fuel statistics in the UK energy balance. In reality, the source of the carbon emitted from feedstock and NEU of fuels will partly be carbon from imported materials, with UK feedstock carbon also exported and emitted elsewhere.

**A 3.1.4 Aviation (MS 7)**

**Table A 3.1.2 CAA aircraft types assigned to EMEP-EEA Emissions Inventory Guidebook aircraft types**

EMEP/EEA Aircraft Type	CAA Aircraft Types
Airbus A300 – B4	AIRBUS A300 ( ALL FREIGHTER ); AIRBUS A300-600; AIRBUS A300-600F ( ALL FREIGHTER ); AIRBUS A300B1/B2; AIRBUS A300B4-100/200; AIRBUS A300F4
Airbus A310	AIRBUS A310; AIRBUS A310-202; AIRBUS A310-300
Airbus A318	AIRBUS A318
Airbus A319	AIRBUS A319; AIRBUS A319 CJ (EXEC); BOMBARDIER CSERIES CS100
Airbus A320	AIRBUS A320-100/200
Airbus A321	AIRBUS A321
Airbus A330-200	AIRBUS A330-200
Airbus A330-300	AIRBUS A330-300
Airbus A340-200/300	AIRBUS A340-200; AIRBUS A340-300; AIRBUS A350-900
Airbus A340-500	AIRBUS A340-500
Airbus A340-600	AIRBUS A340-600
Airbus A380-800	AIRBUS A380-800
Antonov 26	ANTONOV AN-24; ANTONOV AN26B/32; DOUGLAS DC4 SKYMASTER; NAMC YS11; VICKERS VISCOUNT 700
ATR 42 - 320	ATR42-300; BRISTOL 170 FREIGHTER; CONVAIR 240/340/440; GULF AMERICAN GULFSTREAM I; ILYUSHIN IL12/IL14
ATR 42 - 45	ATR42-500
ATR 72 - 200	ATR72; ATR72 200/500; ATR72 200/500/600; HANDLEY PAGE HERALD 200; HANDLEY PAGE HERALD 700; NORD 2501 NORTALAS
Avro RJ85	AVROLINER RJ100/115; AVROLINER RJ70; AVROLINER RJ85/QT
BAe 1-11	AEROSPATIALE CARAVELLE 10B/10R; AEROSPATIALE CARAVELLE 12; AEROSPATIALE CARAVELLE 6/6R; BAE(BAC)111-200; BAE(BAC)111-300/400/475; BAE(BAC)111-500; GA GULFSTREAM 3; GULF AMERICAN GULFSTREAM II; TUPOLEV TU124

EMEP/EEA Aircraft Type	CAA Aircraft Types
Bae Jetstream 31	BAE JETSTREAM 31/32
Bae Jetstream 41	BAE JETSTREAM 41
BAe146 -100/200/300	BAE 146-100; BAE 146-200/QT; BAE 146-300
Beech 1900C airline	AEROSPATIALE (NORD)262; BEECHCRAFT 1900C/D AIRLINER; BEECHCRAFT STARSHIP MODEL 2000; DOUGLAS DC3 C47 DAKOTA
Beech Super King Air 200B	BEECHCRAFT 200 SUPERKING AIR; BEECHCRAFT B200 SUPERKING AIR; PIAGGIO P.180 AVANTI
Beech Super King Air 350	BEECHCRAFT 300 / 350 SUPER KING AIR; PIPER PA42 CHEYENNE III/IV
Boeing 727-100	BOEING 727-100/100C
Boeing 727-200	BOEING 727-200/200 ADVANCED; TUPOLEV TU154A/B; TUPOLEV TU154M
Boeing 737 100	ANTONOV 148/158; ANTONOV AN72; ANTONOV AN72 / 74; BOEING 737-100; CONVAIR 880; GULF AMERICAN GULFSTREAM 500-550; GULF AMERICAN GULFSTREAM IV; TUPOLEV TU134
Boeing 737-200	BOEING 737-200; DASSAULT-BREGUET MERCURE; GULFSTREAM G650
Boeing 737-300	BOEING 737-300
Boeing 737-400	BOEING 737-400
Boeing 737-500	BOEING 737-500
Boeing 737-600	BOEING 737-600
Boeing 737-700	BOEING 737-700; BOEING BBJ
Boeing 737-800	BOEING 737-800; BOEING 737-900; BOEING 737-900 ER
Boeing 747-100/300/800	BAC/AEROSPATIALE CONCORDE; BOEING 747-100/100F; BOEING 747-300(STRETCH UP DK); BOEING 747-300M (COMBI); BOEING 747-8 (FREIGHTER); BOEING 747-8 (I); BOEING 747SP
Boeing 747-200	ANTONOV AN-124; ANTONOV AN-225 MRIYA; BOEING 747-200B; BOEING 747-200B (COMBI); BOEING 747-200C/200F
Boeing 747-400	BOEING 747-400; BOEING 747-400F; BOEING 747-400M (COMBI)
Boeing 757-200	BOEING 757-200
Boeing 757-300	BOEING 757-300
Boeing 767 200	BOEING 767-200; BOEING 767-200ER
Boeing 767 300 ER	BOEING 767-300; BOEING 767-300ER/F; BOEING 767-400ER; BOEING 787-800 DREAMLINER; BOEING 787-900 DREAMLINER
Boeing 777-200 ER	BOEING 777-200; BOEING 777-200ER
Boeing 777-200 LRF	BOEING 777 FREIGHTER; BOEING 777 FREIGHTER SERIES; BOEING 777-200LR
Boeing 777-300	BOEING 777-300
Boeing 777-300 ER	BOEING 777-300ER
Canadair Regional Jet CRJ-200	BOMBARDIER CHALLENGER 850; BOMBARDIER REGIONAL JET 100/200; DASSAULT FALCON 7X

EMEP/EEA Aircraft Type	CAA Aircraft Types
Canadair Regional Jet CRJ-900	BOMBARDIER GLOBAL 5000; BOMBARDIER GLOBAL EXPRESS; BOMBARDIER GLOBAL EXPRESS (BD700 EXEC); BOMBARDIER REGIONAL CRJ 1000/1000 ER; BOMBARDIER REGIONAL JET CRJ900; BOMBARDIER REGIONAL JET CRJ900 ER; BOMBARDIER REGIONAL JET CRJ900 ER/LR; BOMBARDIER REGIONAL JET RJ700; BOMBARDIER REGIONAL JET RJ700ER
Cessna 208 Caravan	Other small piston aircraft
Cessna Citation II	Other small jets
Dash 8 A	BOMBARDIER DASH 8 Q100/200; DE HAVILLAND DASH 8-100
Dash 8 C	DE HAVILLAND DASH 8-300/Q300
Dash 8 D	ARMSTRONG WHITWORTH ARGOSY; BOMBARDIER DASH 8 Q400; DE HAVILLAND DASH 8 Q400
Dornier 328-110	DE HAVILLAND DHC-7 DASH-7; DORNIER 328
Embraer 110P2A Bandeirante	Other small turboprops
Embraer EMB 120 Brasilia	EMBRAER EMB 120 BRASILIA
Embraer ERJ145	BOMBARDIER CHALLENGER 300; BOMBARDIER CHALLENGER 300/350; EMBRAER LEGACY 600 (BJ135); EMBRAER LEGACY 600/650 (BJ135); EMBRAER RJ135; EMBRAER RJ145; LOCKHEED JETSTAR II
Embraer ERJ170-ERJ175	EMB ERJ170 (170-100); EMB ERJ175 (170-200); EMBRAER ERJ 170; EMBRAER ERJ170; EMBRAER ERJ175
Embraer ERJ190	CANADAIR CL-600-604 CHALLENGER; EMBRAER 195; EMBRAER ERJ190; EMBRAER ERJ195
Falcon 2000	BAE125-1000; CESSNA 680 CITATION SOVEREIGN; CESSNA 750 CITATION X; DASSAULT BREGUET FALCON 50; DASSAULT MYSTERE-FALCON 2000; DASSAULT MYSTERE-FALCON 900; DASSAULT MYSTERE-FALCON 900EX; DORNIER 328 JET; EMBRAER LEGACY 500 (EMB-550); GULFSTREAM G200 (IAI GALAXY); HAWKER 4000; LEARJET 60; RAYTHEON HAWKER HORIZON; YAKOVLEV YAK-40
Fokker F100	FOKKER 100; FOKKER 70
Fokker F27	BAE (HS) 748; FAIRCHILD HILLER FH 227B; FOKKER F27 100-400/600; FOKKER F27-500
Fokker F28	FOKKER F28-1000; FOKKER F28-2000; FOKKER F28-3000; FOKKER F28-4000/6000
Fokker F50	BAE ATP; FOKKER 50
Let L-410 Turbolet	LET 410; MITSUBISHI MU2; SHORTS SC7 SKYLINER; SHORTS SC7 SKYVAN
Lockheed C-130H Hercules	AEROSPACELINES B377SUPER GUPPY; ANTONOV AN-12; CANADAIR CL-44; ILYUSHIN IL18; LOCKHEED L100 HERCULES; LOCKHEED L188 ELECTRA; SHORTS BELFAST; V953C MERCHANTMAN
McDonnell Douglas DC-10	LOCKHEED L1011-1/100 TRISTAR; LOCKHEED L1011-200 TRISTAR; LOCKHEED L1011-500 TRISTAR; MCDONNELL-DOUGLAS DC10-10; MCDONNELL-DOUGLAS DC10-30; MCDONNELL-DOUGLAS DC10-40

EMEP/EEA Aircraft Type	CAA Aircraft Types
McDonnell Douglas DC8-50	BOEING 707 ALL SERIES; BOEING 707-120/121B; BOEING 720B; MCDONNELL-DOUGLAS DC8-10/50; MCDONNELL-DOUGLAS DC8F 54/55
McDonnell Douglas DC8-60/70	ILYUSHIN IL62; MCDONNELL-DOUGLAS DC861/3 71/3; MCDONNELL-DOUGLAS DC8-62/72; MCDONNELL-DOUGLAS DC8-71/73
McDonnell Douglas DC-9-10	MCDONNELL-DOUGLAS DC9-10/15
McDonnell Douglas DC-9-20/30/40/50	MCDONNELL-DOUGLAS DC9-20; MCDONNELL-DOUGLAS DC9-30; MCDONNELL-DOUGLAS DC9-40; MCDONNELL-DOUGLAS DC9-50
McDonnell Douglas MD-11	MCDONNELL-DOUGLAS MD11
McDonnell Douglas MD-82/87/88	BOEING 717-200; MCDONNELL-DOUGLAS MD80-MD83; MCDONNELL-DOUGLAS MD87; MCDONNELL-DOUGLAS MD88; SUKHOI RRJ95
McDonnell Douglas MD-83	Mc DONNELL DOUGLAS MD90; TUPOLEV TU104; YAKOVLEV YAK-42
Saab 2000	CONVAIR 580/600/640; DOUGLAS DC6/6A/6B/6C; SAAB 2000; VICKERS VISCOUNT 800
Saab 340B	SAAB FAIRCHILD 340
Shorts 360-300	SHORTS 330; SHORTS 360
Swearingen Metro III	FAIRCHILD SA-227 METRO 23; FAIRCHILD SA-227 METRO III; SWEARINGEN MERLIN IIA/IIB/IIIB; SWEARINGEN MERLIN IVA; SWEARINGEN METRO II
Tupolev TU 204	ILYUSHIN 76 90VD (PERM); ILYUSHIN IL76; ILYUSHIN IL86; ILYUSHIN IL96-300; TUPOLEV TU204

### A 3.1.5 Gas leakage

An overview of the time series of estimates of gas leakage at the point of use, together with overall gas use by economic sector and appliance type is presented in **Table A 3.1.3** below.

**Table A 3.1.3 Activity data and methane leakage estimates for Gas leakage at Point of Use, including cooking appliances, gas fires and boilers**

Source / Appliance type	Units	1990	1995	2000	2005	2010	2015	2016	2017
<b>Annual Gas Use</b>									
Domestic gas fires	ktoe	462	520	621	650	673	475	498	474
Domestic manual ignition hobs / cookers	ktoe	590	530	511	496	444	459	485	487
Domestic auto-ignition hobs / cookers	ktoe	211	190	183	177	159	164	174	174
Domestic auto-ignition space and water heating	ktoe	24572	26796	30491	31512	32223	24490	25617	24404

Source / Appliance type	Units	1990	1995	2000	2005	2010	2015	2016	2017
Service sector catering (ovens and hobs)	ktoe	596	762	772	774	705	570	590	572
Other service sector appliances (boilers)	ktoe	6646	8508	9818	9290	8643	8106	8412	8143
<b>Methane Leakage</b>									
Domestic cooking and gas fires	ktCH <sub>4</sub>	1.02	0.94	0.86	0.85	0.8	0.78	0.83	0.84
Domestic boilers and water heating	ktCH <sub>4</sub>	0.76	0.83	0.94	0.98	1	0.76	0.79	0.76
Service sector (all sources)	ktCH <sub>4</sub>	0.83	1.06	1.09	1.05	1	0.91	0.95	0.93
<b>Total</b>	<b>ktCH<sub>4</sub></b>	<b>2.61</b>	<b>2.83</b>	<b>2.9</b>	<b>2.88</b>	<b>2.8</b>	<b>2.46</b>	<b>2.57</b>	<b>2.52</b>

### A 3.2 INDUSTRIAL PROCESSES (CRF SECTOR 2)

There is currently no additional information for this sector in this Annex.

### A 3.3 AGRICULTURE (CRF SECTOR 3)

Note that the references for this section are included in **Section 17.4**.

**Table A 3.3.1 Livestock Population Data by Animal Type ('000 animal places)**

Livestock Category	1990	1995	2000	2005	2010	2015	2016	2017
Total cattle	12,125	11,760	11,048	10,698	10,014	9,785	9,886	9,837
- dairy cows	2,848	2,603	2,336	2,003	1,839	1,906	1,910	1,901
- all other cattle	9,277	9,157	8,713	8,695	8,175	7,879	7,977	7,936
Sheep	45,380	44,174	43,117	36,138	31,727	34,034	34,663	35,577
Pigs	7,548	7,627	6,482	4,862	4,468	4,739	4,866	4,969
Total poultry	138,381	142,267	169,773	173,909	163,842	167,579	172,607	181,811
- laying hens	33,624	31,837	28,687	29,544	28,751	28,311	29,184	30,193
- broilers	73,944	77,177	105,689	111,475	105,309	107,056	110,639	117,612
Total horses	570	684	1006	1036	1024	978	963	954
- horses kept on agricultural holdings	202	273	287	346	312	283	268	258

Livestock Category	1990	1995	2000	2005	2010	2015	2016	2017
- professional horses	62	62	70	91	91	87	87	87
- domestic horses	305	348	649	599	621	608	608	608
Goat	98	75	74	95	93	101	102	105
Deer	47	37	36	33	31	31	31	31

### A 3.3.1 Enteric Fermentation (3A)

**Table A 3.3.2 Methane Emission Factors for Livestock Emissions for 2017**

Animal type		Enteric methane kg CH <sub>4</sub> /head/year	Methane from manures kg CH <sub>4</sub> /head/year
Cattle	Dairy cows	122.70	37.24
	Dairy heifers	52.25	6.90
	Dairy replacements >1 year	49.25	6.79
	Dairy calves <1 year	42.59	5.42
	Beef cows	76.49	11.48
	Beef females for slaughter	49.74	6.81
	Bulls for breeding	59.67	9.15
	Cereal fed bull	50.49	10.27
	Heifers for breeding	48.89	6.86
	Steers	50.51	6.87
Pigs		1.50	5.20
Sheep	Ewes	6.48	0.18
	Rams	7.00	0.19
	Lambs	2.74	0.07
Goats		5.0	0.13
Horses		18.0	1.56
Deer		20.0	0.22
Poultry	Laying hens	NA	0.028
	Growing pullets	NA	0.013
	Broilers	NA	0.013
	Turkeys	NA	0.091
	Breeding flock	NA	0.013

Animal type		Enteric methane kg CH <sub>4</sub> /head/year	Methane from manures kg CH <sub>4</sub> /head/year
	Ducks	NA	0.289
	Geese	NA	0.289
	All other poultry	NA	0.013

### A 3.3.2 Manure Management (3B)

#### A 3.3.2.1 Methane emissions from animal manures

**Table A 3.3.3 Methane conversion factors for Manure Management Systems in the UK**

<sup>a</sup>No differentiation is made between crusted and non-crusted slurry storage

Manure Handling System	Methane Conversion Factor %
Liquid <sup>a</sup>	17
Daily spread	0.1
Deep bedding/farm yard manure – cattle, pigs	17
Deep bedding/farm yard manure – sheep	2.0
Pasture range and paddock	1.0
Poultry manure	1.5

**A 3.3.2.2 Nitrous Oxide emissions from Animal Waste Management Systems**

**Table A 3.3.4 Nitrogen Excretion Factors, kg N animal place<sup>-1</sup> year<sup>-1</sup> for livestock in the UK (1990-2017)**

Livestock Category	1990	1995	2000	2005	2010	2015	2016	2017
Dairy cows	86.6	88.2	94.5	102.2	105.5	109.6	107.0	109.9
Other cattle <sup>a</sup>	45.3	45.6	46.4	45.7	45.5	45.3	45.1	44.8
Sows	23.6	22.5	21.6	20.1	18.1	18.1	18.1	18.1
Gilts	15.5	15.5	15.5	15.5	15.5	15.5	15.5	15.5
Boars	28.7	27.4	26.1	24.5	21.8	21.8	21.8	21.8
Fatteners > 80 kg	20.2	19.3	18.4	17.2	15.4	15.4	15.4	15.4
Fatteners 20-80 kg	14.6	13.9	13.2	12.4	11.1	11.1	11.1	11.1
Weaners (<20 kg)	4.6	4.4	4.2	3.9	3.4	3.4	3.4	3.4
Ewes	6.7	6.8	6.8	6.8	6.8	7.0	6.8	6.8
Rams	9.1	9.1	9.1	8.9	8.9	9.0	8.9	8.9
Lambs	3.0	2.9	3.0	3.2	3.2	3.4	3.3	3.3
Goats	20.6	20.6	20.6	20.6	20.6	20.6	20.6	20.6
Horses								
– horses kept on agricultural holdings	50	50	50	50	50	50	50	50
– professional horses	129	129	129	129	129	129	129	129
– domestic horses	50	50	50	50	50	50	50	50
Deer	13.0	13.0	13.0	13.0	13.0	13.0	13.0	13.0
Laying hens	0.87	0.83	0.80	0.77	0.70	0.70	0.70	0.70
Broilers	0.64	0.59	0.55	0.49	0.40	0.40	0.40	0.40

Livestock Category	1990	1995	2000	2005	2010	2015	2016	2017
Turkeys	1.50	1.59	1.68	1.76	1.82	1.82	1.82	1.82
Pullets	0.42	0.39	0.36	0.34	0.33	0.33	0.33	0.33
Breeding flock	1.16	1.13	1.10	1.07	1.02	1.02	1.02	1.02
Ducks	1.30	1.41	1.52	1.62	1.71	1.71	1.71	1.71
Geese	1.30	1.41	1.52	1.62	1.71	1.71	1.71	1.71
Other poultry	1.30	1.41	1.52	1.62	1.71	1.71	1.71	1.71

<sup>a</sup>Weighted average for all other cattle categories

**Table A 3.3.5 Distribution of Animal Waste Management Systems (%) used for Different Animal types, 2017**

Animal Type		Liquid System	Daily Spread	Solid storage/Deep litter/Poultry litter <sup>b</sup>	Pasture Range and Paddock
Cattle	Dairy cows	61.0	8.2	9.3	21.5
	All other cattle	18.5	11.7	21.7	48.2
Pigs	All pigs	35.2	14.7	39.4	10.7
Sheep	Ewes	0.0	0.0	8.9	91.1
	Rams	0.0	0.0	0.0	100.0
	Lambs	0.0	0.0	0.6	99.4
Goats		0.0	0.0	8.2	91.8
Deer		0.0	0.0	24.9	75.1
Horses		0.0	0.0	30.1	69.9
Poultry	All poultry	0.0	35.8	61.4	2.9

**Table A 3.3.6 Nitrous Oxide Emission Factors for Animal Waste Handling Systems**

Emission source	EF (% of total N)	Uncertainty limits (95% CI)	Data source
Cattle housing			
Slurry – solid floor	0	N/A	IPCC 2006
Slurry – slatted floor	0.2	Factor of 2	IPCC 2006
FYM systems	2.0	Factor of 2	UK measurement (at storage)
Outdoor yards	0	N/A	IPCC 2006
Cattle manure storage			
Slurry – without crust	0	N/A	IPCC 2006
Slurry – with crust	0.5	Factor of 2	IPCC 2006
Weeping wall store	0.5	Factor of 2	IPCC 2006
FYM heap	0.0	N/A	Included in housing
Pig housing			
Slurry – slatted floor	0.2	Factor of 2	IPCC 2006
FYM systems	2.0	Factor of 2	UK measurement (at storage)
Pig manure storage			
Slurry (no crust)	0	N/A	IPCC 2006
FYM heap	0	N/A	Included in housing
Sheep housing (FYM)	2.0	Factor of 2	Based on cattle/pig
Sheep FYM storage	0	N/A	Included in housing
Layer housing	0.5	Factor of 2	UK measurement (at storage)
Layer manure storage	0	N/A	Included in housing
Broiler housing	0.5	Factor of 2	UK measurement (at storage)
Broiler litter storage	0	N/A	Included in housing
Duck housing	2.0	Factor of 2	Based on cattle/pig
Duck manure storage	0	N/A	Included in housing

**A 3.3.3 Agricultural Soils (3D)**
**A 3.3.3.1 Inorganic Fertiliser**
**Table A 3.3.7 EF for direct N<sub>2</sub>O emissions from managed soils in the UK inventory**

Emission source	EF (% of total N)	Uncertainty	Data source
Urea fertiliser	Non-linear function of application rate (see <b>Section 5.5.2.1</b> )		Topp et al., 2016
Other mineral fertilisers	Non-linear function of application rate and annual rainfall (see <b>Section 5.5.2.1</b> )		Topp et al., 2016
Livestock slurry	0.7475	SE 0.17328	Topp et al., 2016
Livestock solid manure (FYM, poultry manure)	0.3635	SE 0.06622	Topp et al., 2016
Sewage sludge	1.0	0.3 – 3.0	IPCC 2006
Crop residues	1.0	0.3 – 3.0	IPCC 2006
N mineralisation	1.0	0.3 – 3.0	IPCC 2006
Histosols	8 kg N <sub>2</sub> O-N/ha	0 - 24	IPCC 2006
Cattle urine	0.629	SE 0.0930	Topp et al., 2016
Cattle dung	0.193	SE 0.0212	Topp et al., 2016
Sheep, goat, horse and deer urine	0.629	SE 0.0930	Cattle value assumed
Sheep, goat, horse and deer dung	0.193	SE 0.0212	Cattle value assumed
Outdoor pig and poultry			IPCC 2006

**Table A 3.3.8 Areas of UK Crops and quantities of fertiliser applied for 2017**

Crop Type	Crop area, ha	Fertiliser, ktN	Crop Type	Crop area, ha	Fertiliser, ktN
Oats	128,425	13.5	Potatoes (maincrop)	120,557	16.2
Spring oats	23,456	2.0	Potatoes (seed or earlies)	24,343	3.3
Winter oats	9,168	0.8	Sugar beet	111,269	10.5
Spring barley	14,031	1.4	Maize	70,196	4.9
Spring barley (malting)	392,939	43.6	Grain maize	7,759	0.5
Spring barley (non-malting)	346,924	33.7	Forage maize	119,475	7.4
Winter barley	54,623	8.5	Rootcrops for stockfeed	28,509	2.2

## Other Detailed Methodological Descriptions **A3**

Crop Type	Crop area, ha	Fertiliser, ktN	Crop Type	Crop area, ha	Fertiliser, ktN
Winter barley (malting)	76,614	10.5	Leafy forage crops	4,153	0.3
Winter barley (non-malting)	291,610	45.9	Other fodder crops	48,823	4.0
Wheat	8,730	1.5	Vegetables (not-differentiated)	1,717	0.1
Wheat (milling)	637,374	126.3	Vegetables (brassicas)	4,855	0.4
Wheat (non-milling)	1,145,725	199.3	Vegetables (legumes)	39,941	0.0
Minor cereals	51,631	5.7	Vegetables (other non-legumes)	72,558	5.7
Oilseed rape	5,086	0.9	Other horticultural crops	14,690	1.2
Spring oilseed rape	8,674	1.6	Soft Fruit	8,558	0.6
Winter oilseed rape	548,637	100.0	Top Fruit	24,634	1.8
Linseed	26,375	2.2	Miscanthus	7,366	0.0
Field beans (harvested dry)	191,711	0.2	Willow (short rotation coppice)	3,610	0.0
Field peas (harvested dry)	39,864	0.0	Other field crops	25,365	2.1
Field beans and peas combined not Vining peas	0.0	0.0	Wine grapes	1,992	0.1
Fruit (mixed top & soft fruit)	11	0.0			
Permanent grass	6,135,379	297.5	Temporary grass	1,144,430	112.3

**Table A 3.3.9 Trends in area grown ('000 ha) and N fertiliser applied (kg/ha) for the major UK crops, 1990-2017**

Year	Wheat		Spring barley		Winter barley		Main crop potatoes		Oilseed rape		Grass leys (<5yrs)		Permanent grassland	
	'000 ha	kg/ha N	'000 ha	kg/ha N	'000 ha	kg/ha N	'000 ha	kg/ha N	'000 ha	kg/ha N	'000 ha	kg/ha N	'000 ha	kg/ha N
1990	2,014	183	635	90	882	140	148	184	390	225	1,606	166	5,316	108
1991	1,980	187	552	89	841	140	148	185	440	221	1,603	168	5,334	107
1992	2,067	185	515	89	784	141	151	175	421	196	1,579	157	5,286	94
1993	1,759	185	518	91	650	136	143	189	377	179	1,567	146	5,278	100
1994	1,811	186	481	94	628	143	138	191	404	179	1,456	170	5,375	110
1995	1,859	193	504	97	689	144	144	176	354	187	1,407	170	5,375	108
1996	1,977	185	518	93	749	140	149	171	356	190	1,396	166	5,347	104
1997	2,036	192	518	94	839	143	133	166	445	199	1,394	147	5,290	103
1998	2,045	182	484	91	769	135	131	186	507	192	1,302	156	5,365	99
1999	1,847	185	631	99	548	142	148	153	495	196	1,226	180	5,449	102
2000	2,086	188	539	106	589	146	138	157	395	189	1,226	142	5,363	90
2001	1,635	185	783	109	462	144	137	153	446	196	1,205	130	5,584	84
2002	1,996	189	555	110	546	150	129	152	436	194	1,243	135	5,519	77
2003	1,836	197	621	107	455	148	118	149	549	194	1,200	128	5,683	75
2004	1,990	190	587	101	420	144	121	164	498	171	1,246	117	5,620	71
2005	1,870	188	553	98	384	140	113	164	588	186	1,193	111	5,711	66
2006	1,836	181	494	100	388	134	117	142	568	178	1,137	106	5,967	60
2007	1,830	183	515	97	383	134	112	136	674	179	1,176	98	5,965	56

Year	Wheat		Spring barley		Winter barley		Main crop potatoes		Oilseed rape		Grass leys (<5yrs)		Permanent grassland	
	'000 ha	kg/ha N	'000 ha	kg/ha N	'000 ha	kg/ha N	'000 ha	kg/ha N	'000 ha	kg/ha N	'000 ha	kg/ha N	'000 ha	kg/ha N
2008	2,080	179	616	92	416	134	114	150	598	180	1,141	92	6,036	45
2009	1,814	188	749	98	411	137	118	169	581	167	1,262	88	6,081	48
2010	1,939	192	539	96	382	140	114	137	642	186	1,231	98	5,925	53
2011	1,969	194	611	99	359	140	120	157	705	186	1,278	91	5,877	51
2012	1,992	194	618	99	385	144	123	141	756	183	1,357	92	5,799	50
2013	1,615	183	903	108	310	143	114	160	715	149	1,390	94	5,802	55
2014	1,936	192	651	108	429	145	115	149	675	186	1,396	97	5,824	51
2015	1,832	193	659	105	442	148	105	162	652	190	1,135	96	5,886	49
2016	1,821	189	683	107	439	147	114	141	579	182	1,144	94	6,118	50
2017	1,792	183	754	104	423	154	121	134	562	182	1,144	98	6,135	48

**A 3.3.3.2 Crop Residues**

**Table A 3.3.10 Parameter values for crop residue management.**

Crop	Crop Harvest Index <sup>a</sup>	Above Ground Residue Retained after harvest	IPPC Crop Yield To Above Ground Residue Slope <sup>b</sup>	IPPC Crop Yield To Above Ground Residue Intercept <sup>b</sup>	IPCC Above To Below Ground Residue ratio
Oats	0.46	0.5	NA	NA	0.25
Spring oats	0.46	0.5	NA	NA	0.25
Winter oats	0.46	0.5	NA	NA	0.25
Spring barley	0.52	0.5	NA	NA	0.22
Spring barley (malting)	0.52	0.5	NA	NA	0.22
Spring barley (non-malting)	0.52	0.5	NA	NA	0.22
Winter barley	0.52	0.5	NA	NA	0.22
Winter barley (malting)	0.52	0.5	NA	NA	0.22
Winter barley (non-malting)	0.52	0.5	NA	NA	0.22
Wheat	0.50	0.5	NA	NA	0.23
Wheat (milling)	0.50	0.5	NA	NA	0.23
Wheat (non-milling)	0.50	0.5	NA	NA	0.23
Minor cereals	0.49	0.5	NA	NA	0.23
Oilseed rape	0.30	1	NA	NA	0.35
Spring oilseed rape	0.30	1	NA	NA	0.35
Winter oilseed rape	0.30	1	NA	NA	0.35
Linseed and Flax	0.38	0.5	NA	NA	0.35
Linseed	0.38	1	NA	NA	0.35
Flax	0.38	0.2	NA	NA	0.35
Field beans (harvested dry)	NA	1	1.13	0.85	0.19
Field peas (harvested dry)	NA	1	1.13	0.85	0.19
Field beans and peas combined not Vining peas	NA	1	1.13	0.85	0.19
Potatoes	NA	1	0.10	1.06	0.20
Potatoes (maincrop)	NA	1	0.10	1.06	0.20
Potatoes (seed or earlies)	NA	1	0.10	1.06	0.20
Sugar beet	NA	1	1.07	1.54	0.20
Maize	NA	1	1.03	0.61	0.22
Grain maize	NA	1	1.03	0.61	0.22
Forage maize	NA	0.15	1.03	0.61	0.22
Rootcrops for stockfeed	NA	0.15	1.07	1.06	0.20
Leafy forage crops	NA	0.15	0.30	0.00	0.35
Other fodder crops	NA	0.1	NA	NA	0.35
Vegetables (not-differentiated)	NA	1	0.30	0.00	0.35
Vegetables (brassicas)	NA	1	0.30	0.00	0.35
Vegetables (legumes)	NA	1	0.30	0.00	0.35
Vegetables (other non-legumes)	NA	1	0.30	0.00	0.35
Other horticultural crops	NA	1	0.30	0.00	0.35
Soft Fruit	NA	1	0.20	0.00	0.35

Crop	Crop Harvest Index <sup>a</sup>	Above Ground Residue Retained after harvest	IPPC Crop Yield To Above Ground Residue Slope <sup>b</sup>	IPPC Crop Yield To Above Ground Residue Intercept <sup>b</sup>	IPCC Above To Below Ground Residue ratio
Top Fruit	NA	1	0.20	0.00	0.35
Miscanthus	NA	1	1.00	0.00	0.35
Willow (short rotation coppice)	NA	1	1.00	0.00	0.35
Other field crops	0.52	0.5	NA	NA	0.22
Wine grapes	NA	1	0.20	0.00	0.35
Fruit (mixed top & soft fruit)	NA	1	0.20	0.00	0.35
Field beans and peas combined	NA	1	1.13	0.00	0.19

<sup>a</sup>Where 'NA' appears in the Harvest Index column, it indicates that the IPCC 2006 method was used; <sup>b</sup>where 'NA' appears in the IPCC slope or intercept column, it means that the Harvest Index approach was used

**Table A 3.3.11 N concentrations in above and below ground biomass**

Crop	Below Ground N, kg N/[t DM]	Crop Residue Above Ground N, kg N/[t DM]
Oats	8.0	5.4
Spring oats	8.0	5.4
Winter oats	8.0	5.4
Spring barley	14.0	6.7
Spring barley (malting)	14.0	6.7
Spring barley (non-malting)	14.0	6.7
Winter barley	14.0	6.7
Winter barley (malting)	14.0	6.7
Winter barley (non-malting)	14.0	6.7
Wheat	9.0	6.2
Wheat (milling)	9.0	6.2
Wheat (non-milling)	9.0	6.2
Minor cereals	9.0	6.6
Oilseed rape	11.0	9.9
Spring oilseed rape	11.0	9.9
Winter oilseed rape	11.0	9.9
Linseed and Flax	11.0	9.9
Linseed	11.0	9.9
Flax	11.0	9.9
Field beans (harvested dry)	8.0	8.0
Field peas (harvested dry)	8.0	8.0
Field beans and peas combined not Vining peas	8.0	8.0
Potatoes	14.0	17.3
Potatoes (maincrop)	14.0	17.3
Potatoes (seed or earlies)	14.0	17.3
Sugar beet	14.0	24.6
Maize	7.0	6.0
Grain maize	7.0	6.0

<b>Crop</b>	<b>Below Ground N, kg N/[t DM]</b>	<b>Crop Residue Above Ground N, kg N/[t DM]</b>
Forage maize	7.0	6.0
Rootcrops for stockfeed	14.0	12.6
Leafy forage crops	12.0	26.3
Other fodder crops	14.0	6.7
Vegetables (not-differentiated)	12.0	26.1
Vegetables (brassicas)	12.0	38.4
Vegetables (legumes)	22.0	23.2
Vegetables (other non-legumes)	22.0	16.7
Other horticultural crops	22.0	26.1
Soft Fruit	11.0	17.7
Top Fruit	11.0	3.9
Miscanthus	11.0	0.3
Willow (short rotation coppice)	11.0	0.3
Other field crops	11.0	6.7
Wine grapes	11.0	3.3
Fruit (mixed top & soft fruit)	11.0	8.1
Field beans and peas combined	8.0	8.0

**A 3.3.3.3 Mineralisation****Table A 3.3.12 Mineralised N from soils**

	1990	1995	2000	2005	2010	2013	2014	2015	2016	2017
N in mineral soils that is mineralised as a result of historical land use change to Cropland (kt N/y)	39.34	44.12	52.04	72.85	86.44	87.53	87.88	88.21	88.54	88.54
N in mineral soils that is mineralised as a result of Cropland Management (kt N/y)	0.0002287	0.0002600	0.0002423	0.0002411	0.0002338	0.0002277	0.0002219	0.0002314	0.0002169	0.0002169
Direct N <sub>2</sub> O emissions from mineralised N as a result of historical land use change to Cropland, kt N <sub>2</sub> O/y	0.62	0.69	0.82	1.14	1.36	1.38	1.38	1.39	1.39	1.39

	1990	1995	2000	2005	2010	2013	2014	2015	2016	2017
Direct N <sub>2</sub> O emissions from mineralised N as a result of Cropland Management, kt N <sub>2</sub> O/y	0.000004	0.000004	0.000004	0.000004	0.000004	0.000004	0.000003	0.000004	0.000003	0.000003
Indirect N <sub>2</sub> O emissions from mineralised N as a result of historical land use change to Cropland and Cropland management (kt N <sub>2</sub> O/y)	0.14	0.16	0.18	0.26	0.31	0.31	0.31	0.31	0.31	0.31
Total N <sub>2</sub> O emissions from Mineralisation (kt N/y)	0.76	0.85	1.00	1.40	1.66	1.69	1.69	1.70	1.70	1.70

**A 3.3.3.4 Histosols**


The total area used to calculate emissions from histosols is 2857 km<sup>2</sup>.


### **A 3.4 LAND USE, LAND USE CHANGE AND FORESTRY (CRF SECTOR 4)**


The following section describes in detail the methodology used in the LULUCF sector described in **Chapter 6**.


The flow chart (**Figure A 3.1**) shows the interrelationships between different data sources and the main calculation steps.


Figure A 3.1 Data flow diagrams for each land use sub-category, showing cross-linkages between sectors: (i) 4A and 4G, (ii) 4B, (iii) 4C, (iv) 4D, (v) 4E














### A 3.4.1 Carbon stock changes due to afforestation and forest management (4A)

#### A 3.4.1.1 The Forest carbon accounting model CARBINE

Carbon uptake by the forests planted in the UK is calculated by a carbon accounting model, CARBINE, as gains and losses in pools of carbon in standing trees, litter and soil in conifer and broadleaf forests and in harvested wood products. Restocking is assumed in all forests. The method is Tier 3, as defined by IPCC (2006). Matthews *et al.* (2014) gives an overview of the CARBINE model and a comparison of its use in the 1990-2012 LULUCF inventory with the C-Flow model previously used model to forest carbon stock changes.

A key concept underlying the net exchanges of C between forests and the other C reservoirs is that it can be inferred from the changes in C stocks. So CARBINE simulates forest C stock changes represented by tree biomass growth, mortality and subsequent loss. The CARBINE model is primarily dedicated to reproducing the UK forest conditions.

The model as used for this inventory consists of three sub-models or 'compartments' which estimate carbon stocks in the forest biomass, soil, and harvested wood products. The forest biomass carbon sub-model is further compartmentalised to represent fractions due to tree stems, branches, foliage, and roots.

The main driving module of CARBINE consists of a set of computerised mathematical functions and algorithms describing the accumulation (and loss) of carbon in tree biomass of different forestry systems at the per-hectare scale. Different functions and algorithms are used to represent distinct forestry systems, defined in terms of:

- Tree species composition
- Tree growth rate (yield class)
- Management regime applied.

The tree species and growth rates represented are based on yield models originally produced by the British Forestry Commission (Matthews et al., 2016a, 2016b). The tree species covered include examples for coniferous species of spruces, pines, firs, larches, cedars, cypresses and all the major temperate and boreal broadleaf tree species. Growth rates in terms of mean annual increment (MAI) of stem volume can be represented in the range from 2 m<sup>3</sup> ha<sup>-1</sup> yr<sup>-1</sup> up to 30 m<sup>3</sup> ha<sup>-1</sup> yr<sup>-1</sup>.

The CARBINE model uses standard estimates for wood density wood carbon content to derive stem biomass from the stem volume predictions simulated by the M1 model (Lavers and Moore, 1983; Jenkins et al., 2011; Matthews, 1993). Wood and bark density along with the carbon content differences are not taken into account. The density of bark is lower than that of wood (Aaron, 1970), but the carbon content is usually higher (Matthews, 1993), hence it is assumed that the two effects cancel out. The volume, biomass and carbon in tree foliage, branches, and coarse and fine roots are derived from the results for the stem by applying expansion factors. Species-specific biomass expansion factors are applied for these calculations.

The biomass of a component of interest is calculated by multiplying stem biomass by a corresponding expansion factor. The UK species-specific crown and root biomass expansion factors were derived from the report of Jenkins *et al.*, (2011) report. Branch biomass is calculated by subtracting foliage biomass from crown biomass. The coarse root biomass expansion factor includes an allowance for stump material. Robust information on foliage expansion factors relevant to UK conditions were not available, hence these were obtained from scientific literature. The ratio of foliage to stem changes over time, but approaches an asymptote (Matthews et al., 1991; Matthews and Duckworth, 2005). However, the asymptote in general is more suited to older, larger trees and as such is considered not to be representative of typical forests under regular management. It was decided to use a biomass expansion relationship for trees of approximately 20 cm diameter in order to better represent managed forests. It is likely that this will ultimately underestimate foliage biomass in smaller trees, and conversely over-estimate in older, larger trees. Finally, fine root biomass is calculated with a uniform expansion factor  $\beta_r=0.02$  from a Liski et al., (2002) study. The expansion factors are not sensitive to stand age, management regime or growth rate. This approach was adopted for the simplicity and ease of implementation on the large scale simulations.

The mass of carbon in a forest was calculated from biomass by multiplying by the fraction of carbon in wood (0.5 assumed). As an example, the values used for these parameters for Sitka spruce (*P. Sitchensis*) are given in **Table A 3.4.1**. Sitka spruce is the most common species in UK forests (c. 30%); parameters for other tree species are given in Matthews *et al.* (2014).

**Table A 3.4.1 Main parameters for forest carbon flow model used to estimate carbon uptake by planting of forests of Sitka spruce (*P. Sitchensis*), yield class 12.**

Parameter	Value
Time of maximum mean annual increment (years)	60
Initial spacing (m)	2
First table age (years)	20
Age at first thinning (years)	25
Stemwood density (oven dried tonnes m <sup>-3</sup> )	0.33

<b>Parameter</b>	<b>Value</b>
Stemwood conversion loss	10%
% Branchwood left in forest	100%
% Branchwood harvested for fuel	0%
% fuel from bark	30%
% non-fuel products from bark	70%
% small roundwood (underbark) used as fuel	20%
% Pallets and fencing from small roundwood (under bark)	20%
% Paper from small roundwood (under bark)	35%
% Particleboard etc. from small roundwood (under bark)	25%
% Fuel from sawlogs (under bark)	30%
% Pallets and fencing from sawlogs (under bark)	0%
% Particleboard from sawlogs (under bark)	40%
% Structural timber from sawlogs (under bark)	30%
Root:Stem ratio	0.49
Crown:Stem ratio	0.32
Foliage:stem ratio	0.13
Fine root:stem ratio	0.02
Foliage turnover rate (annual)	0.2
Branchwood turnover rate (annual)	0.04
Coarse Root Turnover rate (annual)	0.02
Fine Root turnover rate (annual)	0.8
Underbark/overbark ratio at 15cm DBH (varies with DBH)	0.9
Ratio of thinned stem volume that is sawlog at 15cm DBH (varies with DBH)	0.05

CARBINE includes a sub-model for representing accumulation and loss of carbon in dead wood and litter. Inputs of litter are related to the standing biomass of trees and also to rates of tree mortality. Levels of tree mortality are represented implicitly in the standard Forestry Commission growth models, and explicit estimates are included in models for stands subject to no thinning, where mortality levels are high. Root and branch wood volume associated with dead trees is estimated in the same way as for living stemwood, by reference to allometric relationships. Deadwood and litter are assumed to decay according to a first order process, with rate constants that are normally set to be consistent with boreal and temperate conditions but can be adjusted

for Mediterranean and tropical conditions. The other significant input of carbon to the dead wood and litter pool is due to harvesting operations (as part of either thinning or clearfelling). The carbon in roots of harvested trees is assumed to enter the litter pool. The harvesting of stem wood is assumed to involve a conversion loss equivalent to 10% of standing stem volume, which also enters the litter pool. It is difficult to make robust assumptions about the fate of branch wood and foliage at time of harvesting. In many situations, this material will be left on-site to deteriorate and decay. Sometimes it is possible that branch wood remaining after clearfelling may be deliberately burned. There has also been an increasing interest in active harvesting of branch wood (or at least some proportion of it) to supply biomass to the Energy sector. However, currently, such practice remains very limited. For this inventory the assumption has been made that no branch wood is harvested but is left to degrade and decay on site as part of the litter pool.

The branch ATR was fixed at 4% in accordance to Canadian forest carbon accounting model CBM-CFS (Kurz et al., 2009). Deciduous species foliage turnover is assumed to be 100% (Kurz et al., 2009; Ľupek et al., 2015). Conifer species foliage ATRs were obtained by referring to relevant scientific literature. If insufficient empirical literature and data was available the species were mapped to an allometrically similar species. Coarse root annual turnover was assumed to be 2% as in the CBM-CFS (Kurz et al., 2009; Kurz and Beukema, 1996; Li et al., 2003). Fine root ATRs were mapped from the available scientific literature and the UK specific datasets provided by Vanguelova (pers. com.). The UK ATRs for fine roots were derived from Kielder forest for Sitka spruce and Alice Holt forest for oak. Lastly, root exudate ATR was set to 160% of fine root dry biomass, the upper quartile of reported exudate mass from grassland was adopted (Jones et al., 2009), because of limited understanding about forest rhizodeposition. Aboveground shed litter, foliage and branches, are accumulated in a litter layer and after partial degradation passed to the Fermenting (F) layer. Residues that are left after thinning or felling can be set to enter a litter layer. If the crop is not a forest, it is assumed that the litter and F layers are zero. The litter layer decomposition is modelled using modified ForClim-D model version (Liski et al., 2002; Perruchoud et al., 1999). Below ground litter is not included in this simulation, while the annual transfer rates are applied to foliage ( $C_f$ ) and branch ( $C_b$ ) litter biomass. They are expressed as a proportion relocated annually.

Branch and foliage litter transfer are set according to the model proposed by Liski et al. (2002). The transferred biomass is pooled and degraded by a fixed constant of 0.5, which is the average of constants given in the Liski et al. (2002) study

The new CARBINE Soil Carbon Accounting model (CARBINE-SCA; **Figure A 3.2**), is based on a simplified version of the ECOSSE model (Smith *et al.*, 2011), coupled with a litter decomposition model derived from the ForClim-D model (Perruchoud *et al.*, 1999; Liski *et al.*, 2002). Above-ground turnover of material such as foliage, branches and dead stemwood enters the litter pool, which is then broken down to F-material (Fermenting) as a function of temperature and rainfall, releasing CO<sub>2</sub>. Within the soil, a number of layers exist, each with its own set of texture (Sand, Silt, Clay) characteristics. Carbon from decayed litter, dead roots, and root exudates enters each layer and is assigned to four active pools; resistant plant material (RPM), readily decomposable plant material (DPM), biological material (BIO) and humic material (HUM). A proportion of organic carbon is also assumed to be inert, and unavailable for further activity. The active pools undergo decomposition and transference, releasing CO<sub>2</sub>. Decomposition (aerobic and anaerobic) within each pool and layer is influenced by response functions to water saturation in the soil, temperature, pH, and the presence (or not) of plant cover on the soil surface. The availability of water within each layer, and the level of saturation are largely defined from soil texture following Saxton and Rawls (2006) coupled with inputs from rainfall, (or drainage) and removal of water through evapotranspiration. In any soil layer, water above field capacity can drain to lower soil

layers, complete with any dissolved organic carbon (DOC). The rates of potential decomposition of each carbon pool and the response functions follow ECOSSE (Smith et al., 2011).

New carbon input to the soil arises from four sources:


- Recently dead root material (according to a rooting profile depth),
- Transfer from the F-material arising from the decomposition of above-ground litter,
- Secretions and exudates from the roots,
- DOC; this carbon can become available to the biological pool and enter the 'reactive material cycle'.

Turnover rates for mortality of tree components (roots, foliage etc.) are species dependent and obtained from scientific literature (see **Table A 3.4.1** for example). A full description of the model will be presented in a separate technical report.

An improved version of the soil sub-model was implemented for the 1990-2016 inventory. This included work on parameterisation of litter input from ground flora and other non-forest vegetation, assuming a decrease in the contribution of non-tree litter from that assumed in Ecosse for pasture to zero contribution at canopy closure.

A more comprehensive description of the soil sub-model will be described in a technical report.

Figure A 3.2 The CARBINE-SCA model



The harvested wood products sub-model represents wood products as long-lived and short-lived sawn timber, particleboard, paper and fuel (see also **Section 6.8**). Carbon in harvested stemwood

is allocated to these wood product categories using an assortment forecasting model that accounts for variation in product out-turn due to tree species and tree size class distribution at time of harvest (Rollinson and Gay, 1983). Wood products in primary use are assumed to decay over time with no account taken of carbon stocks in landfill or greenhouse gas emissions (due to wood products) from landfill (these are taken into account in the Waste sector).

### **A 3.4.1.2 Forest activity data: management**

Information from the Sub-compartment database (SCDB, the Forestry Commission database of information on the growth rate and management of the Forestry Commission (FC) and Natural Resources Wales (NRW) forest estate) was used to create a distribution of species and yield class (an indication of growth rate) for the FC/NRW forest estate. For the non-FC/NRW forest estate information from the NFI survey of woodlands was analysed to estimate yield class and species by age class, and scaled to represent the whole private forest estate. Data from the Forestry Commission's new planting and wood production statistics were used to assign the areas in an age class to individual years, either as areas restocked or areas newly planted.

Management of forests is represented as one of four options: Clearfell with thinnings, clearfell without thinnings, managed but not clearfelled, and not used for timber production. For the clearfell forests restocking occurs after the rotation period. For non-clearfell productive woodlands it is assumed there is a 30 year overlap of restocking and non-restocked trees. The area of land felled each year was estimated from the wood production statistics separately for both FC/NRW and private forests. The rotation periods for forests were estimated based on information on the intended management of the FC/NRW estate. This analysis gave a target rotation period for each modelled species and yield class.

The actual rotations historically applied to the forest estate are unknown and for the private forests the area of woodland used for timber production is also unknown. In order to match production, given the age class distribution of the forest, an algorithm was implemented to adjust the assumed rotations and the percentage of private sector woodland not used for timber production. This algorithm adjusts these assumptions in order to match the modelled wood production with the timber production statistics separately for the FC/NRW and private forests. It was assumed that the forests would be felled evenly over a period +/-7 years from the target rotation period. A comprehensive description of this algorithm will be presented in a separate technical report.

Information on the management of privately owned forests that is used to inform the inventory estimates, as well as a description of how forest land AD (forest land remaining forest land; land converted to forest land) are derived is included in the National Forest Accounting Plan, which can be found at:

<https://www.gov.uk/government/publications/uk-national-forestry-accounting-plan-2021-to-2025>

### **A 3.4.1.3 Forestry activity data: historical and current afforestation rates**

Irrespective of species assumptions, the variation in CO<sub>2</sub> removals from 1990 to the present is determined by the afforestation rate in earlier decades, the effect this has on the age structure in the present forest estate, and hence the average growth rate. Afforestation is assumed to occur on ground that has not been wooded for many decades, based on the assumption that if it had previously been woodland it would be in the restocking statistics rather than the new planting statistics as a result of the regulatory framework that applies to forestry in the UK.

A comparison of historical forest census data and the historical annual planting rates has been undertaken. Forest censuses were taken in 1924, 1947, 1965, 1980 and the late 1990s. The latest census (National Forest Inventory) has only just been completed. The comparison of data sources

showed that discrepancies in annual planting rates and inferred planting/establishment date (from woodland age in the forest census) are due to restocking of older (pre-1920) woodland areas and variations in the harvesting rotations. However, there is also evidence of shortened conifer rotations in some decades and transfer of woodland between broadleaved categories (e.g. between coppice and high forest). It is difficult to incorporate non-standard management in older conifer forests and broadleaved forests into the Inventory because it is not known whether these forests are on their first rotation or subsequent rotations (which would affect carbon stock changes, particularly in soils). The area of afforestation in a given year is predicted based on applying the yearly distribution from the new planting and restocking statistics to the age class inventory. Age classes prior to the availability of new planting statistics are assigned evenly to individual years. For this inventory submission the assumption was made that we can estimate the area felled for recent years based on the timber production in the year of felling. It is assumed that woodland felled is immediately restocked. As we have an estimate of the area restocked for these years, the remainder of the area for each year was assumed to be restocking or natural regeneration. For years prior to the timber production statistics (i.e. prior to 1976), an estimated ratio between restocking and afforestation was used based on the earliest data. For restocked woodland the forest area was assumed to have been restocked twice and had been managed in the same fashion and on the same rotation.

The planting data used as input to the CARBINE model come from national planting statistics from 1921 to the present (provided by the Forestry Commission) for England, Scotland and Wales and from 1900 to the present (provided by the Northern Ireland Forest Service). For England, Scotland and Wales estimates of area of woodland by species, yield class and broad age class came from analysis of the NFI (for private woodland) and the SCDB (for FC/NRW woodland).

The NFI provides woodland statistics for Great Britain, (England, Wales and Scotland), broken down by region. It comprises a digital woodland map based on comprehensive aerial photography and a field survey using 15,000 one-hectare sample squares. The digital map and field survey cover all woodland areas down to 0.5 hectares. An initial digital woodland map was published in spring 2011. The NFI woodland field survey provides direct assessments of woodland growing stock including species composition, stand structure, tree age (distribution) productivity indices, numbers of trees, and diameter and height distribution. Standing biomass (and carbon) in trees, including above and below ground biomass, can be derived from these assessments using GB-specific conversion factors and allometric equations. A complete 5-year cycle of ground survey has now been completed. NFI data do not allow the carbon stocks of deadwood or litter to be estimated. The NFI has been supplemented by an assessment of the area of small woods (woodland between 0.1 ha and 0.5 ha) to align with the minimum woodland area for UNFCCC reporting as set out in CMP.7 (Forestry Commission, 2017). The analysis of small woods area included no characterisation of the resource. Since there is currently no information on the age-distribution of the area of small woods, it was assumed to have established evenly between 1900 and 1970.

The NFI uses a lower integral open space threshold of 0.5 ha (as opposed to 1 ha), which requires a downward adjustment to areas. However, the main differences in 2010 GB woodland cover between the NFI (2982 kha) and previous estimates (2757 kha, Forestry Statistics 2010) arise from identified errors in the previous woodland survey, particularly the under-estimate of woodland areas between 0.5 and 2 hectares. Estimates of woodland loss have been assessed, which affect the total estimated woodland area in the GHGI (but are not yet reflected in the national Forestry Statistics).

We assumed that the NFI survey gives a distribution of all the non FC/NRW forest area for a base year of 2011, and the SCDB gives a distribution of all the FC/NRW forest area for a base year of 2014.

The main NFI survey includes areas of woodland >0.5 ha. An adjustment was made to the areas of woodland to account for woods between 0.1 ha and 0.5 ha. For England and Wales, the estimates are derived from a calibration of tree cover plotted in the National Tree Map (NTM) product across England and Wales<sup>6</sup>, using a comparison of manual photographic interpretation with the NTM product within a sample of 1 km square tiles. For Scotland, the estimates are derived from a direct evaluation of polygons in the map constructed for the Native Woodlands of Scotland Survey (NWSS)<sup>7</sup>, which mapped all woodland polygons in Scotland down to 0.1 hectares in size by photographic interpretation. The areas of small woods used in this inventory were based on data published in 2017 by the Forestry Commission in the report “Tree cover outside woodland in Great Britain”<sup>8</sup>.

An algorithm was used to obtain the area of woodland afforested each year by removing the area of felling from the age class distribution. The species were then allocated to this “residual distribution’ by starting in the base year and allocating the shortest rotations first. The planting years for all restocked woodland are assigned by the algorithm to give two rotations of the same length as the assigned rotation, and are thus notional. This approach was undertaken to spin up the model in terms of soil and litter in order to reach a state consistent with land that has been forest for a long period. This algorithm will be described in detail in the same technical report as the description of allocation of the management of forests.

Conifer planting on organic soil is a subset of total conifer planting. All broadleaf planting is assumed to be on non-organic soil. As explained above, the planting rates given in **Table A 3.4.2** are derived from administrative records, information on forest age class distribution from NFI field assessments and interim assumptions about the age distribution of ‘small woods’. The planting rates given in **Table A 3.4.2** are therefore significantly different to those reported as official planting statistics supported by grant-aid. The afforestation rates for each planting type in the UK have been calculated from the data and are shown in **Table A 3.4.2**.

**Table A 3.4.2      Afforestation rate of conifers and broadleaves in the United Kingdom since 1500 based on estimates of woodland area by age from the NFI and administrative records.**

Period	Planting rate (kha annum <sup>-1</sup> )		
	Conifers on all soil types	Conifers on organic soil	Broadleaves
1501-1600	0.00	0.00	0.02
1601-1700	0.06	0.00	0.68
1701-1750	0.14	0.00	2.31

<sup>6</sup> <http://www.bluesky-world.com/national-tree-map>

<sup>7</sup> <http://scotland.forestry.gov.uk/supporting/strategy-policy-guidance/native-woodland-survey-of-scotland-nwss>

<sup>8</sup> <sup>8</sup> <https://www.forestresearch.gov.uk/tools-and-resources/national-forest-inventory/what-our-woodlands-and-tree-cover-outside-woodlands-are-like-today-8211-nfi-inventory-reports-and-woodland-map-reports/>

Period	Planting rate (kha annum <sup>-1</sup> )		
	Conifers on all soil types	Conifers on organic soil	Broadleaves
1751-1800	0.42	0.00	1.69
1801-1850	1.10	0.00	0.91
1851-1900	5.67	0.02	0.86
1901-1910	4.81	0.65	8.94
1911-1920	2.71	0.32	11.57
1921-1930	2.89	0.30	12.20
1931-1940	4.54	0.47	12.92
1941-1950	7.83	1.06	15.64
1951-1960	16.82	2.65	16.64
1961-1970	24.11	4.61	18.36
1971-1980	24.33	5.44	13.46
1981-1990	20.38	4.79	15.41
1991	12.41	2.93	11.97
1992	10.83	2.60	13.65
1993	8.58	2.06	16.74
1994	9.21	2.20	17.67
1995	8.23	1.98	15.17
1996	7.94	1.85	14.72
1997	7.39	1.64	16.14
1998	6.83	1.48	15.94
1999	6.63	1.39	16.21
2000	5.36	1.08	17.95
2001	6.01	1.15	14.41
2002	5.67	1.06	13.80
2003	5.08	0.93	12.43
2004	7.81	1.20	12.10
2005	6.46	0.95	10.91
2006	7.16	1.00	11.73
2007	6.42	0.87	9.44
2008	6.13	0.75	7.31
2009	5.99	0.66	6.55
2010	7.15	0.74	8.82
2011	9.66	0.93	11.98
2012	6.14	0.54	12.66
2013	6.12	0.54	11.66
2014	6.48	0.58	8.63
2015	2.17	0.23	4.73
2016	3.23	0.34	3.16
2017	4.85	0.52	3.65

The proportion of forest planting on mineral and organic soils was re-assessed in 2012, as part of the work to estimate N<sub>2</sub>O emissions due to drainage on forest soils (Yamulki et al. 2012).

**A 3.4.1.4 Allocation of CARBINE outputs to UNFCCC inventory sub-categories**

The CARBINE model output was post-processed using the IPCC default 20-year transition period for Land converted to Forest to move into the Forest remaining Forest category. The area within the Land converted to Forest Land sub-category is split between cropland, pasture grassland, semi-natural grassland, settlement and other areas. Pasture grassland and semi-natural grassland are combined for Grassland reporting in the CRF. This split is based on the relative proportions of historical land use change from these categories to forest. The proportions for each country change over time because the 20-year transition period has a different start date for each reporting year.

The area and carbon stock changes in the Forest remaining Forest category are adjusted to take account of losses of forest converted to other land use categories (deforestation), as these losses, in their entirety, are not reflected in the statistics published by the Forestry Commission. Implied carbon stock changes per unit area are calculated using the unadjusted forest area and carbon stock changes. The forest area is then adjusted to reflect losses due to forest conversion and multiplied by the implied carbon stock change to obtain the adjusted carbon stock change.

The CARBINE model has not yet been implemented for forest in the Isle of Man and Guernsey (Crown Dependencies of the UK) and instead the C-Flow model is used as it was in previous submissions (UK Greenhouse Gas Inventory, 1990-2011, Annex 3.6). Transition to the use of CARBINE for the Overseas Territories and Crown Dependencies has been delayed to allow for further development and documentation of CARBINE.

**A 3.4.1.5 Nitrogen fertilization of forest land**

Nitrogen fertilization of forest land is assumed to occur only when absolutely necessary, i.e. new planting on 'poor' soils (mining spoil, impoverished brown field sites, or upland organic soils). In terms of the inventory, this means that N fertilisation is assumed for Settlement converted to Forest land and Grassland converted to Forest Land on organic soils. The areas of new planting with these conditions were taken from the same dataset used in the CARBINE model (see **Table A 3.4.2**) for 4.A.2. Land converted to Forest land.

Where fertilisation occurs, an application rate of 150 kg N ha<sup>-1</sup> is assumed based on Forestry Commission fertilisation guidelines (Taylor 1991). The guidelines recommend applying fertiliser on a three-year cycle until canopy closure (at approximately 10 years), but this is thought to be rather high (Skiba 2007) and unlikely to occur in reality, so two applications are adopted as a compromise. These applications occur in year 1 and year 4 after planting. The N<sub>2</sub>O emission factor for applied nitrogen fertiliser is the default value of 1% used in the IPCC 2006 Guidelines. Emissions of N<sub>2</sub>O from N fertilisation of forests are estimated using a Tier 1 methodology and IPCC default emission factors. The emissions have fallen since 1990 due to reduced rates of new forest planting. A GWP of 298 for N<sub>2</sub>O is used.

**A 3.4.1.6 Emissions from drainage on forest soils**

Work on developing this method was undertaken by Forest Research in 2012 (Yamulki *et al.* 2012), using new GIS data on forest planting in England, Wales and Scotland. Comparable data were not available for Northern Ireland. This method was described in the 1990-2012 National Inventory Report.

The calculations use the same data on forest planting on mineral and organic soils as are used by the CARBINE model for the calculation of carbon stock changes. It is assumed that only forests planted since 1920 have been drained. The areas of forest planted on mineral soil, nutrient-rich organic soil, nutrient-poor organic soils, nutrient-rich organo-mineral soils and nutrient-poor

organo-mineral soils are estimated based on the proportion of forest cover on different soil types (Yamulki *et al.* 2012), adjusted by the amount of forest planted since 1920. The area of forest on mineral soil is adjusted further by splitting it between free-draining mineral soils and imperfectly draining (easily waterlogged) mineral soils, which require artificial drainage (based on the current guidance and policy for forest operations and management). The proportion of mineral soils requiring artificial drainage is: 34% in England, 24% in Scotland, 3% in Wales, 68% in Northern Ireland and 26% in the UK as a whole. We assumed all forest on organic and organo-mineral soils is cultivated prior to planting and therefore effectively drained.

N<sub>2</sub>O emissions are estimated using the Tier 1 methodology and the IPCC default emission factors for drained mineral (0.06 kg N<sub>2</sub>O-N ha<sup>-1</sup> yr<sup>-1</sup>), nutrient-rich organic (0.6 kg N<sub>2</sub>O-N ha<sup>-1</sup> yr<sup>-1</sup>) and nutrient-poor organic soils (0.1 kg N<sub>2</sub>O-N ha<sup>-1</sup> yr<sup>-1</sup>) (IPCC, 2006).

### A 3.4.2 Land Use Change and Soils (4B, 4C, 4E)

Changes in soil carbon due to land use change are modelled with a dynamic model of carbon stock change which is driven by matrices of change calculated from land surveys.

#### A 3.4.2.1 Land Use Change Matrices

For Great Britain (England, Scotland and Wales), matrices from the Monitoring Landscape Change (MLC) data from 1947 & 1980 (MLC 1986) and the Countryside Surveys (CS) of 1984, 1990, 1998 (Haines-Young *et al.* 2000) and 2007 (Smart *et al.* 2009) are used.

In Northern Ireland, matrices were calculated from the Northern Ireland Countryside Surveys of 1990, 1998 (Cooper and McCann 2002) and 2007 (Cooper, McCann and Rogers 2009). The only data available for Northern Ireland pre-1990 are land use areas from The Agricultural Census and The Forest Service (Cruickshank and Tomlinson 2000). Matrices of land use change were estimated for 1970-79 and 1980-89 using area data. The relationship between the matrix of land use transitions and initial area from recent Countryside Surveys is assumed to be the same as the relationship between the matrix and area data for each of the earlier periods – 1970-79 and 1980-89. The matrices developed in this approach were used to extrapolate areas of land use transition back to 1950 to match the start year in the rest of the UK.

The AFOLU Guidance (IPCC 2006) recommends use of six types of land for descriptive purposes: Forest, Grassland, Cropland, Settlements, Wetlands and Other Land. Areas undergoing active commercial peat extraction and areas of inland water and flooded land are reported under Wetlands in the current inventory, and the remaining land in the UK has been placed into the five other types. The more detailed habitats for the two surveys in Great Britain were combined as shown in **Table A 3.4.3** for the Monitoring Landscape Change dataset and **Table A 3.4.4** for the Countryside Survey Broad Habitats (Jackson, 2000).

**Table A 3.4.3 Grouping of MLC land cover types for soil carbon change modelling**

CROPLAND	GRASSLAND	FORESTLAND	SETTLEMENTS (URBAN)	OTHER
Crops	Upland heath	Broadleaved wood	Built up	Bare rock
Market garden	Upland smooth grass	Conifer wood	Urban open	Sand/shingle
	Upland coarse grass	Mixed wood	Transport	Inland water
	Blanket bog	Orchards	Mineral workings	Coastal water
	Bracken		Derelict	
	Lowland rough grass			
	Lowland heather			
	Gorse			

CROPLAND	GRASSLAND	FORESTLAND	SETTLEMENTS (URBAN)	OTHER
	Neglected grassland			
	Marsh			
	Improved grassland			
	Rough pasture			
	Peat bog			
	Fresh Marsh			
	Salt Marsh			

**Table A 3.4.4 Grouping of Countryside Survey Broad Habitat types (Jackson, 2000) for soil carbon change modelling**

CROPLAND	GRASSLAND	FORESTLAND	SETTLEMENTS (URBAN)	OTHER
Arable and horticulture	Improved grassland	Broadleaved, mixed and yew woodland	Built up areas and gardens	Inland rock
	Neutral grassland	Coniferous woodland	Unsurveyed urban land	Supra littoral rock
	Calcareous grassland		Boundary and linear features	Littoral rock
	Acid grassland			Standing open water and canals
	Bracken			Rivers and streams
	Dwarf shrub heath			Sea
	Fen, marsh, swamp			
	Bogs			
	Montane			
	Supra littoral sediment			
	Littoral sediment			

The area data used between 1947 and the latest inventory year are shown in **Table A 3.4.5** and **Table A 3.4.6**.

The land use change data over the different periods were used to estimate annual changes by assuming that the rates of change were uniform across the period. The full set of annual land use change matrices 1990-latest inventory year are provided in in **Chapter 6** of the main NIR report, **Section 6.1.1** for the UK and **Section 6.9.2** for the Overseas Territories and Crown Dependencies.

**Table A 3.4.5 Sources of land use change data used to estimate changes in soil carbon in Great Britain for different periods.**

Year or Period	Method	Change matrix data
1950-1979	Measured LUC matrix	MLC 1947->MLC1980
1980 - 1984	Interpolated	CS1984->CS1990
1984 - 1989	Measured LUC matrix	CS1984->CS1990
1990 - 1998	Measured LUC matrix	CS1990->CS1998
1999-2007	Measured LUC matrix	CS1998->CS2007

Year or Period	Method	Change matrix data
2008-latest year	<i>Extrapolated</i>	CS1998->CS2007

**Table A 3.4.6 Sources of land use change data used to estimate changes in soil carbon in Northern Ireland for different periods.**


Year or Period	Method	Change matrix data
1950 – 1969	Extrapolation and ratio method	NICS1990->NICS1998
1970 – 1989	Land use areas and ratio method	NICS1990->NICS1998
1990 – 1998	Measured LUC matrix	NICS1990->NICS1998
1999-2007	Measured LUC matrix	NICS1998->NICS2007
2008-latest year	<i>Extrapolated</i>	NICS1998->NICS2007

NICS = Northern Ireland Countryside Survey

The transitions between habitat types in the Countryside Surveys for the latest survey (2007) were calculated with Geographical Information System software (ArcGIS). 544 Countryside Survey squares of Great Britain were identified that coincided between the 1998 and 2007 surveys. Survey square locations are confidential. For each coincident square, the area that changed from one habitat type in 1998 to another in 2007 was calculated. There are 47 broad habitats described by the Countryside Survey. Individual surveyed squares contain a subset of these habitats and changes between habitats are called transitions. Each coincident survey square also has a 'land class' assigned to it that does not change between survey years. There are currently 45 land classes in the Land Classification of Great Britain. Land classes represent the stratification of environments across the UK. A simplified picture of the stratification is shown in **Figure A 3.3**. Northern Ireland is treated as a single uniform unit as its smaller area means that it does not display the climatic or elevation variations evident across the rest of the UK.

Transitions between broad habitats were grouped by land class. The ratio of the total area of each land class to the total area sampled within each land class is calculated so that the transitions can be up-scaled to the land class areas. Transitions can then be extracted at various scales i.e. UK or Devolved Authorities scale or 20 km by 20 km squares. These scales are required by the soil carbon and non-forest biomass models.

**Figure A 3.3 Stratification of environments across the UK with areas 1 to 6 based on the underlying Land Classification (45 classes).**



**A 3.4.2.2 Soils modelling**

A database of soil carbon density for the UK (Milne & Brown 1997, Cruickshank *et al.* 1998, Bradley *et al.* 2005) is used in conjunction with the land use change matrices. There are three soil surveys covering England and Wales, Scotland and Northern Ireland. The field data, soil classifications and laboratory methods for these surveys have been harmonized to reduce uncertainty in the final joint database. The depth of soil considered was also restricted to 1 m at maximum as part of this process. **Table A 3.4.7** shows total stock of soil carbon (1990) for different land types in the four devolved areas of the UK.

**Table A 3.4.7 Soil carbon stock (TgC = MtC) for depths to 1 m in different land types in the UK**

Region Type	England	Scotland	Wales	N. Ireland	UK
Forestland	108	295	45	20	467
Grassland	995	2,349	283	242	3,870
Cropland	583	114	8	33	738
Settlements	54	10	3	1	69

Region Type	England	Scotland	Wales	N. Ireland	UK
Other	0	0	0	0	-
<b>TOTAL</b>	<b>1,740</b>	<b>2,768</b>	<b>340</b>	<b>296</b>	<b>5,144</b>

The dynamic model of carbon stock change requires the change in equilibrium carbon density from the initial to the final land use. The core equation describing changes in soil carbon with time for any land use transition is:

$$C_t = C_f - (C_f - C_0)e^{-kt}$$

where

$C_t$  is carbon density at time  $t$

$C_0$  is the assumed equilibrium carbon density initial land use

$C_f$  is the assumed equilibrium carbon density after change to new land use

$k$  is time constant of change

Differentiating this equation gives the flux  $f_t$  (emission or removal) per unit area:

$$f_t = k(C_f - C_0)e^{-kt}$$

This equation gives, for any inventory year, the land use change effects from any specific year in the past. If  $A_T$  is area in a particular land use transition in year  $T$  considered from 1950 onwards then total carbon lost or gained in an inventory year, e.g. 1990, is given by:

$$F_{1990} = \sum_{T=1950}^{t=1990} kA_T (C_f - C_0)(e^{-k(1990-T)})$$

This equation is used with  $k$ ,  $A_T$  and  $(C_f - C_0)$  chosen by Monte Carlo methods within ranges set by prior knowledge, e.g. literature, soil carbon database, agricultural census, LUC matrices.

In the model, the change in equilibrium carbon density from the initial to the final land use is calculated. These are calculated for each land use category as averages for Scotland, England, Wales and Northern Ireland. These averages are weighted by the area of Land Use Change occurring in four broad soil groups (organic, organo-mineral, mineral, unclassified) in order to account for the actual carbon density where change has occurred. In the UK land use change other than afforestation generally is assumed not to occur on organic soils. Changes in soil carbon stock on afforested land are modelled using the CARBINE model rather than the exponential loss model described above. Other areas of land use change on organic soils are believed to be very small and are currently not separated out from change on mineral soils.

Hence mean soil carbon density change is calculated as:

$$\bar{C}_{ijc} = \frac{\sum_{s=1}^6 (C_{sijc} L_{sijc})}{\sum_{s=1}^6 L_{sijc}}$$

This is the weighted mean, for each country, of change in equilibrium soil carbon when land use changes, where:

$i$  = initial land use (Forestland, Grassland, Cropland, Settlements)

$j$  = new land use (Forestland, Grassland, Cropland, Settlements)

$c$  = country (Scotland, England, N. Ireland & Wales)

$s$  = soil group (organic, organo-mineral, mineral, unclassified)

$C_{sijc}$  is change in equilibrium soil carbon for a specific land use transition

The land use data (1990 to 1998) is used in the weighting. The averages calculated are presented in **Table A 3.4.8 - Table A 3.4.11**.

**Table A 3.4.8 Weighted average change in equilibrium soil carbon density (t ha<sup>-1</sup>) to 1 m deep for changes between different land types in England**

From To	Forestland	Grassland	Cropland	Settlements
Forestland	0	25	32	83
Grassland	-21	0	23	79
Cropland	-31	-23	0	52
Settlements	-87	-76	-54	0

**Table A 3.4.9 Weighted average change in equilibrium soil carbon density (t ha<sup>-1</sup>) to 1 m deep for changes between different land types in Scotland**

From To	Forestland	Grassland	Cropland	Settlements
Forestland	0	47	158	246
Grassland	-52	0	88	189
Cropland	-165	-90	0	96
Settlements	-253	-187	-67	0

**Table A 3.4.10 Weighted average change in equilibrium soil carbon density (t ha<sup>-1</sup>) to 1 m deep for changes between different land types in Wales**

From To	Forestland	Grassland	Cropland	Settlements
Forestland	0	23	57	114
Grassland	-18	0	36	101
Cropland	-53	-38	0	48
Settlements	-110	-95	-73	0

**Table A 3.4.11 Weighted average change in equilibrium soil carbon density (t ha<sup>-1</sup>) to 1 m deep for changes between different land types in Northern Ireland**

From To	Forestland	Grassland	Cropland	Settlements
Forestland	0	94	168	244
Grassland	-94	0	74	150

From To	Forestland	Grassland	Cropland	Settlements
<b>Cropland</b>	-168	-74	0	76
<b>Settlements</b>	-244	-150	-76	0

The rate of loss or gain of carbon is dependent on the type of land use transition (Table A 3.12). For transitions where carbon is lost e.g. transition from Grassland to Cropland, a ‘fast’ rate is applied whilst a transition that gains carbon occurs much more slowly. A literature search for information on measured rates of changes of soil carbon due to land use was carried out and ranges of possible times for completion of different transitions were selected, in combination with modelling and expert judgement (Milne and Brown, 1999; Ashman, *et al*, 2000, Salway *et al*, 2001). These are shown in Table A 3.13.

**Table A 3.4.12 Rates of change of soil carbon for land use change transitions**

		Initial			
		Forestland	Grassland	Cropland	Settlement
Final	Forestland		<i>slow</i>	<i>slow</i>	<i>slow</i>
	Grassland	<i>fast</i>		<i>slow</i>	<i>slow</i>
	Cropland	<i>fast</i>	<i>fast</i>		<i>slow</i>
	Settlement	<i>fast</i>	<i>fast</i>	<i>fast</i>	

(“Fast” & “Slow” refer to 99% of change occurring in times shown in Table A 3.4.13)

**Table A 3.4.13 Range of times for soil carbon to reach 99% of a new value after a change in land use in England (E), Scotland (S) and Wales (W)**

	Low (years)	High (years)
Carbon loss (“fast”) E, S, W	50	150
Carbon gain (“slow”) E, W	100	300
Carbon gain (“slow”) S	300	750

Changes in soil carbon from equilibrium to equilibrium ( $C_t - C_0$ ) were assumed to fall within ranges based on 2005 database values for each transition (Bradley *et al*, 2005) and the uncertainty indicated by this source (up to  $\pm 11\%$  of mean). The areas of land use change for each transition were assumed to fall in a range of uncertainty of  $\pm 30\%$  of the mean.

A Monte Carlo approach is used to vary the rate of change, the area activity data and the values for soil carbon equilibrium (under initial and final land use) for all countries in the UK. The model of change was run 1000 times using parameters selected from within the ranges described above. The mean carbon flux for each region resulting from this imposed random variation is reported as the estimate for the Inventory. An adjustment was made to these calculations for each country to remove increases in soil carbon due to afforestation, as the CARBINE model provides a better estimate of these fluxes in the Land Converted to Forest Land category. Variations from year to year in the reported net emissions reflect the trend in land use change as described by the matrices of change.

**A 3.4.2.2.1 Change in soil carbon stock due to cropland management activities**

Change in soil carbon stocks due to cropland management activities is estimated using the methodology developed in Defra project SP1113 (Moxley *et al*, 2014a) which reviewed UK relevant literature on the effects of cropland management practices on soil carbon stocks and attempted to model UK specific emission factors.

Increases in inputs of fertiliser, manure and crop residues were found to increase soil carbon stocks of tillage land, but changes in the tillage regime from conventional tillage to reduced or zero tillage were found to have no significant effect in a UK context.


Using this methodology, tillage crops are divided into Medium and Low residue groups based on the data on total crop biomass. Where land receives inputs of fertiliser or manure the inputs moved up a class (e.g. cropland producing a Low residue crop which receives manure is considered to receive Medium inputs, while land producing a Medium residue crop which received manure inputs is considered to receive High inputs). If crop residues are removed from land the input level drops. A decision tree for assessing the effect of cropland management on soil carbon stocks is shown in **Figure A 3.4**.

For most cropland management activities there were insufficient UK field data to develop reliable Tier 2 stock change factors, and so Tier 1 factors have been used (for manure and residue inputs, and for differentiating perennial crops, annual crops and set-aside). However, for tillage reduction both a literature review and modelling work suggested that it did not have a significant effect on soil carbon stocks, and that the Tier 1 stock change factors over-estimated its effect under UK conditions. Therefore, a stock change factor of 1 has been used for tillage reduction.

As changes in soil carbon stocks due to changes in cropland management are smaller than changes due to land use change the IPCC default transition time of 20 years is used.

Data on the areas under the main crop types is obtained from the annual June Agricultural Censuses carried out by each UK administration (Defra; Welsh Government; Scottish Government; DAERA). Data on the areas of cropland receiving inputs of manure, fertiliser and crop residues is obtained from the annual British Survey of Fertiliser Practice (Defra).

**Figure A 3.4 Decision tree for assessing the effects of cropland management activities on soil carbon stocks.**



**A 3.4.2.2.2 Change in soil carbon stock due to grassland management activities**

Defra project SP1113 attempted to develop a methodology to allow reporting of changes in soil carbon stocks resulting from grassland management activities. There are reasonable data on the effects of management practices such as liming, reseeding and drainage on improved grassland on mineral soils. However, there are few data on the effect of many management practices if applied to semi-natural grassland or those on organo-mineral or organic soils where there is a

risk that more intensive management could increase carbon losses. As semi-natural grassland makes up a large proportion of grassland in the UK the lack of field data makes it impossible to reliably report changes in soil carbon stocks from grassland management activities. A research project commissioned by BEIS is currently underway to fill this knowledge gap; reporting on the effect of grassland management on soil carbon will be included in the inventory when new data from this project become available.

### **A 3.4.2.3 Future development**

A new vector-based approach to tracking land use change (Levy et al. 2017) has been developed using data from the CORINE land cover map. The approach can be used to produce a set of 100 x 100 m resolution maps, where each 100 m square has an associated vector of land use over time. The maps can be aggregated into a set of distinct representative vectors with their corresponding areas. This has the potential to improve the modelling of change in soil carbon stocks resulting from land use change compared to the current statistically based land use change matrices.


Further implementation of this approach is under discussion, awaiting the outcome of related projects on the potential of earth observation to track LUC in the UK and clarity on the long-term availability of relevant datasets. As there will not be a repeat of the Countryside Survey (last undertaken in 2007), alternative data sources must be used, such as CEH land cover maps (1990,2000, 2007), the NFI map, annual agricultural survey data or the Integrated Administration and Control System (IACS) dataset (used to administer Common Agricultural Policy payments). These all have advantages and disadvantages in terms of their coverage, spatial and temporal resolution, long-term availability and/or processing requirements.

### **A 3.4.3 Changes in stocks of carbon in non-forest biomass due to management and land use change (4B2, 4C2, 4E2)**

#### **A 3.4.3.1 Change in biomass carbon stock due to change in Cropland and Grassland Management.**

Change in Cropland biomass carbon stocks was assessed based on agricultural census data. Areas under different crop types were taken from annual agricultural census data and assigned on one of four categories: annual crops, orchard crops, shrubby perennial crops and set aside and fallow (**Figure A 3.5**). Crop types reported in the agricultural census vary slightly for each administration. **Table A 3.4.14** shows how agricultural census crop types were grouped to assess biomass carbon stocks.

**Figure A 3.5 Crop type area in the UK 1989-2017**



**Table A 3.4.14 Aggregation of Agricultural Census crop types for estimating biomass carbon stock changes from Cropland Management**

Devolved Administration	Annual Crops	Orchard Crops	Shrubby perennial crops	Set Aside and Fallow
<b>England</b>	Cereals, Other arable not stockfeed, Crops for stockfeeding, Vegetables for Human Consumption	Orchard Fruit	Soft fruit, Hardy nursery stock, bulbs and flowers, Area under glass or plastic covered structures.	Uncropped land
<b>Scotland</b>	Cereals, Oilseed rape, Peas for combining, Beans for combining, Linseed, Potatoes, Crops for stockfeeding, Vegetables for human consumption, Other crops	Orchard fruit	Soft fruit	Fallow , Set Aside
<b>Wales</b>	Cereals, Other arable not for stockfeeding, Crops for stockfeeding, Salad and vegetables grown in the open, Total hardy crops	Commercial orchards, Other orchards	Glasshouse	Bare fallow

Devolved Administration	Annual Crops	Orchard Crops	Shrubby perennial crops	Set Aside and Fallow
Northern Ireland	Cereals, Other arable not for stockfeeding, Vegetables	Fruit	Ornamentals	Fallow and set aside

The areas under each aggregated crop type were multiplied by the biomass carbon stock of each crop type using the biomass carbon stock factors in **Table A 3.4.15**. These factors were generated from a literature review. (Moxley *et al.* 2014b).

**Table A 3.4.15 Biomass stock factors for UK Cropland types**

Crop Type	Total biomass Carbon Stock t C/ha	Uncertainty t C/ha (95% CI)	Root: Shoot ratio
Annual	5	1.2	Assume no Below Ground Biomass.
Orchards	10	6.75	0.24
Shrubby perennial crops	3.7	2.0	Assume no Below Ground Biomass.
Set Aside and Fallow	5	1	4.0

Biomass carbon stock change was assumed to occur in the year in which the change in crop type was reported. Cropland biomass stock changes resulting from land use change to or from Cropland were subtracted from the changes due to change in cropland management, as they are accounted for under land use change.

Change in Grassland biomass carbon stocks was assessed based on Countryside Survey data. Grassland was separated into shrubby, non-shrubby and unvegetated Grassland based on Countryside Survey Broad Habitat types. **Table A 3.4.16** shows which Broad Habitats were allocated to which Grassland type.

**Table A 3.4.16 Aggregation of Countryside Survey Broad Habitats for estimating biomass carbon stock changes from Grassland Management**

Shrubby Grassland	Non-shrubby Grassland	Unvegetated Grassland
Dwarf Shrub Heath Bracken Montane	Improved Pasture Improved Pasture Neutral Grassland Calcerous Grassland Acid Grassland Bogs	Littoral sediment Supra littoral sediment

The areas under each aggregated Grassland type were multiplied by the biomass carbon stock of each crop types using the biomass carbon stock factors in **Table A 3.4.17**. These factors were generated from literature reviews (Moxley *et al.* 2014b). Only biomass carbon stock changes resulting from change between shrubby and non-shrubby Grassland were considered, as changes to and from unvegetated littoral and supra-littoral sediments were considered unlikely.

**Table A 3.4.17 Biomass stock factors for UK Grassland types**

Crop Type	Total biomass Carbon Stock t C/ha	Uncertainty t C/ha (95% CI)	Root: Shoot ratio
Non-shrubby Grassland	2.8	1.5	4.0
Shrubby Grassland.	10	3.6	0.53
Unvegetated Grassland	0	0	0
Managed hedge	34.86	68.75	0.3
Unmanaged hedge	175.3	476.6	0.3

Countryside Survey data are only collected on an approximately decadal basis. The annual stock change between survey years was estimated using linear interpolation. Biomass carbon stock change was assumed to occur in the year in which the change in Grassland type occurred. Grassland biomass stock changes resulting from land use change to or from Grassland were subtracted from the changes due to change in grassland management.

Change in Grassland biomass carbon stocks due to change in hedge length are included in the estimate of change in Grassland biomass carbon stock using Countryside Survey data on hedge length and condition. Hedges were divided into managed hedges which are trimmed to prevent the growth of large trees and unmanaged hedges which do not received routine maintenance. Unmanaged hedges do not fall within the UK’s definition of Forest, but may contain isolated trees and may also have some gaps in them. The biomass carbon stocks of managed and unmanaged hedges are estimated as the median of UK-relevant values in published literature, based on a literature review commissioned by BEIS (Moxley *et al.* 2014b) supplemented with more recent data. Full details of these values and data sources are included the Grassland Management Biomass calculation workbook.

**A 3.4.3.2 Change in biomass carbon stock due to land use change.**

Changes in stocks of carbon in biomass due to land use change are based on the same area matrices used for estimating changes in carbon stocks in soils (see previous section). The average biomass carbon density for Cropland, Grassland and Settlement are shown in **Table A 3.4.18**: these were derived from the distribution and biomass densities of the different crop and grassland types in each country of the UK. For Settlements the biomass stocks from Milne and Brown (1997) and land cover data from the 2007 Land Cover Map was used to assess the proportion of gardens, pasture-type grass (including sports pitches, golf courses and parks) and urban (built over) area within areas identified as Settlements.

The average change in biomass carbon density for each country is shown in **Table A 3.4.19 - Table A 3.4.22**. Changes between these equilibrium biomass carbon densities were assumed to happen in a single year.

**Table A 3.4.18 Mean biomass carbon stock densities, tC/ha**

Mean C stock tC/ha	Cropland	Grassland	Settlement
England	5.02	3.37	2.77
Scotland	5.00	4.16	2.91
Wales	5.03	3.61	2.81
Northern Ireland	5.14	2.93	2.64

**Table A 3.4.19 Weighted average change in equilibrium biomass carbon density (kg m<sup>2</sup>) for changes between different land types in England**

From To	Forestland	Grassland	Cropland	Settlements
Forestland				
Grassland		0	-1.66	0.60
Cropland		1.66	0	2.25
Settlements		-0.60	-2.25	0

(Transitions to and from Forestland are considered elsewhere)

**Table A 3.4.20 Weighted average change in equilibrium biomass carbon density (kg m<sup>2</sup>) for changes between different land types in Scotland.**

From To	Forestland	Grassland	Cropland	Settlements
Forestland				
Grassland		0	-0.83	1.25
Cropland		0.83	0	2.08
Settlements		-1.25	-2.08	0

(Transitions to and from Forestland are considered elsewhere)

**Table A 3.4.21 Weighted average change in equilibrium biomass carbon density(kg m<sup>2</sup>) for changes between different land types in Wales.**

From To	Forestland	Grassland	Cropland	Settlements
Forestland				
Grassland		0	-1.42	0.80
Cropland		1.42	0	2.22
Settlements		-0.80	-2.22	0

(Transitions to and from Forestland are considered elsewhere)

**Table A 3.4.22 Weighted average change in equilibrium biomass carbon density (kg m<sup>-2</sup>) for changes between different land types in Northern Ireland.**

From To	Forestland	Grassland	Cropland	Settlements
Forestland				
Grassland		0	-2.21	0.29
Cropland		2.21	0	2.49
Settlements		-0.29	-2.49	0

(Transitions to and from Forestland are considered elsewhere)

Living biomass carbon stocks and Dead Organic Matter (DOM) stocks on Forest Land are modelled using CARBINE and used to calculate changes in carbon stocks due to conversions to and from Forest Land. When land is deforested to another land use, it is assumed that all living biomass and DOM is either converted to Harvested Wood Products or burnt on site in the year in which deforestation takes place. Under KP-LULUCF reporting all HWP from Article 3.3 Deforestation are assumed to be instantaneously oxidised. Increase in biomass carbon and DOM stocks on afforested land is modelled in CARBINE. Full details of CARBINE modelling of carbon stocks on Forest Land are given in **Annex A 3.4.1.1**.

#### **A 3.4.3.3 Future development**

A new vector-based approach to tracking land use change (Levy et al. 2017) as described in **Section A 3.4.2.3**.

#### **A 3.4.4 Carbon stock changes and biomass burning emissions due to Deforestation (4B, 4C, 4E, 4G)**

Deforestation is an activity that cuts across LULUCF categories, affecting net emissions and removals in all the land use categories except 4D Wetlands. The process of land use change affects carbon stock changes in biomass and soil, and the woody material left after felling either moves into the harvested wood products pool or is assumed to be burnt on-site, resulting in immediate biomass burning emissions.

Levy and Milne (2004) discuss methods for estimating deforestation since 1990 using a number of data sources. Their approach of combining Forestry Commission felling licence data for rural areas with Ordnance Survey data for non-rural areas was expanded to include new sources of information and to improve coverage of all countries in the UK. Deforestation before 1990 (which contributes to soil carbon stock change from historical land use change) is estimated from the land use change matrices described in **Section A 3.4.2**.

##### **A 3.4.4.1 Types of deforestation activity in the UK**

In Great Britain, some activities that involve tree felling require permission from the Forestry Commission, in the form of a felling licence, or a felling application within the various forestry grant schemes. There is a presumption that the felled areas will be restocked with trees, usually by replanting but sometimes by natural regeneration. However, some licences are granted without

the requirement to restock – so-called unconditional felling licences. A felling licence is required unless special conditions are met<sup>9</sup>.

Felling for urban development, with no requirement to restock, can be allowed under planning permission but only local planning authorities hold documentation for this. Since 2006, remotely sensed data used in the NFI has included this change, but prior to this, the need for collation of data from local authorities makes estimating the national total difficult. However, in England, the Ordnance Survey (the national mapping agency) makes an annual assessment of land use change from the data it collects for map updating and provides this assessment to the Department of Communities and Local Government (DCLG)<sup>10</sup>. DCLG provides an extract of this dataset, listing annual land use change from Forest to developed land uses (1990-2008 in the latest submission). This dataset comes from a continuous rolling survey programme, both on the ground and from aerial photography. The changes reported each year may have actually occurred in any of the preceding 1-5 years. The survey frequency varies among areas, and can be up to 10 years for moorland/mountain areas. Consequently, for pre-2006 deforestation to Settlement a five-year moving average is applied to the data to smooth out the between-year variation appropriately, to give a suitable estimate with annual resolution.

The Countryside Survey land use change matrix (**Section A 3.4.2**) gives estimates of forest conversion to other land use categories for all countries in the UK for 1990-1998 and 1999-2007. There are known issues with Countryside Survey over-estimating the extent of Forest conversion compared with the extent estimated by the Forestry Commission. Therefore, Forest Commission data is used for Forest areas and the areas of other land uses estimated by Countryside Survey are adjusted to account for this. This is due to differences in Forest definitions, amongst other causes.

### **A 3.4.4.2      Compilation of activity datasets**

For 1990-1999 the deforestation activity dataset is compiled from the felling licence and DCLG datasets as far as possible, using Countryside Survey (CS) data to fill gaps in the time series, to estimate deforestation in Northern Ireland (for which no direct data are available) and to estimate the conversion to different land use categories. The DCLG data are used to estimate the area of Forest Land converted to Settlement (4.E.2.1). The unconditional felling licence data are used to estimate the area of Forest Land converted to Cropland (4.B.2.1) and of Forest Land converted to Grassland (4.C.2.1). Only England has any post-1990 forest to cropland conversion: the estimated areas in Scotland, Wales and Northern Ireland are so small that they are thought to be due to survey classification error rather than genuine land use change.

The CS data are used to estimate the relative split of Forest conversion between Grassland, Cropland and Settlements (**Table A 3.4.23**), using other known data (e.g. felling licences) to correct the CS areas where datasets overlap in time. **Table A 3.4.24** shows the Corrected Forest conversion rates. A correction ratio is used to adjust the estimated deforestation areas, as the Countryside Survey is known to over-estimate deforestation as described in the section above. There are no non-CS data for Northern Ireland so the correction ratios for England or Wales are used, depending on availability. The 1990-98 correction ratios are also applied to the pre-1990 CS land use change estimates.

---

<sup>9</sup> <https://www.gov.uk/guidance/tree-felling-licence-when-you-need-to-apply>

<sup>10</sup> <http://www.communities.gov.uk/planningandbuilding/planningbuilding/planningstatistics/landusechange/>

The annual area of forest converted to other land uses is removed from the area of 4A1 Forest Land remaining Forest Land to maintain consistency in the land area matrix.

**Table A 3.4.23 Countryside Survey data for Forest conversion**

Countryside Survey land use change		Annual rate of change, kha/yr				Grassland/Cropland fractional split		
		England	Scotland	Wales	N Ireland	England	Scotland	Wales
1990-1998	Forest to Natural Grassland	5.600	4.418	1.099	0.171	0.61	0.86	0.72
	Forest to Pasture Grassland	3.081	0.608	0.418	0.086	0.33	0.14	0.28
	Forest to Cropland	0.545	0.097	0.019	0.008	0.06	0.00	0.00
	Forest to Settlements	1.242	0.293	0.132	0.072			
	Forest to Other Land	0.169	0.231	0.058	0.025			
1999-2007	Forest to Natural Grassland	2.656	10.327	0.120	0.209	0.86	0.98	0.42
	Forest to Pasture Grassland	0.277	0.186	0.162	0.102	0.09	0.02	0.58
	Forest to Cropland	0.141	0.006	0.001	0.001	0.05	0.00	0.00
	Forests to Settlements	0.617	0.098	0.095	0.142			
	Forest to Other Land	0.430	0.695	0.374	0.027			

**Table A 3.4.24 Corrected Forest conversion rates**

		Correction ratio			Estimated annual rate of change, kha/yr			
		England	Scotland	Wales	England	Scotland	Wales	N Ireland
1990-1998	Grassland & Cropland	2% <sup>a</sup>			0.159	0.088 <sup>c</sup>	0.026 <sup>c</sup>	0.005 <sup>c</sup>
	Settlements & Other Land	28% <sup>b</sup>			0.390	0.145 <sup>c</sup>	0.052 <sup>c</sup>	0.027 <sup>c</sup>
1999-2007	Grassland & Cropland	20% <sup>a</sup>	2% <sup>a</sup>	15% <sup>a</sup>	0.602	0.262	0.041	0.045 <sup>d</sup>
	Settlements & Other Land	28% <sup>b</sup>			0.296	0.224 <sup>c</sup>	0.133 <sup>c</sup>	0.048 <sup>c</sup>

<sup>a</sup> Unconditional felling licence data used for correction

<sup>b</sup> Land Use Change Statistics used for correction

<sup>c</sup> England correction ratio used

<sup>d</sup> Wales correction ratio used

For 2000 onward, the area and subsequent land-use of deforestation were estimated based on a combination of data sources:

- observations on forest loss by the National Forest Inventory (internal Forestry Commission analysis) by IPCC category. This inventory includes an analysis of deforestation from 2006 to the current inventory year based on a new analysis of woodland maps (Forestry Commission, 2016);
- unconditional felling licences granted (assumed all converted to Grassland);
- analysis of the FC Sub-Compartment Database for restoration of Forest land to open habitats (assumed all converted to Grassland); and

- conversion to non-forest on private sector forest covered by long-term forest plans rather than felling licences (internal Forestry Commission report, assumed all converted to Grassland).

The revision in deforestation was only done from 2000 onwards, partly because there were no suitable data on which to base adjustments for 1990-1999, but also because a number of policy developments came into play in 2000 or shortly beforehand, which affected deforestation to restore open habitats or develop wind-farms. These include the introduction of the UK's climate change policy (2000), and the diversification in relevant forest policies in England, Scotland and Wales following the devolution of forest policy to countries in the late 1990s (Matthews *et al.* 2014). The deforestation information used in this inventory is to the same as the previous inventory, with an assumption of the deforestation rate for 2017.

Soil carbon stock changes associated with deforestation are estimated using the dynamic soil carbon model described in **Section A 3.4.2**. When deforestation occurs it is assumed that 60% of the standing biomass is removed as timber products and the remainder is burnt. In the UNFCCC inventory, KP-LULUCF reporting of deforestation assumes instantaneous oxidation of HWP. Country-specific forest biomass densities for living and dead organic matter from CARBINE are used. These densities change over time in relation to the forest age and species structure. Biomass losses are reported in the relevant carbon stock change tables, assuming a carbon fraction of 0.5 on a dry weight basis. The carbon removed as timber is reported as Harvested Wood Products (HWP) in 4G, using CARBINE to model emission from HWP (described in **Section 6.8**).

Direct and indirect greenhouse gas emissions from associated biomass burning is estimated using the Tier 1 methodology described in the IPCC 2006 guidelines (IPCC 1997 a, b, c) and the emission ratios for CH<sub>4</sub>, CO, N<sub>2</sub>O and NO<sub>x</sub> from Table 3A.1.15 in the IPCC 2003 GPG for LULUCF. Only immediate losses are considered because sites are normally completely cleared, leaving no debris to decay.

### **A 3.4.5 Biomass Burning – Forest and Non-Forest Wildfires (4A, 4B, 4C)**

#### **A 3.4.5.1 Activity dataset**

Until the 2010 submission only wildfires on Forest land were reported due to a lack of activity data for wildfires on other land use categories. Data on Forest wildfires prior to 2010 come from the Forestry Commission and the Forest Service of Northern Ireland.

In 2010 the Fire and Rescue Service (FRS) began recording wildfires in England, Scotland and Wales on a new Incidence and Reporting Systems (IRS) which includes wildfires on all land use categories. The IRS database contains 30 attributes for each fire to which a fire appliance was called, including date, spatial location, property type description (e.g. heathland and moorland, standing crop) and an estimate of the area burnt. This dataset is available from 1<sup>st</sup> April 2009. The original dataset had >126,000 fire records but 99% of these fires were less than 1 ha in size. The IRS database is manually completed by fire service personnel and its use requires some subjective judgement. This is likely to lead to non-systematic differences in the accuracy and precision of the data. The accuracy of the locations is variable, but an assessment of a number of the larger fires suggests that the land cover type attribute is reliable. The accuracy of the FRS burnt area estimates could not be validated using aerial photography as the available imagery was not recent enough, so Landsat images were used to validate the FRS data. However, it was still difficult to find cloud-free, pre- and post-fire images for fires in 2010. In addition Landsat has been affected by image 'striping' since 2003, which affects the quality of the images and causes

some data loss. There are issues with re-ignited fires or additional fires in the same area being logged in the database as separate events. Overall, the uncertainty associated with this dataset is high but should be re-assessed once a longer time series is available.

To provide data on non-Forest wildfires prior to 2010, thermal anomaly data for 2010 from the NASA-operated MODerate Resolution Imaging Spectroradiometer (MODIS) were obtained from the Fire Information Resource Management System (FIRMS) and allocated to land uses using the proportions of fire on each land use type from the Fire and Rescue Service IRS data. The correlation between MODIS data and IRS data breaks down below 25 ha, so for consistency a 25 ha threshold was set for reporting wildfires logged on the IRS.

Thermal anomalies usually represent active fires, but may also detect industrial heat sources, although these are typically masked out by the thermal anomaly processing chain. The IRS dataset records 89 fires > 25ha occurring in 2010. The FIRMS dataset records 335 fire detections for the same period, however, the FIRMS detections may contain multiple detections for a single fire event and the FIRMS detections are for a single 1 km pixel, and do not have a straightforward conversion to burnt area. Searching the IRS and FIRMS data sets for temporally and spatially coincident events, using a 2 km buffer around the IRS data, suggests that 22 fires were recorded by both the IRS and FIRMS systems. There are wide discrepancies between the two datasets, reflecting their different natures. The IRS data set records fires where a fire service response was required, so does not record controlled burning, unless the fire gets out of control. The FIRMS dataset however, responds to anomalous heat signatures, so records controlled and uncontrolled fires. However, in the UK controlled burning, which is primarily carried out for heath management, is only permitted between October and mid-April to reduce the risk of these burns running out of control (Natural England, 2014<sup>11</sup>; Scottish Government, 2011<sup>12</sup>). As the FIRMS thermal anomaly data is only collected between March and August it will not detect most fires from controlled burning. FIRMS is only able to detect fires under cloud-free or light cloud conditions and is also only able to detect fires alight at the time of the satellite overpass. The FIRMS data are more likely to detect larger fires than smaller ones, probably due to the stronger heat signature and the longer burn time that larger fires tend to exhibit.

The IRS and FIRMS thermal anomalies give a very different perspective on the extent, timing and duration of fire events in the UK. However, the datasets did show correlation ( $R^2 = 70-81\%$ ) for fires larger than 25 ha, which enabled an empirical relationship to be derived to extend the burnt area record back to 2001. A burnt area threshold of 25 hectares was used to extract a subset of the IRS database: this captured 75% of the IRS wildfire-burnt area in England, 86% in Scotland and 64% in Wales.

As more IRS data become available confidence should increase in the relationship between fires detected by FIRMS and fires logged in the IRS. This may allow FIRMS data to be extrapolated to fires covering less than 25 ha the inventory in future. However to extend this to small fires there would need to be reasonable confidence that the ratio of large to small fires used was valid, and also some investigation of whether the distribution of small fires across land use classes was the same as that of larger fires.

It was assumed that all fires in the IRS database were wildfires: even if they started as controlled burning, because the need for a fire appliance call-out indicates that they are no longer under

---

<sup>11</sup> <https://www.gov.uk/guidance/heather-and-grass-burning-apply-for-a-licence>

<sup>12</sup> <http://www.gov.scot/Resource/Doc/355582/0120117.pdf>

control. The IRS property type descriptions were assigned to LULUCF sub-categories (**Table A 3.4.25**). There is a very small area of wildfires that occur on Settlement types, and these are included in the Grassland category as the IRS land type classification suggests that they occur on grassy areas within Settlements and there is not a separate reporting field for wildfires in Settlements in the CRF.

**Table A 3.4.25 IRS database property type descriptions by LULUCF sub-category**


LULUCF sub-category	Forest	Cropland	Grassland	Settlement
IRS property type description	Woodland/forest - conifers/softwood	Straw/stubble burning	Heathland or moorland	Domestic garden (vegetation fire)
	Woodland/forest - broadleaf/hardwood	Stacked/baled crop	Grassland, pasture, grazing etc.	Park
		Nurseries, market garden	Scrub land	Roadside vegetation
		Standing crop	Tree scrub	Railway trackside vegetation
				Wasteland
				Canal/riverbank vegetation

A time series of wildfire-burnt areas for each non-forest land use type was constructed for 1990-2015 (**Figure A 3.6**). For non-forest wildfires for England, Scotland and Wales the IRS burnt areas were used for 2010 to the current inventory year and the burnt area estimated from thermal anomalies from 2000 to 2010. For 1990-2000 the average annual burnt area 2001-2011 was used.

In Northern Ireland, where no IRS data were available, it was assumed that the heathland and grassland burning rates were in the same proportions as the Scottish burning rates, using the area of heathland and grassland from the 2007 Northern Ireland Countryside Survey.


Estimates of the forest area burnt in wildfires 1990-2004 are published in different locations (FAO/ECE 2002; FAO 2005) but all originate from either the Forestry Commission (Great Britain) or the Forest Service (Northern Ireland). There is a gap in the time series 2005-2010 for Great Britain but areas of forest wildfires are reported annually for Northern Ireland. The gap was filled using the annual average areas burnt 1995-2005. These areas refer only to fire damage in state forests; no information is collected on fire damage in privately owned forests. The proportion of private-owned forest that was burnt each year was assumed to be the same as the percentage of the state forest that was burnt each year.

**Figure A 3.6** Annual area of FIRMS thermal anomalies for GB for 2001 to the current inventory year (thermal anomalies were filtered to exclude those recorded over urban/industrial areas).



**Figure A 3.7** shows the temporal pattern of FIRMS thermal anomalies, with peaks in hot dry years such as 2003. The FIRMS data used only includes thermal anomalies for March – August for each year, as these are the months where the IRS database recorded fires greater than 25 ha. Some FIRMS thermal anomalies were recorded outside these months due to FIRMS detecting both controlled burns and some fires less than 25 ha in size which are not included in the IRS data.

**Figure A 3.7** Time series of wildfire burnt areas in the UK 1990 to the current inventory year



**A 3.4.5.2 Estimation of emissions**

The IPCC Tier 1 method is used for estimating emissions of CO<sub>2</sub> and non-CO<sub>2</sub> gases from wildfires (IPCC 2006). The *Calluna* heath fuel biomass consumption factor and grassland emission factors are used for heathland and moorland fires, the agricultural residues EFs for cropland and the “savannah and grassland” EFs for other grassland and settlements.

Country-specific biomass and Dead Organic Matter densities from the CARBINE model are used for estimating fuel consumption in forest fires (as discussed in the deforestation methodology section) and the ‘extra tropical forest’ EFs in the 2006 Guidelines. In line with the default value in the IPCC 2006 Guidelines for AFOLU it is assumed that 45% of the biomass is consumed in a wildfire in an unfelled temperate forest.

Emissions from all wildfires are reported under the ‘Land remaining Land’ categories (i.e. 4A1, 4B1 and 4C1) and IE reporting under 4A2, 4B2 and 4C2.

**A 3.4.6 Emissions from drainage (organic soils) (4B1, 4C1)**

Some Wetlands in the UK were drained many years ago for agricultural purposes and continue to emit carbon from the soil. The inventory includes emissions from areas of drained organic soils under Cropland and improved Grassland throughout the UK. The drained areas were generated from work on the UK Agricultural Greenhouse Gas Platform project (Defra project AC0114) (Anthony, ADAS pers. comm). A lack of data on drainage of semi-natural Grassland meant that the area of semi-natural Grassland on drained organic soils could not be estimated. These areas of drained organic soils have also been used for the Agricultural Sector, so there is consistency within the UK Greenhouse Gas Inventory. Work to implement the Wetlands Supplement (IPCC 2013a) guidance is developing a methodology to estimate the area of semi-natural Grassland on drained organic soils. Emissions have been estimated using Tier 1 emissions factors for drained organic soils under Cropland and Grassland taken from the AFOLU Guidelines (IPCC, 2006) applied to all depths of drained organic soil. Results from a BEIS-funded project to implement the Wetland Supplement guidance will allow more detailed estimates of emissions from drained organic soils to be developed in future inventories.

It is assumed that the area of drained organic soils has remained constant as no drainage or rewetting has occurred since 1990 as there have been no policy drivers to encourage drainage or rewetting of cropland or improved grassland. It is also assumed that land on organic soils has not been converted between land uses.

**A 3.4.7 Emissions of N<sub>2</sub>O due to disturbance associated with land use conversion and land management changes (4(III))**

Methodological coverage of this activity has expanded in the IPCC 2006 Guidelines, as previously only N<sub>2</sub>O emissions due to soil disturbance associated with land use conversion to Cropland were included. All land use conversions or land management changes that result in a loss of soil carbon, leading to N mineralization and N<sub>2</sub>O emissions are now reported. Direct emissions from soils, and indirect emissions from nitrogen leached from soil and subsequently oxidised are included in the inventory. The UK now includes emissions resulting from the land use conversions: 4A2 Land converted to Forest Land, 4B2 Forest to Cropland and Grassland to Cropland, 4C1 Grassland remaining Grassland, 4C2 Forest to Grassland, 4E1 Settlement remaining Settlement and 4E2 Land converted to Settlement. Emissions of N<sub>2</sub>O from 4C1 and 4E1 arise from land use change over 20 years before the inventory reporting year where loss of soil organic matter is still ongoing. Emissions of N<sub>2</sub>O from 4B1 Cropland remaining Cropland (resulting from land use

change over 20 years before the inventory year) are calculated in the same way by the LULUCF inventory team but are included in the Agriculture sector (category 3D1).

The Tier 1 methodology described in the IPCC 2006 Guidelines is used. The activity data are the areas and soil carbon stock changes reported in the relevant categories in 4B, 4C and 4E. Some C:N ratios for UK soil/vegetation combinations are published in the Countryside Survey (with values of 11.7 to 13.4) but only for the top 15 cm of soil. However, the soil carbon stock changes reported in the inventory are from the top 1 m of soil, so these C:N ratios were not felt to be applicable. Therefore, the IPCC default C:N ratio of 15 is used for estimating mineralised N. The emission factor of 1% in the 2006 Guidelines was used to estimate N<sub>2</sub>O emissions from mineralised N. Indirect N<sub>2</sub>O emissions from mineralisation are also estimated from carbon stock change using Tier 1 methodology.

### **A 3.4.8 On-site and off-site emissions from peat extraction (4D)**

On-site emissions of CO<sub>2</sub> and N<sub>2</sub>O from peat extraction activities (for energy and horticultural use) and off-site emissions of CO<sub>2</sub> from the decomposition of horticultural peat are reported in category 4D.

#### **A 3.4.8.1 Activity datasets**

Available data sets on peat extraction vary between Northern Ireland and for Great Britain (England, Scotland and Wales). From 2002 onwards Google Earth imagery has been used to estimate the area of peat extraction and sites list in the Directory of Mines and Quarries and the BritPits online database<sup>13</sup>. Prior to the 2002 no Google Earth images are available, and peat extraction site areas have been estimated from other sources. **Table A 3.4.26** shows the sources of activity data used to estimate emissions from peat extraction.

**Table A 3.4.26 Activity data for peat extraction sites in Northern Ireland**

<b>Data set</b>	<b>Information contained</b>	<b>Geographic extent</b>	<b>Time period</b>	<b>Publication frequency</b>
Directory of Mines and Quarries (DMQ)/BritPits database	Location of active peat extraction sites	England, Scotland, Wales, Northern Ireland	1984 - present	Online database is continuously updated
Google Earth	Land use images to estimate area of extraction sites identified from DMQ	UK	2002 - date	Variable
Cruickshank and Tomlinson (1997)	Area with planning consent for peat extraction Local authority planning consents for peat extraction sites	England, Scotland, Wales	1990/91	One off compilation of data

<sup>13</sup> <https://www.bgs.ac.uk/products/minerals/britpits.html>

Data set	Information contained	Geographic extent	Time period	Publication frequency
Tomlinson (2010)	Estimate of peat extraction area in Northern Ireland.  Volume of peat extracted (sod cutting and vacuum harvesting)	Northern Ireland	1990 - 1991	One off compilation of data
Mineral Extraction in Great Britain (Annual Minerals Raised Inquiry)	Volume of peat extracted	England, Scotland, Wales	1947 - date	Annual
Cruickshank et al. 1995	Volume of peat extracted (hand cutting)	Northern Ireland	1990 - 1991	One off compilation of data

The areas of peat extractions sites listed in the BritPits database were assessed using Google Earth. Polygons were drawn around site boundaries and the area covered by the polygons was calculated in Google Earth. Change over time at individual sites was tracked to give an estimate of the extent of conversion to and from extraction sites. This method is repeated annually to incorporate changes in extraction site areas in new Google Earth images.

Any sites abandoned since 2002 where a change of land use cannot be identified are still judged to be producing on-site emissions, in line with good practice guidance. A time series was constructed using linear interpolation. The extraction area (active and abandoned where there has been no change in land use) declined between 1990 and 2015 by 40% in England, 43% on fuel peat sites in Scotland, 6% on horticultural peat sites in Scotland, 99% on fuel peat sites in Northern Ireland and 40 % on horticultural peat sites in Northern Ireland. This area was assumed to be converted to Grassland.

The 2017 version of the BritPits database lists four sites in Wales, most having ceased extraction. Only one of these is visible on Google Earth and the Mineral Extraction in Great Britain report for 2013 does not report any peat production in Wales, so it is assumed that the visible site is currently inactive but has not been converted to another land use.

Annual peat production in Great Britain (**Table A 3.4.27**) is inferred from extractor sales by volume as published in the annual Mineral Extraction in Great Britain report, formerly known as the Minerals Raised Inquiry (ONS). This gives a breakdown for horticultural and other uses of peat, which are assumed to be fuel, for English regions and for Scotland. No peat extraction is reported in Wales. Annual production is highly variable because extraction methods depend on suitable summer weather for drying peat.

**Table A 3.4.27 Annual peat production, m<sup>3</sup> for England and Scotland (from Annual Minerals Raised Inquiry/Mineral Extraction in Great Britain reports)**

Year	England		Scotland	
	Horticultural	Fuel	Horticultural	Fuel
1990	1,116,940	2,727	293,170	93,163
1991	1,202,000	2,000	241,000	115,000
1992	1,079,000	4,000	332,000	91,000
1993	1,069,820	2,180	306,511	73,489
1994	1,375,000	1,000	498,000	108,000
1995	1,578,000	2,000	657,000	44,000
1996	1,313,000	2,000	517,000	53,000
1997	1,227,000	2,000	332,000	59,000
1998	936,000	0	107,000	32,000
1999	1,224,000	0	392,000	37,000
2000	1,258,000	1,000	336,000	31,000
2001	1,459,000	1,000	325,000	30,000
2002	856,000	1,000	107,000	10,000
2003	1,227,000	1,000	741,000	38,000
2004	902,000	1,000	338,000	21,000
2005	927,000	1,000	556,000	21,000
2006	856,000	1,000	712,000	24,000
2007	654,000	0	221,000	10,000
2008	455,000	41,000	243,000	21,000
2009	476,000	0	390,000	21,000
2010	456,000	1,000	527,000	21,000
2011	429,000	0	369,000	26,000
2012	422,000	0	126,000	20,000
2013	661,000	0	570,000	24,000
2014	294,000	0	469,000	32,000
2015*	563,000	0	417,772	22,528
2016*	563,000	0	417,772	22,528
2017*	563,000	0	417,772	22,528

\* 2005 - 2014 average as UK Minerals Yearbook has ceased production.

#### A 3.4.8.2 Estimation of emissions

Default on-site emission factors for Tier 1 reporting (IPCC 2006) are used to estimate emissions. Peat extracted for horticultural use is inferred to be from oligotrophic (nutrient-poor) bogs. Peat for fuel is inferred to be from mineratrophic (nutrient-rich) fens or bogs. On-site emissions of CO<sub>2</sub> and N<sub>2</sub>O from drainage are reported.

A value of 0.0641 tonnes C m<sup>-3</sup> is used for Great Britain to estimate emissions from extracted horticultural peat volumes based on previous work (Thomson *et al*, 2011). This is slightly lower than the IPCC default emission factor of 0.07 tonnes C m<sup>-3</sup> air-dry peat for nutrient-poor peats.

Tomlinson (2010) gives production estimates of horticultural peat production for Northern Ireland for 1990/91 and 2007/2008. These have been interpolated to produce a time series. The total emission from horticultural peat production is the sum of emissions from vacuum harvesting production, sod extraction production and mechanical extraction production.

Emissions from vacuum harvesting production =

area \* annual depth of extraction \* carbon fraction by volume

where

*Annual depth of extraction by vacuum harvesting, m/ha = 0.1*

*Carbon fraction of air-dry peat by volume, tonnes C/m<sup>3</sup> air-dry peat = 0.0641*

Emissions from sod extraction production =

area \* sod extraction rate \* % dry matter for sods \* mean % C

where

*Sod extraction rate, tonnes/ha/yr = 200*

*Sod extraction, mean % dry matter = 35%*

*Mean % carbon = 49%*

Emissions from mechanical extraction production =

area \* extraction rate \* % dry matter for mechanical extraction \* mean % C

where

*The mechanical extraction rate was estimated to be 206.45 tonnes/ha in 1990/91 and 243.06 tonnes/ha in 2007/08 (Tomlinson, 2010).*

*Mechanical extraction, mean % dry matter = 67%*

*Mean % carbon = 49%*

### **A 3.4.9 Flooded Lands (4D)**

Carbon stock changes on land converted to Flooded Land (reservoirs) are included in the inventory, based on the IPCC 2006 Guidance. Data on all reservoirs over 1 km<sup>2</sup> were compiled but only reservoirs established since 1990 were reported (areas of inland water under 1 km<sup>2</sup> are reported under 4D Wetlands remaining Wetlands). Activity data were compiled for England and Wales from the Public Register of Large Raised Reservoirs provided by the Environment Agency, which listed location, surface area and year built. Activity data for Scotland were compiled from the SEPA Water Body Classification database (of water bodies > 0.5 km<sup>2</sup>) and the associated Water Body data sheets. Additional information on the year of building was obtained from:

- the Gazetteer for Scotland <http://www.scottish-places.info>;
- hydro-electric power generators <http://sse.com/whatwedo/ourprojectsandassets/> <http://www.power-technology.com/projects/glendoehydropowerpla/>; and
- local authorities <http://www.argyll-bute.gov.uk>.

It was established through discussion with local experts that no new large reservoirs had been built in Northern Ireland since the 1950s.

Only five large reservoirs have been established in the UK since 1990, three in England and one each in Scotland and Wales (another five in England are sacrificial floodplains and do not fit the criteria of permanent conversion to Flooded Land). These cover a total of 1.995 kha.

The location of each reservoir was examined using the [www.magic.gov.uk](http://www.magic.gov.uk) geographic information portal. All reservoirs were in upland locations and were assumed to be Grassland prior to their conversion to Flooded Land. (Any forest removed as part of the land conversion will have been captured under the deforestation activity methodology). A Tier 1 methodology was followed, so carbon stock changes in living biomass stock in the year of flooding were estimated, but not carbon stock change in soils. A living biomass density of 2 t dry matter/ha was used to estimate carbon stock losses. This will be updated in the next submission to be consistent with the biomass densities used in other parts of the LULUCF inventory.

### **A 3.4.10 Harvested Wood Products (4G)**

The activity data used for calculating this activity are the annual forest planting rates. CARBINE then applies a forest management regime as given in input to the model. For a given forest stand, carbon enters the HWP pool when thinning is undertaken and when harvesting takes place. Depending on the species, first thinning occurs approximately 20 years after planting.

At thinning and harvest, the CARBINE model allocates merchantable stem volume to various wood products, while the remainder is transferred to the harvesting residue pool. The 'end-use' wood products represented are:

- Long-lived sawn timber, e.g. timber used for construction;
- Short-lived sawn timber, e.g. timber used for fencing;
- Particleboard;
- Paper; and
- Fuel.

For reporting purposes the long-lived and short-lived sawn timber are reported together in the Sawnwood category.

During wood extraction, conversion losses are assumed to be left as on-site harvest residue and enter the litter pool. The allocation of carbon to wood product categories is estimated by inputting the merchantable stem carbon (from the forest yield model) to a stand volume assortment forecasting model which estimates the volume allocated to sawn timber, roundwood and waste. This is implemented in CARBINE as a set of functions derived from the output of a more general and flexible assortment forecasting program known as ASORT (Rollinson and Gay, 1983). The CARBINE model uses standard estimates for oven-dried wood density to derive biomass from the harvested volume (Lavers and Moore, 1983; Jenkins et al., 2011). Carbon content of all oven-dried wood is assumed to be 50% (Matthews, 1993). CARBINE assumes the wood of a tree species all has the same oven-dried wood density and carbon content, irrespective of which semi-finished wood product categories it is assigned to.

The proportions of wood produced which are allocated to different product categories are based on proportions derived from FAO data<sup>14</sup> (prior to 1994) and forestry commission data<sup>15</sup> (after 1993) on production of semi-finished wood products. A carbon retention curve is used to estimate product decay and return of carbon to the atmosphere. Each wood product category has its own carbon retention curve using the default half-lives in the IPCC 2013 Revised Supplementary Methods and Good Practice Guidance, taking into account the decay rate of wood products and the service life as influenced by socio-economic factors. The half-lives are: 35 years for sawn wood; 25 years for wood panels; 2 years for paper. Timber used as woodfuel is assumed to instantaneously oxidise.

In implementing the 2006 IPCC guidelines for HWP the UK has elected to report using the production approach B2, which requires disaggregation of HWP into those produced and consumed domestically and those produced and exported. In the annual Forestry Statistics publication there is data on the apparent consumption of wood products in the UK. A consistent dataset is available at the product level (i.e. sawnwood, wood panels and paper & paperboard) for 2002 onwards. The ten year average of 2002-2011 was calculated for each product type and those values were used for the years 1990-2001. This dataset was used to assign the HWP output from the CARBINE model into either consumed domestically or exported.

### **A 3.4.11 Methods for the Overseas Territories (OTs) and Crown Dependencies (CDs)**

The UK LULUCF inventory includes direct GHG emissions from UK Crown Dependencies (CDs) and Overseas Territories (OTs) which have joined, or are likely to join, the UK's instruments of ratification to the UNFCCC and the Kyoto Protocol. Currently, these are: Guernsey, Jersey, the Isle of Man, the Falkland Islands, the Cayman Islands, Bermuda, and Gibraltar. It should be noted that Bermuda will not ratify the 2<sup>nd</sup> Commitment Period of the Kyoto Protocol and is therefore not included in the 'GBK' submission under the Kyoto Protocol.

A web search of statistical publications was undertaken for any updates in datasets in 2017. This work builds on an MSc project to calculate LULUCF net emissions/removals for the OTs and CDs undertaken during 2007 (Ruddock 2007).

The availability of data for the different OTs and CDs is very variable, so that emission estimates can only be made for the Isle of Man, Guernsey, Jersey and the Falkland Islands. These four comprise over 95% of the area in all the OTs and CDs. Gibraltar wished to produce their own inventory and has assessed its LULUCF net emissions/removals as Not Occurring (Annex 3.6) as the country is extremely small (6 km<sup>2</sup>). A lack of suitable data for the Caribbean territories (discussed in the 1990-2006 NIR) makes it impossible to create inventories for them at the present time. Discussions have started with the Cayman Islands government with a view to reporting LULUCF emissions from this OT in the future.

A number of omissions (reported as Not Estimated) for the Overseas Territories and Crown Dependencies were identified in the 2016 UNFCCC Expert Review of the GHG Inventory. These included wildfires on forest land and grassland, flooded land, peat extraction and some carbon pools affected by deforestation. Only very limited further data has been discovered through

---

<sup>14</sup> <http://www.fao.org/faostat/en/#data/FO>

<sup>15</sup> <https://www.forestresearch.gov.uk/tools-and-resources/statistics/statistics-by-topic/timber-statistics/uk-wood-production-and-trade-provisional-figures/>

additional investigations, and so the threshold of insignificance (Decision 24/CP.19, paragraph 37b) has been applied.

- Wildfires on forest land: no information on the occurrence of wildfires on forest land was obtained by further investigation. The average area of forest wildfires in the UK 2008-2017 was 0.632 kha/year, with an average emission of 73.6 Gg CO<sub>2</sub>e (CO<sub>2</sub>, CH<sub>4</sub> and N<sub>2</sub>O). This equates to 0.018% of UK forests burnt every year, and the equivalent area burnt in the Crown Dependencies (the Falkland Islands have no forest cover) would be <0.001 kha/year with estimated emissions falling well below the insignificance threshold of 500 Gg CO<sub>2</sub>e.
- Wildfires on grassland: no information on grassland fires in the Crown Dependencies was obtained by further investigation. There are some reports of wildfires in the Falkland Islands, caused by lightning strikes or pasture burning. The average area of grassland wildfires in the UK 2008-2017 was 10.148 kha/year, with an average emission of 13.5 Gg CO<sub>2</sub>e (CO<sub>2</sub>, CH<sub>4</sub> and N<sub>2</sub>O). This equates to 0.076% of UK grassland burnt every year, and the equivalent area burnt in the Overseas Territories and Crown Dependencies (dominated by the large area of grassland in the Falkland Islands) would be 0.961 kha/year with estimated emissions of 1.28 Gg CO<sub>2</sub>e falling well below the insignificance threshold of 500 Gg CO<sub>2</sub>e. A watching brief will be kept on this potential emission source, particularly in the Falkland Islands.
- Flooded land: An assessment of flooded land was undertaken for the Overseas Territories and Crown Dependencies. No flooded land areas exceed the area threshold of 1 km<sup>2</sup> used for the UK, so the area of Flooded Land remaining Flooded Land has been included with the Inland Water area in the Wetlands remaining Wetlands category (0.095 kha in total).
- Peat extraction: Peat organic soils occur in the Falkland Islands and Isle of Man but not in the other Overseas Territories and Crown Dependencies. Information on the extent and condition of peat soils in these territories has been compiled a part of the research project on implementation of the Wetlands Supplement (see box in Chapter 6, section 6.5.8). Further work is underway to incorporate these results into the GHG Inventory. Further research on the peat soils of the Falkland Islands is also underway and will be used to improve reporting in this category in due course.
- Deforestation: A small area of deforestation (0.002 kha/yr) occurs in Guernsey, obtained from two habitat surveys in 1999 and 2010. The limited data only permits Tier 1 estimation of carbon stock changes in biomass and litter. The change in forest cover is a result of the changed areas losing sufficient tree cover to be reclassified as dense scrub or parkland, rather than conversion to settlement land or agriculture. Any changes in soil and dead wood are very likely to be minimal. As all emissions from deforestation in the Overseas Territories and Crown Dependencies fall well below the insignificance threshold of 500 Gg CO<sub>2</sub>e (estimated to be 0.64 Gg CO<sub>2</sub> in 2017 if a similar rate of deforestation to the UK rate is assumed), the UK will continue to report these carbon stock changes as Not Occurring and will continue to report emissions from this activity that it has reported in previous years.

Information on the area of each IPCC land category, dominant management practices, land use change, soil types and climate types were compiled for each OT/CD from statistics and personal communications from their government departments (**Table A 3.4.28**). This allowed Tier 1 level inventories to be constructed for the four OT/CDs already mentioned, and a Tier 3 approach for Forest Land on the Isle of Man and Guernsey (using the C-Flow model, for information on C-Flow

model please refer to 1990-2011 NIR). The assumptions and factors used for the estimation of emissions are given in **Table A 3.4.29** and **Table A 3.4.30**. The estimates have high uncertainty and may not capture all relevant activities, but given the size of the territories any missing sources are likely to be small.

**Table A 3.4.28 Information sources for estimating LULUCF emissions from the Overseas Territories and Crown Dependencies**

Territory	LULUCF category	Time period	Reference
Isle of Man	4A	1970-2011	Personal communication from Isle of Man Department of Agriculture, Fisheries and Forestry (Peter Williamson) FAO (2010) Global Forest Resources Assessment: Isle of Man
	4B, 4C, 4D	2002-2011	Isle of Man Agricultural and Horticultural Census: completed by all farmland occupiers on an annual basis Isle of Man Digest of Economic and Social Statistics
	4B, 4C	2012 - 2013	Isle of Man Digest of Economic and Social Statistics
	4B, 4C	2014	The Isle of Man in Numbers
	4E	1991-1994	Isle of Man Ecological Habitat Survey, Phase 1 Report (Sayle et al, 1995)
Guernsey	4A	1990-2010	FAO Global Forest Resources Assessment 2010: Guernsey
	4A, 4B, 4C, 4D, 4E	1998/9, 2005, 2010	Guernsey Habitat Survey Sustainable Guernsey 2005, 2009, Guernsey Facts and Figures 2011
Jersey	4A	1990-2010	FAO Global Forest Resources Assessment 2010: Jersey
	4B	1990 - 2014	Jersey Agricultural Statistics
	4A, 4B, 4C, 4D, 4E	2006, 2008, 2012, 2015	Jersey In Figures 2006/2008/2009/2010/2011/2012
Falkland Islands	4A	1990-2011	Department of Mineral Resources, personal communication FAO Global Forest Resources Assessment 2010: Falkland Islands
	4B, 4C	1991-2013	Falkland Islands Agricultural Statistics
	4E	1990-2005	Falkland Islands Environment and Planning Department, personal communication
	4E	1986 – 2001 with projections 2006 - 2016	Stanley Town Plan, Environmental Planning Dept, Falkland Islands Government.

**Table A 3.4.29 Assumptions used in applying the Tier 1 methodology to the Overseas Territories and Crown Dependencies**

Land Use category	Sub-category	Isle of Man	Guernsey	Jersey	Falkland Islands
<b>Forest land fluxes</b>	Living biomass, DOM, Mineral soils, Organic soils	From C-Flow model	From C-Flow model	Assumed in equilibrium	No forest on Falklands
<b>Crop remaining crop</b>	Living biomass	N/A. Only for perennial crops	N/A. Only for perennial crops	N/A. Only for perennial crops	N/A. Only for perennial crops
	Dead organic matter	N/A	Orchards only. 10 tC/ha	N/A	N/A
	Mineral soils	No change in SOC	No change in SOC	No change in SOC	N/A
	Organic soils	N/A	N/A	N/A	Default (-5 tC/ha)
<b>Land converted to Crop</b>	Living biomass	UK shrubby grass to crop values(5 tC/ha)	UK non-shrubby grass to crop value (2.2 tC/ha)	UK non-shrubby grass to crop value (2.2 tC/ha)	UK non-shrubby grassland to crop value (2.2 tC/ha)
	Dead organic matter	N/A	N/A	N/A	N/A
	Mineral soils	Default SOC = 95 tC/ha, assume conversion from natural grassland (-1.7347 tC/ha)	Default SOC = 95 tC/ha, assume conversion from natural grassland (-0.95 tC/ha)	Default . SOC = 95 tC/ha, assume conversion from natural grassland (-0.95 tC/ha)	N/A
	Organic soils	N/A	N/A	N/A	Default (-5 tC/ha)
	N <sub>2</sub> O emissions	Default (0.001817 t N <sub>2</sub> O/ha)	Default (0.000995 t N <sub>2</sub> O/ha)	Default (0.000995 t N <sub>2</sub> O/ha)	Default (0.012571 t N <sub>2</sub> O/ha)
<b>Grass remaining grass</b>	Living biomass	N/A	N/A	N/A	N/A
	Dead organic matter	N/A	N/A	N/A	N/A
	Mineral soils	No change in SOC	No change in SOC	No change in SOC	N/A
	Organic soils	N/A	N/A	N/A	Assume no soil C stock change

Land Use category	Sub-category	Isle of Man	Guernsey	Jersey	Falkland Islands
<b>Land converted to grass</b>	Living biomass	UK crop to non-shrubby grass values (-2.2 tC/ha)	UK settlement to non-shrubby grass value (0 tC/ha)	Crop to Grassland: UK crop to non-shrubby grass value (-2.2 tC/ha) Settlement to Grassland: assume increase from 0 in glasshouses (2.8 tC/ha)	Use crop to non-shrubby grassland value (-2.2 tC/ha)
	Dead organic matter	N/A	N/A	N/A	N/A
	Mineral soils	Default SOC = 95 tC/ha, assume conversion from cropland (1.7347 tC/ha)	Default SOC = 95 tC/ha, assume conversion from settlement, assume same soil C as for cropland (0.95 tC/ha)	Default SOC = 95 tC/ha, Cropland to Grassland: assume conversion from cropland (0.95 tC/ha) Settlement to Grassland: assume no change	N/A
	Organic soils	N/A	N/A	N/A	Default (-0.25 tC/ha)
<b>Settlements remaining Settlements</b>	Living biomass, DOM, Mineral soils, Organic soils	N/A	N/A	N/A	N/A

Land Use category	Sub-category	Isle of Man	Guernsey	Jersey	Falkland Islands
<b>Land converted to Settlements</b>	Living biomass	UK values, shrubby grass to settlement (-7.2 tC/ha)	UK non-shrubby grass to settlement value (0 tC/ha)	Grassland to Settlement, UK non-shrubby grass to settlement value (0 tC/ha) Cropland to Settlement: use cropland to settlement value (-2.2 tC/ha)	Use shrubby grass to settlement value (-7.2 tC/ha)
	Dead organic matter	N/A	N/A	N/A	N/A
	Mineral soils	Default SOC = 95 tC/ha, assume conversion from grassland and all soil C lost (-4.75 tC/ha)	Default SOC = 95 tC/ha, assume 30% of land is paved over and the rest is turf grass (-1.14 tC/ha)	Default . SOC = 95 tC/ha, Grassland to Settlement: assume 30% of land is paved over and the rest is turf grass (-1.14 tC/ha) Cropland to Settlement: assume 30% of land is paved over and the rest is turf grass (0.95 tC/ha)	N/A
	Organic soils	N/A	N/A	N/A	Default - assume cropland (-5 tC/ha)
	N <sub>2</sub> O emissions	Default (0.004976 t N <sub>2</sub> O/ha)	Default (0.00119 t N <sub>2</sub> O/ha)	Default (0.00119 t N <sub>2</sub> O/ha)	N/A
<b>Other land remaining other land</b>	Living biomass, DOM, Mineral soils, Organic soils	N/A	N/A	N/A	N/A

Land Use category	Sub-category	Isle of Man	Guernsey	Jersey	Falkland Islands
<b>Land converted to other land</b>	Living biomass	N/A	Assume loss of grassland to standing water or cliff (-10 tC/ha)	Assumed loss of grassland to standing water (-2.8 t C/ha)	N/A
	Dead organic matter	N/A	N/A	N/A	N/A
	Mineral soils	N/A	Assume no change in soil stocks	N/A	N/A
	Organic soils	N/A	N/A	N/A	N/A
	N <sub>2</sub> O emissions	N/A	0	N/A	N/A
<b>Harvested wood products</b>		From C-Flow model	From C-Flow model	N/A	N/A

**Table A 3.4.30 IPCC Tier 1 factors used for estimating LULUCF emissions from Overseas Territories and Crown Dependencies**

	Factor	Isle of Man/	Guernsey/ Jersey	Falkland Islands
Biomass carbon densities, tC/ha	Cropland	5	5	5
	Grassland (shrubby)	10	10	10
	Grassland (non-shrubby)	2.8	2.8	2.8
	Settlements	2.8	2.8	2.8
	Soil C density	95	95	87
	Grass F <sub>lu</sub>	1	1	1
	Grass F <sub>mg</sub>	1	1	1
	Grass F <sub>i</sub>	1	1	1
	Crop F <sub>lu</sub>	0.69	0.8	0.69
	Crop F <sub>mg</sub>	1	1	1
	Crop F <sub>i</sub>	0.92	1	0.92
	C/N ratio kg N <sub>2</sub> O-N/kg N	15	15	15
	N <sub>2</sub> O EF	0.01	0.01	0.01
	Frac-Leach	0.3	0.3	0.3
		0.0075	0.0075	0.0075
	Cropland Organic soils EF, tC/ha/yr			-5
	Grassland Organic soils EF, tC/ha/yr			-0.25
	EF2 for temperate organic crop and grassland soils, kg N <sub>2</sub> O-N/ha			8
Forest converted to grassland (Guernsey only)	Living biomass, t DM/ha		120	

	Factor	Isle of Man/	Guernsey/ Jersey	Falkland Islands
	Dead organic matter, tC/ha		16	
	Carbon fraction of dry matter		0.47	
	Biomass C density, tc/ha		56.4	
	Change in mineral soil density, tC/ha		0	


### A 3.4.12 Uncertainty analysis of the LULUCF sector

The purpose of carrying out uncertainty analysis within the LULUCF inventory is to quantify where the largest sources of errors lie, and to identify areas to be targeted in future work so as to reduce the uncertainties. In the 1990-2010 inventory report a sensitivity analysis of the whole of the existing inventory methodology was undertaken, applying uncertainty quantification more widely and rigorously to all model parameters and empirical conversion factors, in order to quantify the impact of those uncertainties on the inventory. Although this analysis was carried out for the C-Flow model, which is no longer used, it is broadly applicable to the CARBINE model as both are similar forest carbon accounting models, based on the same underlying yield tables and input data. An update to this uncertainty analysis, which will include the CARBINE model, will be undertaken in 2019 and is likely to be reported in the 2020 submission.

The results of the simulations, including both input and parameter uncertainty, are that the area undergoing land use change is the single biggest uncertainty in the inventory, followed by uncertainty in the forest model parameters and the choice of model for the change in soil carbon following land use change (**Figure A 3.8**). The next five terms are all of a similar magnitude. Full details of the methodology and results are in the 1990-2010 inventory report.

The uncertainty in the land use change areas is being addressed by the development of a new vector-based approach (see **Chapter 7, Section 1.1**), combining multiple sources of land use data.

**Figure A 3.8 The largest uncertainties in the LULUCF inventory, in terms of standard deviation in the output distributions**



Parameterisation of the forest model is the second largest source of uncertainty. This has been addressed with the move to CARBINE, as 19 tree species are now modelled instead of the two used in previous submissions. Results from the National Forest Inventory (NFI) and small woods dataset will also provide additional information on carbon stocks in trees (e.g. Forestry Commission 2015<sup>16</sup>). The choice of soil carbon model and its parameters are also important, because the time course of the flux following land use change may be quite different, depending on the equations used to represent this, and how carbon is distributed between fast- and slow-turnover pools. The choice of forest model is less important, largely because all the UK forest models are based on the same yield table data.

## A 3.5 WASTE (CRF SECTOR 5)

### A 3.5.1 Solid Waste Disposal on Land (5A)

#### A 3.5.1.1 Input data

Because waste sent to landfill is now evaluated using individual waste consignments by EWC code, there is no need to make assumptions regarding waste composition, other than for two waste categories. These EWC codes are 19.12.12 (residues from waste sorting) and 20.03.01 (mixed municipal waste). Wastes with these codes were allocated in accordance with the findings of a survey carried out on behalf of Defra (Resource Futures, 2012), as set out in **Table A 3.5.1**. Data on DOC, DOCf and material compositions are provided in **Table A 3.5.2**.

The model allocates waste to two types of landfill – old, closed sites which last received waste in 1979, and modern engineered landfills that came into operation from 1980. Only these latter sites

<sup>16</sup> This survey is preliminary and the carbon stocks have been estimated using the same relationships and calculation parameters that underlie CARBINE; they are therefore not an independent validation of the LULUCF estimates.

have gas management systems. The old closed sites have no gas control. The distribution of waste between these types of site is the same as used for compiling the previous NIR.

The quantities of waste sent to landfill are shown in **Table A 3.5.3**. The amounts of methane generated, recovered, used for power generation, flared, oxidised and emitted to the atmosphere are shown in **Table A 3.5.4**.

**Table A 3.5.1 Composition of waste sorting residues and mixed municipal waste**

Material	19.12.12 (residues from waste sorting)	20.03.01 (mixed municipal waste)
Paper	10.3%	10.6%
Card	9.1%	7.7%
Plastic film	9.4%	8.4%
Dense plastics	13.2%	9.6%
Sanitary waste	1.3%	3.1%
Wood	10.0%	5.3%
Textiles and shoes	5.9%	5.6%
Glass	1.3%	3.0%
Food waste	8.2%	21.3%
Garden waste	1.8%	3.5%
Other organic	1.3%	2.1%
Metals	3.2%	3.7%
WEEE	1.4%	1.5%
Haz waste and batteries	1.1%	0.9%
Carpet, underlay & furniture	7.0%	5.0%
Other combustibles	2.7%	1.4%
Bricks, plaster and soil	7.9%	4.1%
Other non-combustible	1.7%	1.5%
Fines <10mm	3.3%	1.8%
Total	100%	100%

Table A 3.5.2 DOC, DOCf and composition of waste materials

Component	Lignin biodegradability DOCf	Non-lignin biodegradability DOCf	Moisture, %fresh matter	Lignin, % dm	Hemicellulose, %dm	Cellulose, %dm	Starch, %dm	Sugar, % dm	Fat, %dm	Proteins, %dm	Fibre, %dm	Readily soluble, %dm	Other (inert), % dm
Carbon contents (DOC)			0%	65.1%	44.6%	40.0%	44.4%	42.1%	76.0%	40.0%	45.0%	45.0%	0.0%
<b>Waste composition on a dry matter basis (other than moisture)</b>													
<b>Municipal solid waste</b>													
Paper	5%	65%	15%	15%	9%	61%							15.00%
Card	5%	65%	20%	15%	9%	61%							15.00%
Nappies	5%	65%	65%			47%							52.70%
Textiles and footwear	5%	65%	20%		15%	15%							69.68%
Miscellaneous combustible	5%	65%	20%		25%	25%							50.00%
Wood	5%	65%	17%	26%	12%	42%							21.00%
Food – corrected	15%	70%	70%	6%	4%	27%	13%	7%	14%	15%	14%	0%	0.00%
Garden	10%	65%	55%	20%	16%	20%			2%			26%	17.10%
Soil and other organic	5%	65%	30%		1%	1%							98.60%
Furniture	5%	65%	12%	1%	10%	11%	0%	0%	0%	0%	0%	0%	77.25%
Mattresses	5%	65%	20%		15%	15%							69.68%
Non-inert Fines		50%	40%		25%	25%							50.00%
Other (100% inert)													100.00%

Component	Lignin biodegradability DOCf	Non-lignin biodegradability DOCf	Moisture, %fresh matter	Lignin, % dm	Hemicellulose, %dm	Cellulose, %dm	Starch, %dm	Sugar, % dm	Fat, %dm	Proteins, %dm	Fibre, %dm	Readily soluble, %dm	Other (inert), % dm
<b>Commercial &amp; industrial waste</b>													
Commercial	5%	65%	37%		8%	76%							16.00%
Paper and Card	5%	65%	15%	15%	9%	61%							15.00%
General industrial waste	5%	65%	37%		8%	76%							16.00%
Food and Abattoir	15%	70%	70%	5%	11%	11%	36%	7%	6%	18%			6.00%
Food effluent	15%	70%	65%		55%	7%							37.40%
Construction and demolition	5%	65%	30%		9%	9%							83.00%
Miscellaneous process waste	5%	65%	20%		10%	10%							80.00%
Other waste	5%	65%	20%		25%	25%							50.00%
Miscellaneous Combustible	5%	65%	20%		25%	25%							50.00%
Furniture	5%	65%	12%	1%	10%	11%	0%	0%	0%	0%			77.25%
Garden	10%	65%	55%	20%	16%	20%			2%			26%	17.10%
Sewage sludge	5%	65%	70%		14%	14%							72.00%
Textiles / Carpet and Underlay	5%	65%	20%	0%	15%	15%	0%	0%	0%	0%			69.68%
Wood	5%	65%	17%	26%	12%	42%							21.00%
Sanitary	5%	65%	65%	0%	0%	47%	0%	0%	0%	0%			52.70%
Other	5%	65%											100.00%

**Table A 3.5.3 Amount of waste landfilled (1945 to 2017)**

Year	England	Scotland	Wales	Northern Ireland	Total	Year	England	Scotland	Wales	Northern Ireland	Total
1945	70.9	9.0	4.6	2.3	86.9	1982	77.1	8.5	4.6	2.5	92.7
1946	71.2	9.0	4.6	2.4	87.2	1983	77.1	8.5	4.6	2.6	92.8
1947	71.5	9.0	4.6	2.4	87.4	1984	77.2	8.5	4.6	2.6	92.8
1948	71.7	9.0	4.6	2.4	87.7	1985	77.2	8.4	4.6	2.6	92.8
1949	72.3	9.0	4.6	2.4	88.4	1986	77.3	8.4	4.6	2.6	92.9
1950	72.9	9.1	4.6	2.4	89.1	1987	77.3	8.3	4.6	2.6	92.9
1951	73.6	9.1	4.6	2.5	89.8	1988	77.4	8.3	4.6	2.6	92.9
1952	74.2	9.1	4.7	2.5	90.5	1989	77.4	8.3	4.6	2.6	92.9
1953	74.2	9.1	4.7	2.5	90.4	1990	77.7	8.3	4.7	2.6	93.3
1954	74.2	9.0	4.6	2.4	90.3	1991	77.7	11.3	4.7	2.6	96.3
1955	74.2	9.0	4.6	2.4	90.3	1992	77.7	12.2	4.7	2.6	97.2
1956	75.7	9.1	4.7	2.5	92.0	1993	77.6	14.0	4.6	2.6	98.8
1957	77.2	9.2	4.8	2.5	93.7	1994	77.6	15.9	4.6	2.6	100.7
1958	78.6	9.3	4.8	2.6	95.4	1995	81.8	15.0	4.9	2.8	104.5
1959	80.1	9.5	4.9	2.6	97.1	1996	80.7	15.0	4.8	2.8	103.3
1960	81.5	9.6	5.0	2.6	98.8	1997	81.1	14.0	4.8	2.8	102.7
1961	81.1	9.7	4.9	2.7	98.4	1998	75.0	11.9	4.5	2.6	93.9
1962	80.9	9.6	4.9	2.7	98.0	1999	69.3	10.9	4.1	2.4	86.6
1963	84.5	10.0	5.1	2.8	102.4	2000	67.4	11.2	4.0	2.3	84.9
1964	84.6	9.9	5.1	2.8	102.4	2001	71.4	8.9	4.2	2.4	86.9
1965	85.7	10.0	5.1	2.8	103.6	2002	66.8	8.2	3.9	2.3	81.3
1966	85.3	9.9	5.1	2.8	103.2	2003	65.4	7.9	3.8	2.2	79.4
1967	85.0	9.8	5.1	2.8	102.7	2004	64.9	7.8	3.8	2.2	78.7
1968	84.8	9.7	5.1	2.8	102.4	2005	60.0	7.1	3.5	2.0	72.6
1969	84.0	9.6	5.0	2.8	101.4	2006	61.7	7.1	4.0	2.0	74.8
1970	83.8	9.5	5.0	2.8	101.0	2007	60.7	7.4	3.2	1.9	73.2
1971	82.8	9.3	4.9	2.7	99.8	2008	53.9	6.1	2.9	1.6	64.5
1972	81.8	9.2	4.8	2.7	98.5	2009	44.0	4.7	2.5	1.1	52.3
1973	81.3	9.1	4.8	2.7	97.9	2010	43.6	4.6	2.3	1.0	51.4

Year	England	Scotland	Wales	Northern Ireland	Total	Year	England	Scotland	Wales	Northern Ireland	Total
1974	79.9	9.0	4.8	2.6	96.3	2011	44.7	4.7	2.2	1.0	52.5
1975	80.1	9.0	4.8	2.6	96.5	2012	41.8	4.5	2.2	1.1	49.6
1976	78.8	8.8	4.7	2.6	94.9	2013	41.1	4.1	2.2	1.1	48.4
1977	78.4	8.8	4.7	2.6	94.5	2014	41.3	4.1	1.5	1.3	48.2
1978	78.8	8.8	4.7	2.6	95.0	2015	43.9	4.2	1.3	1.6	51.0
1979	78.7	8.8	4.7	2.6	94.7	2016	44.7	3.7	2.0	1.9	52.3
1980	78.6	8.7	4.7	2.6	94.6	2017	45.4	3.8	1.8	1.7	52.8
1981	77.0	8.5	4.6	2.5	92.7						

### A 3.5.1.2 Methane emissions

The right-most column of **Table A 3.5.4** shows the current estimate of methane emitted from UK landfills, according to the approach outlined in **Chapter 7**, taking account of recovery and oxidation.

**Table A 3.5.4 Amount of waste landfilled and methane generated, captured, utilised, flared, oxidised and emitted**

Year	Waste Landfilled	Methane generated	Methane captured		Methane used for power generation		Methane flared		Residual methane oxidised		Methane emitted	
	Mt	Kt	kt	%	kt	%	kt	%	kt	%	kt	%
1990	93.25	2,709	33	1%	33	1%	0	0%	268	10%	2408	89%
1991	96.32	2,752	50	2%	50	2%	0	0%	270	10%	2432	88%
1992	97.18	2,797	90	3%	90	3%	0	0%	271	10%	2436	87%
1993	98.85	2,837	107	4%	107	4%	0	0%	273	10%	2457	87%
1994	100.75	2,878	124	4%	124	4%	0	0%	275	10%	2479	86%
1995	104.50	2,939	135	5%	135	5%	0	0%	280	10%	2524	86%
1996	103.26	2,983	170	6%	170	6%	0	0%	281	9%	2532	85%
1997	102.70	3,022	218	7%	218	7%	0	0%	280	9%	2524	84%
1998	93.85	3,038	278	9%	278	9%	0	0%	276	9%	2484	82%

Year	Waste Landfilled	Methane generated	Methane captured		Methane used for power generation		Methane flared		Residual methane oxidised		Methane emitted	
	Mt	Kt	kt	%	kt	%	kt	%	kt	%	kt	%
1999	86.63	3,032	394	13%	394	13%	0	0%	264	9%	2374	78%
2000	84.85	3,028	500	17%	500	17%	0	0%	253	8%	2275	75%
2001	86.92	3,040	566	19%	566	19%	0	0%	247	8%	2227	73%
2002	81.28	3,021	599	20%	598	20%	1	0%	242	8%	2180	72%
2003	79.38	2,981	723	24%	723	24%	0	0%	226	8%	2032	68%
2004	78.71	2,937	874	30%	874	30%	0	0%	206	7%	1857	63%
2005	72.59	2,870	926	32%	926	32%	0	0%	194	7%	1750	61%
2006	74.83	2,753	950	35%	944	34%	6	0%	180	7%	1622	59%
2007	73.21	2,645	989	37%	987	37%	2	0%	166	6%	1490	56%
2008	64.51	2,528	1065	42%	980	39%	85	3%	146	6%	1316	52%
2009	52.33	2,400	1112	46%	1015	42%	97	4%	127	5%	1159	48%
2010	51.38	2,278	1200	53%	1066	47%	134	6%	108	5%	971	43%
2011	52.54	2,159	1181	55%	1075	50%	106	5%	98	5%	881	41%
2012	49.55	2,041	1127	55%	1042	51%	85	4%	91	4%	822	40%
2013	48.43	1,929	1152	60%	1035	54%	117	6%	78	4%	699	36%
2014	48.15	1,820	1151	63%	1007	55%	145	8%	67	4%	602	33%
2015	50.98	1,716	1065	62%	974	57%	91	5%	65	4%	586	34%
2016	52.31	1,627	1007	62%	941	58%	67	4%	62	4%	558	34%
2017	52.76	1544	918	59%	857	56%	61	4%	63	4%	563	36%

Notes

- a. Methane generated is based on the MELMod model.
- b. Methane captured is the sum of methane used for power generation and methane flared.

- c. Methane used for power generation is calculated from official figures on landfill gas electricity generation (Digest of UK Energy Statistics (BEIS, 2016), in GWh/year, assuming a net calorific value for methane of 50 GJ/tonnes and a conversion efficiency between methane use and electricity export of 30% rising to 36%, which includes parasitic losses and on-site use of electricity, e.g. for gas blowers, leachate treatment and site offices.
- d. Methane flared is calculated from site-specific data provided by the Environment Agency at regulated sites for 2009 to 2013, from SEPA for 2013, from a study carried out during 2014, and from site-specific data provided voluntarily by site operators.
- e. Methane oxidised is based on the IPCC default oxidation factor of 10%, applied to methane remaining after subtraction of the amount captured.
- f. Methane emitted = (methane generated – methane captured) x (1-oxidation factor).

**A 3.5.1.3 Overseas Territories and Crown Dependencies**

For the overseas territories and crown dependencies, the IPCC landfill model is used. Where available, country-specific waste generation and composition data have been applied and appropriate defaults chosen e.g. taking into account climatic variation. There are no landfill emissions for Gibraltar as waste is exported. **Table A 3.5.5** below gives the parameters used.

**Table A 3.5.5 Parameters used in landfill emission estimates for overseas territories and crown dependencies**

Region	Methodology	Activity data	MCF	DOC
Guernsey	IPCC Landfill Model	2005 onwards: total MSW to landfill data and percentage that is plastics, other inert. Prior to 2005: flat-lined 2005 data	IPCC default values; waste management type is unmanaged, deep (results from expert consultation, 2014)	IPCC default values
Jersey	N/A, all MSW is incinerated for energy from waste	N/A	N/A	N/A
Gibraltar	N/A, all MSW used to be incinerated, now all waste is exported to be landfilled in Spain.	N/A	N/A	N/A
Isle of Man	IPCC Landfill Model	2004 onwards: all waste incinerated for energy from waste. Prior to 2004: population and IPCC default waste per capita for Western Europe	IPCC default values; waste management type is 50% unmanaged, deep and 50% managed, semi-aerobic (results from expert consultation, 2014)	IPCC default values

Region	Methodology	Activity data	MCF	DOC
Bermuda	IPCC Landfill Model	Total MSW to landfill (Environmental Statistics Compendium)	IPCC default values; no information on management system so assume unmanaged deep	IPCC default values
Cayman Islands	IPCC Landfill Model	2000 onwards: Total MSW to landfill (Department of Environmental Health). Prior to 2000: flat-lined 2000 data	IPCC default values; landfill sites are lined and managed to some degree, but with limited information, "Uncategorised" considered appropriate	IPCC default values
Falkland Islands	IPCC Landfill Model	1998: Halcrow Report. Other years: flat-lined after advice in personal communication from environmental officer	IPCC default values; waste management type is unmanaged, shallow (results from expert consultation, 2014)	IPCC default values

**A 3.5.2 Biological Treatment of Solid Waste (5B)****Table A 3.5.6 Activity Data: Inputs in the composting process**

Year	Composting (Non-household) (Mg)	Composting (Household) (Mg)	MBT - Composting (Mg)
1990	0	181,322	0
1991	19,283	181,756	0
1992	33,194	182,146	0
1993	48,000	182,638	0
1994	64,000	183,213	0
1995	140,000	183,654	0
1996	220,000	184,132	0
1997	315,000	184,643	0
1998	675,000	185,364	0
1999	833,044	185,979	0
2000	1,034,000	186,676	0
2001	1,663,852	187,441	67,882
2002	1,828,000	187,373	62,537
2003	1,953,000	187,300	57,192
2004	2,667,000	238,235	37,179
2005	3,424,000	289,058	88,917
2006	4,090,000	339,909	110,618
2007	4,459,000	390,804	542,678
2008	4,285,000	441,622	629,269
2009	5,265,711	492,393	438,011
2010	5,444,092	496,077	1,282,060
2011	6,053,273	500,008	1,898,570
2012	5,850,257	503,005	1,719,118
2013	5,867,640	505,833	2,138,366
2014	5,954,185	509,350	2,439,341

Year	Composting (Non-household) (Mg)	Composting (Household) (Mg)	MBT - Composting (Mg)
2015	6,010,218	513,040	2,740,317
2016	6,135,538	517,007	3,041,292
2017	6,286,717	519,898	3,342,268

**Table A 3.5.7 Activity Data: Inputs in the anaerobic digestion process**

Year	Anaerobic digestion – non agricultural residue (Mg)	Anaerobic digestion - MBT (Mg)
1990	0	0
1991	1,678	0
1992	1,678	0
1993	1,678	0
1994	5,435	0
1995	5,435	0
1996	5,435	0
1997	6,061	0
1998	6,061	0
1999	6,061	0
2000	6,061	0
2001	6,061	16,970
2002	56,155	15,634
2003	56,155	14,298
2004	87,513	37,179
2005	191,153	17,783
2006	203,676	27,655
2007	223,714	40,847
2008	247,633	66,664
2009	370,988	365,570
2010	721,666	72,262

Year	Anaerobic digestion – non agricultural residue (Mg)	Anaerobic digestion - MBT (Mg)
2011	1,286,593	104,815
2012	1,842,418	795,961
2013	2,850,234	600,692
2014	4,286,018	703,455
2015	5,672,507	806,218
2016	7,268,381	908,982
2017	7,772,206	1,011,745

### A 3.5.3 Waste Incineration (5C)

Table A 3.5.8 Activity Data: UK Waste Incineration

Year	Municipal Waste Incineration <sup>a</sup> (Mt)	Clinical Waste Incineration (Mt)	Chemical Waste Incineration (Mt)	Sewage Sludge Incineration (Mt)
1990	2.093	0.350	0.290	0.075
1991	2.069	0.350	0.290	0.069
1992	1.945	0.330	0.290	0.072
1993	1.677	0.310	0.290	0.084
1994	1.148	0.290	0.289	0.072
1995	0.996	0.270	0.289	0.082
1996	1.062	0.250	0.288	0.088
1997	-	0.230	0.287	0.081
1998	-	0.236	0.287	0.185
1999	-	0.242	0.286	0.189
2000	-	0.248	0.285	0.194
2001	-	0.254	0.285	0.198
2002	-	0.260	0.284	0.203
2003	-	0.221	0.258	0.207
2004	-	0.182	0.232	0.212
2005	-	0.143	0.206	0.216
2006	-	0.105	0.180	0.220
2007	-	0.111	0.171	0.215

Year	Municipal Waste Incineration <sup>a</sup> (Mt)	Clinical Waste Incineration (Mt)	Chemical Waste Incineration (Mt)	Sewage Sludge Incineration (Mt)
2008	-	0.115	0.143	0.192
2009	-	0.121	0.135	0.199
2010	-	0.114	0.148	0.231
2011	-	0.107	0.147	0.224
2012	-	0.107	0.145	0.209
2013	-	0.101	0.164	0.204
2014	-	0.104	0.172	0.177
2015	-	0.096	0.166	0.170
2016	-	0.095	0.169	0.148
2017	-	0.095	0.139	0.144

a Note that all MSW incinerators were either closed or converted to extract power by 1997. In the latter case they were then considered to be power generation and so emissions were reported in 1A1a.

**Table A 3.5.9 Emissions Data: UK Waste Incineration**

Year	Chemical Waste Incineration	Accidental Fires	MSW Incineration <sup>a</sup>	Clinical Waste Incineration	Sewage Sludge Incineration	Total
<b>Carbon Dioxide (kt CO<sub>2</sub>)</b>						
1990	381.8	NE	606.0	308.0	NA	1,295.7
1995	380.0	NE	357.0	237.6	NA	974.7
2000	304.3	NE	NO	218.2	NA	522.6
2005	275.9	NE	NO	126.2	NA	402.2
2010	168.6	NE	NO	100.6	NA	269.2
2012	170.6	NE	NO	93.8	NA	264.3
2013	171.7	NE	NO	89.3	NA	260.9
2014	174.0	NE	NO	91.7	NA	265.7
2015	164.0	NE	NO	84.3	NA	248.3
2016	183.3	NE	NO	83.4	NA	266.7
2017	174.5	NE	NO	83.9	NA	258.4

Year	Chemical Waste Incineration	Accidental Fires	MSW Incineration <sup>a</sup>	Clinical Waste Incineration	Sewage Sludge Incineration	Total
<b>Methane (kt CH<sub>4</sub>)</b>						
1990	NE	1.009	4.175	0.007	0.029	5.311
1995	NE	0.984	1.987	0.005	0.032	3.100
2000	NE	0.772	NO	0.005	0.076	0.916
2005	NE	0.704	NO	0.003	0.084	0.834
2010	NE	0.347	NO	0.002	0.090	0.462
2012	NE	0.331	NO	0.002	0.082	0.437
2013	NE	0.290	NO	0.002	0.080	0.399
2014	NE	0.277	NO	0.002	0.069	0.375
2015	NE	0.283	NO	0.002	0.066	0.379
2016	NE	0.287	NO	0.002	0.058	0.374
2017	NE	0.281	NO	0.002	0.056	0.368
<b>Nitrous oxide (kt N<sub>2</sub>O)</b>						
1990	0.029	NE	0.058	0.011	0.060	0.155
1995	0.029	NE	0.033	0.008	0.066	0.129
2000	0.029	NE	NO	0.007	0.155	0.191
2005	0.021	NE	NO	0.004	0.173	0.198
2010	0.015	NE	NO	0.003	0.185	0.203
2012	0.014	NE	NO	0.003	0.167	0.185
2013	0.016	NE	NO	0.003	0.163	0.183
2014	0.017	NE	NO	0.003	0.141	0.162
2015	0.017	NE	NO	0.003	0.136	0.156
2016	0.017	NE	NO	0.003	0.118	0.138
2017	0.014	NE	NO	0.003	0.115	0.132

a Note that all MSW incinerators were either closed or converted to extract power by 1997. In the latter case they were then considered to be power generation and so emissions were reported in 1A1a.

**A 3.5.4 Wastewater Handling (5D)****A 3.5.4.1 5D1 Domestic and Commercial Waste Water Handling and Sludge Disposal****Table A 3.5.10 UK Domestic and Commercial Waste Water Treatment (5D1) Activity Data**

Treatment/disposal route		unit	1990	1995	2000	2005	2010	2013	2014	2015	2016	2017
Total Sludge		kt tds	1634	1657	1682	1768	1666	1598	1596	1597	1659	1669
Population Equivalent		million	68.3	69.2	70.2	70.5	69.3	72.9	72.9	73.0	73.2	73.4
Additional Treatment	Digested	kt tds	402	433	468	810	835	694	710	657	673	660
	Advanced Digested	kt tds	101	107	115	329	373	295	406	454	556	615
	Composted	kt tds	7	8	8	15	25	48	27	7	18	14
Disposal route	Farmland	kt tds	508	547	590	1216	1282	1287	1332	1422	1434	1479
	Landfill	kt tds	160	153	110	131	35	6	4	7	9	8
	Incineration	kt tds	68	80	211	252	238	252	232	161	198	168
	Sea	kt tds	782	721	611	0	0	0	0	0	0	0
	Composted	kt tds	2	2	2	13	23	53	28	7	18	14
	Land Reclamation	kt tds	31	30	30	96	44	0	0	0	0	0
	Other	kt tds	84	124	129	61	44	0	0	0	0	0

Where tds is total dissolvable solids; this is assumed to be comparable to Biochemical Oxygen Demand (BOD)

**Table A 3.5.11 UK Domestic and Commercial Waste Water Treatment (5D1) Implied Emission Factors**

Treatment/disposal route		unit	1990	1995	2000	2005	2010	2013	2014	2015	2016	2017	
Mechanical treatment and storage <sup>1</sup>		kt/Mt tds	2.70	2.70	2.70	2.70	2.70	2.70	2.70	2.70	2.70	2.70	
Additional Treatment	Digested <sup>2</sup>	kt/Mt tds	16.54	16.29	16.47	16.95	16.50	16.01	15.71	15.63	13.30	13.26	
	Advanced Digested	kt/Mt tds	4.54	4.54	4.54	4.54	4.54	4.56	4.52	4.52	4.57	4.57	
	Composted	kt/Mt tds	10.00	10.00	10.00	10.00	10.00	10.00	10.00	10.00	10.00	10.00	
Disposal route	Farmland <sup>3</sup>	kt/Mt tds	1.36	1.36	1.36	1.44	1.41	1.45	1.31	1.30	1.28	1.22	
	Landfill	kt/Mt tds	15.09	15.09	15.09	15.09	15.09	15.09	15.09	15.09	15.09	15.09	
	Incineration	kt/Mt tds	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	
	Sea <sup>4</sup>	kt/Mt tds	60.00	60.00	60.00	60.00	60.00	60.00	60.00	60.00	60.00	60.00	
	Composted	kt/Mt tds	0.60	0.60	0.60	0.60	0.60	0.60	0.60	0.60	0.60	0.60	
	Land Reclamation <sup>5</sup>	kt/Mt tds	1.36	1.36	1.36	1.36	1.36	1.42	1.29	1.29	1.29	1.27	1.22
	Other <sup>6</sup>	kt/Mt tds	1.19	1.19	1.19	1.19	1.19	1.25	1.14	1.22	1.22	1.20	1.16
Total <sup>7</sup>		kt/Mt tds	29.82	28.55	30.99	13.62	13.61	12.04	12.14	11.68	10.94	10.87	

1. All waste is mechanically treated and stored, so the emission factor is applied to total sludge.

2. Implied emission factor after methane capture.

3. Emission factor varies depending on how the waste is treated.

4. Not an IEF, this is the default IPCC factor for sea, river and lake discharge.

5. Land reclamation hasn't got associated reported emissions, so the factor is based on a weighted average of other waste to land (farmland, composting) IEFs.

6. Other hasn't got associated reported emissions, the factor is based on a weighted average of all other disposal IEFs.

7. For information, IEF when dividing total emissions by total activity.

**Table A 3.5.11 UK Domestic and Commercial Waste Water Treatment (5D1) Emission Estimates (kt CH<sub>4</sub>)**

Treatment/disposal route		1990	1995	2000	2005	2010	2013	2014	2015	2016	2017
Mechanical treatment and storage		4.41	4.47	4.54	4.77	4.50	4.31	4.31	4.31	4.48	4.51
Additional Treatment	Digested	6.65	7.06	7.71	13.74	13.77	11.11	11.15	10.27	8.95	8.75
	AdvancedDigested	0.46	0.49	0.52	1.49	1.69	1.34	1.84	2.06	2.55	2.81
	Composted	0.07	0.08	0.08	0.15	0.25	0.48	0.27	0.07	0.18	0.14
Disposal route	Farmland	0.68	0.73	0.79	1.72	1.82	1.90	1.76	1.85	1.85	1.82
	Landfill	2.41	2.30	1.66	1.97	0.53	0.09	0.06	0.10	0.14	0.12
	Incineration	-	-	-	-	-	-	-	-	-	-
	Sea	33.92	31.99	36.63	-	-	-	-	-	-	-
	Composted	0.00	0.00	0.00	0.01	0.01	0.03	0.02	0.00	0.01	0.01
	LandReclamation	0.04	0.04	0.04	0.13	0.06	-	-	-	-	-
	Other	0.10	0.15	0.15	0.07	0.05	-	-	-	-	-
Total		48.74	47.31	52.13	24.09	22.68	19.24	19.39	18.66	18.14	18.14

**Table A 3.5.12 UK Private Waste Water Management System Emission Estimates Parameters (5D1)**

Data	Unit	1990	1995	2000	2005	2010	2013	2014	2015	2016	2017
Estimated population connected to private waste water management systems	population (thousands)	1417.18	1440.63	1438.67	1465.29	1525.78	1562.08	1579.31	1584.43	1598.92	1605.40
BOD value applied	g/person/day	66.33	66.33	66.33	68.73	65.90	60.09	59.96	59.96	62.07	62.30

## A 3.5.4.2 5D2 Industrial Waste Water Handling and Sludge Disposal

Table A 3.5.13 UK Industrial Waste Water Treatment Activity Data (5D2)

Sector	Unit	1990	1995	2000	2005	2010	2013	2014	2015	2016	2017
Organic chemical production	Mt	1.617	1.617	1.751	1.752	1.487	1.539	1.581	1.668	1.575	1.616
Milk-processing	million PE	1.464	1.464	1.464	0.625	0.629	0.733	0.791	0.803	0.788	0.792
Manufacture of fruit and vegetable products	million PE	1.145	1.145	1.145	1.094	1.094	0.988	1.003	1.134	1.275	1.343
Potato-processing	million PE	0.302	0.302	0.302	0.289	0.289	0.261	0.265	0.299	0.336	0.355
Meat industry	million PE	0.623	0.623	0.623	0.618	0.648	0.634	0.696	0.723	0.717	0.760
Breweries	million PE	0.094	0.094	0.094	0.097	0.096	0.098	0.102	0.102	0.103	0.104
Production of alcohol and alcoholic beverages	million PE	1.931	1.931	1.931	1.992	1.967	2.009	2.093	2.091	2.105	2.127
Manufacture of animal feed from plant products	million PE	0.476	0.476	0.476	0.300	0.378	0.349	0.361	0.354	0.359	0.398
Manufacture of gelatine and of glue from hides, skin and bones	million PE	0.013	0.013	0.013	0.013	0.013	0.013	0.013	0.013	0.013	0.013
Malt-houses	million PE	0.207	0.207	0.207	0.213	0.210	0.215	0.224	0.224	0.225	0.228
Fish-processing industry	million PE	0.018	0.018	0.018	0.006	0.006	0.006	0.007	0.007	0.007	0.007
Total Food and Drink	million PE	6.273	6.273	6.273	5.247	5.329	5.306	5.555	5.750	5.929	6.126

Where PE is population equivalent

## A 3.6 UK CROWN DEPENDENCIES AND OVERSEAS TERRITORIES

### A 3.6.1 Overview of Data Sources

Fuel use data for Isle of Man, Guernsey and Jersey are assumed to be included in UK national energy statistics (see **Section 1.1.2.2**), so fuel thought to be used in these territories are split out from UK total consumption unless otherwise stated in **Section A 4.2.1**.

Activity data including fuel use data for other territories are obtained from government departments for those territories, specifically:

- The Cayman Islands Government Department of Environment Sustainable Development Unit;
- The Department of Energy Bermuda; and,
- The Falkland Islands Government Policy Unit.

Activity and emissions data estimates from LULUCF sources and sinks have been researched via the FAOSTAT database, to supplement data available from the OTs. The LULUCF data for Cayman Islands from FAOSTAT (FAO, 2018) indicates zero emissions, using Tier 1 methods. The data sources and methodologies used for other sectors are described in the main methodology sections of the NIR.

### A 3.6.2 Activity and Emissions Data

**Table A 3.6.1 Isle of Man, Guernsey and Jersey – Emissions of Direct GHGs (Mt CO<sub>2</sub> equivalent)**

Sector	1990	1995	2000	2005	2010	2013	2014	2015	2016	2017
1. Energy	1.48	1.64	1.75	1.55	1.47	1.56	1.48	1.37	1.43	1.40
2. Industrial Processes and Product Use	0.0001	0.0050	0.0245	0.0515	0.0824	0.0816	0.0823	0.0812	0.0758	0.0758
3. Agriculture	0.14	0.14	0.15	0.10	0.13	0.12	0.12	0.12	0.12	0.12
4. LULUCF	-0.02	-0.04	-0.04	-0.03	-0.03	-0.02	-0.03	-0.01	-0.03	-0.03
5. Waste	0.16	0.16	0.17	0.17	0.15	0.14	0.14	0.14	0.13	0.13
7. Other	0	0	0	0	0	0	0	0	0	0
<b>Total</b>	1.74	1.91	2.05	1.84	1.81	1.88	1.79	1.69	1.73	1.69

**Table A 3.6.2 Isle of Man, Guernsey and Jersey – Combustion activity data**

Fuel	Fuel Unit	1990	1995	2000	2005	2010	2013	2014	2015	2016	2017
Aviation spirit	Mt	0.00813	0.00907	0.0123	0.0135	0.00715	0.00576	0.00569	0.00364	0.00285	0.00181
Aviation turbine fuel	Mt	0.0803	0.072	0.0931	0.107	0.0935	0.0855	0.0806	0.0855	0.0881	0.0951
Burning oil	Mt	0.225	0.261	0.35	0.357	0.363	0.381	0.373	0.386	0.408	0.416
Coal	Mt	0.0345	0.0218	0.0152	0.0103	0.000891	0.000891	0.000891	0.000972	0.000975	0.000978
DERV	Mt	0.0737	0.0937	0.133	0.118	0.125	0.131	0.132	0.135	0.139	0.139
Fuel oil	Mt	0.469	0.58	0.438	0.0777	0.0889	0.222	0.158	0.0598	0.0777	0.0454
Gas oil	Mt	0.164	0.174	0.162	0.181	0.119	0.0953	0.0873	0.0855	0.088	0.0879
LPG	Mth	11.7	13.5	46.4	25	22.4	19.1	17.4	16.9	16.2	15.3
Lubricants											
MSW	Mt	0.121	0.169	0.216	0.391	0.359	0.321	0.309	0.296	0.293	0.286
Natural gas	Mth	0	0	0	106	126	114	124	113	127	125
Petrol	Mt	0.236	0.228	0.217	0.219	0.191	0.175	0.171	0.169	0.166	0.164
Petroleum wax											
Wood	Mt	0.00765	0.00765	0.00765	0.00765	0.00765	0.00765	0.00765	0.00765	0.00765	0.00765

**Table A 3.6.3 Isle of Man, Guernsey and Jersey – Animal numbers**

Livestock Category	1990	1995	2000	2005	2010	2013	2014	2015	2016	2017
Dairy	15,888	15,729	16,186	13,127	11,455	10,820	10,841	10,704	11,260	11,331
Non-dairy	28,663	28,333	29,176	16,770	28,615	26,097	27,498	26,513	26,338	24,387

Livestock Category	1990	1995	2000	2005	2010	2013	2014	2015	2016	2017
Sheep	151,764	160,228	176,259	87,537	138,251	134,501	134,310	133,666	126,057	128,927
Pigs	4,854	5,411	4,609	1,148	4,086	2,605	2,602	2,861	2,386	2,342
Poultry	84,048	46,481	46,448	58,160	54,400	57,924	58,850	62,916	60,231	62,537
Goats	333	347	376	141	288	416	477	539	352	326
Horses	2,785	2,785	2,785	2,822	3,236	3,265	3,163	2,891	3,258	2,705

**Table A 3.6.4 Isle of Man, Guernsey and Jersey – Total emissions from Agricultural Soils (kg N<sub>2</sub>O)**

Territory	1990	1995	2000	2005	2010	2013	2014	2015	2016	2017
Isle of Man	39,497	40,148	43,721	27,111	37,931	34,889	36,079	35,396	35,610	34,166
Guernsey	5,713	5,759	5,277	4,750	4,624	4,729	4,832	4,641	4,596	4,539
Jersey	6,361	6,444	6,605	5,256	5,394	5,457	5,451	5,278	5,255	5,248

**Table A 3.6.5 Cayman Islands, Falklands Islands, and Bermuda - Emissions of Direct GHGs (Mt CO<sub>2</sub> equivalent)**

Sector	1990	1995	2000	2005	2010	2013	2014	2015	2016	2017
1. Energy	1.23	1.25	1.4	1.59	1.58	1.55	1.64	1.69	1.51	1.61
2. Industrial Processes and Product Use	0.0000957	0.00274	0.0146	0.0308	0.0418	0.0372	0.0369	0.0363	0.0321	0.0321
3. Agriculture	0.389	0.367	0.338	0.312	0.33	0.323	0.318	0.313	0.311	0.319
4. LULUCF	1.62	1.62	1.75	1.93	1.96	1.91	1.99	2.04	1.86	1.96
5. Waste	0	0	0	0	0	0	0	0	0	0
7. Other	0	0	0	0	0	0	0	0	0	0
<b>Total</b>	<b>3.24</b>	<b>3.24</b>	<b>3.5</b>	<b>3.87</b>	<b>3.91</b>	<b>3.82</b>	<b>3.98</b>	<b>4.08</b>	<b>3.71</b>	<b>3.92</b>

**Table A 3.6.6 Cayman Islands, Falklands Islands, and Bermuda – Combustion activity data**

Fuel	Fuel Unit	1990	1995	2000	2005	2010	2013	2014	2015	2016	2017
Aviation spirit	Mt	0.000342	0	0	0	0	0	0	0	0	0
Aviation turbine fuel	Mt	0.0816	0.0373	0.0411	0.0566	0.071	0.0552	0.0511	0.0507	0.0506	0.0504
Biogenic products											
Burning oil	Mt	0.00433	0.00535	0.00616	0.00797	0.0072	0.00692	0.00715	0.00744	0.00827	0.00884
Coal	Mt	0.147	0.127	0.103	0.232	0.0964	0.13	0.0854	0.0856	0.0857	0.0859
DERV	Mt	0.145	0.152	0.266	0.303	0.315	0.314	0.315	0.314	0.314	0.316
Fuel oil	Mt	0.398	0.447	0.428	0.475	0.546	0.595	0.572	0.583	0.592	0.573
Gas oil	Mt	7.92	8.85	9.76	11.3	11.5	11.7	12.1	12.3	12.5	12.6
LPG	Mth	0.00042	0.204	0.204	0.204	0.192	0.163	0.166	0.174	0.167	0.162
MSW	Mt	0.0473	0.0585	0.0673	0.0882	0.105	0.0905	0.0966	0.0971	0.107	0.115
Natural gas	Mth	0.189	0.192	0.214	0.17	0.196	0.174	0.135	0.147	0.147	0.141
Petrol	Mt	0	0	0	0	0.000609	0.00126	0.00123	0.00135	0.00143	0.00149
Lubricants	Mt	0.00025	0.000214	0.000174	0.000413	0.000216	0.000166	0.000187	0.000251	0.000251	0.000268
Petroleum waxes	Mt	0.000342	0	0	0	0	0	0	0	0	0

**Table A 3.6.7 Cayman Islands, Falklands Islands, and Bermuda – Animal numbers**

Livestock Category	1990	1995	2000	2005	2010	2013	2014	2015	2016	2017
Dairy Cattle	2,161	1,862	1,911	1,145	868	765	723	675	634	647
Non-dairy Cattle	5,256	4,861	5,077	7,845	6,360	5,795	5,609	4,748	4,503	4,913
Sheep	739,999	717,571	669,905	580,864	478,625	486,037	483,135	482,131	479,752	490,213

Livestock Category	1990	1995	2000	2005	2010	2013	2014	2015	2016	2017
Goats	405	867	1,286	1,704	2,251	2,080	1,891	1,812	2,031	2,337
Horses	2,217	2,069	1,703	1,417	1,269	1,192	1,202	1,223	1,252	1,244
Swine	1,116	1,174	1,376	1,384	1,233	1,248	1,242	1,058	1,234	1,301
Poultry	15,319	14,664	20,890	27,164	32,293	37,764	27,769	39,458	39,948	26,710
Deer	0	0	0	0	184	243	243	243	243	243

**Table A 3.6.8 Cayman Islands, Falklands Islands, and Bermuda – Total emissions from Agricultural Soils (kg N<sub>2</sub>O)**

Territory	1990	1995	2000	2005	2010	2013	2014	2015	2016	2017
Bermuda	2,409	2,606	2,803	2,786	2,801	2,802	2,802	2,803	2,803	2,798
Cayman Islands	2,698	3,212	3,691	4,173	4,446	4,566	4,342	4,359	3,916	3,862
Falkland Islands	363,328	350,782	328,265	288,461	240,569	245,081	243,913	242,966	241,529	239,827

**Table A 3.6.9 Cayman Islands, Falklands Islands, and Bermuda - Amount of synthetic fertilizer applied**

Country	kg N applied
Cayman Islands	23,080
Falkland Islands	0
Bermuda	1,480

**Table A 3.6.10 Gibraltar – Emissions of Direct GHGs (Mt CO<sub>2</sub> equivalent)**

Sector	1990	1995	2000	2005	2010	2013	2014	2015	2016	2017
1. Energy	0.219	0.216	0.242	0.278	0.288	0.319	0.308	0.331	0.331	0.322

Sector	1990	1995	2000	2005	2010	2013	2014	2015	2016	2017
2. Industrial Processes and Other Product Use	0.000	0.001	0.003	0.006	0.010	0.011	0.011	0.012	0.011	0.011
3. Agriculture	-	-	-	-	-	-	-	-	-	-
4. LULUCF	-	-	-	-	-	-	-	-	-	-
5. Waste	0.007	0.009	0.012	0.001	0.003	0.002	0.002	0.002	0.002	0.002
Total	0.226	0.226	0.257	0.285	0.301	0.331	0.321	0.345	0.344	0.335

**Table A 3.6.11 Gibraltar – Combustion activity data**

Fuel	Unit	1990	1995	2000	2005	2010	2013	2014	2015	2016	2017
Aviation turbine fuel	Mt	7.90	6.06	5.27	8.43	6.55	8.12	8.64	8.73	11.34	11.85
Charcoal	Mt	-	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
Clinical waste	Mt	-	-	-	-	1.81	0.30	0.26	0.27	0.27	0.27
DERV	Mt	0.82	0.41	0.92	2.21	2.45	2.27	2.74	3.19	3.76	3.81
Fuel oil	Mt	11.69	9.72	10.59	9.36	7.17	-	-	-	-	-
Gas oil	Mt	19.88	23.59	26.37	33.22	41.36	63.65	56.69	60.46	56.92	53.31
LPG	Mth	4.13	4.13	4.13	4.13	4.13	4.13	4.13	4.13	4.13	4.34
Lubricant	kt	0.56	0.54	0.47	0.45	0.37	0.29	0.29	0.27	0.27	0.27
MSW	Mt	16.40	19.00	24.09	-	-	-	-	-	-	-
Petrol	Mt	5.95	5.12	7.14	6.93	6.32	4.63	4.77	4.91	5.08	5.08
Petroleum waxes	Mt	0.03	0.02	0.01	0.03	0.02	0.02	0.02	0.02	0.02	0.02

## **ANNEX 4: National Energy Balance**

### **A 4.1 UK ENERGY BALANCE**

The UK energy balance is produced and published annually by the Department of Business, Energy & Industrial Strategy in the Digest of UK Energy Statistics – DUKES. This is available online from:

<https://www.gov.uk/government/collections/digest-of-uk-energy-statistics-dukes>

The aggregate energy balance for the latest year is presented below (Table 1.1 in DUKES). The following sections explain how the energy balance is used for the UK inventory for individual fuel types, and how the data are supplemented with other statistics that may lead to deviations from the DUKES statistics.

The UK energy statistics (detailed breakdown) are presented on a mass basis for liquid and solid fuels, and on a gross energy basis for gaseous fuels (including derived gases). The UK inventory is calculated using these data directly, and for the purposes of reporting in the CRF and NIR, activity data and emission factors are converted to energy units, on a net basis.

The scope of the UK energy balance, as shown below, is fuel use in the United Kingdom and its Crown Dependencies (Jersey, Guernsey and the Isle of Man), as described in the NIR section 1.1.2.2.

The fuel use estimates for Overseas Territories (OTs) are not included within DUKES, and are obtained through direct communications with the respective government contacts in each of the OTs.

**Table A 4.1.1 UK Energy Balance for 2017 (thousand tonnes of oil equivalent, gross energy basis)**

	Coal	Manufactured fuel(1)	Primary oils	Petroleum products	Natural gas(2)	Bioenergy & waste(3)	Primary electricity	Electricity	Heat sold	Total
<b>Supply</b>										
Production	1,934	-	50,944	-	40,019	12,924	20,924	-	-	126,745
Imports	5,807	712	58,480	36,722	45,132	3,475	-	1,562	-	151,891
Exports	-369	-14	-42,040	-25,374	-10,802	-431	-	-293	-	-79,323
Marine bunkers	-	-	-	-2,596	-	-	-	-	-	-2,596
Stock change(4)	+2,098	-2	+361	-113	+1,028	-	-	-	-	+3,373
<b>Primary supply</b>	<b>9,470</b>	<b>696</b>	<b>67,746</b>	<b>8,639</b>	<b>75,377</b>	<b>15,969</b>	<b>20,924</b>	<b>1,269</b>	<b>-</b>	<b>200,090</b>
Statistical difference(5)	-65	+1	-66	-10	+337	-	-	-35	-	+163
<b>Primary demand</b>	<b>9,535</b>	<b>694</b>	<b>67,811</b>	<b>8,649</b>	<b>75,040</b>	<b>15,969</b>	<b>20,924</b>	<b>1,304</b>	<b>-</b>	<b>199,927</b>
Transfers	-	-	-	-	-	-	-	-	-	-
	-	+10	-2,476	+2,483	+224	-237	-5,801	+5,801	-	+4
<b>Transformation</b>	<b>-8,134</b>	<b>379</b>	<b>-65,335</b>	<b>64,539</b>	<b>-27,182</b>	<b>-9,587</b>	<b>-15,124</b>	<b>23,071</b>	<b>1,592</b>	<b>-35,779</b>
Electricity generation	-5,559	-518	-	-533	-24,594	-9,387	-15,124	23,071	-	-32,645
Major power producers	-5,545	-	-	-146	-22,150	-4,404	-15,124	20,358	-	-27,011
Autogenerators	-14	-518	-	-387	-2,445	-4,983	-	2,713	-	-5,634
Heat generation	-4	-1	-	-52	-2,587	-200	-	-	1,592	-1,252

	Coal	Manufactured fuel(1)	Primary oils	Petroleum products	Natural gas(2)	Bioenergy & waste(3)	Primary electricity	Electricity	Heat sold	Total
Petroleum refineries	-	-	-65,795	65,691	-	-	-	-	-	-104
Coke manufacture	-1,435	1,351	-	-	-	-	-	-	-	-84
Blast furnaces	-989	-596	-	-	-	-	-	-	-	-1,585
Patent fuel manufacture	-146	143	-	-66	-	-	-	-	-	-69
Other(7)	-	-	460	-501	-	-	-	-	-	-40
<b>Energy industry use</b>	<b>-</b>	<b>458</b>	<b>-</b>	<b>4,315</b>	<b>4,903</b>	<b>-</b>	<b>-</b>	<b>2,041</b>	<b>322</b>	<b>12,040</b>
Electricity generation	-	-	-	-	-	-	-	1,332	-	1,332
Oil and gas extraction	-	-	-	715	4,244	-	-	51	-	5,010
Petroleum refineries	-	-	-	3,600	92	-	-	375	322	4,390
Coal extraction	-	-	-	-	6	-	-	39	-	45
Coke manufacture	-	182	-	-	-	-	-	1	-	183
Blast furnaces	-	276	-	-	25	-	-	18	-	319
Patent fuel manufacture	-	-	-	-	-	-	-	-	-	-
Pumped storage	-	-	-	-	-	-	-	85	-	85
Other	-	-	-	-	536	-	-	139	-	675
Losses	-	109	-	-	580	-	-	2,283	-	2,972

	Coal	Manufactured fuel(1)	Primary oils	Petroleum products	Natural gas(2)	Bioenergy & waste(3)	Primary electricity	Electricity	Heat sold	Total
<b>Final consumption</b>	<b>1,401</b>	<b>516</b>	-	<b>71,356</b>	<b>42,599</b>	<b>6,145</b>	-	<b>25,851</b>	<b>1,270</b>	<b>149,139</b>
<b>Industry</b>	<b>972</b>	<b>296</b>	-	<b>4,308</b>	<b>8,677</b>	<b>1,162</b>	-	<b>7,964</b>	<b>692</b>	<b>24,071</b>
Unclassified	-	-	-	3,380	1	135	-	-	-	3,515
Iron and steel	23	296	-	5	331	-	-	230	-	885
Non-ferrous metals	19	-	-	8	261	-	-	370	-	659
Mineral products	439	-	-	174	1,222	203	-	524	-	2,562
Chemicals	43	-	-	115	1,748	5	-	1,336	277	3,524
Mechanical engineering etc	8	-	-	0	975	2	-	556	-	1,540
Electrical engineering etc	3	-	-	1	275	-	-	512	-	791
Vehicles	38	-	-	208	595	-	-	405	-	1,246
Food, beverages etc	54	-	-	110	1,624	47	-	945	3	2,783
Textiles, leather etc	44	-	-	43	243	-	-	231	-	561
Paper, printing etc	64	-	-	31	391	611	-	930	-	2,027
Other industries	232	-	-	38	609	159	-	1,801	412	3,251
Construction	4	-	-	195	401	-	-	125	-	725
<b>Transport (6)</b>	<b>11</b>	-	-	<b>55,051</b>	-	<b>997</b>	-	<b>411</b>	-	<b>56,470</b>
Air	-	-	-	12,995	-	-	-	-	-	12,995

	Coal	Manufactured fuel(1)	Primary oils	Petroleum products	Natural gas(2)	Bioenergy & waste(3)	Primary electricity	Electricity	Heat sold	Total
Rail	11	-	-	661	-	-	-	396	-	1,069
Road	-	-	-	40,468	-	997	-	15	-	41,480
National navigation	-	-	-	926	-	-	-	-	-	926
Pipelines	-	-	-	-	-	-	-	-	-	-
<b>Other</b>	<b>419</b>	<b>172</b>	-	<b>4,507</b>	<b>33,496</b>	<b>3,986</b>	-	<b>17,476</b>	<b>578</b>	<b>60,634</b>
Domestic	392	172	-	2,472	25,540	2,216	-	9,062	260	40,116
Public administration	18	-	-	369	3,111	72	-	1,696	97	5,364
Commercial	4	-	-	883	3,868	1,173	-	6,344	220	12,493
Agriculture	-	-	-	474	110	525	-	373	-	1,482
Miscellaneous	5	-	-	308	866	-	-	-	-	1,179
<b>Non energy use</b>	-	<b>48</b>	-	<b>7,490</b>	<b>426</b>	-	-	-	-	<b>7,964</b>

(1) Includes all manufactured solid fuels, benzole, tars, coke oven gas and blast furnace gas.

(2) Includes colliery methane.

(3) Includes geothermal and solar heat.

(4) Stock fall (+), stock rise (-).

(5) Primary supply minus primary demand.

(6) See paragraphs 5.42 regarding electricity use in transport and 6.66 regarding renewables use in transport.

(7) Back-flows from the petrochemical industry.

**A 4.2 FUELS DATA**

The fuels data are taken from DUKES - the Digest of UK Energy Statistics (BEIS, 2018), so the fuel definitions and the source categories used in the NAEI reflect those in DUKES.

IPCC Guidelines (IPCC, 2006) lists fuels that should be considered when reporting emissions. **Table A 4.2.1** lists the fuels that are used in the GHGI (based on DUKES) and indicates how they relate to the fuels listed in the IPCC Guidelines. In most cases the mapping is obvious but there are a few cases where some explanation is required.

**Table A 4.2.1 Mapping of fuels used in IPCC and the NAEI**

	IPCC	NAEI
Category	Subcategory	Subcategory
Liquid	Motor Gasoline	Petrol
	Aviation Gasoline	Aviation Spirit
	Jet Kerosene	Aviation Turbine Fuel <sup>1</sup> (ATF)
	Other Kerosene	Burning Oil
	Gas/Diesel Oil	Gas Oil/ DERV
	Residual Fuel Oil	Fuel Oil
	Orimulsion	Orimulsion
	Liquefied Petroleum Gases	Liquefied Petroleum Gas (LPG)
	Naphtha	Naphtha
	Petroleum Coke	Petroleum Coke
	Refinery Gas	Other Petroleum Gas (OPG)
	Other Oil: Other Petroleum Products	Refinery Miscellaneous
	Lubricants	Lubricants
Solid	Anthracite	Anthracite
	Coking Coal	Coal <sup>2</sup>
	Other Bituminous Coal	Coal
		Slurry <sup>3</sup>
	Coke Oven Coke	Coke
	Patent Fuel	Solid Smokeless Fuel (SSF)
	Coke Oven Gas	Coke Oven Gas
	Blast Furnace Gas	Blast Furnace Gas

	IPCC	NAEI
Category	Subcategory	Subcategory
Gas	Natural Gas	Natural Gas
		Sour Gas <sup>4</sup>
		Colliery Methane <sup>5</sup>
Other Fuels	Municipal Solid Waste	Municipal Solid Waste
	Industrial Waste: Scrap Tyres	Scrap Tyres
	Waste Oils	Waste Oil
Peat	Peat	Peat
Biomass	Wood/Wood Waste	Wood
	Other Primary Solid Biomass	Straw
		Poultry Litter, Meat & bone meal
	Landfill Gas	Landfill Gas
	Sludge Gas	Sewage Gas
	Charcoal	Charcoal
	Other liquid biofuels	Liquid Biofuels
Other biogas	Biogas	

1 Includes fuel that is correctly termed jet gasoline.

2 Used in coke ovens.

3 Coal-water slurry used in some power stations

4 Unrefined natural gas used on offshore platforms and some power stations

5 IPCC Guidelines (IPCC, 2006) specifies coal seam methane is included in Natural Gas.

#### **A 4.2.1 Reallocations of energy data and differences from UK energy statistics**

The main source of energy consumption data used in the UK inventory is the Digest of UK Energy Statistics (BEIS, 2018). This annual publication gives detailed sectoral energy consumption broken down by fuel type, and covering the entire time period covered by the inventory. In many cases, these data are used directly in the inventory without modification. However, there are instances where the activity data used in the inventory are not based directly on DUKES data, but where alternative data sources provide supplementary data to inform energy use and emission estimates. In general, the UK inventory totals by fuel are kept consistent with the DUKES national totals for each fuel. There are some exceptions where the UK total may be different to that presented in DUKES due to different scopes and reporting requirements.

The reasons for any deviations from use of DUKES data in the inventory are discussed within the source category methodological descriptions in Section 3 of the main report. The main reasons for reallocations or modifications are:

- To account for differences in geographical scope (e.g. to account for energy use in OTs)
- To make best use of EU ETS data (this data is only used indirectly in producing UK energy statistics)
- To utilise other operator reported data (e.g. direct to the Inventory Agency, or via the various regulator's inventories).

- When bottom-up models are available providing fuel consumption data on a more granular level and are considered to be a higher quality estimate by the Inventory Agency.

The fuel reconciliation tables (**Table A 4.2.2 - Table A 4.2.6**) show how the deviations are applied and how the energy data for the major fuels in the UK inventory are reconciled against the energy demand data from DUKES. The tables show:

1. where fuels are re-allocated between sectors, but the overall annual fuel consumption across all UK sectors is kept consistent with the data in DUKES, and
2. where deviations are made to DUKES figures for total UK consumption of a given fuel, and in which source categories these deviations are made.

The Inventory Agency presents data below for the fuel allocations for coal, natural gas, fuel oil, gas oil (including DERV) and petroleum gases (LPG, OPG) for the latest inventory year. Together these fuels constitute the majority of the UK inventory 1A sector emissions total.

Deviations to the energy balance are made in consultation with the authors of the energy statistics.

**A 4.2.1.1 Coal**

Total industrial coal use within the GHG inventory is consistent with the DUKES total and in the most part, coal use at the sectoral level is consistent with the DUKES data. However, there is an apparent step change in the amount of fuel allocated to other industries within DUKES between 1999 and 2000. In addition, between 1997 and 1999, the total coal use allocated to 1A2f is less than the independent estimates for cement and lime production used within the inventory. Cement and lime production would fall into the 1A2f category for IPCC reporting. Therefore, Inventory Agency estimates have been made to construct a consistent time series for coal use. **Table A 4.2.2** below compares inventory estimates with DUKES estimates for the latest inventory year

**Table A 4.2.2 Fuel reconciliation - coal use in the latest year (Mtonnes)**

DUKES Category	DUKES	GHGI	Difference	GHGI category	CRF	Comment
Major power producers	8.702	8.702	0.000	Power stations	1A1a	
Blast furnaces	1.301	1.301	0.000	Blast furnaces	2C1	
Coal extraction	0.000	0.000	0.000	Collieries - combustion	1A1c	
Autogenerators	0.022					
		0.013		Autogenerators	1A2b	
		0.010		Autogeneration - exported to grid	1A2b	
	0.022	0.022		<b>Subtotal</b>		

DUKES Category	DUKES	GHGI	Difference	GHGI category	CRF	Comment
Patent fuel manufacture etc.	0.207	0.207	0.000	Solid smokeless fuel production	1B1b	
Coke manufacture	1.888	1.826	0.062	Coke Production	1B1b	GHGI uses EU ETS data which suggests fewer emissions than in DUKES
Industry total	1.344					
		0.033		Iron & steel - combustion plant	1A2a	
		0.033		Non-Ferrous Metal (combustion)	1A2b	
		0.069		Chemicals	1A2c	
		0.111		Pulp, Paper and Print	1A2d	
		0.078		Food & drink, tobacco	1A2e	
		0.580		Other industrial combustion	1A2g	
		0.580		Cement production - combustion	1A2f	Operator's data used in preference in the GHGI
		0.051		Lime production - non decarbonising	1A2f	EU ETS used in preference in the GHGI
	<b>1.344</b>	<b>1.533</b>	<b>-0.188</b>	<b>Subtotal</b>		
Rail	0.015	0.015	0.000	Rail	1A3c	

DUKES Category	DUKES	GHGI	Difference	GHGI category	CRF	Comment
<b>Domestic - anthracite</b>	<b>0.193</b>	<b>0.193</b>	<b>0.000</b>	<b>Domestic combustion - anthracite</b>	<b>1A4b</b>	
Domestic - coal	0.342	0.342		Domestic combustion - UK	1A4b	
		0.000		Domestic combustion - crown dependencies	1A4b	
	<b>0.342</b>	<b>0.342</b>	<b>0.000</b>	<b>Subtotal</b>		
<b>Agriculture</b>	<b>0.000</b>	<b>0.000</b>	<b>0.000</b>	<b>Agriculture - stationary combustion</b>	<b>1A4c</b>	
Commercial	0.005					
Miscellaneous	0.007	0.012	0.000	Miscellaneous industrial/commercial combustion	1A4a	
	<b>0.012</b>	<b>0.012</b>	<b>0.000</b>	<b>Subtotal</b>		
<b>Public administration</b>	<b>0.156</b>	<b>0.030</b>	<b>0.126</b>	<b>Public sector combustion</b>	<b>1A4a</b>	<b>Differences offset the deviations in the industrial and coke manufacture sectors</b>
<b>TOTAL</b>	<b>14.183</b>	<b>14.183</b>	<b>0.000</b>			

Notes: Rows are shaded to help illustrate reconciliation between sectors.

#### **A 4.2.1.2 Natural Gas**

Data for natural gas use is largely taken directly from DUKES and the national total is consistent between the inventory and the energy statistics, other than a small additional use of natural gas at a number of (international) gas pipeline inter-connectors and also on the Isle of Man (IoM) which is added to the inventory, as natural gas use on IoM is not included in DUKES demand totals. Operator estimates for ammonia production (both fuel and feedstock), and ETS data for gas separation plant lead to minor reallocations of the DUKES data, these are summarised below in **Table A 4.2.3**. In addition, the NAEI model doesn't include any accounting for losses compared to DUKES tables.

**Table A 4.2.3 Fuel reconciliation – natural gas use in the latest year (Mtherms)**

DUKES Category	DUKES	GHGI	Difference	GHGI Category	CRF	Comment
Major power producers	8790	8822	-33	Power stations	1A1a	Isle of Man gas use not offset by DUKES
Heat generation	0		0			
Autogenerators	954					
		398		Autogenerators	1A2f	
		300		Autogeneration - exported to grid	1A2f	
		0		Railways - stationary combustion	1A4a	
	954	699	255	Subtotal		Offset against refineries
Coal extraction	2	2	0	Collieries - combustion	1A1c	
Oil and gas extraction	1684					
		1186		Upstream oil production - fuel combustion	1A1c	
		499		Upstream gas production - fuel combustion	1A1c	
	1684	1684	0	Subtotal		
Petroleum refineries	200	455	-255	Refineries - combustion	1A1b	Offset against autogeneration
Blast furnaces	10	0	10	Blast furnaces	1A2a	
Other energy industries	213					
		242		Gas production	1A1c	
		0		Nuclear fuel production	1A1c	

DUKES Category	DUKES	GHGI	Difference	GHGI Category	CRF	Comment
	213	242	-30	Subtotal		From discussion with BEIS, interconnector sites are missing from DUKES, so we add the estimates for these across the time series, to 15_19.
Non-Ferrous Metal	112	112		Non-Ferrous Metal (combustion)	1A2b	
Chemicals	939	939		Chemicals (combustion)	1A2c	
Pulp, Paper and Print	213	213		Pulp, Paper and Print (combustion)	1A2d	
Food & drink, tobacco	697	697		Food & drink, tobacco (combustion)	1A2e	
		118		Ammonia production - combustion	1A2c	Operator's data
All other industry (minus iron and steel)	1852	1542		Other industrial combustion	1A2f	
Other industrial combustion (colliery methane)	0			Other industrial combustion (colliery methane)		
		47		Lime production - non decarbonising	1A2f	EU ETS
		3		Cement production - combustion	1A2f	Operator's data
<b>All industry except iron and steel</b>	<b>3814</b>	<b>3671</b>	<b>143</b>	<b>Subtotal</b>		<b>Offset against NEU</b>
Non-energy use	169	102		Non-energy use (stored)		
		210		Ammonia production - feedstock use of gas	2B2	
	<b>169</b>	<b>312</b>	<b>-143</b>			<b>Offset against industry</b>
Iron and steel	147	147	0	Iron and steel - combustion plant	1A2a	
Domestic	10135	10145	-10	Domestic combustion	1A4b	Isle of Man gas use not offset against DUKES
Public administration	1409	1409	0	Public sector combustion	1A4a	

DUKES Category	DUKES	GHGI	Difference	GHGI Category	CRF	Comment
Commercial	1705					
Miscellaneous	344					
		2049		Miscellaneous industrial/commercial combustion	1A4a	
	<b>2049</b>	<b>2049</b>	<b>0</b>	<b>Subtotal</b>		
Agriculture	44	44	0	Agriculture - stationary combustion	1A4c	
Autogenerators - colliery methane	16					
Coal extraction - colliery methane	0					
		16		Collieries - combustion (colliery methane)	1A1c	
	<b>60</b>	<b>60</b>	<b>0</b>	<b>Subtotal</b>		
Losses	230		230			The NAEI model doesn't include any accounting for losses compared to DUKES long term trends
<b>Total</b>	<b>29867</b>	<b>29709</b>	<b>158</b>			

Notes: Rows are shaded to help illustrate reconciliation between sectors.

1 Mtherm = 105.51 TJ

#### A 4.2.1.3 Fuel Oil

For shipping, a major research project was completed in 2017 and the results were incorporated into the 2018 submission. The estimated total fuel oil consumption derived from this research is greater than as reported for shipping in DUKES, and any deviations from the national navigation sector are considered additional and are not reconciled elsewhere in the inventory. Additional sectoral deviations are also made to account for known use of fuel oil in power stations, and the Crown Dependencies.

**Table A 4.2.4 Fuel reconciliation – Fuel oils use in latest year (Mtonnes)**

DUKES Category	DUKES	GHGI	Difference	GHGI category	CRF	Comment
Major power producers	0.101					

DUKES Category	DUKES	GHGI	Difference	GHGI category	CRF	Comment
		0.110		Power Stations - UK	1A1a	EUETS data
		0.015		Power Stations - crown dependencies	1A1a	Local data sets
	<b>0.101</b>	<b>0.124</b>	<b>-0.024</b>	<b>Subtotal</b>		
<b>Autogenerators</b>	<b>0.015</b>		<b>0.015</b>			
Industry	0.240					
		0.011		Iron and steel - combustion plant	1A2a	Reduced to offset increase in 1A1a
		0.000		Non-Ferrous Metal (combustion)	1A2b	
		0.026		Chemicals (combustion)	1A2c	
		0.001		Pulp, Paper and Print (combustion)	1A2d	
		0.052		Food & drink, tobacco (combustion)	1A2e	
		0.205		Other industrial combustion	1A2g	Reduced to offset increase in 1A1a
		0.000		Cement production - combustion	1A2f	Operator's data
	<b>0.240</b>	<b>0.296</b>	<b>-0.055</b>	<b>Subtotal</b>		Reduced to offset increase in 1A1a
<b>Petroleum refineries</b>	<b>0.208</b>	<b>0.208</b>	<b>0.000</b>	<b>Refineries - combustion</b>	<b>1A1b</b>	
<b>Oil and gas extraction</b>	<b>0.064</b>					
<b>Energy sector</b>	<b>0.628</b>	<b>0.628</b>	<b>0.000</b>	<b>Subtotal</b>		
Agriculture	0.017	0.017	0.000	Agriculture - stationary combustion	1A4c	
<b>Domestic</b>	<b>0.000</b>	<b>0.000</b>	<b>0.000</b>	<b>Domestic combustion</b>	<b>1A4b</b>	
<b>Commercial</b>	<b>0.062</b>					
<b>Miscellaneous</b>	<b>0.013</b>					
		0.075		Miscellaneous combustion - UK	1A4a	
		0.000		Miscellaneous combustion - crown dependencies	1A4a	Local data sets

DUKES Category	DUKES	GHGI	Difference	GHGI category	CRF	Comment
	0.075	0.075	0.000	<b>Commercial and Miscellaneous Total</b>		
<b>Public administration</b>	<b>0.025</b>	<b>0.025</b>	<b>0.000</b>	<b>Public sector combustion</b>	<b>1A4a</b>	
National navigation	0.070	0.167		Shipping - coastal	1A3d	Bottom-up AIS approach estimates greater fuel oil and is considered additional to DUKES.
		0.010		Fishing vessels	1A4c	
		0.000		Shipping between UK and CDs	1A3d	
		0.006		Shipping between UK and Gibraltar	1A3d	
		0.000		Shipping between UK and Bermuda	1A3d	
		0.005		Shipping between UK and OTs (excl. Gib and Bermuda)	1A3d	
Marine bunkers	0.777	0.766		Shipping - international IPCC definition		
<b>Shipping</b>	<b>0.777</b>	<b>0.954</b>	<b>-0.177</b>	<b>Subtotal</b>		<b>Deviation due to the implementation of the bottom-up shipping inventory, which is known to deviate from DUKES</b>
<b>TOTAL</b>	<b>1.523</b>	<b>1.700</b>	<b>-0.177</b>			

Notes: Rows are shaded to help illustrate reconciliation between sectors.

#### **A 4.2.1.4 Gas Oil**

Gas oil is used in both off-road transport and machinery diesel engines, and as a fuel for stationary combustion. The varied use of this fuel and the complexity of the supply chain complicates the means of allocating consumption across the wide range of sectors that use the fuel in the inventory. DUKES provides a breakdown of gas oil consumption in different industry and other sectors, but the data resolution in DUKES does not distinguish between use of the fuel for stationary combustion and off-road machinery, a distinction which is necessary for the inventory.

The GHGI estimates consumption of gas oil and emissions for off-road machinery using a bottom-up method based on estimates of population and usage of different types of machinery. However, this has led to a situation where the total amount of gas oil consumption across sectors exceeds that which is available as given in DUKES. Therefore, consumption figures, mainly for stationary combustion in industry sectors, have had to be adjusted to obtain a total fuel balance.

The problem is extended when new sources of gas oil consumption are found. For example, the recent development of an inventory for the UK's inland waterways requires the allocation of gas oil to this sector (Walker et al, 2011). During the process of compiling the inland waterways inventory, it became clear that not all vessels with diesel engines use gas oil, but use road diesel and that this may also be the case for other off-road machinery sources, especially those that consume small amounts of fuel on an irregular basis, e.g. for private or recreational use rather than commercial use. There are also inconsistencies in terminology used to define types of fuel; it became apparent that the terms "gas oil", "red diesel" and "diesel" are used interchangeably by fuel suppliers and consumers and this confuses the situation when considering fuel allocations across different sectors.

In light of this, Task 5 of the 2011 UK GHG Inventory Improvement Programme aimed to address the allocation of gas oil and DERV in the GHGI (Murrells et al., 2011). The methodology outlined in Murrells et al. (2011) was used in the compilation of the 2011 inventory, and is summarised here. The same approach has been used in this inventory.

Several fuel suppliers and experts in the petroleum industry and at the Department for Transport were consulted to understand terminologies used, the physical differences between gas oil and DERV, and to gauge opinions on what determines where the fuels are mainly used where it is possible to use either gas oil or DERV. The study concluded that while the majority of agricultural and industrial machinery will be using low tax gas oil (red diesel), a small amount of DERV is likely to be used by private recreational boat users and by equipment with small engines used for private or small-scale commercial use on an irregular basis and the gas oil fuel supply infrastructure makes it more convenient to use DERV.

The study provided new estimates of the amount of DERV and petrol consumed by non-road transport sources with small internal combustion engines. This reduces the overestimation of gas oil consumption and relieves the pressure on how much gas oil consumption by other sources has to be adjusted to match the total amount available as given in DUKES.

The study also considered the allocation of gas oil given in DUKES to different industry and other sectors and how these can be mapped to inventory reporting categories. The detailed bottom-up method is used to estimate gas oil consumption by different off-road machinery and marine vessel types. Independent sources were used to estimate gas oil used by the rail sector while data provided by industrial sites reporting under emission trading schemes (EU ETS) were used to derive an allocation of gas oil consumption by stationary combustion sources in different industry, commercial and other sectors. Also, the UK energy statistics now include an allocation of gas oil for consumption by the oil and gas sector, but since only a partial time series was made available, the study included making estimates of gas oil for this category back to 1990.

A method of re-allocation was developed using an over-arching condition that the total sum of gas oil consumption across all sectors was consistent with the total consumption figures given in DUKES across all years. The method allowed the consumption estimates for industrial off-road machinery and stationary combustion by industry, commercial and public sector activities to vary in order to align the total consumption estimates with DUKES on the basis that the estimates for these sources are the most uncertain.

As with fuel oil, the introduction of the results of a major research project into the shipping sector in the 2018 submission, whereby Automatic Identification System (AIS) data was used to calculate shipping movements along the coast of the UK and the Crown Dependencies, however suggested that gas oil consumption reported by DUKES is an underestimate. As a result, total gas oil use

(not including DERV) deviates from DUKES as any further consumption in the national navigation sector are considered additional to DUKES and are not reconciled elsewhere in the inventory.

**Table A 4.2.5** below summarised the DUKES and GHGI allocations for the latest inventory year.

**Table A 4.2.5 Fuel reconciliation – Gas oil use in latest year (Mtonnes)**

DUKES Category	DUKES	GHGI	Difference	GHGI Category	CRF	Comment
Petroleum refineries	0.000	0.000	0.000	Refineries - combustion	1A1b	
Major power producers	0.039	0.039		Power stations - UK		
		0.002		Power stations - crown dependencies		
	0.039	0.041	-0.002	Subtotal		
Autogenerators	0.056		0.056			
Industry	1.740					
		0.002		Iron and steel - combustion plant	1A2b	
		0.000		Non-Ferrous Metal	1A2c	
		0.003		Chemicals	1A2d	
		0.001		Pulp, Paper and Print	1A2e	
		0.001		Food & drink, tobacco		
		0.047		Other industrial combustion - UK	1A2g	
		0.000		Other industrial combustion - crown dependencies	1A2g	
		0.008		Cement production - combustion	1A2f	Operator's data

DUKES Category	DUKES	GHGI	Difference	GHGI Category	CRF	Comment
		0.177		Aircraft - support vehicles	1A3e	Inventory agency estimates
		1.526		Industrial off-road mobile machinery	1A2g	Inventory agency estimates
	<b>1.740</b>	<b>1.766</b>	<b>-0.026</b>	<b>Subtotal</b>		
Commercial	0.378					
Miscellaneous	0.271	0.029		Miscellaneous combustion - UK	1A4a	Reduced to offset higher consumption elsewhere
		0.008		Miscellaneous combustion - crown dependencies	1A4a	Reduced to offset higher consumption elsewhere
	<b>0.271</b>	<b>0.037</b>	<b>0.235</b>	<b>Subtotal</b>		
<b>Public administration</b>	<b>0.299</b>	<b>0.019</b>	<b>0.280</b>	<b>Public sector combustion</b>	<b>1A4a</b>	Reduced to offset higher consumption elsewhere
<b>Agriculture</b>	<b>0.328</b>	<b>1.250</b>	<b>-0.922</b>	<b>Agriculture - mobile machinery</b>	<b>1A4c</b>	<b>Inventory Agency estimates which far exceed DUKES</b>
<b>Rail</b>	<b>0.610</b>	<b>0.608</b>	<b>0.002</b>	<b>Railways</b>	<b>1A3c</b>	<b>Reduced to offset higher consumption elsewhere</b>
<b>Subtotal (of all above)</b>	<b>3.721</b>	<b>3.721</b>	<b>0.000</b>			
Oil & gas extraction	0.599			Upstream Gas Production - fuel combustion		
		0.059		Upstream Oil Production - fuel combustion	1A1c	
		0.539			1A1c	
	<b>0.599</b>	<b>0.599</b>	<b>0.000</b>	<b>Subtotal</b>		
National navigation	0.853	1.217		Shipping - coastal		Bottom-up AIS approach estimates greater fuel oil and is considered additional to DUKES.
		0.175		Fishing vessels	1A3d	

DUKES Category	DUKES	GHGI	Difference	GHGI Category	CRF	Comment
		0.006		Shipping between Uks and CDs	1A3d	
		0.137		Shipping - naval		
		0.046		Motorboats / workboats	1A5b	
		0.002		Inland goods-carrying vessels	1A3d	
Marine bunkers	<b>1.653</b>	1.653		Shipping - international IPCC definition	1A3d	Bottom-up AIS approach estimates greater fuel oil and is considered additional to DUKES.
<b>Shipping total</b>	<b>2.506</b>	<b>3.236</b>	<b>-0.730</b>	<b>Subtotal</b>		<b>Deviation due to the implementation of the bottom-up shipping inventory, which is known to deviate from DUKES</b>
Road	24.911					
		0.011		House and garden machinery - DERV	1A4b	
		0.326		Industrial off-road mobile machinery - DERV	1A2gvii	
		0.002		Sailing boats with auxiliary engines	1A3d	
		0.112		Motorboats / workboats (e.g. canal boats, dredgers, service boats, tourist boats, river boats)	1A3d	
		24.413		Road transport - UK	1A3b	
		0.046		Road transport -	1A3b	

DUKES Category	DUKES	GHGI	Difference	GHGI Category	CRF	Comment
	24.911	24.911	0.000	Subtotal		
Domestic	0.144	0.137	0.137	Domestic combustion - UK	1A4b	0.000
		0.007	0.007	Domestic combustion - crown dependencies	1A4b	
	0.144	0.144	0.000	Subtotal		
<b>Total</b>	<b>31.880</b>	<b>32.610</b>	<b>-0.730</b>			

Notes: Rows are shaded to help illustrate reconciliation between sectors

#### A 4.2.1.5 Petroleum gases

For petroleum gases (LPG, OPG), the total fuel use in the inventory is greater than the national statistics in several years, to reflect information from other sources (such as EU ETS data) that indicate potential under-reports in the UK energy statistics. These modifications to the energy balance are set out in **Table A 4.2.6**. They mostly relate to refineries, use of feedstock as fuel in the petrochemicals sector, and fuel use for upstream oil and gas production.

**Table A 4.2.6 Fuel reconciliation – Use of petroleum gases in the latest year (Mtherms)**

DUKES Category	DUKES	GHGI	Difference	GHGI Category	CRF	Comment
Petroleum refineries, other gases	943	1115		Refineries, OPG	1A1b	
Autogenerators, other gases	123					

DUKES Category	DUKES	GHGI	Difference	GHGI Category	CRF	Comment
<b>Sub-total</b>	<b>1066</b>	<b>1115</b>	<b>-49</b>			<b>OPG total inferred from EU ETS dataset. This is larger than the reported DUKES value</b>
<b>Petroleum refineries, propane</b>	<b>7</b>	<b>7</b>	<b>0</b>	<b>Refineries, LPG</b>	<b>1A1b</b>	
<b>Iron &amp; steel, propane</b>	<b>0</b>	<b>0</b>	<b>0</b>	<b>Iron &amp; steel combustion, LPG</b>	<b>1A1b</b>	
Industry, propane	133					
Industry, butane	37					
Agriculture, propane	40					
Agriculture, butane	0					
Commercial, propane	162					
Commercial, butane	1					
Public administration, propane	7					
		379		Industrial combustion, LPG - UK	1A2g	
		0		Industrial combustion, LPG - crown dependencies	1A2g	
<b>Sub-total</b>	<b>380</b>	<b>379</b>	<b>1</b>			
<b>Industry, other gases</b>	<b>0</b>	<b>523</b>	<b>-523</b>	<b>Chemicals, OPG</b>	<b>2B8g</b>	<b>EU ETS higher than DUKES</b>

DUKES Category	DUKES	GHGI	Difference	GHGI Category	CRF	Comment
Road, propane	32	32	0	Road transport - all vehicles LPG use	1A3b	
Domestic, propane	94	88		Domestic combustion, LPG - UK	1A4b	
Domestic, butane	0	5		Domestic combustion, LPG - crown dependencies	1A4b	
<b>Sub-total</b>	<b>94</b>	<b>93</b>	<b>1</b>			
(excluded from DUKES)		145		Gas separation plant, OPG	1A1c	EEMS. Outside scope of DUKES
(excluded from DUKES)		7		Gas separation plant, LPG	1A1c	EEMS. Outside scope of DUKES
<b>Total</b>	<b>1579</b>	<b>2301</b>	<b>-722</b>			

Notes: Sequences of shaded rows indicate categories which are grouped for purposes of data reconciliation, and should be considered together.

1 Mtherm = 105.51 TJ

# **ANNEX 5: Additional Information to be Considered as Part of the Annual Inventory Submission and the Supplementary Information Required Under Article 7, paragraph 1, of the Kyoto Protocol Other Useful Reference Information.**

## **A 5.1 ANNUAL INVENTORY SUBMISSION**

No additional information.

## **A 5.2 SUPPLEMENTARY INFORMATION UNDER ARTICLE 7, PARAGRAPH 1**

No additional information.



# ANNEX 6: Comparison of Emission Estimates Using Atmospheric Observations

This Annex discusses the verification of the UK estimates of the Kyoto Gases.

## A 6.1 MODELLING APPROACH USED FOR COMPARISON WITH THE UK GHGI

Comparison of the UK GHGI with emission estimates made using atmospheric observations is considered to be best practice by the UNFCCC as it allows for an independent assessment of the GHG emissions from the UK using a comprehensively different approach. Significant differences in the emissions estimated using the two methods are a means of identifying areas worthy of further investigation, for example as occurred with a re-assessment of the emissions of HFC-134a from mobile air conditioning.

In order to provide a comparison with the UK Greenhouse Gas Inventory (GHGI), BEIS (Department of Business, Energy and Industrial Strategy) supported the establishment and maintenance of a high-quality remote observation station at Mace Head (MHD) on the west coast of Ireland. The station reports high-frequency concentrations of the key greenhouse gases and is under the supervision of Prof. Simon O'Doherty of the University of Bristol (O'Doherty *et al.* 2004, 2014). BEIS extended the measurement programme in 2012 with three new tall tower stations across the UK (collectively called the UK DECC (Deriving Emissions linked to Climate Change) network): Tacolneston (TAC) near Norwich; Ridge Hill (RGL) near Hereford; and Tall Tower Angus (TTA) near Dundee, Scotland which relocated to Bilsdale (BSD) in North Yorkshire in Sept 2015. Methane (CH<sub>4</sub>), carbon dioxide (CO<sub>2</sub>), nitrous oxide (N<sub>2</sub>O) and sulphur hexafluoride (SF<sub>6</sub>) are measured at all stations across the UK DECC network. The hydrofluorocarbons (HFCs) and perfluorocarbons (PFCs) are measured at MHD and TAC, and nitrogen trifluoride (NF<sub>3</sub>) is only measured at MHD.


The Met Office, under contract to BEIS, employs the Lagrangian dispersion model NAME (Numerical Atmospheric dispersion Modelling Environment) (Ryall *et al.* 1998, Jones *et al.* 2007) driven by three-dimensional modelled meteorology to interpret the observations. NAME determines the history of the air arriving at Mace Head at the time of each observation. By estimating the underlying background trend (Northern Hemisphere mid-latitude atmospheric concentrations where the short-term impact of regional pollution has been removed from the data) and by modelling where the air has passed over on route to the observation stations on a regional scale, estimates of UK emissions are made. A methodology called Inversion Technique for Emission Modelling (InTEM) has been developed that uses a minimisation technique, Non-Negative Least Squares (NNLS) (Lawson and Hanson, 1974) to determine the emission map that most accurately reproduces the observations (Manning *et al.* 2003, 2011 and Arnold *et al.* 2017).

For each reported gas the Northern Hemisphere background trend and the UK emission estimates are presented. InTEM estimates using only Mace Head (MHD) data are presented along with the estimates made using the full UK DECC network. In 2014 two additional tower sites, Heathfield (HFD) south of London and Bilsdale (BSD) in North Yorkshire, were established

through the NERC GAUGE (Global And UK Greenhouse gas Emissions) programme and have been included in the emission estimates. HFD is currently operated by NPL, both stations are now in the UK DECC Network. For CH<sub>4</sub> only, a further dataset from the tall tower station at Cabauw in The Netherlands has been included courtesy of ECN, The Netherlands. When only MHD data are used a three-year inversion window is assumed (an inversion is performed for a three year period and then the period is incremented by one-year e.g. 1989 – 1991, 1990 – 1992 etc., from which a median for each year is estimated), however with the additional data from the other stations the inversion time window has been shortened to one year or smaller (2 months for CH<sub>4</sub>, N<sub>2</sub>O) depending on data. The geographical spread of the UK DECC (and other stations) network allows the spatial distribution of the emissions across the UK to be better constrained within InTEM. The InTEM estimates of UK emissions using the atmospheric observations are compared to the reported GHGI estimates.


## A 6.2 METHANE

**Figure A 6.1** Monthly Northern Hemisphere trend in methane estimated from Mace Head observations (blue line). Red line denotes the annual trend. Data under grey shading are not yet ratified.



**Figure A 6.1** shows the Northern Hemisphere background atmospheric concentration of methane from 1990 onwards. As with all of the background plots for the different gases, it shows how the overall atmospheric concentration of the gas in question is changing in the atmosphere as a result of global emissions and atmospheric loss processes. For CH<sub>4</sub>, the underlying background trend is positive but there is strong year-to-year variability and a strong seasonal cycle. The growth rate since 2006 has been unusually and consistently positive (average +7.2 ppb/yr).

**Figure A 6.2** Verification of the UK emission inventory estimates for methane in Gg yr<sup>-1</sup> from 1990. GHGI estimates are shown in black. InTEM (MHD+CBW, 3-year) estimates are shown in blue (±1σ). InTEM (DECC, 12-mth) estimates are shown in orange (±1σ).




The emission estimates made for the UK using the InTEM methodology are compared to the GHGI emission estimates for the period 1990 onwards. It is important to note that although the UK methane emissions are estimated to have fallen over the last 28 years, the global atmospheric concentration of methane has increased, indicating that global emissions of methane are still outperforming the global natural removal of methane from the atmosphere.

Methane has a natural (biogenic) component and it is estimated that 22% of the annual global emission is released from wetlands (Nilsson *et al.* 2001). Usually natural emissions are strongly dependent on a range of meteorological factors such as temperature and also growth and decay cycles. Such non-uniform emissions will add to the uncertainties in the modelling, although in North West Europe the natural emissions are thought to be small compared to the anthropogenic emissions (<5%, Bergamaschi *et al.* 2005). Due to the relatively strong local (within 20km) influence of biogenic emissions at some of the stations the influence of observations taken when local emissions are thought to be significant (low boundary layer heights, low wind speeds, stable atmospheres) has been reduced within InTEM by removing all observations taken during such meteorological conditions.

The GHGI trend is monotonically downwards whereas the InTEM estimates, after an initial fall, shows little change (**Figure A 6.2**). The InTEM estimates using all of the available observations (MHD+CBW+DECC+GAUGE) are consistent with the MHD+CBW estimates and agree with the reported GHGI. The InTEM 2-month estimates using the full DECC Network (**Figure A 6.3**) do not show any strong seasonal cycle in UK methane emissions.


**Figure A 6.3** Verification of the UK emission inventory estimates for methane in Gg yr<sup>-1</sup> from 2013. GHGI estimates are shown in black. InTEM (DECC 2-month) estimates are shown in blue ( $\pm 1\sigma$ ).



**A 6.3 NITROUS OXIDE**


**Figure A 6.4** shows the Northern Hemisphere background atmospheric concentration of nitrous oxide (N<sub>2</sub>O) from 1990 onwards. The background trend is monotonic and positive. In 2018, the background increased by 1.1 ppb.

**Figure A 6.4** Monthly Northern Hemisphere trend in nitrous oxide estimated from Mace Head observations (blue line). Red line denotes the annual trend. Data under grey shading are not yet ratified.



The main activities in Europe resulting in the release of nitrous oxide are; agricultural practices resulting in emissions from soils (~60%), chemical industry (~20%) and combustion (~15%) (UNFCCC 1998 figures). The amount emitted from soils has significant uncertainty and has a diurnal and seasonal release cycle. It is driven by the availability of nitrogen, temperature and the soil moisture content.


**Figure A 6.5** Verification of the UK emission inventory estimates for nitrous oxide in Gg yr<sup>-1</sup> from 1990. GHGI estimates are shown in black. InTEM (MHD-only, 3-year) estimates are shown in blue ( $\pm 1\sigma$ ). InTEM (DECC, 12-mth) estimates are shown in orange ( $\pm 1\sigma$ ).



**Figure A 6.5** shows the InTEM and GHGI emission estimates for nitrous oxide for the period 1990 onwards for the UK. The annual InTEM estimates are similar to the GHGI estimates, with both showing declining UK totals. The InTEM estimates are higher than the GHGI post-1998 although within the  $1\sigma$  uncertainty. The GHGI estimates show a sharp decline (40 Gg) between 1998 and 1999 in line with the introduction of clean technology at an adipic acid plant in Wilton, north east England. It is estimated to have cut its emissions of N<sub>2</sub>O by 90%, from 46 thousand tonne yr<sup>-1</sup> to around 6 thousand tonne yr<sup>-1</sup>. The InTEM estimates show a more gradual decline over this period (expected due to the long 3-year inversion time periods) but the overall reduction is similar. The estimates using all available observations are very similar to the MHD-only estimates. Post-2012 the InTEM estimates are notably higher than the GHGI estimates. The improved network of observations from 2013 onwards allows a strong seasonal cycle in emissions to be highlighted, Figure A 6.6 shows there is a peak in UK emissions in spring-summer and a minimum in the winter months. This is aligned with the traditional fertiliser application period.

The nature of the nitrous oxide emissions challenges the InTEM assumption of uniformity of release both in time and space. Also the point of release to the atmosphere may not be coincident with the activity generating the nitrous oxide e.g. the nitrous oxide may be transported from its source, for example by rivers, prior to its release to the atmosphere.


**Figure A 6.6** Verification of the UK emission inventory estimates for nitrous oxide in Gg yr<sup>-1</sup> from 2013. GHGI estimates are shown in black. InTEM (DECC 2-month) estimates are shown in blue ( $\pm 1\sigma$ ).



**A 6.4 HYDROFLUOROCARBONS**


**A 6.4.1 HFC-134a**

**Figure A 6.7** Monthly Northern Hemisphere trend in HFC-134a estimated from Mace Head observations (blue line). Red line denotes the annual trend. Data under grey shading are not yet ratified.



**Figure A 6.7** shows the Northern Hemisphere background atmospheric concentration of HFC-134a from 1994 onwards. The background trend is monotonic and positive, in 2018 the background increased by 6.2 ppt.


**Figure A 6.8** Verification of the UK emission inventory estimates for HFC-134a in Gg yr<sup>-1</sup> from 1990. GHGI estimates are shown in black. InTEM (MHD-only, 3-year) estimates are shown in blue ( $\pm 1\sigma$ ). InTEM (DECC, 12-mth) estimates are shown in orange ( $\pm 1\sigma$ ).



**Figure A 6.8** shows the InTEM and GHGI emission estimates for the UK for HFC-134a for the period 1990 onwards. The GHGI shows a stronger increase in emission compared to the InTEM estimates. The InTEM estimates have risen at about 60% of the rate of the GHGI. Throughout the time series the trend agreement between the GHGI and InTEM is good but the InTEM estimates are consistently about 50-60% of the GHGI estimates, a difference well outside both uncertainty ranges. A similar InTEM result is obtained when the TAC observations are included (from 2013).


### A 6.4.2 HFC-152a

**Figure A 6.9** Monthly Northern Hemisphere trend in HFC-152a estimated from Mace Head observations (blue line). Red line denotes the annual trend. Data under grey shading are not yet ratified.



**Figure A 6.9** shows the Northern Hemisphere background atmospheric concentration of HFC-152a from 1995 onwards. The background trend shows a strong rise from the mid-1990s until 2008, then a much reduced annual increase until 2012. From 2012 onwards a small decline is observed, a result seen globally (Simmonds et al, 2016).


**Figure A 6.10** Verification of the UK emission inventory estimates for HFC-152a in Gg yr<sup>-1</sup> from 1990. GHGI estimates are shown in black. InTEM (MHD-only, 3-year) estimates are shown in blue (±1σ). InTEM (DECC, 12-mth) estimates are shown in orange (±1σ).




**Figure A 6.10** shows the InTEM and the GHGI emission estimates for the UK for HFC-152a for the period 1990 onwards. Between 2002-2007 and from 2011 onwards the GHGI estimates are significantly larger than those estimated through inverse modelling. The InTEM estimates are consistently on the lower side compared to the GHGI estimates with larger uncertainties. It is also interesting to note the positive trend from 2011 until 2017 in the UK GHGI, conflicting with a much flatter InTEM trend. The InTEM UK DECC network estimates are consistent with the InTEM MHD-only estimates.

**A 6.4.3 HFC-125**

**Figure A 6.11** Monthly Northern Hemisphere trend in HFC-125 estimated from Mace Head observations (blue line). Red line denotes the annual trend. Data under grey shading are not yet ratified.



**Figure A 6.12** Verification of the UK emission inventory estimates for HFC-125 in Gg yr<sup>-1</sup> from 1990. GHGI estimates are shown in black. InTEM (MHD-only, 3-year) estimates are shown in blue ( $\pm 1\sigma$ ). InTEM (DECC, 12-mth) estimates are shown in orange ( $\pm 1\sigma$ ).




**Figure A 6.11** shows the Northern Hemisphere background atmospheric concentration of HFC-125 from 1998 onwards. The background trend is monotonic and exponentially increasing, in 2018 the background increased by 3.2 ppt.

InTEM emission estimates for the UK for HFC-125 for the period 1999 onwards are shown in **Figure A 6.12**. Both the InTEM and UK GHGI estimates suggest that the emissions of HFC-125 from the UK have increased significantly from the 1990s. The InTEM estimates are consistently lower than those from the GHGI (by ~50%) and the uncertainties do not overlap. The introduction of the TAC data in the InTEM estimates shows consistency with the MHD-only estimates.


**A 6.4.4 HFC-143a**

**Figure A 6.13** shows the Northern Hemisphere background atmospheric concentration of HFC-143a from 2004 onwards. The background trend is positive, in 2018 it increased by 1.6 ppt.

**Figure A 6.13** Monthly Northern Hemisphere trend in HFC-143a estimated from Mace Head observations (blue line). Red line denotes the annual trend. Data under grey shading are not yet ratified.



**Figure A 6.14** Verification of the UK emission inventory estimates for HFC-143a in Gg yr<sup>-1</sup> from 1990. GHGI estimates are shown in black. InTEM (MHD-only, 3-year) estimates are shown in blue (±1σ). InTEM (DECC, 12-mth) estimates are shown in orange (±1σ).




InTEM emission estimates for the UK for HFC-143a for the period 2005 onwards are shown in **Figure A 6.14** and are compared to the GHGI estimates. UK emissions, as estimated by the GHGI, have increased year on year from the early 1990s until 2010 when a sharp decline is estimated. The InTEM estimates show a slow rise during this period but are consistently lower than the GHGI estimates. The estimates with the DECC network (i.e. including TAC data) are consistent with those from MHD-only and estimate a sharp decline in 2018, a decline that is estimated in GHGI to have started in 2016.

#### A 6.4.5 HFC-23


**Figure A 6.15** shows the Northern hemisphere background atmospheric concentration of HFC-23 from 2008 onwards. The background trend is monotonic and positive, in 2018 the background increased by 1.2 ppt.


InTEM emission estimates for the UK for HFC-23 from 2009 are higher than the low emissions estimated by the GHGI for this period, although the InTEM uncertainties are large and mostly extend down to zero. The introduction of the TAC emissions in 2013 re-enforces this finding, albeit the emissions are very small (**Figure A 6.16**). There are clearly some intermittent emissions of HFC-23 remaining in the UK.

**Figure A 6.15** Monthly Northern Hemisphere trend in HFC-23 estimated from Mace Head observations (blue line). Red line denotes the annual trend. Data under grey shading are not yet ratified.



**Figure A 6.16** Verification of the UK emission inventory estimates for HFC-23 in Gg yr<sup>-1</sup> from 1990. GHGI estimates are shown in black. InTEM (MHD-only, 3-year) estimates are shown in blue ( $\pm 1\sigma$ ). InTEM (DECC, 12-mth) estimates are shown in orange ( $\pm 1\sigma$ ). The second plot is plotted from the year 2000 on an expanded y-axis.






**A 6.4.6 HFC-32**


Figure A 6.17 shows the Northern Hemisphere background atmospheric concentration of HFC-32 from 2004 onwards. The background trend is monotonic, positive and the growth rate is increasing, in 2018 the background increased by 3.4 ppt.

**Figure A 6.17 Monthly Northern Hemisphere trend in HFC-32 estimated from Mace Head observations (blue line). Red line denotes the annual trend. Data under grey shading are not yet ratified.**



InTEM emission estimates for the UK for HFC-32 for 2004 onwards are shown in Figure A 6.18. The InTEM emission estimates are lower than the GHGI estimates. Both trends are positive but the rate of increase of the GHGI is significantly larger than the InTEM estimates. By 2017 the GHGI estimated emissions are significantly (more than 100%) larger than those estimated by InTEM. The DECC network InTEM estimates show a slight decline and then rise but are broadly consistent with the MHD-only InTEM estimates.


**Figure A 6.18** Verification of the UK emission inventory estimates for HFC-32 in Gg yr<sup>-1</sup> from 1990. GHGI estimates are shown in black. InTEM (MHD-only, 3-year) estimates are shown in blue ( $\pm 1\sigma$ ). InTEM (DECC, 12-mth) estimates are shown in orange ( $\pm 1\sigma$ ).




**A 6.4.7 HFC-43-10mee**

Figure A 6.19 shows the Northern Hemisphere background atmospheric concentration of HFC-43-10mee from 2011 onwards. There is a slight positive trend in the background with an overall growth rate of  $\sim 0.01$  ppt yr<sup>-1</sup>. The UK emissions of this gas are small. The GHGI estimates are in agreement with those estimated by InTEM. The DECC network InTEM estimates are consistent with the MHD-only InTEM estimates.

**Figure A 6.19** Monthly Northern Hemisphere trend in HFC-43-10mee estimated from Mace Head observations (blue line). Red line denotes the annual trend. Data under grey shading are not yet ratified.




**Figure A 6.20** Verification of the UK emission inventory estimates for HFC-43-10mee in Gg yr<sup>-1</sup> from 1990. GHGI estimates are shown in black. InTEM (MHD-only, 3-year) estimates are shown in blue ( $\pm 1\sigma$ ). InTEM (DECC, 12-mth) estimates are shown in orange ( $\pm 1\sigma$ ).




**A 6.4.8 HFC-227ea**

Figure A 6.21 shows the Northern Hemisphere background atmospheric concentration of HFC-227ea from 2007 onwards. There is a positive trend in the background; in 2018 it increased by 0.13 ppt. The GHGI estimates are significantly (more than 100%) higher than those obtained through inverse modelling. Both the GHGI and InTEM estimate increasing UK emissions.

**Figure A 6.21** Monthly Northern Hemisphere trend in HFC-227ea estimated from Mace Head observations (blue line). Red line denotes the annual trend. Data under grey shading are not yet ratified.




**Figure A 6.22** Verification of the UK emission inventory estimates for HFC-227ea in Gg yr<sup>-1</sup> from 1990. GHGI estimates are shown in black. InTEM (MHD-only, 3-year) estimates are shown in blue ( $\pm 1\sigma$ ). InTEM (DECC, 12-mth) estimates are shown in orange ( $\pm 1\sigma$ ).




**A 6.4.9 HFC-365mfc**

**Figure A 6.23** shows the Northern Hemisphere background atmospheric concentration of HFC-365mfc from 2005 onwards. There is positive trend in the background; in 2016 it increased by 0.05 ppt. The GHGI shows a sharp decline in emissions in 2008, the MHD-only InTEM estimates, with the 3-yr inversion time periods, shows the same but understandably slower response. Post-2012 the GHGI estimates rising UK emissions, a trend reproduced by InTEM, however the InTEM estimates then show a decline starting in 2017 that is not seen in the GHGI.

**Figure A 6.23** Monthly Northern Hemisphere trend in HFC-365mfc estimated from Mace Head observations (blue line). Red line denotes the annual trend. Data under grey shading are not yet ratified.




**Figure A 6.24** Verification of the UK emission inventory estimates for HFC-365mfc in Gg yr<sup>-1</sup> from 1990. GHGI estimates are shown in black. InTEM (MHD-only, 3-year) estimates are shown in blue ( $\pm 1\sigma$ ). InTEM (DECC, 12-mth) estimates are shown in orange ( $\pm 1\sigma$ ).




**A 6.4.10 HFC-245fa**

**Figure A 6.25** shows the Northern Hemisphere background atmospheric concentration of HFC-245fa from 2007 onwards. There is a positive trend in the background; in 2018 it increased by 0.21 ppt. The InTEM estimates have significant uncertainty and are consistently lower than the GHGI estimates. The GHGI estimates show a significant decline in 2008 and then a steady annual increase. The DECC network InTEM estimates show a strong rise in emissions from 2014 to 2018.

**Figure A 6.25** Monthly Northern Hemisphere trend in HFC-245fa estimated from Mace Head observations (blue line). Red line denotes the annual trend. Data under grey shading are not yet ratified.



**Figure A 6.26** Verification of the UK emission inventory estimates for HFC-245fa in Gg yr<sup>-1</sup> from 1990. GHGI estimates are shown in black. InTEM (MHD-only, 3-year) estimates are shown in blue (±1σ). InTEM (DECC, 12-mth) estimates are shown in orange (±1σ).




**A 6.5 PERFLUOROCARBONS**

**A 6.5.1 PFC-14**


**Figure A 6.27** shows the Northern Hemisphere background atmospheric concentration of PFC-14 from 2004 onwards. The background trend is positive; in 2018 it increased by 0.95 ppt.

The drop in emissions in 2012 in the GHGI reflects the closure of the last significant aluminium production plant in the UK. The InTEM uncertainty ranges are large and generally extend down to zero, probably because the majority of emissions come from intermittently emitting point sources. Overall there is good agreement between the GHGI and the InTEM estimates.

**Figure A 6.27** Monthly Northern Hemisphere trend in PFC-14 estimated from Mace Head observations (blue line). Red line denotes the annual trend. Data under grey shading are not yet ratified.




**Figure A 6.28** Verification of the UK emission inventory estimates for PFC-14 in Gg yr<sup>-1</sup> from 1990. GHGI estimates are shown in black. InTEM (MHD-only, 3-year) estimates are shown in blue (±1σ). InTEM (DECC, 12-mth) estimates are shown in orange (±1σ).




**A 6.5.2 PFC-116**

Figure A 6.29 shows the Northern Hemisphere background atmospheric concentration of PFC-116 from 2004 onwards. The background trend is monotonic and positive; in 2018 the background increased by 0.1 ppt. The UK InTEM estimates are consistent with those reported in the GHGI (Figure A 6.30) given the uncertainties in both estimates.

**Figure A 6.29** Monthly Northern Hemisphere trend in PFC-116 estimated from Mace Head observations (blue line). Red line denotes the annual trend. Data under grey shading are not yet ratified.



**Figure A 6.30** Verification of the UK emission inventory estimates for PFC-116 in Gg yr<sup>-1</sup> from 1990. GHGI estimates are shown in black. InTEM (MHD-only, 3-year) estimates are shown in blue ( $\pm 1\sigma$ ). InTEM (DECC, 12-mth) estimates are shown in orange ( $\pm 1\sigma$ ).




**A 6.5.3 PFC-218**


Figure A 6.31 shows the Northern Hemisphere background atmospheric concentration of PFC-218 from 2004 onwards. The background trend is monotonic and positive; in 2018 the background increased by 0.02 ppt.

The InTEM estimates are consistent with those reported in the GHGI (Figure A 6.32). Interestingly the dip in UK GHGI estimates in 2008 and 2009 is shown by the InTEM estimates, although it is estimated to have occurred later in 2011. The DECC network InTEM estimates show a decline in emissions in 2016 and then a rise.

**Figure A 6.31** Monthly Northern Hemisphere trend in PFC-218 estimated from Mace Head observations (blue line). Red line denotes the annual trend. Data under grey shading are not yet ratified.




**Figure A 6.32** Verification of the UK emission inventory estimates for PFC-218 in Gg yr<sup>-1</sup> from 1990. GHGI estimates are shown in black. InTEM (MHD-only, 3-year) estimates are shown in blue (±1σ). InTEM (DECC, 12-mth) estimates are shown in orange (±1σ).




**A 6.5.4 PFC-318**

Figure A 6.33 shows the Northern Hemisphere background atmospheric concentration of PFC-318 from 2010 onwards. The background trend is monotonic and positive; in 2018 the background increased by 0.07 ppt. The UK InTEM estimates are significantly higher than the very small emissions reported in the GHGI (Figure A 6.34). However the estimated quantities have large uncertainties extending down to zero emissions.

**Figure A 6.33** Monthly Northern Hemisphere trend in PFC-318 estimated from Mace Head observations (blue line). Red line denotes the annual trend. Data under grey shading are not yet ratified.



**Figure A 6.34** Verification of the UK emission inventory estimates for PFC-318 in Gg yr<sup>-1</sup> from 1990. GHGI estimates are shown in black. InTEM (MHD-only, 3-year) estimates are shown in blue (±1σ). InTEM (DECC, 12-mth) estimates are shown in orange (±1σ).




**A 6.6 SULPHUR HEXAFLUORIDE**


**Figure A 6.35** shows the Northern Hemisphere background atmospheric concentration of sulphur hexafluoride (SF<sub>6</sub>) from 2004 onwards. The background trend is monotonic and positive; in 2018 the background increased by 0.35 ppt.

The UK MHD-only InTEM estimates show a rise from 2005 until 2009 and then a decline until 2015. From 2005 until 2009 the GHGI shows a steady decline from ~0.05 Gg yr<sup>-1</sup> to ~0.02 Gg yr<sup>-1</sup>, a small rise in 2010 and then a slow decline until 2015. There is a significant difference between the GHGI and InTEM estimates between 2008 and 2013. Between 2014 and 2017 the InTEM estimates with the DECC network agree with the MHD-only InTEM estimates and the GHGI.

**Figure A 6.35** Monthly Northern Hemisphere trend in SF<sub>6</sub> estimated from Mace Head observations (blue line). Red line denotes the annual trend. Data under grey shading are not yet ratified.




**Figure A 6.36** Verification of the UK emission inventory estimates for SF<sub>6</sub> in Gg yr<sup>-1</sup> from 1990. GHGI estimates are shown in black. InTEM (MHD-only, 3-year) estimates are shown in blue (±1σ). InTEM (DECC, 12-mth) estimates are shown in orange (±1σ).



### A 6.7 NITROGEN TRIFLUORIDE

**Figure A 6.37** shows the Northern Hemisphere background atmospheric concentration of nitrogen trifluoride (NF<sub>3</sub>) from 2013 onwards. The background trend is monotonic and positive; the growth rate in 2018 was estimated to be 0.22 ppt yr<sup>-1</sup>. NF<sub>3</sub> is only measured at MHD, they started in May 2013. The InTEM emission estimates for the UK are 1800 kg but with an uncertainty that extends down to zero. The GHGI estimate for 2017 is 31 kg.

**Figure A 6.37** Monthly Northern Hemisphere trend in NF<sub>3</sub> estimated from Mace Head observations (blue line). Red line denotes the annual trend. Data under grey shading are not yet ratified.




### A 6.8 CARBON DIOXIDE

High precision, high frequency measurements of carbon dioxide (CO<sub>2</sub>) are made across the UK DECC network. The Northern Hemisphere background trend is positive; in 2018 it increased by 2.2 ppm. It has a strong seasonal cycle due to the influence of the biosphere.

The CO<sub>2</sub> observed has three principle components:

1. Northern hemisphere background (**Figure A 6.38**).
2. Anthropogenic (man-made)
3. Non-anthropogenic (natural)

**Figure A 6.38 Monthly Northern Hemisphere trend in CO<sub>2</sub> estimated from Mace Head observations (blue line). Red line denotes annual trend. Data under grey shading are not yet ratified.**



Plants both respire CO<sub>2</sub> and absorb it through photosynthesis. Therefore the CO<sub>2</sub> flux from vegetation has a strong diurnal and seasonal cycle and switches from positive to negative on a daily basis. This unknown natural (biogenic) component of the observed CO<sub>2</sub> is significant when compared to the anthropogenic (man-made) component and cannot be assumed negligible (minimised during the winter months). From direct CO<sub>2</sub> observations it is not possible to distinguish between biogenic and anthropogenic CO<sub>2</sub>. Therefore it is difficult to use the CO<sub>2</sub> observations directly in an inversion to estimate anthropogenic emissions. This is because the diurnally varying biogenic CO<sub>2</sub> flux is at odds with a key assumption of the inversion method, namely that emissions do not strongly vary in time over the inversion time-window (monthly). Methods are under development to attempt to over-come these challenges, such as: the use of isotopic observations; through ratios with respect to anthropogenic carbon monoxide (CO); and tracking at what time of day air passes over the ground and using biogenic process models. The uncertainties associated with each of these methods are predicted to be significant.

The estimated uncertainties in the CO<sub>2</sub> GHGI are very small compared to inversion results. Work is on-going to seek to improve our methods of verifying inventory CO<sub>2</sub> emission estimates.



## ANNEX 7: Analysis of EU ETS Data

### A 7.1 INTRODUCTION

This annex summarises the analysis of the 2017 European Union Emissions Trading System (EU ETS) energy and emissions data that is used within the compilation of the UK GHG inventory. The EU ETS data are used to inform activity data estimates for heavy industry sectors, carbon dioxide emission factors of UK fuels within those sectors, and for comparison of fuel allocations to specific economic sectors against data presented in the *Digest of UK Energy Statistics* (DUKES), published by the Department of Business, Energy and Industrial Strategy (BEIS).

The EU ETS data are used in the UK GHGI compilation as follows:

- EU ETS raw data on energy and emission estimates are processed and checked to enable integration of the activity data, implied emission factors and installation emission estimates as far as practicable within the UK GHG inventory compilation. Emission sources reported in EU ETS are allocated to inventory fuels and source codes, outliers are identified and clarifications of data inconsistencies are sought with the regulatory agencies;
- EU ETS activity data are closely compared against the UK national energy balance (DUKES) published by BEIS, and any inconsistencies are researched, seeking to resolve these through consultation with BEIS wherever possible;
- The verified EU ETS data provides up to date high quality fuel compositional analysis of UK fuels, and these data are used to improve inventory emission estimates across the highly energy intensive sources such as power stations, refineries, cement kilns, and oil and gas sources;
- The EU ETS dataset for offshore oil and gas installations are checked to assess data consistency in emissions reporting between the EU ETS and the (more comprehensive) EEMS dataset that is used within the UK GHGI compilation;
- Overall, the Inventory Agency approach seeks to minimise data discrepancies between EU ETS and the GHGI as far as practicable, in order that the derivation of traded and non-traded emission estimates from the UK GHGI are as accurate as possible. Close consistency between the EU ETS and GHGI is an important aspect of the development of a complete and consistent evidence base for policy development and tracking progress towards UK GHG reduction targets in the non-traded sector under the EU Effort Sharing Decision.

The scope of reporting under EU ETS increased from the 2013 dataset onwards. Phase II of the EU ETS ran from 2008-2012 inclusive. Phase III reporting began in 2013, with some new emission sources and new installations reporting for the first time on their GHG emissions; in particular, the definition of combustion has now been extended to cover installations such as furnaces, driers, and other plant where heat is used directly. A handful of industrial process sources of CO<sub>2</sub> are also included from 2013, such as soda ash production. In the UK, the changes in reporting in Phase III are most significant for the chemicals sector, where the scope of reporting is larger than previously and now encompasses both new industrial process emission sources, and additional energy use. There is also a notable shift towards estimation methods that are based on mass balance calculations (e.g. for chemical manufacturing) within the UK operator reporting to EU ETS. Other sectors with significant increases in reporting are food and drink manufacture, where installations such as driers, ovens etc. were included for the first time thus adding to the emissions

from boilers and CHP plant that were reported in previous years, and roadstone coating, a sector which has not been present in the dataset before.

Analysis of the phase III data enabled the Inventory Agency to improve estimates of emissions from the combustion of waste residues and process off-gases within the chemical and petrochemical sectors (which are all reported under IPPU sector 2B10), as well as to generate improved estimates for the IPPU component of several specific manufacturing processes, such as for soda ash (2B7), and titanium dioxide (2B6). In addition, following a review of methodology for all IPPU sources, EU ETS data for phase II onwards has been used to improve emission estimates for glass production (2A3), brickmaking (2A4) and reductant use in electric arc furnaces (2C1).

The key findings from the analysis and use of the EU ETS data include:

- In the 2017 EU ETS dataset, a very high coverage of Tier 3 emissions data is evident for all fuel use in the power sector, and for solid, liquid and waste-derived fuels used in the cement and lime sectors. The proportion of Tier 3 data is somewhat lower for refinery fuel use, but still sufficiently high for the ETS to be considered the most reliable data available. All of the fuel quality data for these sources and fuels are therefore used within the UK GHGI, as the EU ETS fuel quality data is the most representative dataset available to inform UK carbon dioxide emission factors in the inventory;
- EU ETS emissions data from refineries are higher than estimates derived from DUKES activity data for all but two years within the time series, with a discrepancy evident in OPG emissions. Consultation with the industry trade association, UKPIA, and cross-checking with their data shows that the EU ETS data are felt to more accurately reflect estimates of CO<sub>2</sub>, and therefore UK GHGI estimates are based on EU ETS data rather than refinery fuel use data reported in the UK energy balance;
- There are a range of other activity data discrepancies when compared to DUKES within the oil & gas, cement and lime, other industry and iron and steel sectors. Revisions to fuel allocations within the UK GHGI have been implemented for a number of sources, whilst further research is needed in some instances to clarify the issues where the reporting format of EU ETS does not map explicitly to energy balance and GHG inventory reporting requirements;
- EU ETS data for fuel use at chemical and petrochemical production facilities has helped to identify and quantify the combustion of process off-gases that are derived from Natural Gas Liquid (NGL) feedstock to petrochemical production processes, and from combustion of carbon-containing process residues. Analysis of “fuel gas” calorific values and carbon content informs the calculations to estimate emissions from NGL-derived gases and other residues.

The use of EU ETS data in the UK GHG inventory is summarised in **Table A 7.1.1**.

Table A 7.1.1 Summary of the use of EU ETS data in the UK inventory

Category	Sub-categories	EU ETS data used			Comments
		Factors	Activity	Emissions	
1A1a	Power stations - coal, fuel oil natural gas, sour gas	✓			
1A1a	Power stations – pet coke			✓	Some additional data is sourced from process operators.
1A1b	Refineries – pet coke & OPG			✓	EU ETS figures only used where higher than DUKES-based emissions.
1A1b	Refineries – natural gas	✓			
1A1c 1B2	Upstream oil and gas production – Gas oil, natural gas, LPG, OPG			✓	
1A1c	Gas industry – natural gas		✓		
1A1c 1B1b 1A2a 2C1	Integrated steelworks	✓	✓		Use of various EU ETS data in complex carbon balance – factors for some fuels, activity data for others
1A1c	Collieries – Colliery methane	✓			
1A2b	Autogenerators - coal	✓			
1A2f	Lime - coal			✓	
1A2f	Lime – natural gas		✓		
1A2g	Industry - pet coke & waste solvents			✓	No alternative data available for this emission source.
1A2g	Industry – colliery methane	✓			
2A1	Cement			✓	Data used is actually from industry trade-association, but this is based on EU ETS returns
2A2	Lime			✓	
2A3	Glass			✓	
2A4	Bricks			✓	

Category	Sub-categories	EU ETS data used			Comments
		Factors	Activity	Emissions	
2B7	Soda ash			✓	
2B8g	Ethylene & other petrochemicals			✓	
2C1	Electric arc furnaces - reductants			✓	

## A 7.2 BACKGROUND

### A 7.2.1 EU ETS Data and GHG Inventories

The European Union Emissions Trading System (EU ETS) data provides annual estimates of fuel use and fuel quality data from the most energy intensive sites in the UK, and provides a source of data, or can be used to cross-check data held in the UK Greenhouse Gas Inventory (GHGI), and to inform the carbon contents of current UK fuels. The EU ETS has operated since 2005, and there are now 13 years' worth of data on fuel use and emissions across major UK industrial plant, for 2005-2017.

The data reported under the EU ETS includes quantities of fuels consumed (or other activity data for process sources of CO<sub>2</sub>), carbon contents of fuels and other inputs, calorific values (fuels only) and emissions of carbon dioxide, all presented by installation and by emission source. Activity data are also given for many biofuels, although emissions of CO<sub>2</sub> from these fuels are not included in the emissions data. This is useful though, since PI/SPRI/WEI/NIPI emissions data for CO<sub>2</sub> often include biocarbon as well as fossil carbon, and the EU ETS data on biofuels helps to explain differences between CO<sub>2</sub> emissions reported in EU ETS and in those regulator inventories. EU ETS data for individual installations are treated as commercially confidential by the UK regulatory authorities and so only aggregated emissions data are reported in inventory outputs.

As part of the UK's annual reporting requirements to the MMR and UNFCCC, the UK must include a comparison of the EU ETS data against the national inventory dataset within the National Inventory Report. Furthermore, the analysis of the inventory against the EU ETS dataset is coming under increasing scrutiny due to the development of domestic GHG reduction targets that are based on non-traded<sup>17</sup> emissions data only, and the growing need to understand the UK non-traded sector emissions for future reporting under the Effort Sharing Decision.

The EU ETS dataset helps to improve the UK GHG inventory in a number of ways:

- Identifying new sources, therefore improving completeness;
- Helping assess true levels of uncertainty in fuel- and sector-specific data;
- Providing fuel quality data and oxidation factors for complex processes;

<sup>17</sup> All GHG emissions that are regulated within the EU ETS are defined as "traded" emissions, whilst all other GHG emissions are defined as "non-traded". The EU Effort Sharing Decision will lead to the UK adopting a new target for GHG reductions by 2020 for all of the non-traded emissions (i.e. everything outside of EU ETS), and progress towards this target will be monitored through the UK GHG inventory.

- Providing information on process-specific emissions that are not apparent from the national energy balances;
- Reducing uncertainty in the GHGI; and
- Acting as a source of quality assurance to inventory data.

In the 1990-2017 inventory cycle, the Inventory Agency has updated and extended the EU ETS analysis conducted for inventory compilation, using the 2017 EU ETS dataset, which is the fifth year of reporting under the Phase III EU ETS scope. This annex presents a comprehensive review of the thirteen years' of EU ETS data, indicating where the data have already been used in the improvement of the GHGI, as well as highlighting outstanding issues which could be investigated further, with potential for further revision and improvement of the GHGI.

The Inventory Agency has also been provided with details of the 2011-2017 EU ETS data for all offshore oil and gas installations, which are regulated by the BEIS Offshore Inspectorate. Access to these detailed data has enabled a more thorough review of the fuel/gas quality and reported emissions from combustion and flaring sources at offshore installations, and has directly improved the completeness and accuracy of the sector estimates within the UK GHGI.

The analysis of the EU ETS data for use in the UK GHGI necessitates a detailed review of the available data, in order to ensure correct interpretation and application of the available data. The study team prioritises effort to the sources and sites that are the most significant in UK GHGI terms, and/or where data reporting discrepancies have been identified from previous work. For those sectors where EU ETS data are used in the GHGI, it is important to review emission factors from all major installations to ensure that any outliers are identified and checked prior to their inclusion in inventory calculations.

Wherever possible, consistent assumptions are made when interpreting data across all years of the EU ETS. For instance ensuring that each site is allocated to the same inventory sector in each year (unless there is reason to change it – some industrial combustion plants in recent years have been converted into power stations, and so these sites do need to be allocated to different sectors in different years), and that there is consistency in the way in which site-specific names for fuels are interpreted across the entire period. The information on the EU ETS method “Tier” used for each of the data dictates whether they are used in inventory compilation. The highest tier EU ETS data are assumed to be subject to the lowest level of uncertainty, and so only tier 3 and tier 4 emission factor data are used. Occasionally there are internal inconsistencies in the EU ETS data between the data on consumption of a given fuel and emissions from the use of that fuel. These need to be resolved before the data can be used in the UK GHGI. As emissions data are verified, we cross-check the detailed emissions data against the final verified emissions for each site. As a general rule it is found that the most appropriate solution to inconsistencies is to assume that the EU ETS emissions data are correct as EU ETS reporting requirements are well regulated, and that it is the activity data that need to be amended instead.

### **A 7.2.2 Scope of the UK EU ETS and Implications for the GHG Inventory**

There are a number of limitations to the EU ETS data that affect the data usefulness in GHG inventory compilation, including:

- The EU ETS data are only available from 2005 onwards, whilst the UK GHG inventory reports emission trends back to 1990. The additional information that EU ETS provides (e.g. year-specific emission factors for many fuels in energy intensive sectors) helps to reduce the uncertainties in inventory emission estimates for the later years, but care is needed where revisions to the time series are made back to 2005. A consistent approach

to inventory compilation across the time series is a key tenet of IPCC good practice guidance, and care is needed to ensure that the use of EU ETS data does not introduce a systematic reporting step-change in the UK GHGI;

- Further to this point, it is important to note that the scope of EU ETS reporting has evolved through the years, from Phase I (2005 to 2007) into Phase II (2008 to 2012 data) and now to Phase III (2013 onwards). The comparability of EU ETS data for many sectors is poor between these three phases. For example, many cement kilns did not report to EU ETS until Phase II: several sectors including cement were reporting under Climate Change Agreements and were opted-out of EU ETS during Phase I. Therefore in several sectors, more complete coverage of EU ETS reporting is evident in Phase II and data from 2008 onwards are therefore much more useful for UK GHGI reporting. The scope of coverage of chemical industry emissions has gone through two step changes – in 2008, and again in 2013, and some sectors (such as roadstone coating) only appear for the first time in the 2013 EU ETS data. Less significantly for the GHGI, many small installations, mostly in the public sector, were removed from EU ETS at the end of phase I. It is vital that the GHGI takes full account of such changes and that UK inventory data do not include trends that merely result from the increase (or decrease) in scope of EU ETS. The changes in EU ETS scope have made the data set increasingly useful, and there are now five years' worth of Phase II data and four years of data under Phase III, hence the EU ETS dataset is now an important source of information for the UK inventory;
- In the UK during EU ETS Phases I and II, the regulators adopted a “medium” definition of the term “combustion”, and as a result there were many sectors where fuel use in specific types of combustion unit were not included in the EU ETS reporting scope until the start of Phase III (2013 onwards). Examples of this include flaring on chemical sites, and fuel use in heaters, dryers, fryers and stenters in industry sectors such as: chemicals, food and drink, textiles, paper and pulp. Hence the total fuel use and GHG emissions from these sectors have typically been under-reported within the EU ETS historically compared with the UK inventory, with many sites and sources excluded from the scope of EU ETS. However, the EU ETS data for these sectors is also incomplete both in Phase II and Phase III because small installations are not covered by EU ETS. Therefore, while the change in scope for combustion installations in phase III is a positive step, it has relatively little impact on the data used in GHGI compilation. Some Phase III data has been used to improve the estimates of emissions from combustion of process wastes / off-gases in the chemical and petrochemical sector in the recent submissions.
- Phase III also brought an increased scope for industrial process sources of CO<sub>2</sub>, and data appeared for the first time for soda ash production, and titanium dioxide manufacture.
- When using the EU ETS data, assumptions and interpretations are required to be made regarding the fuel types used by operators; assumptions are made on a case by case basis depending on knowledge of the site or industry and expert judgement. Operators are free to describe fuels as they wish in their returns, rather than choosing from a specific list of fuels, and so assumptions occasionally need to be made where the fuel type used is not clear from the operator's description. This issue was more significant in the earlier years of EU ETS reporting, with operators often using terms such as “Fuel 1”. The assumption then made about fuel type was based both on the other data the operator provided on the fuel such as calorific value, but also by comparison with later data for the same site, since operators now tend to use more recognisable fuel names, and the use of wholly ambiguous terms is now very rare.

Note that:

- The direct use of EU ETS data (e.g. fuel use data by sector) to inform UK GHGI estimates is limited to where the EU ETS is known to cover close to 100% of sector installations. For example, the EU ETS is regarded as representative and 100% comprehensive in coverage of refineries, power stations (except in the case of some small power stations burning biomass, gas oil, or burning oil as the main fuel), integrated steelworks, cement and lime kilns, soda ash plant, titanium dioxide plant, petrochemical works and glassworks (container, flat, wool & continuous filament fibre only – small lead glass and frit producers are not included). Coverage is very close to 100% for brickworks and tileries. For many other industrial sectors (such as chemicals, non-ferrous metals, food and drink, engineering) the EU ETS is not comprehensive and therefore the data are of more limited use, mainly providing a de-minimis fuel consumption figure for these sectors;
- EU ETS Implied Emissions Factors (IEFs) can be used within the UK GHGI, but only where the evidence indicates that EU ETS data are representative of the sector as whole and provides more comprehensive and accurate data than alternative sources. The key criteria to consider in the assessment of EU ETS IEF usefulness is the percentage of annual fuel use by sector where operator estimates use Tier 3 emission factors.
- Review of the EU ETS IEFs for different fuels across different sites provides a useful insight into the level of Tier 3 reporting within different sectors, the progression of higher-Tier reporting within EU ETS through the time series and the level of variability in fuel quality for the different major fuels in the UK. As a general rule, those energy-intensive sectors with near 100% coverage in EU ETS also report a very high proportion of emission factor data at Tier 3. Those sectors with incomplete coverage tend to report most emission factor data below Tier 3. As a result, in all cases where the level of sectoral coverage is high, the quality of reported data is also sufficiently high to be used with confidence in the UK inventory.

### **A 7.2.3 Limitations of EU ETS Data Integration with GHG Inventory: Autogeneration**

Despite detailed research there remain some fundamental limitations in the use of EU ETS data within national inventories where the sector allocation of energy use and emissions cannot be resolved against the national energy statistics that underpin the GHG inventory compilation. One key example is that of the division between fuel use in autogeneration (or heat generation) and direct fuel use within a specific sector. For example, based on the data available from EU ETS, it is impossible to differentiate between gas use in autogeneration on a chemical installation, and gas used directly to heat chemical production processes. In this example, the allocation of EU ETS energy use and emissions between 1A2c (chemicals) and 1A2f (autogenerators) is uncertain, and therefore comparison of EU ETS and GHGI estimates is uncertain. The EU ETS data are not sufficiently detailed and transparent to enable accurate allocation, and so in all cases fuels and emissions are allocated to the industry sector, and not to autogeneration.

It is worth noting here that the UK energy statistics are also subject to some uncertainty, however small, and that there is likely to be more uncertainty in estimates at industrial sector-level, rather than at more aggregated levels. For example, while fuel producers and suppliers will be able to quantify total fuel demand with a high level of certainty, it would be far more difficult for them to estimate fuel use by specific industrial sectors. This will be reflected in the quality of UK energy statistics which are used to estimate emissions from 1A2c etc. We consider that a high proportion of fossil fuel use by the UK chemical industry will be included in the EU ETS, on the basis that most industrial chemical processes will require sufficiently large combustion installations to

exceed the threshold for EU ETS. Therefore, we consider that it is reasonable to assume that EU ETS emissions for the chemical sector should cover most of the sector and therefore be similar in magnitude to those estimated from UK energy statistics and even, given the uncertainty in fuel allocation for autogeneration, to exceed them. For other sectors such as metals, paper, and food and drink, we would assume that the level of sectoral coverage by the EU ETS would be lower, so that emission estimates based on EU ETS would probably be lower than those based on energy statistics, even taking into account the uncertainty regarding autogeneration.

### **A 7.3 DATA PROCESSING**

BEIS provided the detailed EU ETS regulator data from the Environment Agency, Natural Resources Wales, Scottish Environment Protection Agency and Northern Ireland Environment Agency during April & May 2018, and the Inventory Agency industrial emissions experts progressed the analysis, combining the datasets to generate a UK-wide EU ETS dataset. The work built on analysis conducted in previous years, as the EU ETS has been in place since 2005, but this latest analysis, while focussing on the 2017 submission, did involve review of the data for earlier years, to ensure a consistent approach to the interpretation of energy and emissions data across the time series.

The initial step in the analysis is the allocation of all sites in the dataset to one of the economic sectors as reported within the DUKES Commodity Balance tables. Next, the reported fuels for every UK installation have to be allocated to one of the GHGI fuel names, which are also aligned with the fuel types reported within DUKES. This enables a direct comparison of EU ETS fuel totals against sector fuel allocations within DUKES and therefore used within the GHGI.

Most of the allocations have been made as part of previous years' work, and do not need to be revisited. However, several new installations included in the 2017 EU ETS data had to be allocated to DUKES' sectors, and all of the fuel data for 2017 also has had to be allocated to DUKES/GHGI fuel types. In a very small number of cases, we have revised data for earlier years, for example when it has become apparent that existing assumptions are likely to be incorrect. The allocation process does rely upon some expert judgement, with the Ricardo team using the reported EU ETS fuel names as well as the reported fuel quality data such as calorific values and carbon emission factors in order to make the fuel-type allocation for each entry in the EU ETS spreadsheet. The allocation is, occasionally, quite uncertain, particularly with the allocation of petroleum-based fuels such as the GHGI fuel categories LPG, OPG, gas oil and fuel oil, often because of the use of abbreviations or other slightly ambiguous names for fuels within the EU ETS reporting system. Cross-checking of data across the time series for each installation has been used to ensure as much consistency in fuel allocations as possible, although in some cases, operators of installations use different fuel terminology in different years, and the possibility of the use of different fuels in different years at a site cannot always be ruled out.

The quality checking and allocation process is very resource-intensive and essentially an open-ended task for such a large dataset, and hence the Inventory Agency focuses on the highest emitters and the known "problem" sites and fuel types. Where uncertainties arise in allocations, the most important allocation decisions are copied across to the BEIS DUKES team, for their information and input, as ultimately the EU ETS analysis by the Inventory Agency is taken into account to some degree within the compilation of DUKES for the following year.

As a data verification step, the installation emissions (broken down by fuel) from the EU ETS regulator spreadsheets are then compared against the total installation emissions for 2017 on the European Union Transaction Log (EUTL) which is a central website that holds the verified EU

ETS emissions totals for all EU installations in the scheme. Each year we have noted that for some sites the regulator data does not match the EUTL dataset, and therefore some “residual” emissions allocations are generated, from the difference between EUTL and regulator information. In cases where these residual emissions are large, then these are fed back to the regulator contacts, for their consideration and to request any insights into the likely fuels that the residual emissions should be allocated against. Minor residual emissions are ignored for the purposes of the analysis reported here.

A final data set is then available for fuel combustion emission sources, which includes the following data fields:

- GHGI Source Category;
- GHGI Fuel Category;
- Fuel Consumed;
- Fuel Calorific Value;
- Fuel Carbon Emission Factor; and
- Related Emissions of CO<sub>2</sub>

The Inventory Agency then combines the data by sector and/or fuel category to provide data for comparison against GHGI emissions data, and energy statistics published in DUKES. In this way, the analysis can:

- provide improved CO<sub>2</sub> emission factors for highly energy-intensive industrial sectors covered by the GHGI through the use of verified data;
- provide a comparison with UK energy statistics, allowing the identification of inconsistencies between EU ETS and DUKES;
- Identify any emission sources that are not contained in the GHGI.

The analysis of the EU ETS data for all onshore facilities was completed by May 2018 and provided to the BEIS team of energy statisticians, in time for them to consider the EU ETS dataset during compilation of the UK energy balance for 2017, as published within DUKES 2018 (which was published in July 2018).

The EU ETS data for offshore oil and gas installations was provided in May 2018 and were used directly in the compilation of emission estimates for the upstream oil and gas sector, after the UK energy balance had been compiled by BEIS. Access to these EU ETS data for offshore facilities provided more fuel-specific information (GCV, carbon content) to help improve completeness and accuracy of the upstream oil and gas estimates in the UK GHGI, augmenting the EEMS dataset which is a more comprehensive dataset (i.e. EEMS covers more emission sources than EU ETS) but does not provide the same level of fuel-specific data.

## A 7.4 EU ETS DATA COVERAGE

The coverage of the EU ETS data has changed over the 13 years for which data are available. Major changes have been outlined in **Section A 7.1**, and these changes in scope have an impact on the usefulness of data for some sectors, with data generally being more complete for Phase II (2008-12) and Phase III (2013-17) of EU ETS. In addition, smaller combustion installations in the industrial, commercial and public sectors are outside the scope of EU ETS, and in fact coverage was decreased after 2007 due to the exemption of certain 'small emitters' from the UK EU ETS. For some source sectors in the GHGI, the EU ETS data therefore only includes a small proportion of the sector and the EU ETS data are not useful to directly inform the GHGI.

The following GHGI source sectors are well represented in the EU ETS data sets in the UK, with all UK installations included:

- Power stations burning coal, gas, and fuel oil as the principal fuel;
- Oil refineries;
- Coke ovens & Integrated steelworks;
- Cement kilns (from Phase II onwards); and
- Lime kilns (from Phase II onwards)
- Glassworks - container, flat, wool & continuous filament glass fibre subsectors only (from phase II onwards)
- Brickworks and other sites manufacturing heavy ceramic goods (from Phase II onwards)
- Titanium dioxide and soda ash manufacture (from Phase III onwards).

However, GHGI sectors such as industrial combustion, autogeneration, and public sector combustion are only partially represented in the EU ETS data. An indication of the actual level of coverage of the EU ETS data can be seen in . The number of sites in each sector which are included in the ETS dataset for 2005 and 2017 are given, together with the Inventory Agency's estimate of the total number of installations in that sector throughout the UK in those years.

**Table A 7.4.1 Numbers of installations included in the EU ETS data**

Sector	Number of installations			
	2005		2017	
	EU ETS	UK total	EU ETS	UK total
Power stations (fossil fuel, > 75MWe)	60	60	51	51
Power stations (fossil fuel, < 75MWe)	23	27	30	35
Power stations (nuclear)	12	12	9	9
Coke ovens	4	4	2	2
Sinter plant	3	3	2	2
Blast furnaces	3	3	2	2
Cement kilns	8	15	11	11
Lime kilns	4	15	11	11
Refineries	12	12	8	8
Combustion – iron & steel industry	11	200 <sup>a</sup>	24	200 <sup>a</sup>
Combustion – other industry	171	5000 <sup>a</sup>	469	5000 <sup>a</sup>
Combustion – commercial sector	28	1000 <sup>a</sup>	108	1000 <sup>a</sup>
Combustion – public sector	169	1000 <sup>a</sup>	124	1000 <sup>a</sup>
Glassworks (flat, special, container & fibre)	6	32	23	23
Brickworks	18	80 <sup>b</sup>	52	52 <sup>b</sup>

Sector	Number of installations			
	2005		2017	
	EU ETS	UK total	EU ETS	UK total
Soda ash & titanium dioxide	0	4	1	1

<sup>a</sup> These estimates are 'order of magnitude' figures, based on expert judgement of the inventory team, to show that the number of installations in the UK is likely to be considerably higher than the number of installations reporting in the EU ETS.

<sup>b</sup> Numbers of brickworks are not certain in 2005 but will have been significantly higher than in 2008 (when there were about 70) since many brickworks were closed or mothballed in the second half of 2007. All brickworks are believed to be covered by EU ETS in 2017.

Data are included in EU ETS for all coke ovens, refineries, sinter plant and blast furnaces. Power stations are divided into three categories in the table in order to show that, although a few stations are not included in the EU ETS data for 2017, these are all small (in most cases, very small diesel-fired plant supplying electricity to Scottish islands). In comparison, coverage is quite poor in 2005 for cement and lime kilns (due to CCA participants opting out during Phase I) and for combustion processes (due to CCA/UKETS opt-outs and the fact that numerous combustion plant are too small to be required to join the EU ETS). All cement kilns and all lime kilns are included in 2017. Coverage of glassworks and brickworks was very limited during Phase I, but since 2008 has been very good: all large glassworks have been included since 2008, and all but one brickworks were included in Phase II, with that remaining site being added for Phase III. UK totals for brickworks are subject to some uncertainty however, and may be revised in future should more data be obtained. Both soda ash plant and both plants manufacturing titanium dioxide via the chloride process have only been included in EU ETS since the start of Phase III.

For most emission sources the level of detail given in the EU ETS data matches well with the structures of the GHGI, allowing comparison of like with like. Only in the case of coke ovens and integrated steelworks is this not the case, since the EU ETS reporting format does not provide a breakdown of emissions for the sectors reported within the GHGI: i.e. estimates of emissions from coke ovens, blast furnaces and sinter plants are not provided explicitly. However, for these sectors, additional detailed analysis, including the collection of other industry data, has allowed for far greater use of EU ETS data for the inventory.

## A 7.5 EU ETS DATA USE IN THE UK GHGI

The use of EU ETS data in the UK GHGI may conveniently be divided into two classes:

- Instances where activity data and, in most cases, emission totals as well are taken from EU ETS;
- Instances where emission factors only are taken from EU ETS and then used in the UK GHG Inventory with activity data from other sources such as DUKES.

### A 7.5.1 Activity and Emissions Data

#### A 7.5.1.1 Crude Oil Refineries

The comparison of EU ETS emissions data against GHGI data based on DUKES fuel use allocations for petcoke, natural gas, fuel oil and OPG use is inconsistent to varying degrees in different years. Previous EU ETS analysis indicated that petcoke data in DUKES were too low; the BEIS energy statistics team have investigated this matter with the refinery operators and have

revised data for a number of sites that had been misreporting through the DORS system used to compile DUKES. In recent years, therefore, the EU ETS and DUKES data are closely consistent for petcoke use by refineries.

Data inconsistencies between DUKES and EU ETS remain for other fuels, however. In some cases, this will be due to misallocation of fuel use data within the EU ETS analysis, where fuel names are unclear, e.g. “fuel gas” could be interpreted solely as refinery use of OPG or to also cover the use of natural gas as a back-up fuel within the refinery fuel gas system.

The fuel oil activity data in most years is around 10% higher in EU ETS than in DUKES. Natural gas is a relatively minor fuel in the sector; whilst the EU ETS allocations indicate an over-report in DUKES, there is considerable uncertainty over the allocations of gases in the EU ETS dataset, as noted above. However, DUKES data for natural gas used in autogeneration includes some fuel burnt at refineries, thus the difference between refinery fuel use as given in EU ETS, and that derived from DUKES data can be reduced by taking this into account. Consumption of naphtha reported in DUKES as “unclassified industry” is allocated to refineries as the only known consumers in the UK. However, in the case of OPG, there is typically an under-report in DUKES, although the data in DUKES is higher in two years. **Table A 7.5.1** below presents the emissions allocated to OPG for those years (2004 onwards except 2005, 2012) where UKPIA and EU ETS data indicates that DUKES data are too low. Note that the GHGI estimates also include the assumption that all of the OPG allocation to “autogenerators” within the DUKES commodity balance tables (in the column “Other gases”) is used within the refinery sector. Consultation with the BEIS DUKES team has indicated (Personal Communication, Evans, 2010) that the “Other gases” column in the Commodity Balance tables is the OPG on the refinery basis, with CHP plant on site allocated to the autogeneration line. We have therefore retained this assumption in the current analysis, including the autogenerator allocation of “other gases” within the refinery sector.

To resolve the refinery sector under-report, we have compared DUKES data against EU ETS data, and also considered the total carbon dioxide emissions for the refinery sector provided annually by UKPIA. At the installation level, the UKPIA and EU ETS data show very close consistency for recent years (typically within 1%). The close consistency of the EU ETS and UKPIA data further strengthens the case for using EU ETS data as the primary dataset to inform the UK GHG inventory, in preference to the DUKES energy statistics.

At the fuel-specific level, the greatest disparity is evident in the reporting of OPG use at refineries; the reporting disparity has therefore been resolved through a top-down emissions comparison between DUKES-derived data and the best available operator data from EU ETS (2005 onwards) and UKPIA (pre-2005), with the difference between the two then allocated to OPG use in the UK GHGI. UK inventory estimates of emissions for the sector are therefore aligned with EU ETS totals back to 2005, and with UKPIA data prior to 2005, unless the estimates derived from DUKES data are higher than those from UKPIA or EU ETS (i.e. in 2005 and 2012).

No deviations from UK energy statistics have been made prior to 2004, as the data from UKPIA and GHGI estimates based on DUKES are closely consistent with the DUKES-derived data being slightly higher; therefore a conservative approach is adopted, using DUKES-derived GHG estimates.

The time series of emissions data and the additional OPG and petroleum coke emissions data (where DUKES data are low) for the sector are shown below.

**Table A 7.5.1 Refinery Emissions Data Comparison and Revision to OPG Activity**

Year	Best Operator Data <sup>1</sup>	Refinery emissions total (if based on DUKES)	Additional emissions assumed from OPG	Additional emissions assumed from Pet Coke
	kt C	kt C	kt C	kt C
2000	4,599	4,717	0	0
2001	4,535	4,664	0	0
2002	4,767	5,243	0	0
2003	4,772	5,085	0	0
2004	4,999	4,925	74	0
2005	5,007	5,422	0	150
2006	4,910	4,751	83	76
2007	4,857	4,781	26	50
2008	4,709	4,470	118	121
2009	4,492	4,126	239	126
2010	4,632	4,426	130	76
2011	4,739	4,491	248	0
2012	4,287	4,300	0	0
2013	4,002	3,856	145	2
2014	3,678	3,559	118	0
2015	3,682	3,657	0	47
2016	3,708	3,586	65	56
2017	3,698	3,629	56	13

<sup>1</sup> For 2005 onwards, the EU ETS data are verified by third parties and regarded as the best available sector estimates; prior to 2005 the best available operator emissions data are from the trade association, UKPIA.

<sup>2</sup> For 2005, DUKES activity data for petroleum coke are somewhat lower than the corresponding figure in the EU ETS, so even though CO<sub>2</sub> emission estimates based on DUKES figures for all fuels exceed the CO<sub>2</sub> figure given in the EU ETS, we use the higher (EU ETS) figure for petroleum coke, with the result that for 2005, the UK inventory figure for refinery CO<sub>2</sub> is higher (at 5422 kt C) higher than either the operator or DUKES based totals.

There is some level of uncertainty in the allocation of fuels in EU ETS to specific “DUKES” fuels, although the OPG use in refineries seems to be reported quite consistently as “Refinery Gas”, “Refinery Off-Gas”, or “OPG/RFG”. The BEIS DUKES team have reviewed the year to year consistency of OPG use in refineries through the DORS system.

#### **A 7.5.1.2 Oil & Gas Terminal OPG and LPG Use**

The allocation of fuel use reported within EU ETS to UK energy balance fuel nomenclature is uncertain in some cases. Analysis of the EU ETS fuel use data does indicate that there are small amounts of LPG and OPG being used in the upstream oil & gas sector that are not evident within DUKES.

The DUKES team have noted previously (Personal communication, DECC, 2010) that some LPG and OPG fuels are abstracted from upstream oil and gas exploration and production sources, rather than purchased from other sources, and that no data have been collected for this source since DUKES last published data, for the year 2002.

Therefore, the data from the EU ETS from oil and gas processing terminals on LPG and OPG combustion are used directly within the UK GHG inventory for the Phase II and Phase III years of 2008 to 2017, with estimates for 2003 to 2007 derived by interpolation between the EU ETS 2008 data and the DUKES 2002 data.

#### **A 7.5.1.3 Natural Gas Use by Downstream Gas Supply Installations**

The EU ETS data includes natural gas use by large gas compressor and storage sites that operate on the UK gas transmission and distribution network, as well as the three operational LNG terminals and a small number of other downstream gas industry sites.

The gas use reported in EU ETS for these sites throughout Phase II and III has been notably higher than the allocation of gas within DUKES Commodity Balance table 4.2 (Energy Industry Use, Other). This has been evident in the traded / non-traded analysis for the gas supply sector in the UK and DA GHGI.

As this gas use arises from the downstream network, the Inventory Agency and the BEIS DUKES team consider that the DUKES data indicate a small misallocation of gas use, rather than a gap in reported gas use. For 2005 to 2017, therefore, the EU ETS data for this source are used within the UK GHG inventory, and the overall gas use data are balanced by reducing the allocation of gas use to “other industrial combustion” (IPCC source 1A2g); the EU ETS data since 2005 shows good consistency with the data from DUKES for earlier years.

The increased gas use for this sector based on EU ETS data is expected to still be a small under-report for the sector as a whole, as the EU ETS scope only includes around 35 of the larger gas compressors, LNG terminals and storage sites on the UK network, and it is likely that additional gas use on smaller sites also occurs. However, the Inventory Agency has no data to inform such estimates.

#### **A 7.5.1.4 Other Industry OPG use**

There are a number of “other industry” sites where OPG use has been allocated by the Inventory Agency from EU ETS data, where the fuel is defined as either a specific gas (e.g. ethane, propane, butane) or more generic terms such as “OPG”, “High Pressure Refinery Gas”, “Low Pressure Refinery gas”, “fuel gas” or “RFG/OPG/ROG” within the EU ETS forms.

In refinery complexes, the use of RFG for autogeneration (for the refinery and/or for co-located plant) is reported within the energy balance (allocated to “OPG”). At a number of other UK installations, commodities that are used initially as feedstocks in chemical and petrochemical production (e.g. naphtha, ethane, LPG, gas oil) are allocated to “non-energy use” in the UK energy statistics; any subsequent use of process off-gases (derived from these NEU feedstocks) as a fuel is not reflected in DUKES. Therefore the inventory agency uses other data from industry, primarily from EU ETS, to generate estimates of the use of such secondary fuels. For a small number of sites, consultation with the DUKES team, regulators and operators has clarified that the EU ETS energy and emissions data are the best available dataset for use in the UK GHGI.

In the 1990-2017 inventory cycle, EU ETS data for fuel use at petrochemical production facilities helped to identify and quantify the combustion of process off-gases that are derived from Natural

Gas Liquid (NGL) feedstock to petrochemical production processes. Analysis of “fuel gas” calorific values and carbon content informs the calculations to estimate emissions from NGL-derived gases and other residues.

#### **A 7.5.1.5 Industrial Processes**

The EU ETS dataset contains data on a number of industrial processes for which alternative data sources are either unavailable or of low quality. The EU ETS data therefore constitute the most reliable set of emissions data for these processes and are used in the UK inventory. In almost all cases, the EU ETS activity data are difficult to use directly, largely because different operators provide activity data and emission factors on a different basis (e.g. some may provide input material and emission factors on a consumption basis, others will provide production data and emission factors on a production basis). Therefore, for all of the industrial process sources, the EU ETS emissions data are adopted, and activity data are generally back-calculated from the emissions using a suitable IPCC emission factor. The industrial process sources where EU ETS data have been used to generate estimates of emissions included within the UK GHGI in this submission, include:

- Emissions from the manufacture of lime. UK activity data for limestone and dolomite consumption in lime production would yield much lower emission estimates than is suggested by EU ETS returns therefore, as a conservative approach, the EU ETS data are used instead. Activity data are back-calculated using the IPCC default factor for lime production. See **Section 4.3** for further details.
- Emissions from the use of carbonates in the manufacture of glass. As with lime production, the available data on consumption of limestone and dolomite for glass production are suspect, being very inconsistent across the time series, and so EU ETS data are used in the generation of the inventory time series, as detailed in **Section 4.4**.
- Emissions from the use of clays, carbonate minerals and other additives in the manufacture of bricks and roofing tiles, as detailed in **Section 4.5**. The EU ETS data are very detailed, with separate lines for different input materials such as different types of clay, carbonate minerals used in the bricks or in scrubbers used to abate fluoride emissions, and coke oven coke/petroleum coke used as an additive in certain bricks. UK brick production data are used as activity data.
- Estimates for emissions from the use of limestone in flue-gas desulphurisation (FGD) plant for the years 2005-2017 are taken from EU ETS data, because UK activity data (for gypsum produced from the FGD plant) are incomplete for those years. Activity data for 1990-2004 are available from non-EU ETS sources, and are back-calculated from the EU ETS CO<sub>2</sub> emissions for 2005 onwards assuming an emission factor of 253 kg CO<sub>2</sub> per tonne gypsum produced (which is based on an assumed 100% conversion of limestone and SO<sub>2</sub> into gypsum and CO<sub>2</sub>).
- EU ETS Phase III saw the introduction of data for soda ash manufacturing sites and EU ETS data, and CO<sub>2</sub> emissions reported for earlier years in the PI, are used as the basis of UK inventory emissions data for that sector. See **Section 4.12** for more details.
- Titanium dioxide production was also included in Phase III of the EU ETS, but full data for the UK plant are not included in the data set provided, and so emission estimates are generated using an alternative, more conservative method.
- Petroleum coke is added to some electric arc furnaces as a reductant, and emissions and activity data for the period 2005-2017 are taken from EU ETS data, with emissions in earlier years being extrapolated on the basis of plant production. See **Section 4.16** for further details.

## A 7.5.2 Implied Emission Factors

### A 7.5.2.1 Power Stations

**Table A 7.5.2** summarises EU ETS data for fuels burnt by major power stations and coal burnt by autogenerators. The percentage of emissions based on Tier 3 emission factors is given (Tier 3 factors are based on fuel analysis, and are therefore more reliable than emission factors based on default values), as well as the average emission factor for EU ETS emissions based on Tier 3 factors.

**Table A 7.5.2 EU ETS data for Fuels used at Power Stations and Autogenerators (Emission Factors in kt / Mt for Coal & Fuel Oil, kt / Mth for Gases)**

Year	Fuel	% Tier 3	Average Carbon Emission Factor (Tier 3 only)
2005	Coal	99	615.3
2006		100	615.0
2007		100	614.7
2008		100	612.4
2009		100	607.2
2010		100	609.0
2011		100	608.9
2012		100	611.7
2013		100	612.5
2014		100	611.8
2015		100	607.9
2016		94	488.5
2017		100	613.0
2005	Fuel oil / Waste oil <sup>a</sup>	59	860.3
2006		66	873.0
2007		68	871.1
2008		91	869.5
2009		94	872.7
2010		95	873.2
2011		94	873.9
2012		95	873.1
2013		94	869.2
2014		93	868.5
2015		90	870.3
2016		86	872.4
2017		87	871.5
2005	Natural gas	52	1.443
2006		76	1.465

Year	Fuel	% Tier 3	Average Carbon Emission Factor (Tier 3 only)
2007		95	1.464
2008		97	1.467
2009		100	1.464
2010		99	1.460
2011		99	1.458
2012		100	1.461
2013		99	1.464
2014		100	1.461
2015		100	1.462
2016		99	1.462
2017		99	1.466
2005	7Coal - autogenerators	100	594.3
2006		100	596.3
2007		100	594.5
2008		100	581.3
2009		100	600.6
2010		100	599.9
2011		100	594.9
2012		100	598.3
2013-17		0 <sup>b</sup>	N/A

<sup>a</sup> It is not possible to distinguish between fuel oil and waste oil in the EU ETS data, so all emissions have been reported under fuel oil.

<sup>b</sup> Plant operated as a power station after 2012 and included in the figures for power stations burning coal

The EU ETS data shown are regarded as good quality data, since a high proportion of emissions are based on Tier 3 emission factors (i.e. verified emissions based on fuel analysis to ISO17025). The factors are also very consistent across the time-series, which would be expected for this sector. As shown in Section 3, the EU ETS data for power stations also cover almost all UK installations in this sector, and certainly cover all of the installations which burn coal, fuel oil and natural gas.

A few power stations burn small quantities of petroleum coke as well as coal. One supplies data to ETS for coal/petroleum coke blends i.e. there are no separate emissions data or carbon factors for the coal and the petroleum coke at that site. We therefore back-calculate the coal IEF in those blends by using an assumed default for the petcoke carbon content and more detailed activity data on the constituents of the fuel blends, obtained directly from the operator.

The EU ETS based emission factors presented above for power stations are used directly as the emission factors in the GHGI, with the exception of the 2005 figure for gas, where Tier 3 factors were only used for about half of the sector's emissions reported in EU ETS. Small quantities of sour gas were burnt at one power station in 2005-2007 and 2009 and EU ETS Tier 3 emission factors are available and therefore used. [Due to the confidentiality of the data, the emission factors are not shown]. Prior to 2005, the emission factors for these sectors are based on the

methodology established by Baggott *et al*, 2004, since it has been concluded that this represents the most reliable approach.

The EU ETS factors for coal-fired autogenerators are slightly different to the factors for the power stations in that, although the EU ETS data are exclusively Tier 3, they only represent about 80-90% of total fuel used by the sector.

### A 7.5.2.2 Crude Oil Refineries

**Table A 7.5.3** below summarises the EU ETS data for the major fuels burnt by refineries in the UK.

The main fuels in refineries are fuel oil and OPG and emissions also occur due to the burning off of 'petroleum coke' deposits on catalysts used in processes such as catalytic cracking. In the latter case, emissions in the EU ETS are not generally based on activity data and emission factors but are instead based on direct measurement of carbon emitted. This is due to the technical difficulty in measuring the quantity of petroleum coke burnt and the carbon content. Refineries also use natural gas, although it is a relatively small source of emissions compared to other fuels.

**Table A 7.5.3 Refinery EU ETS Data for Fuel Oil, OPG and Natural Gas (Emission Factors in kt / Mt for Fuel Oil and kt / Mth for OPG and Natural Gas)**

Year	Fuel	% Tier 3	Average Carbon Emission Factor (Tier 3 sites only)
2005	Fuel Oil	25	860.9
2006		65	873.8
2007		78	877.2
2008		91	871.6
2009		91	876.2
2010		97	878.2
2011		85	877.6
2012		82	887.1
2013		95	874.3
2014		96	875.8
2015		61	876.7
2016		66	876.1
2017		90	864.8
2005		OPG	56
2006	54		1.468
2007	65		1.587
2008	78		1.482
2009	78		1.494
2010	79		1.509
2011	68		1.445
2012	61		1.463
2013	77		1.493

Year	Fuel	% Tier 3	Average Carbon Emission Factor (Tier 3 sites only)
2014		64	1.508
2015		62	1.492
2016		61	1.470
2017		66	1.481
2005	Natural Gas	0	N/A
2006		43	1.460
2007		45	1.462
2008		98	1.475
2009		98	1.480
2010		97	1.465
2011		81	1.447
2012		63	1.442
2013		89	1.459
2014		87	1.459
2015		87	1.465
2016		81	1.456
2017		84	1.462

There has been some variation in the proportion of Tier 3 reporting for all three fuels, which will adversely affect the quality of the emission factors, although coverage is still in excess of 50% for all fuels.

Emission factors for **fuel oil** generated from EU ETS data have been adopted in the GHGI, with the exception of data for 2005, where Tier 3 methods were used for only 25% of fuel.

Carbon factors can be derived for **OPG** based on moderate levels of Tier 3 reporting for 2005-2007 and 2011-2017 but 80% for 2008-2010, which gives us a high confidence in the representativeness of the carbon factors for 2008-10. There is some uncertainty regarding the allocation of EU ETS fuels to the OPG fuel category, and the derived emission factors do cover a wider spread of values than for many other fuels in EU ETS. However, this reflects the nature of this fuel, and the data for all years have been used in the inventory.

Carbon factors for natural gas are based on a low % of Tier 3 reporting until 2008; in 2008 to 2010 over 90% of gas use is reported at Tier 3 and over 80% in 2011 and 2013-2017. Within the UK GHGI, the EU ETS factors for 2008 to 2016 are used directly, whilst emission factors for earlier years are derived from gas network operator gas compositional analysis.

EU ETS emission data for **petroleum coke** are higher in 2005-2010, when compared against the estimates derived from DUKES activity data and the industry-recommended emission factor. This is especially noticeable for 2005, where the petroleum coke consumption given in DUKES would have to be more than 100% carbon in order to generate the carbon emissions given in the EU ETS. Consultation with BEIS energy statisticians has identified that the figures given in DUKES are subject to uncertainty and hence the EU ETS data are used directly within the UK GHGI for those years.

### A 7.5.2.3 Integrated Steelworks & Coke Ovens

**Table A 7.5.4** summarises EU ETS data for the major fuels burnt at integrated steelworks and coke ovens. The data exclude one independent coke oven which calculated emissions using a detailed mass balance approach which makes it more difficult to assess the data in the same way as the other installations. This site closed at the end of 2014.

**Table A 7.5.4 EU ETS data for fuels used at integrated steelworks & coke ovens (Emission Factors in kt/Mt for solid & liquid fuels, kt/Mth for gases)**

Year	Fuel	% Tier 3	Average Carbon Emission Factor (Tier 3 sites only)
2005	Blast furnace gas	0	N/A
2006		100	6.873
2007		90	6.920
2008		92	6.945
2009		92	7.029
2010		100	6.949
2011		94	6.990
2012		96	6.815
2013		91	6.766
2014		91	6.776
2015		100	7.653
2016		100	7.578
2017		90	7.219
2005	Coke oven gas	0	N/A
2006		0	N/A
2007		0	N/A
2008		53	1.093
2009		96	1.140
2010		96	1.117
2011		96	1.089
2012		96	1.094
2013		96	1.103
2014		100	1.143
2015		100	1.216
2016		48	1.659
2017		100	1.068
2005	Natural gas	0	N/A
2006		3	1.479
2007		2	1.478
2008		0	N/A
2009		58	1.425

Year	Fuel	% Tier 3	Average Carbon Emission Factor (Tier 3 sites only)
2010		68	1.441
2011		64	1.441
2012		64	1.443
2013		27	1.447
2014		23	1.445
2015		0	N/A
2016		12	1.445
2017		33	1.446
2005	Fuel oil	0	N/A
2006		0	N/A
2007		0	N/A
2008		84	878.3
2009		89	884.7
2010		83	887.6
2011		88	888.7
2012		67	877.3
2013		33	846.2
2014		30	844.7
2015		32	845.1
2016		0	N/A
2017		0	N/A

Most of the ETS data for coke ovens and steelworks are now used in the GHGI, although not the emission factors shown above. Instead, the Inventory Agency have used the EU ETS data and other detailed, site-specific and fuel-specific data, provided by the process operators to refine the carbon balance model used to generate emission estimates for the sector. Details of the revisions to the carbon balance model can be found in the research report from the 2013-2014 inventory improvement programme (Ricardo-AEA, 2014)

#### A 7.5.2.4 Cement Kilns

**Table A 7.5.5** summarises EU ETS data for the major fuels burnt at cement kilns.

**Table A 7.5.5 EU ETS data for Fuels used at Cement Kilns (kt / Mt)**

Year	Fuel	% Tier 3	Average Carbon Emission Factor (Tier 3 sites only)
2005	Coal	100	678.6
2006		100	546.2
2007		100	664.3
2008		100	655.8
2009		97	658.8

Year	Fuel	% Tier 3	Average Carbon Emission Factor (Tier 3 sites only)
2010		97	637.8
2011		100	645.8
2012		100	662.4
2013		100	694.2
2014		100	673.9
2015		100	675.3
2016		100	683.6
2017		100	683.3
2005	Petroleum coke	0	810.7
2006		100	820.8
2007		100	830.2
2008		100	819.1
2009		73	812.6
2010		69	791.9
2011		100	738.4
2012		100	770.2
2013		100	811.1
2014		100	793.4
2015		100	824.6
2016		100	822.2
2017		100	823.1

The EU ETS dataset also provides a detailed breakdown of cement sector process emissions from the decarbonisation of raw materials during the clinker manufacturing process. These data are useful to compare against statistics provided by the Mineral Products Association (MPA) regarding clinker production and the non-combustion emissions associated with UK cement production. The MPA data on clinker production are commercially confidential.

The two data sets show significant differences for 2005-2007; however the EU ETS data cover only a fraction of the sector, so differences might be expected. From 2008 onwards, there is close agreement (average of 0.5% difference) between the two data sets. The coal IEF data across the time series are also fairly consistent, other than in 2006 where the ETS value is very much lower than in other years. Because of the good agreement in both activity data and emission factors for 2008 onwards, the industry-wide estimates provided by the MPA and used within the GHGI show very close comparison with the EU ETS estimates. The difference between the EU ETS and those reported to the GHGI are consistently less than 1%, as outlined below in **Table A 7.5.6**.

**Table A 7.5.6 Comparison of Cement Sector Carbon Dioxide Emissions\* within the UK GHGI and the EU ETS**

	2008	2009	2010	2011	2012	2013	2014	2015	2016	2017
GHGI CO <sub>2</sub> emissions (kt)	8,296	5,687	5,789	6,148	5,602	6,009	6,212	6,622	6,819	6,569
Sum of EU ETS CO <sub>2</sub> emissions (kt)	8,259	5,645	5,722	6,087	5,556	5,972	6,205	6,534	6,800	6,560
EU ETS / GHGI	99.5%	99.3%	98.8%	99.0%	99.2%	99.4%	99.9%	98.7%	99.7%	99.9%

\*The data in this table include fuel combustion emissions (reported under IPCC 1A2f) and process emissions (reported under IPCC sector 2A1) from UK cement kilns.

### A 7.5.2.5 Lime Kilns

**Table A 7.5.7** summarises data given in the EU ETS datasets for the major fuels burnt at lime kilns. Unlike cement kilns, which often burn a variety of fuels, many lime kilns burn just a single fuel, often natural gas. The data below exclude coke oven coke used in lime kilns at soda ash plant since these kilns were not covered by EU ETS until Phase III, and the small number of sites make the data confidential in any case.

**Table A 7.5.7 EU ETS data for Fuels used at Lime Kilns (Emission Factors in kt / Mt for Solid Fuels and kt / Mth for Gases)**

Year	Fuel	% Tier 3	Average Carbon Emission Factor (Tier 3 sites only)
2005	Coal*	-	N/A
2006		-	N/A
2007		34	846.9
2008		79	701.4
2009		100	698.9
2010		100	634.4
2011		100	703.9
2012		100	725.6
2013		100	689.1
2014		100	680.2
2015		100	693.1
2016		100	688.8
2017		100	677.1

\*Coal used in the lime industry in the UK includes a proportion of anthracitic coal, and hence some of these IEFs are notably higher than for coal used in other sectors of UK industry.

The EU ETS data for lime kilns vary across the time series, both in terms of the proportion of emissions based on Tier 3 factors, and in the emission factors themselves. EU ETS based factors are currently used for coal and petroleum coke from 2008 onwards, as the EU ETS data do include all lime kilns burning those fuels and almost all of those data are Tier 3 and hence are regarded as highly reliable.

EU ETS data for natural gas use in the lime industry does cover all installations burning this fuel, however the proportion of emissions based on Tier 3 factors is very low. Therefore the EU ETS emission factors are not used in the UK GHGI, and the emission factors for natural gas continue to be based on the methodology given in Baggott *et al*, 2004.

**Table A 7.5.8** shows implied emission factors for process-related emissions from lime kilns that are used within the UK GHG inventory. The lime industry can be sub-divided into those installations where lime is the primary product, and carbon dioxide is an unwanted by-product; and those installations where both lime and carbon dioxide are utilised. The latter include kilns in the sugar industry (where carbon dioxide is used in the purification stages) and soda ash production (where carbon dioxide is combined with other chemicals to produce sodium carbonate), and in these kilns, the carbon dioxide from decarbonisation of the limestone or dolomite feedstock is assumed to be fully consumed in the process, rather than emitted to atmosphere. **Table A 7.5.8** therefore does not cover these installations. None of the emission factors in EU ETS are Tier 3, so the table shows the overall emission factors for all tiers of data.

**Table A 7.5.8 EU ETS emission factor data for production of lime (kt / Mt lime produced)**

Year	Activity	EU ETS
2005	Lime production	192.4
2006		192.0
2007		192.9
2008		180.5
2009		178.1
2010		182.1
2011		178.9
2012		180.2
2013		178.5
2014		179.5
2015		180.0
2016		178.4
2017		179.4

These factors compare with a theoretical emission factor based on the stoichiometry of the lime manufacturing process of 214 kt / Mt lime, assuming use of pure limestone. We note that the EU ETS factors are all lower than the theoretical emission factor and this is despite some use of dolomitic limestone in the UK industry which would be expected to further increase the emission

factor above the 214 kt/Mt lime factor. The EU ETS data are subject to third party verification, and therefore the emissions data are assumed to be accurate. It is assumed that the reason for this deviation from the theoretical emission factor is due to the production activity data being inflated by either the products containing some proportion of slaked lime (i.e. hydrated product and hence containing a lower proportion of carbon than pure lime) and/or other additives to the lime product which decrease the % carbon content of the lime product.

#### A 7.5.2.6 Other Industrial Combustion

**Table A 7.5.9** summarises EU ETS data for coal, fuel oil and natural gas used by industrial combustion installations.

At first sight, the data for coal looks like it should be reliable enough to be used in the GHGI with 92% or more of emissions based on Tier 3 factors in each year. However, it must be recalled that numerous smaller industrial consumers will not be represented in EU ETS and that the EU ETS data are not fully representative of UK fuels as a whole – see **Section A 7.4** for details. This is also true for EU ETS data for fuel oil and natural gas but here, in addition, very little of the EU ETS data are based on Tier 3 factors. Therefore, none of these data have been used directly in the compilation of the GHGI estimates.

**Table A 7.5.9 EU ETS data for Coal, Fuel Oil and Natural Gas used by Industrial Combustion Plant (Emission Factors in kt / Mt for Coal & Fuel Oil, kt / Mth for Natural Gas)**

Year	Fuel	% Tier 3	Average Carbon Emission Factor	GHGI Carbon Emission Factor
			(Tier 3 sites only)	
2005	Coal	98	607.1	647.8
2006		98	603.0	647.8
2007		99	615.7	662.4
2008		94	598.6	656.6
2009		92	595.4	668.5
2010		88	576.5	674.3
2011		91	589.0	653.4
2012		90	599.2	653.7
2013		95	653.4	653.3
2014		98	654.3	651.2
2015		100	645.8	652.1
2016		100	624.9	650.9
2017		100	647.4	651.2
2005	Fuel oil	48	864.7	879.0
2006		74	865.3	879.0
2007		50	872.3	879.0
2008		35	871.4	879.0
2009		39	871.3	879.0

Year	Fuel	% Tier 3	Average Carbon Emission Factor (Tier 3 sites only)	GHGI Carbon Emission Factor
2010		40	873.0	879.0
2011		51	874.2	879.0
2012		49	875.1	879.0
2013		44	871.3	879.0
2014		48	875.0	879.0
2015		55	872.1	879.0
2016		63	876.2	879.0
2017		65	880.0	879.0
2005	Natural gas	16	1.593	1.477
2006		37	1.470	1.476
2007		42	1.466	1.476
2008		29	1.496	1.475
2009		43	1.499	1.473
2010		40	1.503	1.472
2011		39	1.466	1.470
2012		40	1.469	1.470
2013		43	1.473	1.473
2014		42	1.472	1.472
2015		42	1.480	1.470
2016		41	1.474	1.468

Emission factors can also be derived from EU ETS where a high percentage of Tier 3 analysis is evident, for a number of other minor fuels. Due to the very low number of sites that report data for each fuel type, these EU ETS-derived emission factors are confidential and are not tabulated here. The source/activity combinations for which EU ETS emission factor data are used within the inventory are:

- Other industrial combustion / petroleum coke
- Other industrial combustion / waste solvents
- Other industrial combustion / colliery methane

The EU ETS-derived emission factors for colliery methane for each year (2005-2017) are also applied to all other sources using these fuels.

## **ANNEX 8: UK Domestic Emissions Reporting Requirements**

In addition to the reporting requirements of the UNFCCC, Kyoto Protocol (KP) and EU MMR, UK Greenhouse Gas Inventory statistics are published annually in a Department for Business, Energy and Industrial Strategy National Statistics release<sup>18</sup>. The geographical coverage of these estimates differs from the UNFCCC, KP and EU MMR coverage, with the totals mainly covering emissions from the UK only (i.e. excluding overseas territories and crown dependencies), although progress towards the Kyoto Protocol is still reported.

As part of the Climate Change Act 2008, the UK committed to reducing greenhouse gas emissions by at least 80 percent by 2050 (relative to the base year<sup>19</sup>), with an interim target of reducing greenhouse gas emissions by at least 34 percent by 2020, also relative to the base year. These targets are accompanied by legally binding five-year carbon budgets, which set the trajectory to reaching the targets by placing a restriction on the total amount of greenhouse gases the UK can emit over the five-year period.

Summary tables of the National Statistics release data are presented below. The data are presented in the nine categories used for the UK's National Communications to the UNFCCC and in UK Official Statistics (NC Categories). Note that the scope of emissions used for calculating Carbon Budgets differs slightly from those presented here – for example Carbon Budgets currently exclude NF<sub>3</sub>. The 2019 UK GHG emissions statistical release included an update of the UK's performance against the second carbon budget<sup>20</sup>. Note that performance against the first carbon budget was set in May 2014<sup>21</sup>, updated inventories do not update the first carbon budget or our performance against it.

---

<sup>18</sup> <https://www.gov.uk/government/collections/final-uk-greenhouse-gas-emissions-national-statistics>

<sup>19</sup> Under the Kyoto Protocol, the UK uses 1990 as the base year for carbon dioxide, methane and nitrous oxide emissions, and 1995 as the base year for the fluorinated gases (or F-gases: hydrofluorocarbons, perfluorocarbons and sulphur hexafluoride). To ensure consistency with our international obligations, the same base year for each greenhouse gas is used under the Climate Change Act.

<sup>20</sup> <https://www.gov.uk/government/statistics/final-uk-greenhouse-gas-emissions-national-statistics-1990-2017>

<sup>21</sup> <https://www.gov.uk/government/statistics/final-statement-for-the-first-carbon-budget-period>

**A 8.1 NATIONAL STATISTICS****Table A 8.1.1 Summary table of GHG emissions by NC Category, including net emissions/removals from LULUCF (Mt CO<sub>2</sub>eq) – National Statistics coverage (UK only)**

NC category	1990	1995	2000	2005	2010	2013	2014	2015	2016	2017
Energy supply	277.9	238.0	221.6	231.3	207.2	190.1	165.2	145.3	121.8	112.6
Business	114.0	111.8	115.4	108.9	94.1	88.6	86.6	85.1	81.4	80.1
Transport	128.1	129.7	133.3	136.0	124.5	120.0	121.3	123.5	125.9	125.9
Public	13.5	13.3	12.1	11.2	9.5	9.1	7.8	8.0	8.2	7.8
Residential	80.1	81.6	88.7	85.7	87.5	77.5	64.8	67.4	69.8	66.9
Agriculture	54.0	52.9	50.3	47.9	44.6	44.2	45.6	45.1	45.2	45.6
Industrial processes	59.9	50.8	27.1	20.6	12.6	12.9	13.0	12.7	10.6	10.8
LULUCF <sup>22</sup>	0.3	-1.7	-3.9	-7.1	-9.1	-9.4	-9.6	-9.7	-9.8	-9.9
Waste management	66.6	69.1	62.9	49.0	29.7	23.1	20.9	20.6	20.0	20.3
Total	794.2	745.6	707.5	683.7	600.9	556.2	515.6	497.9	473.1	460.2

**Table A 8.1.2 Summary table of GHG emissions by Gas, including net emissions/removals from LULUCF (Mt CO<sub>2</sub>eq) – National Statistics coverage (UK only)**

Gas	1990	1995	2000	2005	2010	2013	2014	2015	2016	2017
CO <sub>2</sub>	596.3	560.1	558.3	557.9	498.3	464.0	424.9	408.3	385.8	373.2
CH <sub>4</sub>	132.5	125.9	108.4	86.9	63.9	55.5	53.3	52.7	51.1	51.5
N <sub>2</sub> O	48.2	38.6	28.5	24.4	21.3	20.3	20.8	20.3	20.2	20.5
HFCs	14.4	19.1	9.8	13.0	16.4	15.7	15.9	15.9	15.1	14.1
PFCs	1.7	0.6	0.6	0.4	0.3	0.3	0.3	0.3	0.4	0.4
SF <sub>6</sub>	1.3	1.3	1.8	1.1	0.7	0.5	0.5	0.5	0.5	0.5
NF <sub>3</sub>	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
Total	794.360	745.6	707.5	683.7	600.9	556.2	515.6	498.0	473.1	460.2

<sup>22</sup> Land use, land use change and forestry

---

## ANNEX 9: End User Emissions

### A 9.1 INTRODUCTION

This Annex explains the concept of an end user emissions (sometimes also referred to a “final user emissions”, summarises the end user calculation methodology with examples, and contains tables of greenhouse gas emissions according to the end user from 1990 to 2017.

The end user sectoral categories used are consistent with those used in the National Communications (NC) to the UNFCCC. The sectoral categories in the NC are derived from the UNFCCC reporting guidelines on national communications<sup>23</sup>.

The purpose of the end user calculations is to allocate emissions from fuel and electricity producers to the energy users - this allows the emission estimates for a consumer of energy to include the emissions from the production of the fuel or electricity they use.

The UNFCCC does not require end user data to be included in the UK’s National Inventory Report. These data have been included to provide BEIS with information for their policy support needs.

The tables in this Annex present summary data for UK greenhouse gas emissions for the years 1990-2017, inclusive. These data are updated annually to reflect revisions in the methods used to estimate emissions, and the availability of new information within the inventory. These recalculations are applied retrospectively to earlier years to ensure a consistent time series and this accounts for any differences in data published in previous reports.

Emissions presented in this chapter show emissions from the UK only, consistent with the BEIS UK statistical release.

### A 9.2 DEFINITION OF END USERS

The end user<sup>24</sup> or calculations allocate emissions from fuel producers to fuel users. The end user calculation therefore allows estimates to be made of emissions for a consumer of fuel, which also include the emissions from producing the fuel the consumer has used.

The emissions included in the end user categories can be illustrated with an example of two end users - the residential sector and road transport:

- Emissions in the **residential** end user category include:
  1. All direct emissions from domestic premises, for example, from burning gas, coal or oil for space heating.
  2. A portion of indirect emissions used by domestic consumers from: power stations generating electricity; emissions from refineries including refining, storage, flaring and

---

<sup>23</sup> See page 84 of UNFCCC Guidelines contained in FCCC/CP/1999/7 available at: <http://unfccc.int/resource/docs/cop5/07.pdf>

<sup>24</sup> An end user is a consumer of fuel for useful energy. A ‘fuel producer’ is someone who extracts, processes or converts fuels for the end use of end users. Clearly there can be some overlap of these categories but here the fuel uses categories of the UK BEIS publication DUKES are used, which enable a distinction to be made.

extraction; emissions from coal mines (including emissions due to fuel use in the mining industry itself and fugitive emissions of methane from the mines); and emissions from the extraction, storage and distribution of mains gas.


- Emissions in the **road transport** end user category include:
  1. Direct emissions from motor vehicle exhausts.
  2. A portion of indirect emissions from: refineries producing motor fuels, including refining, storage, flaring and extraction of oil; the distribution and supply of motor fuels; and power stations generating the electricity used by electric vehicles.

### A 9.3 OVERVIEW OF THE END USER CALCULATIONS

Fuel and electricity producers also require the use of energy which comes from other producers. Therefore, in the process of reallocating emissions to the end user, emissions are allocated from one to the other and then are reallocated to end users. This circularity results in an iterative approach being used to estimate emissions from categories of end users.

**Figure A 9.1** shows a simplified view of the energy flows in the UK (the fuels used in the greenhouse gas inventory have hundreds of uses). This figure shows that while end users consuming electricity are responsible for a proportion of the emissions from power stations they are also responsible for emissions from collieries, and some of these emissions in turn come from electricity generated in power stations and from refineries.

**Figure A 9.1** Simplified fuel flows for an end user calculation.



---

The approach for estimating end user emissions is summarised in the three steps below:

1. Emissions are calculated for each sector for each fuel.
2. Emissions from fuel and electricity producers are then distributed to those sectors that use the fuel according to the energy content<sup>25</sup> of the fuel they use (these sectors can include other fuel producers). This distribution is based on inventory fuel consumption data and DUKES electricity consumption data.
3. By this stage in the calculation, emissions from end users will have increased and those from fuel and electricity producers will have decreased. The sum of emissions from fuel producers and power stations in a particular year as a percentage of the total emissions is then calculated. If this percentage, for any year, exceeds a predetermined value (In the model used to determine emissions from end users, the value of this percentage can be adjusted. The tables presented later in this Annex were calculated for a convergence at 0.001%) the process continues at Step 2. If this percentage matches or is less than the predetermined value, the calculation is finished.

Convergence occurs as the fuel flows to the end users are much greater than fuel flows amongst the fuel producers.

While a direct solution could possibly be used it was decided to base the calculation on an iterative approach because:

- This can be implemented in the database structures already in existence for the UK greenhouse gas inventory;
- It can handle a wide range of flows and loops that occur without any of the limits that other approaches may incur; and
- The same code will cover all likely situations and will be driven by tabular data stored in the database.

## A 9.4 EXAMPLE END USER CALCULATION

The following example illustrates the methodology used to calculate emissions according to end users. The units in this example are arbitrary.


The example in **Figure A 9.2** has two fuel producers, *power stations* and *collieries*, and three end users, *residential*, *industry* and *commercial*. The following assumptions have been made for simplicity:

- The only fuels used are coal and electricity;
- Coal is the only source of carbon emissions (released from burning coal in power stations to produce electricity and from burning coal in the home for space heating); and
- Commerce uses no coal and so has zero 'direct' emissions.

---

<sup>25</sup> If calorific data for the fuels is not available then the mass of fuel is used instead. This is the case for years prior to 1990.

Figure A 9.2 Fuel use in the example calculation



In **Figure A 9.2**, the tonnes refer to tonnes of coal burnt (black arrows), and the units refer to units of electricity consumed (blue arrows).

In this example the coal extracted by the colliery is burnt in the power station to produce electricity for the end users. Industrial and residential users also directly burn coal. Although the colliery uses electricity produced by the power station, it is not considered to be an end user. The colliery is a ‘fuel producer’ as it is part of the chain that extracts, processes and converts fuels for the end users.

**Table A 9.4.1** summarises the outputs during this example end user calculation.

**Table A 9.4.1 Example of the outputs during an end user calculation**

		Sector					Unallocated emissions as percentage of total emission	Total emission of carbon (tonnes)	
		Colliery	Power Station	Residential	Industrial	Commercial			
Coal use (tonnes)	Mass	100	100,000	50,000	100,000	0			
	Energy content	25,000	25,000,000	12,500,000	25,000,000	0			
Electricity use (arbitrary units)	Energy units	10,000		50,000	100,000	100,000			
	Initial	70	70,000	35,000	70,000	0	40.02	175,070	
Emissions of carbon (tonnes)	Emissions after iteration step	1	2,692	28	48,476	96,951	26,923	1.55	175,070
		2	1	1077	49,020	98,039	26,934	0.62	175,070
		3	41	1	49,227	98,454	27,348	0.02	175,070
		4	0	17	49,235	98,470	27,348	0.01	175,070
		5	1	0	49,238	98,477	27,355	0	175,070
		6	0	0	49,239	98,477	27,355	0	175,070


The initial carbon emissions are 70% of the mass of coal burnt. The emissions from the power stations are distributed to the other sectors by using the factor:

- $(\text{Electricity used by that sector}) / (\text{total electricity used minus own use by power stations})$ ;
- Similarly for the colliery emissions the following factor is used; and
- $(\text{Energy of coal used by that sector}) / (\text{total energy of coal consumed used minus own use by collieries})$ .

At the end of iteration step one, the commerce sector has 26923 tonnes of carbon emissions allocated to it, mainly derived from power stations. Emissions allocated to the residential and industry sectors have also increased over their initial allocations. However collieries and power stations still have some emissions allocated to them (these come from each other) and so the reallocation process is repeated to reduce these allocations to zero – these two sectors are not end users. The total unallocated (in this example, equal to the total emissions from collieries and power stations) falls in each iteration until the emissions are consistently allocated across the sectors. In this example, six iterations are needed to achieve a consistent allocation across the sectors.

The sum of emissions allocated to the sectors (175,070 tonnes of carbon) remains unchanged from the initial allocation to the allocation in the sixth iteration. This check is an important quality control measure to ensure all emissions are accounted for during the end user calculations.

**Figure A 9.3** Comparison of 'direct' and end user emissions of carbon according the sectors considered in the end user example



**Figure A 9.3** compares the quantities of direct and end user carbon emitted from each sector at the end of the end user calculation. The direct emissions of carbon are from the combustion of coal in the sectors. The direct and end user emissions are from two distinct calculations and must be considered independently – in other words, the direct and end user emissions in each sector must not be summed. The sum of all the direct emissions and the sum of the end user emissions, are identical.

There are relatively large direct emissions of carbon from power stations, residential and industry sectors. The end user emissions from the power stations and the colliery are zero because these two sectors are not end users. The carbon emissions from these two sectors have been reallocated to the residential, industrial and commercial sectors. This reallocation means the end user emissions for the residential and industrial sectors are greater than their 'direct' emissions.

## A 9.5 END USER CALCULATION METHODOLOGY FOR THE UK GREENHOUSE GAS INVENTORY

The approach divides fuel user emissions into 8 categories (see column 1 of **Table A 9.5.1**). For each of these groups, source categories are distributed by the total energy consumption of a group of fuels. For example, for the coal group, the emissions of four source categories are distributed to end users according to the energy use of anthracite and coal combined.

Table A 9.5.1 Sources reallocated to end users and the fuels used

End user group	Emission sources to be reallocated to end users	Fuels used for redistribution
1. Coke	Gasification processes	Coke
	Coke production	Blast furnace gas
	Iron and steel – flaring	
2. Coal	Closed Coal Mines	Coal
	Coal storage and transport	Anthracite
	Collieries – combustion	
	Deep-mined coal	
	Open-cast coal	
3. Natural gas	Gas leakage	Natural gas
	Gas production	
	Upstream Gas Production – flaring	
	Upstream Gas Production – fuel combustion	
	Upstream Gas Production – Gas terminal storage	
	Upstream Gas Production – Offshore Well Testing	
	Upstream Gas Production - Onshore Oil Loading	
	Upstream Gas Production – process emissions	
	Upstream Gas Production – venting	
	Upstream Gas production – combustion at gas separation plant	
4. Electricity	Nuclear fuel production	Electricity
	Power stations	
	Autogeneration – exported to grid	
	Power stations – FGD	

End user group	Emission sources to be reallocated to end users	Fuels used for redistribution
5. Petroleum	Upstream Oil Production – fuel combustion	Aviation spirit
	Upstream Oil Production –flaring	Aviation turbine fuel
	Upstream Oil Production –venting	Biodiesel
	Upstream Oil Production – Offshore Oil Loading	Bioethanol
	Upstream Oil Production – Offshore Well Testing	Burning oil
	Upstream Oil Production – Oil terminal storage	Burning oil (premium)
	Upstream Oil Production – Onshore Oil Loading	DERV
	Upstream Oil Production – process emissions	Fuel oil
	Petrol stations – petrol delivery	Gas oil
	Petrol stations – vehicle refuelling	LNG
	Petrol terminals – storage	LPG
	Petrol terminals – tanker loading	Naphtha
	Petroleum processes	OPG
	Refineries – combustion	Petrol
	Refineries – drainage	Petroleum coke
	Refineries – flares	Refinery miscellaneous
	Refineries – general	Vaporising oil
	Refineries – process	
	Refineries – road/rail loading	
	Refineries – tankage	
Sea going vessel loading		
Ship purging		
6. Solid Smokeless Fuels	Solid Smokeless fuel production	Solid Smokeless Fuels
8. Charcoal	Charcoal production	Charcoal

Comments on the calculation methodology used to allocate emissions according to the end users are listed below:

- Emissions are allocated to end users on the basis of the proportion of the total energy produced that is used by a given sector. This approach is followed to allow for sectors such as petroleum where different products are made in a refinery;

- 
- Some emissions are allocated to an “exports” category. This is for emissions within the UK from producing fuels, (for example from a refinery or coal mine), which are subsequently exported or sent to bunkers for use outside the UK. Therefore these emissions are part of the UK inventory even if the use of the fuel produces emissions that cannot be included in the UK inventory because it takes place outside the UK;
  - No allowance is made for the emission from the production of fuels or electricity outside the UK that are subsequently imported;
  - Some of the output of a refinery is not used as a fuel but used as feedstock or lubricants. This is not currently treated separately and the emissions from their production (which are small) are allocated to users of petroleum fuels. This is partly due to lack of data in the database used to calculate the inventory, and partly due to the lack of a clear, transparent way of separating emissions from the production of fuels and from the production of non-fuel petroleum products; and
  - End user emissions are estimated for aviation in four categories: domestic take-off and landing, international take-off and landing, domestic cruise and international cruise. This enables both IPCC and UNECE categories to be estimated from the same end user calculation.

Our exact mapping of end user emissions to IPCC categories is shown in **Table A 9.5.2**. The NAEI source sectors and activity names are also shown, as it is necessary to subdivide some IPCC categories. This classification has been used to generate the end user tables for the greenhouse gases given in this section. As this table is for end users, no fuel producers are included in the table.

**Table A 9.5.2 End user category, IPCC sectors, and NAEI source names and activity names used in the emission calculation**

National Communication Category	IPCC Sector	Source Name	Activity Name
Agriculture	1A4ci_Agriculture/Forestry/Fishing:Stationary	Agriculture - stationary combustion	Burning oil
			Coal
			Fuel oil
			Natural gas
			Straw
	1A4cii_Agriculture/Forestry/Fishing:Off-road	Agriculture - mobile machinery	Gas oil
			Petrol
	2D1_Lubricant_Use	Agricultural engines	Lubricants
	3A1a	Enteric	Dairy - Dairy Cows
	3A1b	Enteric	Other cattle - Beef females for slaughter
			Other cattle - Bulls for breeding
			Other cattle - Cereal fed bull
			Other cattle - Cows
			Other cattle - Dairy Calves Female
			Other cattle - Dairy In Calf Heifers
Other cattle - Dairy Replacements Female			
Other cattle - Heifers for breeding			

National Communication Category	IPCC Sector	Source Name	Activity Name
	3A2	Enteric	Other cattle - Steers
			Sheep - Ewe
			Sheep - Lamb
	3A3	Enteric	Sheep - Ram
			Pig - Boar
			Pig - Fattening Pig < 20 kg
			Pig - Fattening Pig > 80 kg
			Pig - Fattening Pig 20 to 80 kg
			Pig - Gilt
	3A4	Enteric	Pig - Sow
			Deer
			Goats
	3B11a	Excreta	Horses
			Dairy - Dairy Cows
	3B11b 3B12 3B13	Excreta	Managed Manure
			Dairy - Dairy Cows
			Other cattle - Beef females for slaughter
			Other cattle - Bulls for breeding
			Other cattle - Cereal fed bull

National Communication Category	IPCC Sector	Source Name	Activity Name
			Other cattle - Cows
			Other cattle - Dairy Calves Female
			Other cattle - Dairy In Calf Heifers
			Other cattle - Dairy Replacements Female
			Other cattle - Heifers for breeding
			Other cattle - Steers
		Managed Manure	Other cattle - Beef females for slaughter
			Other cattle - Bulls for breeding
			Other cattle - Cereal fed bull
			Other cattle - Cows
			Other cattle - Dairy Calves Female
			Other cattle - Dairy In Calf Heifers
			Other cattle - Dairy Replacements Female
			Other cattle - Heifers for breeding
		Excreta	Sheep - Ewe
			Sheep - Lamb

National Communication Category	IPCC Sector	Source Name	Activity Name
		Managed Manure	Sheep - Ram
			Sheep - Ewe
			Sheep - Lamb
		Excreta	Pig - Boar
			Pig - Fattening Pig < 20 kg
			Pig - Fattening Pig > 80 kg
			Pig - Fattening Pig 20 to 80 kg
			Pig - Gilt
			Pig - Sow
		Managed Manure	Pig - Boar
			Pig - Fattening Pig < 20 kg
			Pig - Fattening Pig > 80 kg
			Pig - Fattening Pig 20 to 80 kg
	Pig - Gilt		
	Pig - Sow		
	3B14	Excreta	Poultry - Breeding flock
			Poultry - Broilers
			Poultry - Ducks
			Poultry - Geese

National Communication Category	IPCC Sector	Source Name	Activity Name
			Poultry - Growing Pullets
			Poultry - Laying Hens
			Poultry - Other
			Poultry - Turkeys
		Managed Manure	Poultry - Breeding flock
			Poultry - Broilers
			Poultry - Ducks
			Poultry - Geese
			Poultry - Growing Pullets
			Poultry - Laying Hens
			Poultry - Other
			Poultry - Turkeys
		Wastes	Deer
			Goats
			Horses
		3B21a	Dairy - Dairy Cows - Direct
3B21b	Other cattle - Beef females for slaughter - Direct	Housing	
	Other cattle - Bulls for breeding - Direct	Housing	
	Other cattle - Cereal fed bull - Direct	Housing	

National Communication Category	IPCC Sector	Source Name	Activity Name
		Other cattle - Cows - Direct	Housing
		Other cattle - Dairy Calves Female - Direct	Housing
		Other cattle - Dairy In Calf Heifers - Direct	Housing
		Other cattle - Dairy Replacements Female - Direct	Housing
		Other cattle - Heifers for breeding - Direct	Housing
		Other cattle - Steers - Direct	Housing
	3B22	Sheep - Ewe - Direct	Storage
		Sheep - Lamb - Direct	Storage
	3B23	Pig - Boar - Direct	Housing
		Pig - Fattening Pig < 20 kg - Direct	Housing
		Pig - Fattening Pig > 80 kg - Direct	Housing
		Pig - Fattening Pig 20 to 80 kg - Direct	Housing
		Pig - Gilt - Direct	Housing
		Pig - Sow - Direct	Housing
	3B24	Deer Wastes - Direct	Excreta N managed as manure
		Goats Wastes - Direct	Excreta N managed as manure
		Horses Wastes - Direct	Excreta N managed as manure
		Poultry - Breeding Flock - Direct	Housing
		Poultry - Broilers - Direct	Housing

National Communication Category	IPCC Sector	Source Name	Activity Name	
		Poultry - Ducks - Direct	Housing	
		Poultry - Geese - Direct	Housing	
		Poultry - Growing Pullets - Direct	Housing	
		Poultry - Laying Hens - Direct	Housing	
		Poultry - Other - Direct	Housing	
		Poultry - Turkeys - Direct	Housing	
	3B25	Dairy - Dairy Cows - Indirect Deposition		Housing
				Storage
				Yarding
		Dairy - Dairy Cows - Indirect Leach	Storage	
		Deer Wastes - Indirect Leaching	N leached from manure management	
		Deer Wastes - Indirect Volatilisation	N volatilised from manure management	
		Goats Wastes - Indirect Leaching	N leached from manure management	
		Goats Wastes - Indirect Volatilisation	N volatilised from manure management	
		Horses Wastes - Indirect Leaching	N leached from manure management	
		Horses Wastes - Indirect Volatilisation	N volatilised from manure management	
	Housing			

National Communication Category	IPCC Sector	Source Name	Activity Name
		Other cattle - Beef females for slaughter - Indirect Deposition	Storage
			Yarding
		Other cattle - Beef females for slaughter - Indirect Leach	Storage
		Other cattle - Bulls for breeding - Indirect Deposition	Housing
			Storage
			Yarding
		Other cattle - Bulls for breeding - Indirect Leach	Storage
		Other cattle - Cereal fed bull - Indirect Deposition	Housing
			Storage
			Yarding
		Other cattle - Cereal fed bull - Indirect Leach	Storage
		Other cattle - Cows - Indirect Deposition	Housing
			Storage
			Yarding
		Other cattle - Cows - Indirect Leach	Storage
Other cattle - Dairy Calves Female - Indirect Deposition	Housing		
	Storage		
	Yarding		
Other cattle - Dairy Calves Female - Indirect Leach	Storage		

National Communication Category	IPCC Sector	Source Name	Activity Name
		Other cattle - Dairy In Calf Heifers - Indirect Deposition	Housing
			Storage
			Yarding
		Other cattle - Dairy In Calf Heifers - Indirect Leach	Storage
		Other cattle - Dairy Replacements Female - Indirect Deposition	Housing
			Storage
			Yarding
		Other cattle - Dairy Replacements Female - Indirect Leach	Storage
		Other cattle - Heifers for breeding - Indirect Deposition	Housing
			Storage
			Yarding
		Other cattle - Heifers for breeding - Indirect Leach	Storage
		Other cattle - Steers - Indirect Deposition	Housing
			Storage
			Yarding
Other cattle - Steers - Indirect Leach	Storage		
Pig - Boar - Indirect Deposition	Housing		
	Storage		
Pig - Boar - Indirect Leach	Storage		

National Communication Category	IPCC Sector	Source Name	Activity Name
		Pig - Fattening Pig < 20 kg - Indirect Deposition	Housing
			Storage
		Pig - Fattening Pig < 20 kg - Indirect Leach	Storage
		Pig - Fattening Pig > 80 kg - Indirect Deposition	Housing
			Storage
		Pig - Fattening Pig > 80 kg - Indirect Leach	Storage
		Pig - Fattening Pig 20 to 80 kg - Indirect Deposition	Housing
			Storage
		Pig - Fattening Pig 20 to 80 kg - Indirect Leach	Storage
		Pig - Gilt - Indirect Deposition	Housing
			Storage
		Pig - Gilt - Indirect Leach	Storage
		Pig - Sow - Indirect Deposition	Housing
			Storage
		Pig - Sow - Indirect Leach	Storage
		Poultry - Breeding Flock - Indirect Deposition	Housing
Storage			
Poultry - Breeding Flock - Indirect Leach	Storage		
Poultry - Broilers - Indirect Deposition	Housing		

National Communication Category	IPCC Sector	Source Name	Activity Name
			Storage
		Poultry - Broilers - Indirect Leach	Storage
		Poultry - Ducks - Indirect Deposition	Housing
			Storage
		Poultry - Ducks - Indirect Leach	Storage
		Poultry - Geese - Indirect Deposition	Housing
			Storage
		Poultry - Geese - Indirect Leach	Storage
		Poultry - Growing Pullets - Indirect Deposition	Housing
			Storage
		Poultry - Growing Pullets - Indirect Leach	Storage
		Poultry - Laying Hens - Indirect Deposition	Housing
			Storage
		Poultry - Laying Hens - Indirect Leach	Storage
		Poultry - Other - Indirect Deposition	Housing
			Storage
		Poultry - Other - Indirect Leach	Storage
		Poultry - Turkeys - Indirect Deposition	Housing
			Storage

National Communication Category	IPCC Sector	Source Name	Activity Name
		Poultry - Turkeys - Indirect Leach	Storage
		Sheep - Ewe - Indirect Deposition	Housing
			Storage
		Sheep - Ewe - Indirect Leach	Storage
		Sheep - Lamb - Indirect Deposition	Housing
			Storage
	Sheep - Lamb - Indirect Leach	Storage	
	3D11	Arable - Direct	Ammonium Nitrate Application
			Ammonium Sulphate and Diammonium Phosphate Application
			Calcium Ammonium Nitrate Application
			Other Nitrogen Including Compounds Application
			Urea Ammonium Nitrate Application
			Urea Application
		Grass - Direct	Ammonium Nitrate Application
			Ammonium Sulphate and Diammonium Phosphate Application
Calcium Ammonium Nitrate Application			

National Communication Category	IPCC Sector	Source Name	Activity Name
			Other Nitrogen Including Compounds Application
			Urea Ammonium Nitrate Application
			Urea Application
	3D12a	Dairy - Dairy Cows - Direct	Spreading
		FAM - direct	Manure N applied to soil
		Goats FAM - Direct	Manure N applied to soil
		Horses FAM - direct	Manure N applied to soil
		Other cattle - Beef females for slaughter - Direct	Spreading
		Other cattle - Bulls for breeding - Direct	Spreading
		Other cattle - Cereal fed bull - Direct	Spreading
		Other cattle - Cows - Direct	Spreading
		Other cattle - Dairy Calves Female - Direct	Spreading
		Other cattle - Dairy In Calf Heifers - Direct	Spreading
		Other cattle - Dairy Replacements Female - Direct	Spreading
		Other cattle - Heifers for breeding - Direct	Spreading
		Other cattle - Steers - Direct	Spreading
		Pig - Boar - Direct	Spreading
Pig - Fattening Pig < 20 kg - Direct	Spreading		

National Communication Category	IPCC Sector	Source Name	Activity Name	
		Pig - Fattening Pig > 80 kg - Direct	Spreading	
		Pig - Fattening Pig 20 to 80 kg - Direct	Spreading	
		Pig - Gilt - Direct	Spreading	
		Pig - Sow - Direct	Spreading	
		Poultry - Breeding Flock - Direct	Spreading	
		Poultry - Broilers - Direct	Spreading	
		Poultry - Ducks - Direct	Spreading	
		Poultry - Geese - Direct	Spreading	
		Poultry - Growing Pullets - Direct	Spreading	
		Poultry - Laying Hens - Direct	Spreading	
		Poultry - Other - Direct	Spreading	
		Poultry - Turkeys - Direct	Spreading	
		Sheep - Ewe - Direct	Spreading	
		Sheep - Lamb - Direct	Spreading	
		3D12b	Sewage Sludge Application - Direct	Sewage sludge N applied to soil
		3D13	All Horses PRP - Direct	Excreta N returned at grazing
			Dairy - Dairy Cows - Direct	Grazing
Deer PRP - Direct	Excreta N returned at grazing			
Goats PRP - Direct	Excreta N returned at grazing			

National Communication Category	IPCC Sector	Source Name	Activity Name
		Other cattle - Beef females for slaughter - Direct	Grazing
		Other cattle - Bulls for breeding - Direct	Grazing
		Other cattle - Cereal fed bull - Direct	Grazing
		Other cattle - Cows - Direct	Grazing
		Other cattle - Dairy Calves Female - Direct	Grazing
		Other cattle - Dairy In Calf Heifers - Direct	Grazing
		Other cattle - Dairy Replacements Female - Direct	Grazing
		Other cattle - Heifers for breeding - Direct	Grazing
		Other cattle - Steers - Direct	Grazing
		Pig - Boar - Direct	Grazing
		Pig - Fattening Pig < 20 kg - Direct	Grazing
		Pig - Fattening Pig > 80 kg - Direct	Grazing
		Pig - Fattening Pig 20 to 80 kg - Direct	Grazing
		Pig - Gilt - Direct	Grazing
		Pig - Sow - Direct	Grazing
		Poultry - Breeding Flock - Direct	Grazing
		Poultry - Broilers - Direct	Grazing
		Poultry - Ducks - Direct	Grazing
		Poultry - Geese - Direct	Grazing

National Communication Category	IPCC Sector	Source Name	Activity Name	
		Poultry - Growing Pullets - Direct	Grazing	
		Poultry - Laying Hens - Direct	Grazing	
		Poultry - Other - Direct	Grazing	
		Poultry - Turkeys - Direct	Grazing	
		Sheep - Ewe - Direct	Grazing	
		Sheep - Lamb - Direct	Grazing	
		Sheep - Ram - Direct	Grazing	
	3D14	Arable - Direct		Ammonium Nitrate Residue
				Ammonium Sulphate and Diammonium Phosphate Residue
				Calcium Ammonium Nitrate Residue
				Other Nitrogen Including Compounds Residue
				Urea Ammonium Nitrate Residue
				Urea Residue
		Grass - Direct		Ammonium Nitrate Residue
	Ammonium Sulphate and Diammonium Phosphate Residue			
	Calcium Ammonium Nitrate Residue			
	No Nitrogen Fertiliser Applied Residue			

National Communication Category	IPCC Sector	Source Name	Activity Name
			Other Nitrogen Including Compounds Residue
			Urea Ammonium Nitrate Residue
			Urea Residue
	3D15	Cropland management	Mineralisation
	3D16	Managed Histosols	Land area
	3D21	Arable - Indirect Deposition	Ammonium Nitrate Application
			Ammonium Sulphate and Diammonium Phosphate Application
			Calcium Ammonium Nitrate Application
			Other Nitrogen Including Compounds Application
			Urea Ammonium Nitrate Application
			Urea Application
		Dairy - Dairy Cows - Indirect Deposition	Grazing
			Spreading
		Deer FAM - Indirect Volatilisation	N volatilised from manure applications
		Deer PRP - Indirect Volatilisation	N volatilised from grazing excreta
		Goats FAM - Indirect Volatilisation	N volatilised from manure applications

National Communication Category	IPCC Sector	Source Name	Activity Name
		Goats PRP - Indirect Volatilisation	N volatilised from grazing excreta
		Grass - Indirect Deposition	Ammonium Nitrate Application
			Ammonium Sulphate and Diammonium Phosphate Application
			Calcium Ammonium Nitrate Application
			Other Nitrogen Including Compounds Application
			Urea Ammonium Nitrate Application
			Urea Application
		Horses FAM - Indirect Volatilisation	N volatilised from manure applications
		Horses PRP - Indirect Volatilisation	N volatilised from grazing excreta
		Other cattle - Beef females for slaughter - Indirect Deposition	Grazing
			Spreading
		Other cattle - Bulls for breeding - Indirect Deposition	Grazing
			Spreading
		Other cattle - Cereal fed bull - Indirect Deposition	Grazing
			Spreading
		Other cattle - Cows - Indirect Deposition	Grazing
			Spreading

National Communication Category	IPCC Sector	Source Name	Activity Name
		Other cattle - Dairy Calves Female - Indirect Deposition	Grazing
			Spreading
		Other cattle - Dairy In Calf Heifers - Indirect Deposition	Grazing
			Spreading
		Other cattle - Dairy Replacements Female - Indirect Deposition	Grazing
			Spreading
		Other cattle - Heifers for breeding - Indirect Deposition	Grazing
			Spreading
		Other cattle - Steers - Indirect Deposition	Grazing
			Spreading
		Pig - Boar - Indirect Deposition	Grazing
			Spreading
		Pig - Fattening Pig < 20 kg - Indirect Deposition	Grazing
			Spreading
		Pig - Fattening Pig > 80 kg - Indirect Deposition	Grazing
			Spreading
		Pig - Fattening Pig 20 to 80 kg - Indirect Deposition	Grazing
			Spreading
Pig - Gilt - Indirect Deposition	Grazing		

National Communication Category	IPCC Sector	Source Name	Activity Name
			Spreading
		Pig - Sow - Indirect Deposition	Grazing
			Spreading
		Poultry - Breeding Flock - Indirect Deposition	Grazing
			Spreading
		Poultry - Broilers - Indirect Deposition	Grazing
			Spreading
		Poultry - Ducks - Indirect Deposition	Grazing
			Spreading
		Poultry - Geese - Indirect Deposition	Grazing
			Spreading
		Poultry - Growing Pullets - Indirect Deposition	Grazing
			Spreading
		Poultry - Laying Hens - Indirect Deposition	Grazing
			Spreading
		Poultry - Other - Indirect Deposition	Grazing
			Spreading
		Poultry - Turkeys - Indirect Deposition	Grazing
			Spreading

National Communication Category	IPCC Sector	Source Name	Activity Name
		Sewage Sludge Application - Indirect Volatilisation	N volatilised from sewage sludge applications
		Sheep - Ewe - Indirect Deposition	Grazing
			Spreading
		Sheep - Lamb - Indirect Deposition	Grazing
			Spreading
		Sheep - Ram - Indirect Deposition	Grazing
		3D22	Arable - Indirect Leach
	Ammonium Sulphate and Diammonium Phosphate Application		
	Calcium Ammonium Nitrate Application		
	Other Nitrogen Including Compounds Application		
	Urea Ammonium Nitrate Application		
	Urea Application		
		Arable - Residue Indirect Leach	Ammonium Nitrate Application
			Ammonium Sulphate and Diammonium Phosphate Application
Calcium Ammonium Nitrate Application			

National Communication Category	IPCC Sector	Source Name	Activity Name
			Other Nitrogen Including Compounds Application
			Urea Ammonium Nitrate Application
			Urea Application
		Cropland management	Mineralisation - indirect leach
		Dairy - Dairy Cows - Indirect Leach	Grazing
			Spreading
		Deer FAM - Indirect Leaching	N leached from manure applications
		Deer PRP - Indirect Leaching	N leached from grazing excreta
		Goats FAM - Indirect Leaching	N leached from manure applications
		Goats PRP - Indirect Leaching	N leached from grazing excreta
		Grass - Indirect Leach	Ammonium Nitrate Application
			Ammonium Sulphate and Diammonium Phosphate Application
			Calcium Ammonium Nitrate Application
			Other Nitrogen Including Compounds Application
			Urea Ammonium Nitrate Application
			Urea Application
		Grass - Residue Indirect Leach	Ammonium Nitrate Application

National Communication Category	IPCC Sector	Source Name	Activity Name
			Ammonium Sulphate and Diammonium Phosphate Application
			Calcium Ammonium Nitrate Application
			No Nitrogen Fertiliser Applied
			Other Nitrogen Including Compounds Application
			Urea Ammonium Nitrate Application
			Urea Application
		Horses FAM - Indirect Leaching	N leached from manure applications
		Horses PRP - Indirect Leaching	N leached from grazing excreta
		Other cattle - Beef females for slaughter - Indirect Leach	Grazing
			Spreading
		Other cattle - Bulls for breeding - Indirect Leach	Grazing
			Spreading
		Other cattle - Cereal fed bull - Indirect Leach	Grazing
			Spreading
		Other cattle - Cows - Indirect Leach	Grazing
			Spreading
Other cattle - Dairy Calves Female - Indirect Leach	Grazing		

National Communication Category	IPCC Sector	Source Name	Activity Name
			Spreading
		Other cattle - Dairy In Calf Heifers - Indirect Leach	Grazing
			Spreading
		Other cattle - Dairy Replacements Female - Indirect Leach	Grazing
			Spreading
		Other cattle - Heifers for breeding - Indirect Leach	Grazing
			Spreading
		Other cattle - Steers - Indirect Leach	Grazing
			Spreading
		Pig - Boar - Indirect Leach	Grazing
			Spreading
		Pig - Fattening Pig < 20 kg - Indirect Leach	Grazing
			Spreading
		Pig - Fattening Pig > 80 kg - Indirect Leach	Grazing
			Spreading
		Pig - Fattening Pig 20 to 80 kg - Indirect Leach	Grazing
			Spreading
		Pig - Gilt - Indirect Leach	Grazing
			Spreading

National Communication Category	IPCC Sector	Source Name	Activity Name
		Pig - Sow - Indirect Leach	Grazing
			Spreading
		Poultry - Breeding Flock - Indirect Leach	Grazing
			Spreading
		Poultry - Broilers - Indirect Leach	Grazing
			Spreading
		Poultry - Ducks - Indirect Leach	Grazing
			Spreading
		Poultry - Geese - Indirect Leach	Grazing
			Spreading
		Poultry - Growing Pullets - Indirect Leach	Grazing
			Spreading
		Poultry - Laying Hens - Indirect Leach	Grazing
			Spreading
		Poultry - Other - Indirect Leach	Grazing
			Spreading
		Poultry - Turkeys - Indirect Leach	Grazing
			Spreading

National Communication Category	IPCC Sector	Source Name	Activity Name	
		Sewage Sludge Application - Indirect Leaching	N leached from sewage sludge applications	
		Sheep - Ewe - Indirect Leach	Grazing	
			Spreading	
		Sheep - Lamb - Indirect Leach	Grazing	
	Spreading			
		Sheep - Ram - Indirect Leach	Grazing	
		3F11_Field_burning	Field burning	Wheat residue
		3F12_Field_burning	Field burning	Barley residue
		3F14_Field_burning	Field burning	Oats residue
		3F5_Field_burning	Field burning	Linseed residue
		3G1_Liming - limestone	Liming	Limestone
		3G2_Liming - dolomite	Liming	Dolomite
		3H	Fertiliser Application	Urea Application
	non-IPCC	Agriculture - stationary combustion	Electricity	
Business	1A1ai_Public_Electricity&Heat_Production	Autogenerators	Biogas	
	1A2a_Iron_and_steel	Blast furnaces	Blast furnace gas	
			Coke oven gas	
			LPG	

National Communication Category	IPCC Sector	Source Name	Activity Name
			Natural gas
		Iron and steel - combustion plant	Blast furnace gas
			Coal
			Coke
			Coke oven gas
			Fuel oil
			Gas oil
			LPG
	Natural gas		
	1A2b_Non-Ferrous_Metals	Autogeneration - exported to grid	Coal
		Autogenerators	Coal
		Non-Ferrous Metal (combustion)	Coal
			Fuel oil
			Gas oil
			Natural gas
	1A2c_Chemicals	Chemicals (combustion)	Coal
			Fuel oil
			Gas oil
Natural gas			

National Communication Category	IPCC Sector	Source Name	Activity Name
	1A2d_Pulp_Paper_Print	Pulp, Paper and Print (combustion)	Coal
			Fuel oil
			Gas oil
			Natural gas
	1A2e_food_processing_beverages_and_tobacco	Food & drink, tobacco (combustion)	Coal
			Fuel oil
			Gas oil
			Natural gas
	1A2f_Non-metallic_minerals	Cement production - combustion	Coal
			Fuel oil
			Gas oil
			Natural gas
			Petroleum coke
			Scrap tyres
			Waste
Waste oils			
Waste solvent			
Lime production - non decarbonising		Coal	
	Coke		

National Communication Category	IPCC Sector	Source Name	Activity Name	
			Natural gas	
		Other industrial combustion	Scrap tyres	
	1A2gvii_Off-road_vehicles_and_other_machinery	Industrial off-road mobile machinery		DERV
				Gas oil
				Petrol
	1A2gviii_Other_manufacturing_industries_and_construction	Other industrial combustion	Autogeneration - exported to grid	Natural gas
			Autogenerators	Natural gas
				Biomass
				Burning oil
				Coal
				Coke
				Coke oven gas
				Colliery methane
				Fuel oil
				Gas oil
				LPG
	Lubricants			
	Natural gas			
	OPG			

National Communication Category	IPCC Sector	Source Name	Activity Name
			Petroleum coke
			SSF
			Waste solvent
			Wood
	1A4ai_Commercial/Institutional	Miscellaneous industrial/commercial combustion	Coal
			Fuel oil
			Gas oil
			Landfill gas
			MSW
			Natural gas
	2B1_Chemical_Industry:Ammonia_production	Ammonia production - combustion	Natural gas
	2B8a_Methanol_production	Methanol production – combustion	Natural gas
	2B8g_Petrochemical_and_carbon_black_production:Other	Chemicals (combustion)	OPG
	2C1b_Pig_iron	Blast furnaces	Coal
	2D1_Lubricant_Use	Industrial engines	Lubricants
	2D4_Other_NEU	Non Energy Use: petroleum coke	Petroleum coke
2E1_Integrated_circuit_or_semiconductor	Electronics - HFC	Non-fuel combustion	
	Electronics - NF3	Non-fuel combustion	

National Communication Category	IPCC Sector	Source Name	Activity Name
	2F1a_Commercial_refrigeration	Commercial Refrigeration	Refrigeration and Air Conditioning - Disposal
			Refrigeration and Air Conditioning - Lifetime
			Refrigeration and Air Conditioning - Manufacture
	2F1b_Domestic_refrigeration	Domestic Refrigeration	Refrigeration and Air Conditioning - Disposal
			Refrigeration and Air Conditioning - Lifetime
			Refrigeration and Air Conditioning - Manufacture
	2F1c_Industrial_refrigeration	Industrial Refrigeration	Refrigeration and Air Conditioning - Disposal
			Refrigeration and Air Conditioning - Lifetime
			Refrigeration and Air Conditioning - Manufacture
	2F1d_Transport_refrigeration	Refrigerated Transport	Refrigeration and Air Conditioning - Disposal
			Refrigeration and Air Conditioning - Lifetime
			Refrigeration and Air Conditioning - Manufacture

National Communication Category	IPCC Sector	Source Name	Activity Name
	2F1e_Mobile_air_conditioning	Mobile Air Conditioning	Refrigeration and Air Conditioning - Disposal
			Refrigeration and Air Conditioning - Lifetime
			Refrigeration and Air Conditioning - Manufacture
	2F1f_Stationary_air_conditioning	Stationary Air Conditioning	Refrigeration and Air Conditioning - Disposal
			Refrigeration and Air Conditioning - Lifetime
			Refrigeration and Air Conditioning - Manufacture
	2F2a_Closed_foam_blowing_agents	Foams	Non-fuel combustion
		Foams HFCs for the 2006 GLs	Non-fuel combustion
	2F2b_Open_foam_blowing_agents	One Component Foams	Non-fuel combustion
	2F3_Fire_Protection	Firefighting	Non-fuel combustion
	2F5_Solvents	Precision cleaning - HFC	Non-fuel combustion
	2F6b_Other_Applications:Contained-Refrigerant_containers	Refrigerant containers	Non-fuel combustion
2F6b_Other_Applications:Contained-Refrigerant_Processing	F-gas handling	Non-fuel combustion	
2G1_Electrical_equipment	Electrical insulation	Non-fuel combustion	

National Communication Category	IPCC Sector	Source Name	Activity Name
	2G2_Military_applications	AWACS	Non-fuel combustion
	2G2_Particle_accelerators	Particle accelerators	Non-fuel combustion
	2G2e_Electronics_and_shoes	Electronics - PFC	Non-fuel combustion
		Electronics - SF6	Non-fuel combustion
		Sporting goods	Non-fuel combustion
	2G2e_Tracer_gas	SF6 used as a tracer gas	Non-fuel combustion
	2G3a_Medical_applications	N2O use as an anaesthetic	Population
	5C2.2b_Non-biogenic:Other	Accidental fires - other buildings	Mass burnt
	non-IPCC	Chemicals (combustion)	Electricity
		Food & drink, tobacco (combustion)	Electricity
		Iron and steel - combustion plant	Electricity
		Miscellaneous industrial/commercial combustion	Electricity
		Non-Ferrous Metal (combustion)	Electricity
		Other industrial combustion	Electricity
Pulp, Paper and Print (combustion)		Electricity	
Energy Supply	1A1ai_Public_Electricity&Heat_Production	Power stations	Burning oil
			Coal
			Fuel oil
			Gas oil

National Communication Category	IPCC Sector	Source Name	Activity Name
			Natural gas
			Petroleum coke
	1A1b_Petroleum_Refining	Refineries - combustion	Natural gas
	1A1ci_Manufacture_of_solid_fuels	Coke production	Natural gas
		Solid smokeless fuel production	Coke
	1A1cii_Oil_and_gas_extraction	Upstream Gas Production - fuel combustion	Gas oil
		Upstream oil and gas production - combustion at gas separation plant	LPG
			OPG
	1A1ciii_Other_energy_industries	Upstream Oil Production - fuel combustion	Natural gas
		Collieries - combustion	Natural gas
		Gas production	LPG
	1B1b_Solid_Fuel_Transformation	Nuclear fuel production	Natural gas
		Coke production	Coal
	non-IPCC	Coke production	Coal
			Petroleum coke
		Collieries - combustion	Electricity
Exports	Aviation_Bunkers	Gas production	Electricity
		Refineries - combustion	Electricity
		Aircraft - international cruise	Aviation spirit

National Communication Category	IPCC Sector	Source Name	Activity Name
			Aviation turbine fuel
		Aircraft - international take off and landing	Aviation spirit
			Aviation turbine fuel
		Aircraft between UK and Bermuda - Cruise	Aviation turbine fuel
		Aircraft between UK and Bermuda - TOL	Aviation turbine fuel
		Aircraft between UK and CDs - Cruise	Aviation spirit
			Aviation turbine fuel
		Aircraft between UK and CDs - TOL	Aviation spirit
			Aviation turbine fuel
		Aircraft between UK and Gibraltar - Cruise	Aviation spirit
			Aviation turbine fuel
		Aircraft between UK and Gibraltar - TOL	Aviation spirit
			Aviation turbine fuel
		Aircraft between UK and other OTs (excl Gib. and Bermuda) - Cruise	Aviation turbine fuel
	Aircraft between UK and other OTs (excl Gib. and Bermuda) - TOL	Aviation turbine fuel	
Marine_Bunkers	Shipping - international IPCC definition	Fuel oil	
		Gas oil	
	Shipping between UK and Bermuda	Fuel oil	

National Communication Category	IPCC Sector	Source Name	Activity Name	
		Shipping between UK and CDs	Fuel oil	
			Gas oil	
		Shipping between UK and Gibraltar	Fuel oil	
		Shipping between UK and OTs (excl. Gib and Bermuda)	Fuel oil	
	non-IPCC	Exports		Aviation turbine fuel
				Burning oil
				Coke
				DERV
				Electricity
				Fuel oil
				Petrol
	SSF			
	Industrial Process	2A1_Cement_Production	Cement - decarbonising	Clinker production
2A2_Lime_Production		Lime production - decarbonising	Limestone	
2A3_Glass_production		Glass - general	Dolomite	
			Limestone	
			Soda ash	
2A4a_Other_process_uses_of_carbonates:ceramics		Brick manufacture - all types	Bricks	
			Brick manufacture - Fletton	Fletton bricks

National Communication Category	IPCC Sector	Source Name	Activity Name
	2B1_Ammonia_Production	Ammonia production - feedstock use of gas	Natural gas
	2B10_Chemical_Industry:Other	Chemical industry - general	Process emission
	2B2_Nitric_Acid_Production	Nitric acid production	Acid production
	2B3_Adipic_Acid_Production	Adipic acid production	Adipic acid produced
	2B6_Titanium_dioxide_production	Chemical industry - titanium dioxide	Coke
			Petroleum coke
	2B7_Soda_Ash_Production	Chemical industry - soda ash	Soda ash produced
	2B8a_Methanol_production	Chemical industry - methanol	Methanol
			Natural gas
	2B8b_Ethylene_Production	Chemical industry - ethylene	Ethylene
	2B8c_Ethylene_Dichloride_and_Vinyl_Chloride_Monomer	Chemical Industry - ethylene dichloride	Ethylene dichloride
	2B8d_Ethylene_Oxide	Chemical industry - ethylene oxide	Ethylene oxide
	2B8e_Acrylonitrile	Chemical industry - acrylonitrile	Acrylonitrile
	2B8f_Carbon_black_production	Chemical industry - carbon black	Carbon black capacity
	2B9a1_Fluorchemical_production:By-product_emissions	Halocarbons production - by-product	Non-fuel combustion
	2B9b3_Fluorchemical_production:Fugitive_emissions	Halocarbons production - fugitive	Non-fuel combustion
	2C1a_Steel	Basic oxygen furnaces	Dolomite

National Communication Category	IPCC Sector	Source Name	Activity Name
		Electric arc furnaces	Petroleum coke
			Steel production (electric arc)
		Ladle arc furnaces	Steel production (electric arc)
			Steel production (oxygen converters)
	2C1b_Pig_iron	Blast furnaces	Coke
			Fuel oil
	2C1d_Sinter	Sinter production	Coke
			Dolomite
			Limestone
	2C3_Aluminium_Production	Primary aluminium production - general	Primary aluminium production
		Primary aluminium production - PFC emissions	Primary aluminium production
	2C4_Magnesium_production	Magnesium cover gas	Non-fuel combustion
	2C6_Zinc_Production	Non-ferrous metal processes	Coke
	2G3b_N2O_from_product_uses:_Other	Other food - cream consumption	Process emission
		Recreational use of N2O	Process emission
2G4_Other_product_manufacture_and_use	Chemical Industry – other process sources	Process emission	
non-IPCC	Blast furnaces	Electricity	
Land Use Change	4_Indirect_N2O_Emissions	LULUCF Indirect N2O - Atmospheric Deposition	Non-fuel combustion
		LULUCF Indirect N2O - Nitrogen Leaching and Run-off	Non-fuel combustion

National Communication Category	IPCC Sector	Source Name	Activity Name
	4A_Forest Land_Emissions_from_Drainage	Forest Land - Drainage and rewetting and other management of organic and mineral soils	Non-fuel combustion
	4A1_ Forest Land remaining Forest Land	Forest Land remaining Forest Land - Biomass Burning - Wildfires	Biomass
		Forest Land remaining Forest Land - Carbon stock change	Non-fuel combustion
	4A2_1_Cropland converted to Forest Land	Cropland converted to Forest Land - Carbon stock change	Non-fuel combustion
	4A2_2_Grassland converted to Forest Land	Grassland converted to Forest Land - Carbon stock change	Non-fuel combustion
	4A2_4_Settlements converted to Forest Land	Settlements converted to Forest Land - Carbon stock change	Non-fuel combustion
	4A2_Cropland converted to Forest Land	Cropland converted to Forest Land - Direct N2O emissions from N Mineralization/Immobilization	Non-fuel combustion
	4A2_Grassland converted to Forest Land	Grassland converted to Forest Land - Direct N2O emissions from N Mineralization/Immobilization	Non-fuel combustion
	4A2_Land converted to Forest Land_Emissions_from_Fertilisation	Direct N2O emission from N fertilisation of forest land	Non-fuel combustion
	4A2_Settlements converted to Forest Land	Settlements converted to Forest Land - Direct N2O emissions from N Mineralization/Immobilization	Non-fuel combustion
	4B1_Cropland Remaining Cropland	Cropland remaining Cropland - Biomass Burning - Wildfires	Biomass
		Cropland remaining Cropland - Carbon stock change	Non-fuel combustion
	4B1_Cropland_Remaining_Cropland	Carbon stock change	Non-fuel combustion
	4B2_1_Forest Land converted to Cropland	Forest Land converted to Cropland - Biomass Burning - Controlled Burning	Biomass

National Communication Category	IPCC Sector	Source Name	Activity Name
		Forest Land converted to Cropland - Carbon stock change	Non-fuel combustion
		Forest Land converted to Cropland - Direct N2O emissions from N Mineralization/Immobilization	Non-fuel combustion
	4B2_2_Grassland converted to Cropland	Grassland converted to Cropland - Carbon stock change	Non-fuel combustion
		Grassland converted to Cropland - Direct N2O emissions from N Mineralization/Immobilization	Non-fuel combustion
	4B2_4_Settlements converted to Cropland	Settlements converted to Cropland - Carbon stock change	Non-fuel combustion
	4C_Grassland_Emissions_from_Drainage	Grassland - Drainage and rewetting and other management of organic and mineral soils	Non-fuel combustion
	4C1_Grassland Remaining Grassland	Grassland remaining Grassland - Biomass Burning - Wildfires	Biomass
		Grassland remaining Grassland - Carbon stock change	Non-fuel combustion
		Grassland remaining Grassland - Direct N2O emissions from N Mineralization/Immobilization	Non-fuel combustion
	4C2_1_Forest Land converted to Grassland	Forest Land converted to Grassland - Biomass Burning - Controlled Burning	Biomass
		Forest Land converted to Grassland - Carbon stock change	Non-fuel combustion
		Forest Land converted to Grassland - Direct N2O emissions from N Mineralization/Immobilization	Non-fuel combustion
	4C2_2_Cropland converted to Grassland	Cropland converted to Grassland - Carbon stock change	Non-fuel combustion
	4C2_3_Wetlands converted to Grassland	Wetlands converted to Grassland - Carbon stock change	Non-fuel combustion

National Communication Category	IPCC Sector	Source Name	Activity Name
	4C2_4_Settlements converted to Grassland	Settlements converted to Grassland - Carbon stock change	Non-fuel combustion
	4D_Wetlands_Emissions_from_Drainage	Wetlands - Drainage and rewetting and other management of organic and mineral soils	Non-fuel combustion
	4D1_Wetlands remaining wetlands	Peat Extraction Remaining Peat Extraction - Carbon stock change	Non-fuel combustion
	4D2_Land converted to Wetlands	Grassland converted to flooded land - Carbon stock change	Non-fuel combustion
		Land converted for Peat Extraction - Carbon stock change	Non-fuel combustion
	4E1_Settlements remaining settlements	Settlements remaining Settlements - Carbon stock change	Non-fuel combustion
		Settlements remaining Settlements - Direct N2O emissions from N Mineralization/Immobilization	Non-fuel combustion
	4E2_1_Forest Land converted to Settlements	Forest Land converted to Settlements - Biomass Burning - Controlled Burning	Biomass
		Forest Land converted to Settlements - Carbon stock change	Non-fuel combustion
		Forest Land converted to Settlements - Direct N2O emissions from N Mineralization/Immobilization	Non-fuel combustion
	4E2_2_Cropland converted to Settlements	Cropland converted to Settlements - Carbon stock change	Non-fuel combustion
		Cropland converted to Settlements - Direct N2O emissions from N Mineralization/Immobilization	Non-fuel combustion
	4E2_3_Grassland converted to Settlements	Grassland converted to Settlements - Carbon stock change	Non-fuel combustion
		Grassland converted to Settlements - Direct N2O emissions from N Mineralization/Immobilization	Non-fuel combustion

National Communication Category	IPCC Sector	Source Name	Activity Name
	4G_Harvested Wood Products	HWP Produced and Consumed Domestically - Carbon stock change	Non-fuel combustion
		HWP Produced and Exported - Carbon stock change	Non-fuel combustion
Public	1A4ai_Commercial/Institutional	Public sector combustion	Burning oil
			Coal
			Coke
			Fuel oil
			Gas oil
			Natural gas
			Sewage gas
	non-IPCC	Public sector combustion	Electricity
Residential	1A4bi_Residential_stationary	Domestic combustion	Anthracite
			Burning oil
			Charcoal
			Coal
			Coke
			Fuel oil
			Gas oil
			LPG

National Communication Category	IPCC Sector	Source Name	Activity Name
			Natural gas
			Peat
			Petroleum coke
			SSF
			Wood
	1A4bii_Residential:Off-road	House and garden machinery	DERV
			Petrol
	2D2_Non-energy_products_from_fuels_and_solvent_use:Paraffin_wax_use	Non-aerosol products - household products	Petroleum waxes
	2F4a_Metered_dose_inhalers	Metered dose inhalers	Non-fuel combustion
	2F4b_Aerosols:Other	Aerosols - halocarbons	Non-fuel combustion
5B1a_composting_municipal_solid_waste	Composting (at household)	Biological waste	
5C2.2b_Non-biogenic:Other	Accidental fires - dwellings	Mass burnt	
5C2.2b_Non-biogenic:Other_Accidental fires (vehicles)	Accidental fires - vehicles	Mass burnt	
non-IPCC	Domestic combustion	Electricity	
Transport	1A3a_Domestic_aviation	Aircraft - domestic cruise	Aviation spirit
			Aviation turbine fuel
		Aircraft - domestic take off and landing	Aviation spirit

National Communication Category	IPCC Sector	Source Name	Activity Name	
			Aviation turbine fuel	
		1A3bi_Cars	Road transport - cars - cold start	DERV
				Petrol
			Road transport - cars - motorway driving	DERV
				Petrol
			Road transport - cars - rural driving	DERV
				Petrol
		Road transport - cars - urban driving	DERV	
			Petrol	
		1A3bii_Light_duty_trucks	Road transport - LGVs - cold start	DERV
				Petrol
			Road transport - LGVs - motorway driving	DERV
				Petrol
			Road transport - LGVs - rural driving	DERV
				Petrol
			Road transport - LGVs - urban driving	DERV
				Petrol
		1A3biii_Heavy_duty_trucks_and_buses	Road transport - buses and coaches - motorway driving	DERV
Road transport - buses and coaches - rural driving	DERV			

National Communication Category	IPCC Sector	Source Name	Activity Name
		Road transport - buses and coaches - urban driving	DERV
		Road transport - HGV articulated - motorway driving	DERV
		Road transport - HGV articulated - rural driving	DERV
		Road transport - HGV articulated - urban driving	DERV
		Road transport - HGV rigid - motorway driving	DERV
		Road transport - HGV rigid - rural driving	DERV
		Road transport - HGV rigid - urban driving	DERV
	1A3biv_Motorcycles	Road transport - mopeds (<50cc 2st) - urban driving	Lubricants
			Petrol
		Road transport - motorcycle (>50cc 2st) - urban driving	Petrol
		Road transport - motorcycle (>50cc 4st) - motorway driving	Petrol
		Road transport - motorcycle (>50cc 4st) - rural driving	Petrol
	Road transport - motorcycle (>50cc 4st) - urban driving	Petrol	
	1A3bv_Other_road_transport	Road transport - all vehicles biofuels use	Biodiesel
			Bio-MTBE
		Road transport - all vehicles LPG use	LPG
	1A3c_Railways	Rail - coal	Coal
		Railways - freight	Gas oil

National Communication Category	IPCC Sector	Source Name	Activity Name	
		Railways - intercity	Gas oil	
		Railways - regional	Gas oil	
	1A3d_Domestic_navigation	Inland goods-carrying vessels	Inland goods-carrying vessels	Gas oil
			Motorboats / workboats (e.g. canal boats, dredgers, service boats, tourist boats, river boats)	DERV
				Gas oil
				Petrol
			Personal watercraft e.g. jet ski	Petrol
			Sailing boats with auxiliary engines	DERV
			Shipping - coastal	Fuel oil
	Gas oil			
	1A3eii_Other_Transportation	Aircraft - support vehicles	Gas oil	
	1A4ai_Commercial/Institutional	Railways - stationary combustion	Burning oil	
			Fuel oil	
			Natural gas	
	1A4ciii_Fishing	Fishing vessels	Fuel oil	
			Gas oil	
	1A5b_Other:Mobile	Aircraft - military	Aviation spirit	
			Aviation turbine fuel	
		Shipping - naval	Gas oil	

National Communication Category	IPCC Sector	Source Name	Activity Name
	2D1_Lubricant_Use	Marine engines	Lubricants
		Road vehicle engines	Lubricants
	2D3_Non-energy_products_from_fuels_and_solvent_use:Other	Road transport - urea	Urea consumption
	non-IPCC	Railways - regional	Electricity
		Road vehicle engines	Electricity
Waste Management	5A1a_Managed_Waste_Disposal_sites_anaerobic	Landfill	Non-fuel combustion
	5B1a_composting_municipal_solid_waste	Mechanical Biological Treatment - Composting	Biological waste
		Total composting (non-household)	Biological waste
	5B2a_Anaerobic_digestion_municipal_solid_waste	Anaerobic Digestion (other)	Biological waste
		Mechanical Biological Treatment - Anaerobic Digestion	Biological waste
	5C1.1b_Biogenic:Sewage_sludge	Incineration - sewage sludge	Sewage sludge combustion
	5C1.2a_Non-biogenic:municipal_solid_waste	Incineration	MSW
	5C1.2b_Non-biogenic:Clinical_waste	Incineration - clinical waste	Clinical waste
	5C1.2b_Non-biogenic:Other_Chemical_waste	Incineration - chemical waste	Chemical waste
		5D1_Domestic_wastewater_treatment	Sewage sludge decomposition
	Sewage sludge decomposition in private systems	Non-fuel domestic	
5D2_Industrial_wastewater_treatment	Industrial Waste Water Treatment	Non-fuel combustion	

---

**A 9.6 DETAILED EMISSIONS ACCORDING TO END USER CATEGORIES**

The end user categories in the data tables in this summary are those used in National Communications. The end user reallocation includes emissions from the UK, this is the coverage used for the UK statistical release, where the end users data are presented in more detail.

The base year for hydrofluorocarbons, perfluorocarbons, nitrogen trifluoride, and sulphur hexafluoride is 1995. For carbon dioxide, methane and nitrous oxide, the base year is 1990.

**Table A 9.6.1 End user emissions from all National Communication categories, MtCO<sub>2</sub> equivalent**

End user category	Base Year	1990	1995	2000	2005	2010	2013	2014	2015	2016	2017
Agriculture	57.8	57.8	56.1	53.0	50.6	47.2	46.6	47.7	47.1	47.0	47.2
Business	248.6	247.8	218.1	217.1	212.0	186.6	175.7	161.0	148.4	131.7	125.6
Energy Supply	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
Exports	9.1	9.1	13.0	12.8	16.7	15.9	13.9	12.2	11.9	11.9	12.1
Industrial Process	65.5	63.3	53.6	29.2	21.3	13.5	13.5	13.6	13.3	11.1	11.2
Land Use Change	0.3	0.3	-1.7	-3.9	-7.1	-9.1	-9.4	-9.6	-9.7	-9.8	-9.9
Public	31.5	31.5	28.9	24.4	22.4	19.0	17.8	15.2	14.5	13.4	12.4
Residential	171.9	171.3	157.1	158.2	162.4	155.9	139.8	118.0	113.3	107.3	100.4
Transport	146.6	146.6	151.4	153.9	156.3	142.1	135.2	136.4	138.5	140.5	140.9
Waste Management	66.6	66.6	69.1	62.9	49.0	29.7	23.1	20.9	20.6	20.0	20.3
<b>Total greenhouse gas emissions</b>	<b>798.0</b>	<b>794.4</b>	<b>745.6</b>	<b>707.5</b>	<b>683.7</b>	<b>600.9</b>	<b>556.2</b>	<b>515.6</b>	<b>498.0</b>	<b>473.1</b>	<b>460.2</b>

**Table A 9.6.2 End user CO<sub>2</sub> emissions from all National Communication categories, MtCO<sub>2</sub> equivalent**

End user category	Base Year	1990	1995	2000	2005	2010	2013	2014	2015	2016	2017
Agriculture	10.0	10.0	9.4	8.0	8.8	7.9	7.5	7.6	7.4	7.2	7.1
Business	229.8	229.8	203.3	202.7	194.9	166.8	157.2	142.3	129.8	114.4	109.1
Energy Supply	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
Exports	8.4	8.4	12.1	12.1	16.0	15.3	13.2	11.6	11.4	11.4	11.5

End user category	Base Year	1990	1995	2000	2005	2010	2013	2014	2015	2016	2017
Industrial Process	20.9	20.9	18.9	18.1	16.7	11.2	12.7	12.8	12.5	10.4	10.5
Land Use Change	-2.0	-2.0	-3.9	-6.0	-8.9	-10.7	-10.9	-11.1	-11.2	-11.3	-11.3
Public	29.2	29.2	27.1	23.4	21.7	18.5	17.3	14.8	14.1	13.0	12.0
Residential	156.2	156.2	145.2	149.0	154.5	148.9	133.4	112.1	107.5	101.9	95.2
Transport	142.5	142.5	147.0	150.5	153.8	140.2	133.3	134.4	136.6	138.5	138.8
Waste Management	1.3	1.3	1.0	0.5	0.4	0.3	0.3	0.3	0.2	0.3	0.3
<b>Total greenhouse gas emissions</b>	<b>596.3</b>	<b>596.3</b>	<b>560.1</b>	<b>558.3</b>	<b>557.9</b>	<b>498.3</b>	<b>464.0</b>	<b>424.9</b>	<b>408.3</b>	<b>385.8</b>	<b>373.2</b>

**Table A 9.6.3 End user CH<sub>4</sub> emissions from all National Communication categories, MtCO<sub>2</sub> equivalent**

End user category	Base Year	1990	1995	2000	2005	2010	2013	2014	2015	2016	2017
Agriculture	30.6	30.6	29.8	28.8	26.9	25.3	25.1	25.5	25.7	25.6	25.8
Business	15.5	15.5	11.7	7.4	4.8	3.7	2.9	2.9	2.7	2.3	2.3
Energy Supply	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
Exports	0.7	0.7	0.8	0.6	0.5	0.5	0.5	0.5	0.4	0.4	0.5
Industrial Process	2.2	2.2	1.7	1.1	0.5	0.4	0.3	0.3	0.2	0.2	0.2
Land Use Change	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
Public	2.1	2.1	1.7	0.9	0.6	0.5	0.4	0.4	0.4	0.3	0.3
Residential	14.3	14.3	10.7	6.7	4.9	4.5	4.0	3.6	3.6	3.2	3.1

End user category	Base Year	1990	1995	2000	2005	2010	2013	2014	2015	2016	2017
Transport	2.5	2.5	2.2	1.5	1.0	0.8	0.7	0.7	0.7	0.6	0.7
Waste Management	64.5	64.5	67.3	61.4	47.6	28.2	21.5	19.3	19.0	18.4	18.6
<b>Total greenhouse gas emissions</b>	<b>132.5</b>	<b>132.5</b>	<b>125.9</b>	<b>108.4</b>	<b>86.9</b>	<b>63.9</b>	<b>55.5</b>	<b>53.3</b>	<b>52.7</b>	<b>51.1</b>	<b>51.5</b>

**Table A 9.6.4 End user N<sub>2</sub>O emissions from all National Communication categories, MtCO<sub>2</sub> equivalent**

End user category	Base Year	1990	1995	2000	2005	2010	2013	2014	2015	2016	2017
Agriculture	17.2	17.2	16.8	16.2	14.9	14.0	14.0	14.6	14.1	14.1	14.3
Business	1.5	1.5	1.4	1.3	1.3	1.2	1.3	1.2	1.2	1.1	1.1
Energy Supply	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
Exports	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1
Industrial Process	23.9	23.9	14.4	5.4	3.1	1.5	0.3	0.3	0.3	0.3	0.3
Land Use Change	2.3	2.3	2.2	2.1	1.8	1.6	1.5	1.5	1.4	1.4	1.4
Public	0.1	0.1	0.1	0.1	0.1	0.0	0.1	0.1	0.0	0.0	0.0
Residential	0.7	0.7	0.6	0.5	0.5	0.5	0.6	0.5	0.5	0.4	0.4
Transport	1.6	1.6	2.2	1.9	1.5	1.1	1.2	1.2	1.3	1.3	1.4
Waste Management	0.8	0.8	0.8	0.9	1.0	1.2	1.3	1.3	1.4	1.4	1.4
<b>Total greenhouse gas emissions</b>	<b>48.2</b>	<b>48.2</b>	<b>38.6</b>	<b>28.5</b>	<b>24.4</b>	<b>21.3</b>	<b>20.3</b>	<b>20.8</b>	<b>20.3</b>	<b>20.2</b>	<b>20.5</b>

