

# Trends Report 2019: Trends in critical load and critical level exceedances in the UK

**Ed Rowe<sup>1</sup>, Kasia Sawicka<sup>1</sup>, Zak Mitchell<sup>1</sup>, Ron Smith<sup>2</sup>, Tony Dore<sup>2</sup>,  
Lindsay F. Banin<sup>2</sup> & Peter Levy<sup>2</sup>**

**<sup>1</sup>CEH, Environment Centre Wales, Bangor, Gwynedd, LL57 2UW**

**<sup>2</sup>CEH, Bush Estate, Penicuik, Midlothian, EH26 0QB**



**Centre for  
Ecology & Hydrology**  
NATURAL ENVIRONMENT RESEARCH COUNCIL



**NATURAL  
ENVIRONMENT  
RESEARCH COUNCIL**

Cite as: Rowe E, Sawicka K, Mitchell Z, Smith R, Dore T, Banin LF & Levy P (2019) Trends Report 2019: Trends in critical load and critical level exceedances in the UK. Report to Defra under Contract AQ0843, CEH Project NEC05708. <https://uk-air.defra.gov.uk/library/>

Correction: This version (13<sup>th</sup> Dec 2019) corrects an error in ammonia concentration data – see note at the start of Section 4. This version supersedes the version published in July 2019.

## Table of Contents

Executive Summary.....	3
Technical Summary.....	4
Report structure.....	7
Note on rounded numbers .....	7
Section 1. Critical loads and their exceedances.....	7
1.1 Overview of UK critical loads .....	7
1.1.1 Acidity critical loads .....	9
1.1.2 Nutrient nitrogen critical loads .....	9
1.2 Overview of UK deposition data .....	10
1.2.1 Summary of trends in CBED deposition .....	12
1.3 Overview of the calculation of critical load exceedances.....	13
1.3.1 Exceedance and damage.....	15
1.3.2 Critical load exceedance metrics .....	15
1.3.3 Critical load exceedance maps for all habitats combined .....	16
Section 2: Trends in critical loads exceedance by habitat and country.....	18
2.1 Trends by country .....	20
2.1.1 Acidity results.....	20
2.1.2 Nutrient nitrogen results .....	25
2.2 Trends by habitat .....	29
2.2.1 Acidity results.....	29
2.2.2 Nutrient nitrogen results .....	29
Section 3: Site-relevant critical loads and their exceedances.....	38
3.1 Overview of site-relevant critical loads .....	38
3.2 Overview of SRCL exceedance metrics .....	39
3.2.1 Acidity results.....	40
3.2.2 Nutrient nitrogen results .....	46
Section 4: Critical levels and their exceedances .....	51
4.1 Critical levels of ammonia.....	51
4.2 Concentrations of ammonia .....	52
4.3 Calculation of critical levels exceedance.....	52
4.4 Trends in ammonia critical levels exceedance.....	52
4.4.1 UK land area with exceedance of ammonia critical levels.....	52
4.4.2 Nitrogen-sensitive habitats with exceedance of ammonia critical levels.....	55
4.4.3 Designated sites with exceedance of ammonia critical levels.....	57
References .....	61
Annex: Critical load exceedances by habitat and country.....	64

## Executive Summary

This contract provides key information to track the effects on ecosystems of policies aimed at meeting national and international air pollution targets (e.g. CLRTAP, NECD). It also provides the means to help develop targeted action for these emission reduction policies to get the maximum improvement. Exceedance of critical load or level indicates that an ecosystem is at risk from potential harmful effects. When pollution is reduced to below the critical load or level, there may be delays to recovery, but the risk of harm is reduced.

Reduced pollution emissions to meet CLRTAP (Convention on Long-range Transboundary Air Pollution) targets have mainly reduced the extent and magnitude of critical load and level exceedances across the UK. New targets have been developed to reduce impacts further.

- For UK habitats at risk from acidification, the area of acid-sensitive habitats in the UK with exceedance of acidity critical loads has fallen by more than one third, from 72.6% in 1996, to 44.5% in 2012, to 37.6% (29,337 km<sup>2</sup>) in 2016 due mainly to decreases in sulphur deposition.
- For UK habitats at risk from excess nitrogen (eutrophication), the area of N-sensitive habitats in the UK with exceedance of nutrient N critical loads fell from 75.0% in 1996, to 62.5% in 2012, to 58.1% (42,434 km<sup>2</sup>) in 2016.
- Just over 4% (9,793 km<sup>2</sup>) of the UK land area is currently exposed to ammonia concentrations above the critical level set to protect higher plants (3 µg m<sup>-3</sup>), and just over 60% (146,900 km<sup>2</sup>) to ammonia at concentrations above the critical level set to protect lichens and mosses (1 µg m<sup>-3</sup>). The area where the critical level for higher plants is exceeded increased by 0.5% (1,224km<sup>2</sup>) of UK land area since 2010. The area where the critical level for lichens and mosses is exceeded decreased by 1% (2,448km<sup>2</sup>) of UK land area since 2010.

## Technical Summary

Critical loads define the rates of acid or nitrogen (N) deposition (e.g. in kiloequivalents per hectare per year) below which significant harmful effects are not expected to occur in sensitive habitats. Critical load exceedance is the amount by which acid or N deposition exceeds the critical load. Critical levels refer to the concentrations of gaseous pollutants (e.g. in micrograms per cubic metre) below which direct adverse effects on sensitive vegetation are not expected to occur. Critical level exceedance is the amount by which concentration exceeds the critical level. Critical load and critical level exceedances are calculated using rolling 3-year mean data sets for deposition rates and gaseous concentrations, which are updated annually. This report describes critical loads and their exceedances for the period from 1996 to 2016<sup>1</sup>, and critical levels and their exceedances for the period from 2010 to 2015.

This report presents trends in: a) acidity critical load exceedances for UK habitats at risk from acidification by excess sulphur (S) and/or N; b) nutrient-nitrogen critical load exceedances for UK habitats at risk of eutrophication by excess N; c) critical load exceedances for acid- and N-sensitive habitat features of UK designated sites (Special Areas of Conservation: SACs; Special Protected Areas: SPAs; Sites of Special Scientific Interest: SSSIs); and d) exceedances of ammonia critical levels across the UK.

Summary statistics monitor progress in the areas of habitats in the UK at risk from acidification and eutrophication from air pollution over time, and are reported in the annual “UK Biodiversity Indicators” summary produced by the Joint Nature Conservation Committee (<http://jncc.defra.gov.uk/page-4233>; indicator B5a for assessing the pressures from air pollution).

### ***UK habitats at risk from acidification***

- The area of acid-sensitive habitats in the UK with exceedance of acidity critical loads continues to decline due mainly to decreases in sulphur (S) deposition, having fallen from 72.6% in 1996 to 37.6% (29,337 km<sup>2</sup>) in 2016.
- The magnitude of the acidity exceedance (expressed as the Average Accumulated Exceedance) for all UK habitats combined fell by more than two thirds between 1996 and 2016, from 0.78 to 0.21 keq ha<sup>-1</sup> year<sup>-1</sup>.
- The largest reduction in the area of acid-sensitive habitats with exceedance of acidity critical loads has been in Scotland, where it has fallen by two thirds from 68.2% (32774 km<sup>2</sup>) in 1996 to 22.7% (10,933 km<sup>2</sup>) in 2016.
- The smallest reduction in the area of acid-sensitive habitats with exceedance of acidity critical load has been in Northern Ireland, falling from 76.8% (2,718 km<sup>2</sup>) in 1996 to 61.2% (2,164 km<sup>2</sup>) in 2016.
- Of the terrestrial acid-sensitive habitats mapped, dwarf shrub heath occupies the largest area across the UK (10%); the area of this habitat with exceedance of acidity critical loads has more than halved, from 70.3% (17,370km<sup>2</sup>) in 1996 to 20.7% (5,119 km<sup>2</sup>) in 2016.

---

<sup>1</sup> In this report, time periods are usually (for brevity) referred to using the middle year of the 3 years used to calculate the mean. The example here, “from 1996 to 2015”, means the period from 1995-1997 to 2014-2016.

- The terrestrial habitat with the smallest decrease in acidity critical load exceedance, falling by less than a third between 1996 and 2016, has been acid grassland, where 92.0% (14,106 km<sup>2</sup>) was exceeded in 1996 to and 65.4% (10,026 km<sup>2</sup>) in 2016.

#### ***UK habitats at risk from eutrophication (i.e. excess nitrogen)***

- The reduction in the areas of N-sensitive habitats in the UK with exceedance of nutrient N critical loads, decreased by less than one-fifth, from 75.0% (54,785 km<sup>2</sup>) in 1996 to 58.1% (42,434 km<sup>2</sup>) in 2016. This smaller reduction (in comparison to acidity) is due to the smaller proportional reductions in N deposition over the time period, compared to S.
- The Average Accumulated Exceedance for all UK habitats combined for nutrient N has declined by more than one-third, from 9.5 kg N ha<sup>-1</sup> year<sup>-1</sup> in 1996 to 5.6 kg N ha<sup>-1</sup> year<sup>-1</sup> in 2016.
- The largest reduction in the area of N-sensitive habitats with critical load exceedance is in Scotland, falling by almost one-third from 59.4% (25,675 km<sup>2</sup>) in 1996 to 34.7% (15,002 km<sup>2</sup>) in 2016.
- The smallest reduction in the area of N-sensitive habitats with critical load exceedance is in England, falling from 98.3% (19,199 km<sup>2</sup>) in 1996 to 94.8% (18,504 km<sup>2</sup>) in 2016.
- The nutrient N critical loads are exceeded for more than 80% of the areas of six N-sensitive habitats in all years: calcareous grasslands, unmanaged beech woodland, unmanaged oak woodland, other unmanaged woodland, managed conifer and managed broadleaved woodland.
- Almost 100% of the area of unmanaged beech woodland has exceedance of nutrient N critical loads in all years, however, the magnitude of exceedance (Average Accumulated Exceedance) has nearly halved, from 22.7 kg N ha<sup>-1</sup> year<sup>-1</sup> in 1996 to 12.1 kg N ha<sup>-1</sup> year<sup>-1</sup> in 2016.
- There is virtually no exceedance of nutrient N critical loads for saltmarsh in any year, due to a combination of the high critical load for this habitat and lower deposition in coastal areas.

#### ***Designated sites with acid-sensitive feature habitats***

- The percentage of SACs and SPAs in the UK with exceedance of acidity critical loads for one or more features decreased from more than 90% (SACs: 443 sites, SPAs: 165 sites) in 1996 to 71.5% (SACs: 348 sites) and 65.7% (SPAs: 115 sites) in 2016; the percentage of SSSIs with exceedance fell from 77.6% (3,632 sites) in 1996 to 57.1% (2,674 sites) in 2016.
- Scotland had the largest reductions (>35%) in the percentage of designated sites with exceedance of acidity critical loads between 1996 and 2016.
- Roughly 50% of designated sites in England and Scotland, and more than 70% of designated sites in Wales and Northern Ireland, currently have exceedance of acidity critical loads for one or more features.

#### ***Designated sites with nitrogen-sensitive feature habitats***

- Decreases in the percentage of designated sites in the UK with exceedance of nutrient N critical loads (for one or more features) between 1996 and 2016 are small (5.2% for SACs, 9.0% for SSSIs, 14.2% for SPAs), reflecting the smaller reductions in N deposition, compared to acid deposition, over this time period.
- Scotland had the largest decreases (ca. 9-17%) in the percentage of SACs and SSSIs with exceedance of nutrient N critical loads between 1996 and 2016; Wales had the largest reduction

(21.4%) in the percentage of SPAs with nutrient N critical load exceedance over the same time period, however the total number of SPAs in Wales with N critical loads is only 14.

- 56.5-80.6% (73.8% by maximum area) of designated sites in Scotland, and 78.6-98.0% (86.3% by maximum area) of designated sites in England, Wales and Northern Ireland currently have exceedance of nutrient N critical loads for one or more features.

### ***Exceedance of ammonia critical levels***

The trends in ammonia critical levels exceedance are only available for the period 2009 to 2014; there have only been small reductions in ammonia concentrations over this time period.

#### ***UK land area***

- Over 60% of the UK land area currently receives ammonia concentrations above the critical level set to protect lichens and bryophytes ( $1 \mu\text{g m}^{-3}$ ); this represents 87.2% of England, 57.5% of Wales and 18.9% of Scotland and 89.3% of Northern Ireland.
- The UK land area with ammonia concentrations above  $1 \mu\text{g m}^{-3}$  has slightly decreased, from 63.9% in 2010 to 62.8% in 2015.
- Just over 4% of the UK land area receives ammonia concentrations above the critical level set to protect higher plants ( $3 \mu\text{g m}^{-3}$ ); this ranges from 0.1% of Scotland to 22.8% of Northern Ireland.
- The UK land area with ammonia concentrations above  $3 \mu\text{g m}^{-3}$  has slightly increased, from 3.7% in 2010 to 4.2% in 2015.

#### ***Nitrogen-sensitive habitats***

- One-quarter of the mapped area of N sensitive habitats in the UK receive ammonia concentrations above the critical level of  $1 \mu\text{g m}^{-3}$ ; this ranges from 3.6% for Scotland to 71.3% for Northern Ireland.
- Over 85% of the area of calcareous grassland, and ~65-80% of the area of some woodland habitats (unmanaged, unmanaged beech, managed broadleaved) are in areas that receive ammonia concentrations above the critical level of  $1 \mu\text{g m}^{-3}$ . Only 19.1% of acid grassland is in areas exceeding this critical level, but this equates to 2,916 km<sup>2</sup> which is similar to the 86.2% (3,086 km<sup>2</sup>) of calcareous grassland exceeded.
- Less than 1% of the area of N-sensitive habitats in the UK receives ammonia concentrations above  $3 \mu\text{g m}^{-3}$ ; this ranges from <0.1% in Wales and Scotland to 7.1% in Northern Ireland.

#### ***Designated sites***

- 52.7% (129) of SPAs, 60.6% (372) of SACs and 70.2% (4,798) of SSSIs in the UK currently receive ammonia concentrations above  $1 \mu\text{g m}^{-3}$  anywhere across the site. The percentage of sites with exceedance of the  $1 \mu\text{g m}^{-3}$  critical level has fallen by <5% between 2010 and 2015, for all site types.
- 86% - 95% of designated sites in England and Northern Ireland, 44-72% of sites in Wales, and 19-26% of sites in Scotland, currently receive ammonia concentration above  $1 \mu\text{g m}^{-3}$ .
- 4.9% (30) of SACs, 5.7% (14) of SPAs and 3.7% (256) of SSSIs in the UK currently receive ammonia concentrations above  $3 \mu\text{g m}^{-3}$  anywhere across a site. The percentage of designated sites with exceedance of this critical level has fallen by less than 0.1% for SACs and risen by <1% for SPAs and SSSIs between 2010 and 2015.

- Less than 0.5% of sites that have a conservation designation in Scotland currently receive ammonia concentrations above the critical level of  $3 \mu\text{g m}^{-3}$  anywhere across a site, compared with up to 1.7 % of sites in Wales, up to 13.2 % of sites in England and up to 15.4 % of sites in Northern Ireland.

## Report structure

Section 1 of this report provides an overview of critical loads for acidity and for nutrient N, deposition data, and exceedance calculations and metrics for habitats across the whole area of the UK. This is followed in Section 2 by summaries of these trends for specific habitats and countries, i.e. England, Scotland, Wales, and Northern Ireland. Section 3 focuses on designated sites, the application of “site-relevant critical loads” (SRCL) to these sites, and trends in their exceedances. Finally, Section 4 addresses critical levels for ammonia and their exceedances.

## Note on rounded numbers

Numbers in tables are shown to one decimal place. Differences were rounded after calculation, so may not always reflect differences between the numbers in the table. For example, if there is a change of +0.08 units from 72.06 to 72.14, the real change is  $72.14 - 72.06 = +0.08$ , which rounds to +0.1, and the rounded numbers are reported as  $72.1 - 72.1 = 0.1$ . This is not an error.

## Section 1. Critical loads and their exceedances

### 1.1 Overview of UK critical loads

Critical loads are thresholds for effects from atmospheric deposition and defined as “a quantitative estimate of the exposure to one or more pollutants below which significant harmful effects on specified sensitive elements of the environment do not occur according to present knowledge” (Nilsson & Grennfelt 1988).

The methods used to calculate and map UK critical loads are described in detail in Hall et al. (2015). Critical loads are calculated and mapped for UK habitats sensitive to acidification and/or eutrophication (Table 1.1). Here and in Section 2, results are presented for the entire mapped extent of habitats; results for designated sites are presented in Section 3. The critical load methods applied in the UK are based on methods approved at international workshops held under the Convention on Long-Range Transboundary Air Pollution (CLRTAP) and published in the “Mapping Manual” (CLRTAP 2017) .

**Table 1.1: Habitat distributions mapped for acidity and for nutrient nitrogen critical loads**

Habitat	EUNIS habitat class(es) assigned <sup>1</sup>	Mapped for acidity	Mapped for nutrient nitrogen
Acid grassland (wet & dry)	E1.7 & E3.52	Yes	Yes
Calcareous grassland	E1.26	Yes	Yes
Dwarf shrub heath (wet & dry)	F4.11 & F4.2	Yes	Yes
Montane	E4.2	Yes	Yes
Bog	D1	Yes	Yes
Managed coniferous woodland	G3	Yes	Yes
Managed broadleaved woodland	G1	Yes	Yes
Beech woodland (unmanaged)	G1.6	Yes <sup>3</sup>	Yes
Acidophilous oak woodland (unmanaged)	G1.8	Yes <sup>3</sup>	Yes
Scots Pine woodland (unmanaged)	G3.4	Yes <sup>3</sup>	Yes
Other unmanaged woodland	G4	Yes <sup>3</sup>	Yes
Freshwaters <sup>2</sup>	C1 & C2	Yes	No
Dune grassland	B1.4	No	Yes
Saltmarsh	A2.53/54/55	No	Yes

<sup>1</sup> EUNIS class closest to broad habitat and critical loads habitat; class used for assigning empirical nutrient nitrogen critical loads and for classifying UK critical loads data for submission to the CCE.

<sup>2</sup> Critical loads are calculated for 1752 freshwater sites across the UK (see Section 1.1.1 below); habitat areas are based on the catchment areas of these sites.

<sup>3</sup> Unmanaged woodland classes are mapped together.

Published correspondence tables (available from: <http://jncc.defra.gov.uk/page-1425>) are used to relate broad habitats to the European Nature Information System (EUNIS: Moss & Davies 2002) hierarchical habitat classification scheme, developed for pan-European applications.

Habitat distribution maps are based on the CEH Land Cover Map 2000 (LCM2000: Fuller et al. 2002) and additional data sets such as species distribution data and altitude. Habitat areas, used for assessing the areas of habitats at risk from acidification and or eutrophication, are based on the LCM2000 data. *It should be noted that the habitat distribution maps and areas used for UK critical loads (acidity and nutrient N) research (a) only include areas where data exist for the calculation or derivation of critical loads; (b) may differ from other national habitat distribution maps or estimates of habitat areas. This may also result in a difference in the total habitat areas mapped for acidity and for nutrient N critical loads.*

### **1.1.1 Acidity critical loads**

Two methods are used in the UK for calculating acidity critical loads for terrestrial habitats: the empirical approach is used to provide estimates for non-woodland habitats and a simple mass balance equation used for woodland habitats.

In the empirical approach used for non-woodland mineral and organomineral soils, critical loads are assigned to each 1km grid square of the UK based on the amount of acid deposition that could be neutralised by the base cations produced by mineral weathering of the dominant soil type in the grid square (Hornung et al. 1995). This approach is inappropriate for peat soils because of the absence of inputs of alkalinity from mineral weathering (Smith et al. 1992, Gammack et al. 1995). Critical loads of acidity for peat soils are set to the value corresponding to the amount of acid deposition that would give rise to an effective rain pH value of 4.4 (Skiba & Cresser 1989, Calver 2003, Calver et al. 2004); the choice of threshold pH value reflects the buffering effects of organic acids on peat drainage water pH. This method is applicable to upland and lowland acid peat soils, but not to peats in lowland arable fen areas that are less sensitive to acidification, where a higher critical load is set than would be applied to acid peats (Hall et al. 2015).

Acidity critical loads for non-woodland habitats are calculated using the soil acidity critical loads outlined above, together with additional habitat-specific data to derive the three acidity critical load values (CL<sub>maxS</sub>, CL<sub>minN</sub>, CL<sub>maxN</sub>, see Section 2) for each habitat, needed for the calculation of acidity critical load exceedances.

For woodland habitats a simple mass balance (SMB) equation, based on balancing the acidic inputs to and outputs from the ecosystem, is used to derive a critical load that ensures that a specified critical chemical limit is not exceeded (Sverdrup et al. 1990, Sverdrup & De Vries 1994). In the UK, the SMB equation is parameterised using different chemical criteria for woodlands on mineral or organo-mineral soils, and woodlands on peat soils (Hall et al. 2015). Critical loads are calculated for both managed (productive) and unmanaged woodlands to protect the long-term ecosystem function of these woodland habitats; this approach also aims to protect the land under managed conifer forest for possible future non-forest use and reversion to semi-natural land uses. These SMB critical loads are used with additional habitat-specific data to derive the three acidity critical load input values (CL<sub>maxS</sub>, CL<sub>minN</sub>, CL<sub>maxN</sub>) for each woodland type, for use in calculating habitat-specific exceedances (Section 2).

Acidity critical loads for freshwaters are calculated using the catchment-based First-Order Acidity Balance (FAB: Henriksen & Posch 2001) model. FAB is currently applied to 1752 sites across the UK, comprising a mixture of mainly upland, lakes, reservoirs and first-order streams (i.e. streams that feed into other larger streams, but do not have any other streams draining into them). The critical load calculations are based on the water chemistry of samples collected in the 1990s to provide an estimate of the annual mean water chemistry. The FAB model generates the acidity critical load values CL<sub>maxS</sub>, CL<sub>minN</sub> and CL<sub>maxN</sub> (see Part 2).

### **1.1.2 Nutrient nitrogen critical loads**

Empirical and mass balance methods also exist for calculating critical loads for eutrophication (i.e. an excess of nutrients, in this case N). The empirical critical loads are based on experimental or field

evidence of thresholds for changes in species composition, plant vitality or soil processes. The empirical approach is suited to semi-natural communities for which the long-term protection of biodiversity and/or ecosystem function is the key concern. In the UK the empirical approach is applied to natural and semi-natural habitats, including unmanaged (non-productive) woodland, based on critical load values agreed at international workshops (Bobbink & Hettelingh 2011, Hall et al. 2015).

In the mass balance approach the long-term inputs and outputs of N from the ecosystem are calculated, with the critical load being exceeded when any excess N input is calculated to lead to an exceedance of a specified critical rate of N leaching. This approach is suited to managed ecosystems with relatively low biodiversity, in which the inputs and outputs can be quantified with some confidence and in which the key concern is nitrate leaching. In the UK, this approach is applied to managed (productive) woodlands to ensure that long-term ecosystem functions (e.g. of soils, soil biological resources, trees, linked aquatic systems) are protected.

## **1.2 Overview of UK deposition data**

The S, N and base cation deposition data used in the UK calculations of critical loads and their exceedances are based on the “Concentration Based Estimated Deposition” (CBED) methodology (RoTAP 2012). Site based measurements of air concentrations of sulphur and nitrogen, as gases and particulates, are interpolated to generate 5 x 5 km maps of concentrations for the UK. Ion concentrations in precipitation from the UK Eutrophying and Acidifying Pollutants (UKEAP) network are combined with the Met Office annual precipitation map to generate maps of wet deposition. The wet deposition values include (a) direct deposition of cloud droplets to vegetation (known as “occult” deposition); (b) an orographic enhancement to take account of the “seeder-feeder” effect in upland regions (Fowler et al. 1988). Gas and particulate concentration maps are combined with spatially distributed estimates of vegetation-specific deposition velocities (Smith et al. 2000) to generate dry deposition. Combining these data sets produces 5 x 5 km maps of total (wet + cloud + dry) deposition of S (non-marine), oxidised N and reduced N; two different sets of deposition values are used in critical load and exceedance applications: (i) “moorland”: assumes grassland or moorland vegetation everywhere; (ii) “woodland”: assumes forest everywhere, based on the different deposition velocities to different land cover types.





Figure 1.1: CBED deposition for 2015-17: (a) nitrogen (oxidised plus reduced) deposition to moorland; (b) nitrogen (oxidised plus reduced) deposition to woodland; (c) acid (sulphur + nitrogen) deposition to moorland; (d) acid (sulphur + nitrogen) deposition to woodland. Deposition is mapped for N and acidity using the same units ( $\text{keq ha}^{-1} \text{ yr}^{-1}$ ) and class intervals. For the N maps, deposition rates are also shown in  $\text{kg N ha}^{-1} \text{ yr}^{-1}$ .

Significant inter-annual variations in deposition can occur due to the natural variability in annual precipitation (which influences wet deposition) as well as the general circulation of air which can increase or decrease the amount of polluted air imported from the European continent. The CBED deposition data used to calculate critical load exceedances is therefore averaged over a three-year period; this has been demonstrated to be a suitable time period to smooth out inter-annual variations in deposition. Figure 1.1 shows the CBED data for 2015-17.

Since critical loads for terrestrial habitats are mapped on a 1km grid, for exceedance calculations deposition is assumed to be constant for all 1 km squares within each 5km grid square. For freshwater exceedance calculations catchment-weighted mean S and N deposition values are calculated by overlaying the catchment boundary and land cover information (moorland vs forest) onto the 5km<sup>2</sup> deposition maps.

### **1.2.1 Summary of trends in CBED deposition**

To understand the trends in critical load exceedances it is useful to look at the trends in deposition over time. The CBED deposition data exist for all 3-year rolling intervals from 2002 to 2016 (reminder: in this report, the 3-year time periods are referred to in brief using the *middle year* of the period), plus data for three earlier time periods: 1995-1997, 1998-2000 and 1999-2001. The trends in CBED deposition to moorland and to woodland for the period 1996 to 2016 are summarised in Figure 1.2; this clearly shows that decreases in deposition were generally greater before 2005, with smaller decreases since 2005 as well as some increases, particularly for deposition of reduced N. Deposition to woodland is higher than that to moorland due to greater deposition velocities onto tall vegetation of gases (e.g. NH<sub>3</sub> and HNO<sub>3</sub>) in particular, as well as particulates. Grid-averaged deposition of non-marine S (NMS) for the UK, i.e. deposition weighted by the amounts of moorland and woodland per grid cell (Hall et al. 2015), decreased by 76% over the study period from 1996 to 2016. The NMS deposition rate continues to decline, with a 33% decrease over the most recent five-year period reported, i.e. between 2012 and 2016. Deposition of oxidised N (UK grid-average) decreased by 47% between 1996 and 2016, and also shows an ongoing decline, having decreased by 21% between 2012 and 2016. By contrast, the decline in UK grid-average deposition of reduced N has been smaller (10% between 1996 and 2016) and has slowed in recent years, with only a 2% decrease between 2012 and 2016.



**Figure 1.2: Deposition budgets (kilotonnes S or N per year) calculated for all UK 5x5 km grid squares for CBED deposition to moorland and deposition to woodland. NMS = non-marine sulphur, NOx = oxidised nitrogen, NHx = reduced nitrogen.**

### 1.3 Overview of the calculation of critical load exceedances


Critical load exceedances are the amount of excess deposition above the critical load; for nutrient N the calculation is simply total N deposition (derived from N oxides and ammonia) minus the critical load. For acidification, deposition of both S and N compounds can contribute to the exceedance of critical loads. The Critical Load Function, developed under the UNECE CLRTAP (Hettelingh et al. 1995, Posch et al. 1995, Posch & Hettelingh 1997, Posch et al. 1999), defines combinations of S and N deposition that will not cause harmful effects. In its simplest form, an acidity critical load can be defined graphically by a 45 degree diagonal line on a sulphur-nitrogen deposition plot (Figure 1.3a). The line intercepts the x-axis (representing N deposition) and y-axis (representing S deposition) at chemically equivalent points, each representing the N or S deposition equal to the critical load for acidity. Each point along the diagonal line represents the critical load in terms of some combination of S and N deposition.

To allow for the long-term N removal processes by the soil and through harvesting of vegetation, the simple diagonal line is shifted along the N axis to increase the N values across the entire CLF (Figure 1.3b). More N can then be deposited before the acidity critical load is exceeded. There are no similar removal processes that need to be considered for S.

The intercepts of the CLF on the S and N axes (Figure 1.3c) define the following terms:


- The “maximum critical load of S” (CLmaxS): the critical load for acidity expressed in terms of S only, i.e. when N deposition is zero.
- The “maximum critical load of N” (CLmaxN): the critical load for acidity expressed in terms of N only (when S deposition is zero).
- The “minimum critical load of N” (CLminN): the long-term N removal processes in the soil (e.g. N uptake and immobilisation) and harvesting of vegetation.

These critical loads are calculated from the acidity critical loads described in Section 1.1 and additional soil-specific or habitat-specific data.



**Figure 1.3: Development of the CLF: (a) acidity critical load defined by equal amounts of sulphur and nitrogen deposition; (b) shifting the acidity critical load diagonal line to allow for nitrogen removal processes; (c) the 3 nodes of the CLF: CLmaxS, CLminN, CLmaxN. The area shown in grey represents the combinations of sulphur and nitrogen deposition that are below the critical load (i.e. critical load is not exceeded).**

Exceedances are calculated by comparing the values of CLmaxS, CLminN and CLmaxN to the values of S and N (oxidised + reduced) deposition. The actual calculation depends on where the deposition falls in relation to these critical load values; the CLF is divided into five different regions for this purpose (Figure 1.4). The exceedance is defined by the sum of S and N deposition as shown by the red arrows in Figure 1.4 (i.e. not the length of the diagonal line); this is referred to as the “shortest distance” exceedance. Further details on the calculations are given in Hall et al. (2015).



**Figure 1.4: Example of S and N deposition reductions required depending on the region of the CLF. Deposition that falls in region 5 is below the critical load (i.e. critical loads not exceeded).**

### 1.3.1 Exceedance and damage

The critical loads data on which exceedance calculations are based are derived from empirical or steady-state mass balance methods, which are used to define critical loads for the long term. Exceedance of critical loads is an indication that an ecosystem is at risk from potential harmful effects in the long-term. Therefore, exceedance is not a quantitative estimate of damage to the environment; it does not necessarily mean that harmful or adverse effects have already occurred or may be observed, but that there is a risk of damage in the long-term. Reducing deposition to below the critical load does not mean that ecosystems immediately recover. There are time lags before chemical recovery takes place, and further delays before biological recovery. The timescales for both chemical and biological recovery, could be very long, particularly for the most sensitive ecosystems.

### 1.3.2 Critical load exceedance metrics

Critical load exceedances are calculated for each 1km square of the distributions of each terrestrial habitat, and for each catchment for freshwaters. The results are then summarised by habitat and country using the following exceedance metrics:

(i) Area of habitat exceeded

For terrestrial habitats the area values are based on the LCM2000 data; if the critical load for any individual habitat is exceeded, the exceeded area is set to the habitat area within the 1km square for that particular habitat. For freshwater habitats, if the FAB critical load is exceeded, the whole catchment is assumed to be exceeded and the exceeded area set to the catchment area. The total exceeded areas for individual habitats are summarised by country.

(ii) Percentage area of habitat exceeded

This is calculated from the exceeded areas derived in (i) and the total area of each habitat mapped in each country (Section 1.1). While this is a useful metric, it has its limitations, for example, when comparing exceedance results from one year to another (or one deposition scenario to another), there may be very small (or no) changes in the percentage area of habitat exceeded. This is because the magnitude of the exceedance may have reduced, but the area exceeding the critical load remains the same; the area exceeded will only reduce when the critical load is no longer exceeded.

(iii) Accumulated Exceedance (AE)

AE takes account of both the magnitude of exceedance and the habitat area exceeded:

$$AE (\text{keq year}^{-1}) = \text{exceedance} (\text{keq ha}^{-1} \text{ year}^{-1}) * \text{exceeded area (ha)}$$

AE is calculated for each 1km square for each habitat and then summarised by habitat and country. AE is set to zero where critical loads are not exceeded. This metric can be useful for comparing results for different years or scenarios, but because the results are expressed in keq year<sup>-1</sup> they tend to be very large numbers and not intuitive to understand. It should also be noted that the same AE can arise from a large exceedance and small exceeded area, or a small exceedance and a large area.

(iv) Average Accumulated Exceedance (AAE)

AAE averages the AE across the entire sensitive habitat area:

$$AAE (\text{keq ha}^{-1} \text{ year}^{-1}) = AE (\text{keq year}^{-1}) / \text{total habitat area (ha)}$$

This metric provides an exceedance value averaged across the whole habitat area. In the summary statistics presented (Section 2) it is based on the AE for the habitat (by country)

divided by the total habitat area (by country). AAE is set to zero where critical loads are not exceeded. This metric provides a more intuitive value for comparing the exceedance results for different years or scenarios, and gives an indication of the reduction in the magnitude of exceedance even if there is no change in the percentage area of habitat exceeded.

### **1.3.3 Critical load exceedance maps for all habitats combined**


Critical load exceedances are calculated by habitat; exceedance maps can be generated for individual habitats or for all terrestrial habitats combined. The exceedance data for freshwaters are not incorporated into these combination maps because the data are catchment-based rather than for 1km squares and as such may overlap with other habitat data. This section focuses on maps of AAE for all terrestrial habitats combined (Figure 1.5); other maps are presented and discussed in Hall et al. (2015). Maps of AAE provide a good representation of the summary critical load exceedance statistics since they are based on all the critical load values for all habitats and habitat-specific deposition. The AAE for each 1km square is calculated as:

$$AAE = \sum(AE \text{ for all habitats}) / \sum(\text{area for all habitats})$$

AE (and AAE) is set to zero where the critical loads are not exceeded.

The latest AAE maps for acidity and nutrient N (Figure 1.5) clearly show the lower exceedances in Scotland compared to other regions of the UK. High exceedances of acidity critical loads are focused in upland areas of central and north western England, as well as smaller areas in eastern England and the far south-west, as well as parts of Wales and southern Scotland and Northern Ireland. High exceedances of nutrient N critical loads are widespread across England, Wales and Northern Ireland and parts of southern and eastern Scotland, with many areas having exceedances above 14 kg N ha<sup>-1</sup> year<sup>-1</sup> (1 keq ha<sup>-1</sup> year<sup>-1</sup>).

(a) Acidity



(b) Nutrient nitrogen




Figure 1.5: Average Accumulated Exceedance (AAE) of critical loads by CBED deposition for 2015-17. Although the legends for the two maps are given in different units, the class intervals are equivalent (i.e. 7 kg N ha<sup>-1</sup> year<sup>-1</sup> is equivalent to 0.5 keq ha<sup>-1</sup> year<sup>-1</sup>).

## Section 2: Trends in critical loads exceedance by habitat and country

Acidity and nutrient N exceedances by habitat and country are updated annually using the latest 3-year rolling mean CBED deposition data. The summary statistics as described in Section 1.3.1 are made available to Defra and the Devolved Administrations and JNCC; the trends in the percentage area of habitats exceeded are, or have been, used for the following:

- JNCC: Biodiversity Indicator for assessing the pressures from air pollution
- <http://jncc.defra.gov.uk/page-4233>
- Defra: Environmental Statistics – Key Facts  
<https://www.gov.uk/government/statistics/environment-statistics-key-facts>
- Welsh Government: Sustainable Development Indicators for Wales  
<http://wales.gov.uk/topics/statistics/headlines/sustaindev/120829/?lang=en>
- Scottish Government: Key Scottish Environment Statistics  
<http://www.scotland.gov.uk/Topics/Statistics/Browse/Environment/>

The data used for the trends analysis are described briefly in Section 1 and summarised in Box 1 below; there are a few inconsistencies between years due to changes in methods used to derive deposition estimates, and some minor alterations to the acidity critical loads. This information should be taken into account when interpreting the trends results.

**Box 1:****Data used for critical loads trends analysis****Critical loads data**

Acidity: data as summarised in Section 1.1.1 of this report were used for all years except results prior to 2004-2006 where: (a) the acidity critical loads for the bog habitat were based on the dominant soil in each 1x1km grid square; later results use critical loads data that assume all areas of bog habitat occur on peat soils; (b) freshwater exceedances were based on catchment-weighted grid-average deposition; the later results are based on catchment-weighted ecosystem-specific deposition. Note that the freshwater results are based on critical loads for 1752 lake or stream sites across the UK, and therefore do not represent all waters in the UK.

Nutrient nitrogen: data as summarised in Section 1.1.2 of this report.

**Deposition data**

All results based on 5x5 km resolution "concentration based estimated deposition" (CBED) values averaged over a three year period. All data are based on a consistent methodology except:

(a) Deposition data prior to 2001-2003 exclude nitric acid as the monitoring network for this pollutant was not in operation prior to this time.

(b) Deposition data prior to 2002-2004 excludes aerosol deposition of  $NH_4$ ,  $NO_3$ ,  $SO_4$ .

(c) Data for 2004-06 onwards updated in February 2015 to correct for over-estimate of nitric acid deposition.

CBED moorland values are applied to non-woodland terrestrial habitats, and CBED woodland values are applied to woodland habitats.

**Habitat area data**

These are based on the habitat distribution maps generated for UK critical loads research (see Section 1.1 of this report). There was a small reduction in the area mapped for acidity for the bog habitat as a result of the change to the critical loads in 2008; results using the updated habitat area apply to all results from 2004-06 onwards.

The trends results are shown as both tables and simple plots; it is worth noting that while the percentage area exceeded for some habitats may not alter from one year to another, the AAE values fluctuate reflecting changes in the national deposition data.

## 2.1 Trends by country

Table 2.1 shows the total land area by country and the area of habitats sensitive to acidification and eutrophication to which critical loads have been applied; 31% of the UK land area has habitats mapped for acidity critical loads, and 29% for nutrient N. *Note: throughout Section 1 of this report the ‘percentage area exceeded’ represents the percentage area of the habitats mapped as sensitive to acidification or eutrophication (i.e. not % land area of the UK or countries).*

**Table 2.1: Total land area and habitat areas mapped for critical loads by country**

Country	Land area (km <sup>2</sup> ) <sup>#</sup>	Habitat areas mapped for acidity (km <sup>2</sup> )	Area mapped for acidity as % of country	Habitat areas mapped for nutrient nitrogen (km <sup>2</sup> )	Area mapped for nutrient nitrogen as % of country
England	132938	18635	14	19522	15
Wales	21225	7798	37	6837	32
Scotland	80239	48083	60	43200	54
NI	14130	3541	25	3467	25
UK	248532	78051	31	73027	29

<sup>#</sup>The UK and its countries: facts and figures. Office for National Statistics:

<http://webarchive.nationalarchives.gov.uk/20160105160709/http://www.ons.gov.uk/ons/guide-method/geography/beginner-s-guide/administrative/the-countries-of-the-uk/index.html>

### 2.1.1 Acidity results

The results for acidity (Table 2.2, Figure 2.1) show that the total area of habitats exceeding critical loads in the UK has declined by almost half, from 72.6% in 1996 to 37.6% in 2015-17. However, the area exceeded varies between countries (Table 2.2, Figure 2.2), due to (a) geographic location of different sensitive habitats across the country (see Section 2.2); (b) variability in critical load values across the country – lower critical loads associated with habitats on more acid soils; (c) higher deposition found in central and south-west England, parts of Wales and Northern Ireland and south-west Scotland (Figure 1.1). The percentage area of habitats exceeded is lowest in Scotland in all years; however as shown in Table 2.1, 60% of Scotland has habitats mapped for acidity critical loads, and that means the actual areas exceeded are larger than in the other countries (e.g. 10,933 km<sup>2</sup> exceeded by 2015-17 deposition). Although only 14% of England has habitats mapped for acidity critical loads, 57.9% of their area is exceeded for 2015-17, equivalent to 10,782 km<sup>2</sup>. The magnitude of exceedance across the UK, expressed as AAE (Table 2.3, Figure 2.1), has fallen by two-thirds from 0.78 keq ha<sup>-1</sup> year<sup>-1</sup> in 1996 to 0.21 keq ha<sup>-1</sup> year<sup>-1</sup> in 2016. The data show the largest reductions in the exceedances were in the late 1990s; changes since then have been smaller and fluctuated from one year to another, and although continuing the general downward trend there has been very little overall change in the last decade. This pattern of exceedance is consistent with the trends in deposition (Section 1.2.1, Fig 1.2).

**Table 2.2: Acid-sensitive habitat area and percentage area of habitats where acidity critical loads are exceeded, by country and deposition dataset year.**

Year	Percentage acid-sensitive habitat area exceeded by country: <i>[total area (km<sup>2</sup>) acid-sensitive habitats by country]</i>				
	England	Wales	Scotland	NI	UK
	<i>[18635]</i>	<i>[7798]</i>	<i>[48083]</i>	<i>[3537]</i>	<i>[78051]</i>
1995-1997	75.8	90.0	68.2	76.8	72.6
1998-2000	71.6	83.1	52.6	67.2	60.8
1999-2001	71.9	83.0	51.6	66.8	60.3
2001-2003	72.3	82.4	43.0	67.4	55.0
2002-2004	72.3	82.3	44.8	69.2	56.2
2003-2005	71.8	83.2	44.5	67.1	55.9
2004-2006	66.8	81.2	48.0	68.1	56.7
2005-2007	66.1	81.0	46.1	68.5	55.4
2006-2008	64.3	79.2	40.7	68.6	51.4
2007-2009	63.6	77.4	32.9	69.4	46.3
2008-2010	63.2	74.9	31.5	69.6	45.2
2009-2011	63.8	74.5	33.9	71.0	46.8
2010-2012	62.8	74.2	32.2	67.8	45.3
2011-2013	62.1	74.4	31.0	69.4	44.5
2012-2014	61.3	73.4	31.0	67.1	44.1
2013-2015	60.6	72.4	31.4	66.9	44.1
2014-2016	60.0	72.4	29.6	64.3	42.7
2015-2017	57.9	70.0	22.7	61.2	37.6
<b>Reduction in % area exceeded from 1996-2016</b>	<b>17.9</b>	<b>20.0</b>	<b>45.5</b>	<b>15.6</b>	<b>35.0</b>

**Table 2.3: Acidity Average Accumulated Exceedance (AAE in keq ha<sup>-1</sup> year<sup>-1</sup>) by country and deposition dataset year.**

Year	AAE (keq ha <sup>-1</sup> year <sup>-1</sup> ) by country:				
	England	Wales	Scotland	NI	UK
1995-1997	1.33	1.36	0.47	0.80	0.8
1998-2000	1.00	0.84	0.28	0.46	0.5
1999-2001	0.98	0.82	0.27	0.46	0.5
2001-2003	1.04	0.82	0.23	0.51	0.5
2002-2004	0.94	0.79	0.24	0.46	0.5
2003-2005	0.93	0.84	0.24	0.42	0.5
2004-2006	0.77	0.74	0.24	0.42	0.4
2005-2007	0.74	0.73	0.21	0.45	0.4
2006-2008	0.68	0.61	0.17	0.44	0.4
2007-2009	0.62	0.54	0.12	0.45	0.3
2008-2010	0.59	0.49	0.12	0.47	0.29
2009-2011	0.62	0.48	0.15	0.53	0.31
2010-2012	0.60	0.47	0.14	0.46	0.30
2011-2013	0.59	0.47	0.13	0.46	0.29
2012-2014	0.55	0.46	0.13	0.39	0.28
2013-2015	0.54	0.45	0.15	0.38	0.28
2014-2016	0.51	0.44	0.13	0.36	0.26
2015-2017	0.44	0.39	0.09	0.33	0.21
<b>Reduction in AAE 1996-2015</b>	<b>0.89</b>	<b>0.97</b>	<b>0.38</b>	<b>0.47</b>	<b>0.57</b>




Figure 2.1: Acidity: Percentage area of acid-sensitive habitats with exceedance of acidity critical loads in the UK by year, and AAE in keq ha<sup>-1</sup> year<sup>-1</sup>.




Figure 2.2: Acidity: Percentage area of acid-sensitive habitats with exceedance of acidity critical loads, by country and year, and AAE in keq ha<sup>-1</sup> year<sup>-1</sup>

### 2.1.2 Nutrient nitrogen results

The results for nutrient N (Table 2.4 and Figure 2.3) show a decline in the percentage area of habitats exceeded in the UK, from 75% in 1996 to 58.1% in 2016. The results for England and Wales remained above, or close to, 90% exceeded over the same time period (Table 2.4, Figure 2.4). Scotland shows the smallest percentage habitat area exceeded of all countries, but the area exceeded (15,002 km<sup>2</sup> for 2016) is similar to the area exceeded in England (18,504 km<sup>2</sup> in 2016). The results reflect the smaller reductions in N deposition over the last two decades compared to the reductions in S deposition, which helped reduce the exceedances of acidity critical loads.

**Table 2.4: Nitrogen-sensitive habitat area and percentage area of habitats where nutrient nitrogen critical loads are exceeded, by country and deposition dataset year.**

Year	Percentage habitat area exceeded by country: <i>[total area (km<sup>2</sup>) nitrogen-sensitive habitats by country]</i>				
	England <i>[19522]</i>	Wales <i>[6837]</i>	Scotland <i>[43200]</i>	NI <i>[3467]</i>	UK <i>[73027]</i>
1995-1997	98.3	98.0	59.4	92.6	75.0
1998-2000	97.6	92.5	48.9	80.0	67.5
1999-2001	97.7	91.1	50.9	82.5	68.7
2001-2003	97.8	93.5	47.7	85.4	67.1
2002-2004	97.6	93.3	50.2	86.3	68.6
2003-2005	97.5	94.1	50.6	83.8	68.8
2004-2006	96.7	93.2	52.9	84.8	69.9
2005-2007	96.5	93.6	53.6	86.4	70.4
2006-2008	96.1	92.9	49.0	86.8	67.5
2007-2009	96.4	91.7	41.8	88.7	63.3
2008-2010	96.5	89.7	40.7	89.7	62.6
2009-2011	97.0	89.8	44.5	91.4	65.0
2010-2012	96.5	89.6	41.4	88.5	62.9
2011-2013	96.0	90.3	40.7	89.9	62.5
2012-2014	95.9	89.4	40.9	86.4	62.3
2013-2015	95.8	88.5	42.8	87.4	63.4
2014-2016	95.7	88.5	40.8	85.1	62.1
2015-2017	94.8	88.0	34.7	84.2	58.1
<i>Reduction in % area exceeded 1996-2016</i>	<i>3.5</i>	<i>10.0</i>	<i>24.7</i>	<i>8.4</i>	<i>16.9</i>

The magnitude of the nutrient-nitrogen exceedance (expressed as AAE) across the UK has decreased by more than one-third, from 9.5 kg N ha<sup>-1</sup> year<sup>-1</sup> in 1996 to 5.6 kg N ha<sup>-1</sup> year<sup>-1</sup> in 2016 (Table 2.5, Figure 2.3). The AAE varied among regions, with lowest values in Scotland and highest in England (Table 2.5, Figure 2.4). The UK AAE decreased by 0.9 kg N ha<sup>-1</sup> yr<sup>-1</sup> in the decade between 2000 and 2010, and by 1.2 kg N ha<sup>-1</sup> yr<sup>-1</sup> between 2000 and 2016.

**Table 2.5: Nutrient nitrogen Average Accumulated Exceedance (AAE in kg N ha<sup>-1</sup> year<sup>-1</sup>) by country and deposition dataset year.**

Year	AAE (kg N ha <sup>-1</sup> year <sup>-1</sup> ) by country:				
	England	Wales	Scotland	NI	UK
1995-1997	19.0	15.8	4.1	10.6	9.5
1998-2000	16.8	10.3	2.7	6.5	7.4
1999-2001	17.4	10.6	2.9	6.8	7.7
2001-2003	19.7	12.2	3.1	8.9	8.7
2002-2004	18.0	12.2	3.3	8.7	8.3
2003-2005	18.2	13.2	3.3	8.3	8.4
2004-2006	14.9	11.4	3.1	7.9	7.2
2005-2007	14.9	11.4	2.9	8.8	7.2
2006-2008	14.1	9.9	2.5	8.8	6.6
2007-2009	13.8	9.5	2.1	9.4	6.3
2008-2010	13.9	9.2	2.2	9.8	6.3
2009-2011	14.6	9.2	2.6	10.9	6.8
2010-2012	13.8	8.8	2.4	9.6	6.4
2011-2013	13.3	8.9	2.3	9.5	6.2
2012-2014	12.6	8.6	2.3	8.3	5.9
2013-2015	12.8	8.9	2.7	8.4	6.2
2014-2016	12.7	8.9	2.6	8.2	6.1
2015-2017	12.2	8.6	1.9	7.9	5.6
Reduction (kg N ha <sup>-1</sup> yr <sup>-1</sup> ) in AAE 1996-2016	6.8	7.2	2.2	2.8	3.9




Figure 2.3: Nutrient nitrogen: Percentage area of nitrogen-sensitive habitats with exceedance of nitrogen critical loads in the UK by year, and AAE in kg N ha<sup>-1</sup> year<sup>-1</sup>.




Figure 2.4: Nutrient nitrogen: Percentage area of nitrogen-sensitive habitats with exceedance of nitrogen critical loads, by country and year, and AAE in kg N ha<sup>-1</sup> year<sup>-1</sup>.

## **2.2 Trends by habitat**

This section focuses on the results by habitat for the UK; habitat results for individual countries are also calculated and supplied to Defra and the devolved administrations.

### **2.2.1 Acidity results**

Although acidity critical loads are applied to calcareous grassland there is no exceedance of these critical loads in any year, and therefore this habitat is excluded from Tables 2.6 and 2.7 and Figures 2.5 and 2.6. The habitats with the highest percentage area exceeded are acid grassland, montane, bog and managed woodlands (Table 2.6, Figure 2.5); these habitats also have some of the highest AAE values (Table 2.7, Figure 2.5). Of the habitats mapped for acidity, dwarf shrub heath is the habitat with the largest cover across the UK (10%), and also shows the largest decrease in the area exceeded (more than halved), from 70.3% (17,370 km<sup>2</sup>) in 1996 to 20.7% (5,119 km<sup>2</sup>) in 2016. The largest reductions in AAE over the same timescale are for woodland, acid grassland and montane habitats (Table 2.7).

### **2.2.2 Nutrient nitrogen results**

There are six habitats with more than 80% of their area exceeded for nutrient N in all years (Table 2.8, Figure 2.6): calcareous grasslands and different woodland habitats (beech, oak, managed conifer and broadleaf and other unmanaged woodland). The largest reduction in the percentage area exceeded is for dune grassland from 70.6% in 1996 to 22.4% in 2016, however, this habitat only occupies <1% of the total area of N-sensitive habitats mapped. Another coastal habitat, saltmarsh, has virtually no exceedance in any year, due to a combination of its high critical load and the lower deposition in coastal areas. AAE is generally highest for the woodland habitats (Table 2.9, Figure 2.6), with the exception of Scots Pine, which is only found in Scotland where the magnitude of exceedance is generally lower due to the lower deposition in this region. The beech woodland is virtually 100% exceeded in all years, but the AAE has nearly halved from 22.7 kg N ha<sup>-1</sup> year<sup>-1</sup> in 1996 to 12.1 kg N ha<sup>-1</sup> year<sup>-1</sup> in 2016.


**Table 2.6: Acid-sensitive habitat area in the UK and percentage area of habitats where acidity critical loads are exceeded, by deposition dataset year.**

Acid-sensitive habitat areas in the UK and percentage habitat area with exceedance of acidity critical loads:									
Parameter	Acid grassland	Dwarf shrub heath	Bog	Montane	Coniferous woodland (managed)	Broadleaved woodland (managed)	Unmanaged woodland	Freshwaters	All habitats
<i>Habitat area (km<sup>2</sup>)</i>	15336	24705	5454	3054	8374	7452	4011	7857	78051
1995-1997	92.0	70.3	88.0	95.8	79.4	75.8	69.5	29.9	72.6
1998-2000	84.9	49.5	78.8	91.3	69.9	68.4	57.2	24.2	60.8
1999-2001	84.7	47.9	76.1	93.4	70.2	69.1	58.2	23.9	60.3
2001-2003	79.5	40.7	61.6	82.5	66.5	69.6	58.5	21.9	55.0
2002-2004	80.3	42.3	60.1	89.4	68.4	70.8	60.1	21.3	56.2
2003-2005	80.4	41.5	59.3	92.9	68.2	70.1	59.1	21.7	55.9
2004-2006	82.5	45.1	71.7	96.3	64.2	61.5	48.3	21.7	56.7
2005-2007	81.9	41.5	76.4	94.4	63.6	60.5	46.7	21.3	55.4
2006-2008	78.9	35.4	73.3	85.6	60.5	57.2	43.4	20.6	51.4
2007-2009	73.9	28.5	63.7	71.4	57.1	55.9	42.1	19.0	46.3
2008-2010	72.2	28.1	57.9	70.1	55.6	55.6	42.0	18.5	45.2
2009-2011	74.8	30.6	54.9	71.6	58.0	57.0	43.3	18.9	46.8
2010-2012	73.0	29.3	54.4	65.3	56.7	55.0	41.5	19.0	45.3
2011-2013	73.4	28.5	50.1	62.3	56.2	53.5	40.5	18.8	44.5
2012-2014	72.6	27.8	55.5	62.7	54.9	51.6	38.5	18.6	44.1
2013-2015	73.0	28.1	52.4	65.0	55.8	51.0	37.9	18.2	44.1
2014-2016	71.1	26.4	50.1	62.1	54.7	50.5	36.8	18.5	42.7
2015-2017	65.4	20.7	44.0	43.8	51.1	48.1	34.0	15.6	37.6
Reduction in % area exceeded 1996-2016	26.6	49.6	43.9	52.0	28.3	27.7	35.5	14.3	35.0


**Table 2.7: Acidity AAE (in keq ha<sup>-1</sup> year<sup>-1</sup>) by habitat for the UK by deposition dataset year.**

AAE (keq AAE (keq ha <sup>-1</sup> year <sup>-1</sup> ) by habitat:									
Year	Acid grassland	Dwarf shrub heath	Bog	Montane	Coniferous woodland (managed)	Broadleaved woodland (managed)	Unmanaged woodland	Freshwaters	All habitats
1995-1997	1.15	0.47	0.76	0.81	1.13	1.20	0.87	0.36	0.78
1998-2000	0.80	0.28	0.53	0.57	0.68	0.88	0.58	0.23	0.51
1999-2001	0.77	0.26	0.50	0.59	0.68	0.90	0.61	0.21	0.50
2001-2003	0.70	0.24	0.46	0.60	0.72	1.01	0.68	0.18	0.50
2002-2004	0.67	0.22	0.41	0.64	0.74	0.94	0.65	0.17	0.48
2003-2005	0.68	0.21	0.39	0.62	0.73	0.94	0.65	0.17	0.47
2004-2006	0.68	0.22	0.44	0.66	0.58	0.66	0.44	0.17	0.43
2005-2007	0.64	0.19	0.45	0.53	0.56	0.65	0.43	0.16	0.40
2006-2008	0.57	0.16	0.42	0.39	0.49	0.56	0.36	0.13	0.35
2007-2009	0.49	0.12	0.34	0.28	0.43	0.53	0.34	0.12	0.30
2008-2010	0.47	0.12	0.33	0.28	0.42	0.52	0.34	0.11	0.29
2009-2011	0.51	0.14	0.35	0.31	0.46	0.56	0.36	0.12	0.31
2010-2012	0.50	0.13	0.35	0.26	0.43	0.51	0.32	0.12	0.30
2011-2013	0.51	0.13	0.34	0.25	0.42	0.47	0.30	0.12	0.29
2012-2014	0.48	0.12	0.33	0.25	0.39	0.43	0.27	0.12	0.28
2013-2015	0.49	0.13	0.34	0.26	0.41	0.42	0.26	0.11	0.28
2014-2016	0.47	0.12	0.31	0.23	0.39	0.41	0.25	0.11	0.26
2015-2017	0.38	0.08	0.25	0.14	0.32	0.38	0.23	0.09	0.21
Reduction in AAE 1996-2016	0.77	0.39	0.51	0.67	0.81	0.82	0.64	0.27	0.57


a) Acid grassland




b) Dwarf shrub heath




c) Bog




d) Montane




e) Managed coniferous woodland



f) Managed broadleaved woodland



g) Unmanaged woodland



h) Freshwaters




Figure 2.5: Acidity: Percentage area of habitats where acidity critical loads are exceeded, and acidity AAE, for the UK by deposition dataset year.

**Table 2.8: Nutrient-sensitive habitat area in the UK and percentage area of habitats where nutrient nitrogen critical loads are exceeded, by deposition dataset year.**

Nitrogen-sensitive habitat areas in the UK and percentage habitat area with exceedance of nutrient nitrogen critical loads:

Parameter	Acid grassland	Calcareous grassland	Dwarf shrub heath	Bog	Montane	Coniferous woodland (managed)	Broadleaved woodland (managed)	Beech woodland (unmanaged)	Acidophilous oak (unmanaged)	Scots Pine (unmanaged)	Other unmanaged woodland	Dune grassland	Saltmarsh
<i>Habitat area (km<sup>2</sup>)</i>	15235	3578	24826	5526	3129	8383	7482	719	1434	204	1761	323	427
1995-1997	72.6	97.5	59.1	54.2	96.7	95.4	98.4	100.0	98.9	61.1	96.5	70.6	2.0
1998-2000	61.3	95.5	49.0	45.1	95.7	90.5	97.4	100.0	97.0	38.9	95.1	44.8	1.1
1999-2001	61.4	95.5	51.1	45.0	97.1	92.8	97.8	100.0	98.1	52.3	95.5	46.9	2.1
2001-2003	63.1	95.5	47.8	44.6	89.0	90.6	97.4	100.0	96.1	49.7	95.5	41.9	1.0
2002-2004	64.3	93.9	49.8	44.9	92.6	93.0	98.1	100.0	98.2	66.5	95.7	36.1	1.1
2003-2005	64.8	93.9	50.6	45.2	90.5	92.1	98.0	100.0	98.1	67.8	95.7	33.5	1.1
2004-2006	64.8	90.6	54.5	45.9	96.6	90.2	97.5	100.0	95.6	58.0	95.5	29.3	0.8
2005-2007	64.2	89.4	54.3	54.6	96.2	91.0	97.4	100.0	95.5	52.6	95.6	31.8	0.8
2006-2008	60.0	87.7	49.5	55.4	95.5	89.4	97.1	100.0	93.8	34.2	95.5	31.1	0.8
2007-2009	56.3	89.6	43.9	47.1	82.7	86.9	96.7	100.0	89.8	30.7	95.2	29.2	0.9
2008-2010	55.7	91.2	42.7	45.6	81.0	86.1	96.7	99.9	88.5	30.5	95.1	34.7	0.9
2009-2011	61.1	92.3	45.0	45.8	82.1	88.2	97.0	99.9	91.5	32.4	95.3	37.6	0.9
2010-2012	59.7	90.4	42.2	44.8	74.4	86.5	96.8	99.9	87.7	26.2	94.7	34.0	0.9
2011-2013	60.8	87.6	41.6	43.1	71.2	86.4	96.8	100.0	88.6	24.2	95.0	29.2	0.8
2012-2014	59.1	88.2	41.6	45.4	74.2	85.6	96.6	99.9	87.6	26.0	95.0	25.3	0.7
2013-2015	60.8	87.5	43.0	45.6	78.4	86.0	96.7	100.0	88.5	26.4	95.0	26.9	0.8
2014-2016	59.5	87.2	41.5	44.5	72.3	84.9	96.4	99.8	87.6	24.0	94.8	26.4	0.8
2015-2017	54.4	86.2	36.8	40.9	62.0	82.0	95.9	99.9	83.0	20.8	93.9	22.4	1.0
Reduction in % area exceeded 1996-2016	13.1	10.3	17.4	9.7	22.4	10.2	2.0	0.2	9.9	37.1	1.5	44.2	1.1

**Table 2.9: Nutrient nitrogen: AAE (in kg N ha<sup>-1</sup> year<sup>-1</sup>) by habitat for the UK by deposition dataset year.**

AAE (kg N ha <sup>-1</sup> year <sup>-1</sup> ) by habitat:													
Year	Acid grassland	Calcareous grassland	Dwarf shrub heath	Bog	Montane	Coniferous woodland (managed)	Broadleaved woodland (managed)	Beech woodland (unmanaged)	Acidophilous oak (unmanaged)	Scots Pine (unmanaged)	Other unmanaged woodland	Dune grassland	Saltmarsh
1995-1997	6.3	7.6	4.5	5.3	5.5	16.8	24.5	22.7	19.9	3.3	23.2	2.7	0.0
1998-2000	3.9	7.3	3.1	3.8	4.4	12.1	21.8	19.5	16.4	2.0	21.1	1.6	0.1
1999-2001	4.0	7.7	3.2	3.9	5.0	12.8	22.7	20.3	17.3	2.8	22.0	1.7	0.1
2001-2003	4.5	8.9	3.6	4.4	5.7	14.4	25.8	22.9	18.8	3.2	25.2	1.5	0.0
2002-2004	4.2	6.9	3.4	3.8	6.1	14.7	24.5	22.1	19.1	4.0	23.8	0.9	1.7
2003-2005	4.4	6.9	3.4	3.8	6.1	14.8	24.8	22.6	19.4	3.7	24.1	0.9	1.7
2004-2006	4.4	5.7	3.4	3.9	6.4	12.2	19.3	15.8	15.5	2.6	18.7	0.8	0.0
2005-2007	4.3	5.7	3.3	4.0	5.5	12.3	19.4	15.4	15.4	2.3	19.1	0.8	0.0
2006-2008	3.9	5.2	3.0	4.0	4.3	11.5	18.2	14.0	14.2	1.9	18.1	0.7	0.0
2007-2009	3.5	5.3	2.6	3.5	3.3	10.8	18.3	14.4	13.9	1.6	18.5	0.8	0.0
2008-2010	3.4	5.5	2.6	3.5	3.3	10.9	18.5	14.6	13.9	1.7	18.9	0.9	0.1
2009-2011	3.9	5.9	3.0	3.9	3.6	11.8	19.4	15.2	14.7	1.9	19.9	1.0	0.1
2010-2012	3.7	5.3	2.8	3.7	2.9	11.2	18.1	13.9	13.7	1.6	18.4	0.9	0.1
2011-2013	3.7	4.9	2.8	3.7	2.9	11.0	17.3	13.3	13.5	1.5	17.5	0.8	0.0
2012-2014	3.5	4.7	2.7	3.6	3.0	10.5	16.3	12.4	12.7	1.6	16.4	0.6	0.0
2013-2015	3.8	4.6	2.9	3.8	3.4	11.2	16.5	12.3	13.1	1.7	16.7	0.7	0.0
2014-2016	3.8	4.7	2.8	3.7	3.1	10.9	16.6	12.1	12.8	1.5	16.7	0.7	0.0
2015-2017	3.2	4.4	2.3	3.2	2.1	10.0	16.4	12.1	12.2	1.0	16.7	0.6	0.0
Reduction in AAE from 1996 to 2016	3.1	3.2	2.2	2.1	3.4	6.8	8.1	10.6	7.7	2.3	6.5	2.1	0.0






Figure 2.6: Nutrient nitrogen: Percentage area of habitats where nutrient nitrogen critical loads are exceeded and nutrient nitrogen AAE (in kg N ha<sup>-1</sup> year<sup>-1</sup>) in the UK by deposition dataset year.


g) Managed coniferous woodland




h) Managed broadleaved woodland




i) Unmanaged beech woodland



j) Unmanaged acidophilous oak woodland



k) Unmanaged Scots Pine woodland



l) Other unmanaged woodland




Figure 2.6 (continued): Nutrient nitrogen: Percentage area of habitats where nutrient nitrogen critical loads are exceeded and nutrient nitrogen AAE (in kg N ha<sup>-1</sup> year<sup>-1</sup>) in the UK by deposition dataset year.

### Section 3: Site-relevant critical loads and their exceedances

This section of the report focuses on the application of critical loads to designated sites, hereafter referred to as site-relevant critical loads (SRCL), and their exceedances. The critical loads are based on the same methods applied to UK acid- and N-sensitive habitats described in Section 1 of this report, and are applied to acid- and N-sensitive features within the designated sites. Exceedances of critical loads are also calculated in the same way as the habitats (Section 1) and also based on UK 5x5 km CBED deposition, however, some different metrics are used to describe the exceedance results for SRCL and are explained below.

#### 3.1 Overview of site-relevant critical loads

Site relevant critical loads (SRCL) have been applied to three types of statutory protected sites:

- Special Areas of Conservation (SACs) are protected sites designated under the EC Habitats Directive. Annexes I and II of the Directive identify the habitats and species (excluding birds) to be protected; 78 Annex I habitat types and 41 species are believed to occur in, or be native to the UK.
- Special Protected Areas (SPAs) are sites classified under the EC Birds Directive to protect rare and vulnerable birds (as listed in an Annex to the Directive) and regularly occurring migratory species.
- Sites of Special Scientific Interest (SSSIs in England, Wales and Scotland) and Areas of Special Scientific Interest (ASSIs in Northern Ireland) provide statutory protection to the UK's flora and fauna. There are additional SSSIs designated for geological or physiographic features but these are not included in the SRCL assessments.

Digital boundaries for all sites in the UK have been collated by JNCC, together with tables identifying the designated feature habitats and species associated with each site, but no digital information is currently available on the spatial area of each feature within each site. ***Therefore, for the purposes of the national SRCL work described here, it is assumed that all features recorded for a site, occur across the entire site area. To avoid double counting the area exceeding critical loads for sites with more than one designated feature, the maximum area exceeded for any feature is used when summarising results to the site and country levels (Section 3.2).***

To assign SRCL, the first step is to consider if the interest feature is potentially sensitive to acidification and/or eutrophication. Specialists within Natural England, Scottish Natural Heritage and CEH have used expert judgement to determine this (SNIFFER 2007). For SPAs where the features are bird species, the broad habitats the birds depend upon for feeding, breeding and roosting are considered.

To assign critical loads to the habitat features of designated sites it is necessary to link the different habitat classifications used. Acidity critical loads are mapped by broad habitat and empirical critical loads of N are based on the EUNIS (European Nature Information System: Moss & Davies 2002) habitat classification. Look-up tables developed by Moss & Davies (2002) and published in the National Biodiversity Network (NBN) Habitats Dictionary (<http://habitats.nbn.org.uk/>) and available from the JNCC website (<http://jncc.defra.gov.uk/page-1425>) enable linkages to be made between:

- Annex I habitats and EUNIS classes
- Annex I habitats and broad habitats
- EUNIS habitats and broad habitats

Using the look up tables the most appropriate EUNIS class and broad habitat class can be assigned to each interest feature. It should be noted that some sites may contain features sensitive to acidification and/or eutrophication for which no appropriate critical loads are available.

The critical loads assigned to the habitat features are based on the same methods and data as those outlined in Section 1.1 of this report. However, the national critical load maps are based on national scale data sets appropriate for national scale critical load and critical level assessments. This means they may not include all small areas of sensitive habitats or some coastal habitats; therefore some designated sites and/or feature habitats may not be included in the areas mapped nationally for critical loads. To overcome this, for SRCL a separate database of national critical loads for terrestrial habitats was created, that provides critical loads for every 1x1km square in the UK whether the habitat is known to exist there or not. The appropriate SRCL can then be extracted for terrestrial habitat features of each designated site; the SRCL does not include any acidity critical loads for freshwater habitats. For further information refer to the “Methods” report (Hall et al. 2015).

For nutrient N the empirical critical loads approach is applied to designated feature habitats sensitive to N. The critical load value applied to each habitat are the “Recommended” values agreed by habitat specialists for Article 17 reporting ((for more information refer to <http://www.apis.ac.uk/indicative-critical-load-values> and to Hall et al. 2015).

### 3.2 Overview of SRCL exceedance metrics

Exceedances are calculated separately for SACs, SPAs and SSSIs, for all site features that critical loads and deposition data can be assigned to (Hall et al. 2015). Metrics are calculated by:

(a) Feature (within each site):

- Exceedance
- Exceeded area<sup>#</sup>
- Accumulated Exceedance (AE)(i.e. exceedance \* exceeded area)
- Average Accumulated Exceedance (AAE)(i.e. AE / total site area)

(b) Site:

- Total number of features with SRCL
- Number and percentage of features with exceedance of SRCL.
- Maximum area exceeded<sup>##</sup> for any feature within a site
- Maximum AE for any feature within a site
- Maximum AAE for any feature within a site

(c) Country:

- Total number of sites
- Total number and percentage of sites with SRCL for one or more features

- Total number of features with SRCL
- Total number and percentage of sites with exceedance of SRCL for one or more features
- Total number and percentage of features with exceedance of SRCL
- Total area of all sites
- Total area of all sites with SRCL
- Maximum exceeded area<sup>###</sup>
- Maximum AE calculated as the sum of the maximum AE for all sites
- Maximum AAE; calculated from the country maximum AE and total area of all sites (with SRCL) within a country.

# Feature exceeded area: If the critical load is exceeded and the deposition values are constant across the whole site, the exceeded area equals the site area; if the deposition values vary across the site (e.g. as a result of the site crossing the boundaries between different 5x5km grid squares with different deposition values), then the exceeded area will be the sum of the 1x1 km portions of the site where the deposition exceeds the critical load.

## Site maximum exceeded area: is set to the maximum exceeded area for any feature within a site.

### Country maximum exceeded area: is calculated as the sum of the site maximum exceeded areas for all sites within a country.

The sections below summarise the key results by country, based on the CBED deposition (Section 1.2) for 1996 to 2016. ***Note that the summary statistics and maps may present the “worst” case, as they are based on sites where at least one feature is exceeded; other features within a site may (a) have a smaller exceedance or (b) not be exceeded. In addition, the AAE results are based on the maximum exceedance of any feature within a site.***

### 3.2.1 Acidity results

The trends in acidity critical load exceedances are summarised in Tables 3.1-3.3 and present the percentage of sites (with SRCL) by country, where the SRCL is exceeded for one or more features, together with the maximum AAE.

The percentage of SACs with critical load exceedance decreased between 1996 and 2016 by ~4% (Wales and NI) to 36.3% (Scotland), while the AAE fell by between a half (England and NI) to over three-quarters (Scotland). These trends reflect the changing patterns of acid deposition over this time period. For SPAs the largest reductions in the percentage of exceeded sites were 40.7% (Scotland), 30.8 (Wales) and 30.0% (NI) accompanied by a roughly 80-90% reduction in their maximum AAE values. The reductions in England were slightly less: 12.7% reduction in percentage of exceeded sites, and 59% reduction in maximum AAE.

At the UK level, the trends results show:

(i) for SACs, the percentage of sites with acidity exceedance decreased from 91.0% in 1996 to 71.5% in 2016, and the maximum AAE fell by 66% from 1.51 keq ha<sup>-1</sup> year<sup>-1</sup> to 0.51 keq ha<sup>-1</sup> year<sup>-1</sup> over the same time period;

(ii) for SPAs, the percentage of exceeded sites fell by a third from 94.3% in 1996 to 65.7% in 2016, and the maximum AAE decreased by 68% from 1.11 keq ha<sup>-1</sup> year<sup>-1</sup> in 1996 to 0.36 keq ha<sup>-1</sup> year<sup>-1</sup> in 2016.

(iii) for SSSIs, the percentage of exceeded sites decreased from 77.6% in 1996 to 57.1% in 2016, and the maximum AAE fell by 68% from 1.18 keq ha<sup>-1</sup> year<sup>-1</sup> in 1996 to 0.37 keq ha<sup>-1</sup> year<sup>-1</sup> in 2016.

Maps of the maximum AAE per site (Figure 3.1) based on the latest CBED deposition (2016) show the highest exceedances mainly in northern England and parts of Wales and south-west England and southern Scotland. Some sites in the far north of Scotland, a few SACs and SPAs in southern England, and many small SSSIs across central and eastern England have no exceedance for any site feature.

**Table 3.1: Trends in acidity exceedances for SACs; percentage of sites (with SRCL) with exceedance of SRCL for at least one feature, and maximum AAE of all sites/features: (a) % of sites with SRCL with exceedance of SRCL for at least one feature, (b)[maximum AAE keq ha<sup>-1</sup> year<sup>-1</sup>]. NR = Not recorded.**

	England	Wales	Scotland	NI	Eng/Wales*	Eng/Scot*
Number of sites	231	85	236	54	7	3
Number of sites with SRCL for at least one feature	180	71	182	47	6	1
1995-1997	85.0 [2.36]	97.2 [1.87]	92.3 [0.66]	97.9 [1.32]	100.0 [NR]	100.0 [NR]
1998-2000	82.2 [1.80]	97.2 [1.29]	83.5 [0.42]	95.7 [0.76]	100.0 [NR]	100.0 [NR]
1999-2001	81.7 [1.83]	97.2 [1.31]	83.5 [0.44]	95.7 [0.78]	100.0 [NR]	100.0 [NR]
2001-2003	81.1 [1.89]	94.4 [1.31]	75.3 [0.41]	95.7 [0.87]	100.0 [NR]	100.0 [NR]
2002-2004	82.8 [1.77]	95.8 [1.27]	78.0 [0.43]	95.7 [0.77]	100.0 [NR]	100.0 [NR]
2003-2005	82.8 [1.75]	95.8 [1.33]	76.4 [0.42]	95.7 [0.71]	100.0 [NR]	100.0 [NR]
2004-2006	79.4 [1.50]	95.8 [1.08]	79.7 [0.42]	95.7 [0.70]	100.0 [NR]	100.0 [NR]
2005-2007	79.4 [1.45]	95.8 [1.05]	79.7 [0.38]	95.7 [0.73]	100.0 [NR]	100.0 [NR]
2006-2008	77.2 [1.35]	95.8 [0.90]	75.8 [0.31]	95.7 [0.71]	100.0 [NR]	100.0 [NR]
2007-2009	76.7 [1.21]	95.8 [0.82]	69.2 [0.22]	95.7 [0.72]	100.0 [NR]	100.0 [NR]
2008-2010	75.6 [1.16]	95.8 [0.77]	67.6 [0.22]	95.7 [0.75]	100.0 [NR]	100.0 [NR]
2009-2011	76.1 [1.20]	95.8 [0.75]	70.3 [0.23]	95.7 [0.79]	100.0 [NR]	100.0 [NR]
2010-2012	76.1 [1.17]	93.0 [0.75]	68.1 [0.21]	93.6 [0.72]	100.0 [NR]	100.0 [NR]
2011-2013	75.0 [1.18]	93.0 [0.75]	68.1 [0.19]	95.7 [0.72]	100.0 [NR]	100.0 [NR]
2012-2014	74.4 [1.15]	94.4 [0.73]	68.1 [0.19]	95.7 [0.66]	100.0 [NR]	100.0 [NR]
2013-2015	72.2 [1.13]	93.0 [0.67]	67.6 [0.19]	95.7 [0.63]	100.0 [NR]	100.0 [NR]
2014-2016	72.2 [1.07]	93.0 [0.65]	65.4 [0.17]	93.6 [0.60]	100.0 [NR]	100.0 [NR]
2015-2017	71.7 [0.92]	93.0 [0.58]	56.0 [0.12]	93.6 [0.56]	100.0 [NR]	100.0 [NR]
Reduction in % area exceeded from 1996 to 2016	13.3	4.2	36.3	4.3	0.0	0.0
[Reduction in AAE in same period]	[1.44]	[1.29]	[0.54]	[0.76]	[NR]	[NR]

\* Some sites cross the England/Wales or England/Scotland border and have been assigned to these border areas. However, in calculating AAE each 1x1 km square (or part thereof) within each site has been assigned to a single country, so AAE results are calculated for individual countries only.


**Table 3.2: Trends in acidity exceedances for SPAs; percentage of sites (with SRCL) with exceedance of SRCL for at least one feature, and maximum AAE of all sites/features: (a) % of sites with SRCL with exceedance of SRCL for at least one feature, (b)[maximum AAE keq ha<sup>-1</sup> year<sup>-1</sup>]. NR = Not recorded.**

	England	Wales	Scotland	NI	Eng/Wales*
Number of sites	78	17	145	14	3
Number of sites with SRCL for at least one feature	63	13	86	10	3
1995-1997	98.4 [1.73]	100.0 [1.85]	89.5 [0.45]	100.0 [1.09]	100.0 [NR]
1998-2000	98.4 [1.41]	100.0 [1.14]	72.1 [0.24]	100.0 [0.43]	100.0 [NR]
1999-2001	98.4 [1.39]	100.0 [1.21]	73.3 [0.24]	90.0 [0.41]	100.0 [NR]
2001-2003	98.4 [1.38]	84.6 [1.24]	62.8 [0.23]	70.0 [0.55]	100.0 [NR]
2002-2004	96.8 [1.24]	92.3 [1.25]	66.3 [0.23]	80.0 [0.32]	100.0 [NR]
2003-2005	96.8 [1.21]	92.3 [1.32]	65.1 [0.19]	80.0 [0.26]	100.0 [NR]
2004-2006	88.9 [1.08]	92.3 [1.01]	62.8 [0.19]	90.0 [0.27]	100.0 [NR]
2005-2007	90.5 [1.04]	92.3 [0.98]	66.3 [0.19]	80.0 [0.25]	100.0 [NR]
2006-2008	90.5 [0.99]	92.3 [0.77]	64.0 [0.16]	80.0 [0.25]	100.0 [NR]
2007-2009	90.5 [0.91]	92.3 [0.66]	55.8 [0.10]	80.0 [0.23]	100.0 [NR]
2008-2010	90.5 [0.87]	92.3 [0.62]	54.7 [0.09]	80.0 [0.25]	100.0 [NR]
2009-2011	88.9 [0.90]	92.3 [0.59]	58.1 [0.11]	80.0 [0.32]	100.0 [NR]
2010-2012	88.9 [0.89]	84.6 [0.59]	58.1 [0.09]	80.0 [0.27]	100.0 [NR]
2011-2013	87.3 [0.86]	84.6 [0.59]	54.7 [0.08]	90.0 [0.28]	100.0 [NR]
2012-2014	87.3 [0.86]	76.9 [0.58]	55.8 [0.08]	90.0 [0.21]	100.0 [NR]
2013-2015	87.3 [0.84]	76.9 [0.52]	53.5 [0.09]	90.0 [0.20]	100.0 [NR]
2014-2016	85.7 [0.81]	76.9 [0.50]	50.0 [0.08]	70.0 [0.16]	100.0 [NR]
2015-2017	85.7 [0.71]	69.2 [0.43]	48.8 [0.05]	70.0 [0.13]	100.0 [NR]
Reduction in % area exceeded from 1996 to 2016 [Reduction in AAE in same period]	12.7 [1.02]	30.8 [1.42]	40.7 [0.40]	30.0 [0.96]	0.0 [NR]

\* Some sites that cross the England/Wales have been assigned to this border area. No SPAs cross the England/Scotland border. However, in calculating AAE each 1x1 km square (or part thereof) within each site has been assigned to a single country, so AAE results are calculated for individual countries only.

**Table 3.3: Trends in acidity exceedances for SSSIs (ASSIs in Northern Ireland); percentage of sites (with SRCL) with exceedance of SRCL for at least one feature, and maximum AAE of all sites/features: (a) % of sites with SRCL with exceedance of SRCL for at least one feature, (b) [maximum AAE keq ha<sup>-1</sup> year<sup>-1</sup>]. No SSSIs cross the England/Wales or England/Scotland borders, so all SSSIs have been assigned to a single country.**

	England	Wales	Scotland	NI
Number of sites	4115	1018	1452	291
Number of sites with SRCL for at least one feature	2924	676	905	178
1995-1997	71.8 [1.66]	91.4 [1.63]	85.1 [0.60]	82.0 [1.18]
1998-2000	68.0 [1.24]	80.9 [1.11]	76.1 [0.37]	75.3 [0.65]
1999-2001	68.2 [1.24]	81.2 [1.09]	75.8 [0.36]	75.3 [0.66]
2001-2003	67.9 [1.27]	79.6 [1.07]	69.7 [0.32]	75.3 [0.77]
2002-2004	66.5 [1.16]	81.8 [1.04]	70.1 [0.33]	77.0 [0.65]
2003-2005	66.3 [1.15]	82.0 [1.10]	69.1 [0.32]	75.8 [0.59]
2004-2006	62.7 [0.99]	79.1 [0.95]	69.8 [0.33]	74.7 [0.58]
2005-2007	62.3 [0.95]	79.4 [0.93]	70.1 [0.31]	75.8 [0.60]
2006-2008	60.7 [0.88]	78.6 [0.80]	67.2 [0.25]	75.3 [0.59]
2007-2009	60.2 [0.79]	77.5 [0.72]	62.9 [0.17]	75.3 [0.59]
2008-2010	60.0 [0.76]	76.0 [0.67]	61.2 [0.16]	75.8 [0.61]
2009-2011	60.1 [0.79]	76.0 [0.65]	63.2 [0.19]	78.1 [0.65]
2010-2012	59.3 [0.77]	74.7 [0.65]	61.8 [0.17]	76.4 [0.60]
2011-2013	58.2 [0.77]	74.9 [0.66]	59.1 [0.16]	77.0 [0.60]
2012-2014	57.7 [0.74]	74.0 [0.65]	58.9 [0.16]	74.2 [0.55]
2013-2015	56.6 [0.73]	73.8 [0.61]	58.7 [0.17]	74.2 [0.52]
2014-2016	55.8 [0.69]	72.6 [0.59]	56.7 [0.16]	73.6 [0.50]
2015-2017	55.2 [0.59]	71.0 [0.53]	49.6 [0.11]	73.6 [0.47]
Reduction in % area exceeded from 1996 to 2016 [Reduction in AAE in same period]	16.6 [1.07]	20.4 [1.10]	35.5 [0.49]	8.4 [0.71]



**Figure 3.1: Average Accumulated Exceedance (AAE) of acidity critical loads by CBED deposition for 2015-17; maps show the maximum AAE for any feature within each site (other features may have lower or no exceedance).**

### 3.2.2 Nutrient nitrogen results

The trends in nutrient N critical load exceedances from 1996 to 2016 are summarised in Tables 3.4-3.6. The reductions in the percentage of sites with exceedance of nutrient N critical loads for one or more features, and reductions in AAE, are smaller than the reductions seen for acidity, reflecting the smaller decreases in N deposition over time. Reductions vary by country for the different site types. In Northern Ireland there is no reduction in the percentage of SACs or SPAs with critical load exceedance and a 6.9% reduction in the percentage of SSSIs with exceedance, however the maximum AAE for sites in NI has reduced by roughly 20-30%. The largest reductions in the percentage of sites with exceedance are greatest in Scotland for SACs (9.0% reduction) and SSSIs (17.2% reduction), and in Wales (21.4% reduction) for SPAs, though there are fewer SPAs in Wales compared to Scotland and England. There is at least a 40% reduction in the maximum AAE for SACs, SPAs and SSSIs in Wales and Scotland.

At the UK level the trends results show:

- (i) for SACs, the percentage of sites with nutrient N exceedance decreased from 95.0% in 1996 to 89.7% in 2016, and the maximum AAE declined by 42.5% from 14.1 kg N ha<sup>-1</sup> year<sup>-1</sup> to 8.1 kg N ha<sup>-1</sup> year<sup>-1</sup> over the same time period.
- (ii) for SPAs, the percentage of exceeded sites decreased from 84% in 1996 to 69.8% in 2016, and the maximum AAE fell by 40.6% from 13.3 kg N ha<sup>-1</sup> year<sup>-1</sup> in 1996 to 7.9 kg N ha<sup>-1</sup> year<sup>-1</sup> in 2016.
- (iii) for SSSIs, the percentage of exceeded sites decreased from 94.9% in 1996 to 85.8% in 2016, and the maximum AAE reduced by 40.3% from 14.9 kg N ha<sup>-1</sup> year<sup>-1</sup> in 1996 to 8.9 kg N ha<sup>-1</sup> year<sup>-1</sup> in 2016.

Maps of the maximum AAE per site (Figure 3.2) based on the latest CBED deposition (2015-17) show few sites with no exceedance for any feature. Exceedances are widespread across all countries with generally lower exceedances in Scotland. The maximum AAE is above 7 kg N ha<sup>-1</sup> year<sup>-1</sup> for the majority of sites, with many sites having maximum AAE up to 28 kg N ha<sup>-1</sup> year<sup>-1</sup>, and a few sites in central England with maximum AAE above this value.

**Table 3.4: Trends in nutrient nitrogen exceedances for SACs; percentage of sites (with SRCL) with exceedance of SRCL for at least one feature, and maximum AAE of all sites/features: (a) % of sites with SRCL with exceedance of SRCL for at least one feature, (b)[maximum AAE kg N ha<sup>-1</sup> year<sup>-1</sup>]. NR = Not recorded.**

	England	Wales	Scotland	NI	Eng/Wales*	Eng/Scot*
Number of sites	231	85	236	54	7	3
Number of sites with SRCL for at least one feature	197	79	201	50	7	2
1995-1997	98.5 [20.5]	98.7 [14.1]	89.6 [7.3]	98.0 [14.4]	100.0 [NR]	50.0 [NR]
1998-2000	97.0 [17.4]	96.2 [10.3]	85.1 [5.9]	96.0 [9.8]	100.0 [NR]	50.0 [NR]
1999-2001	97.0 [18.3]	96.2 [10.9]	85.6 [6.3]	96.0 [10.3]	100.0 [NR]	50.0 [NR]
2001-2003	98.0 [19.8]	94.9 [11.9]	84.6 [6.3]	98.0 [12.7]	100.0 [NR]	50.0 [NR]
2002-2004	97.5 [18.4]	93.7 [11.7]	85.6 [6.6]	98.0 [11.3]	100.0 [NR]	50.0 [NR]
2003-2005	97.5 [18.6]	96.2 [12.2]	85.6 [6.4]	98.0 [10.9]	100.0 [NR]	50.0 [NR]
2004-2006	95.9 [15.8]	94.9 [9.7]	84.6 [6.2]	98.0 [11.0]	100.0 [NR]	50.0 [NR]
2005-2007	94.9 [15.7]	94.9 [9.7]	86.1 [6.5]	98.0 [11.9]	100.0 [NR]	50.0 [NR]
2006-2008	94.4 [15.0]	93.7 [8.8]	86.6 [6.1]	98.0 [11.8]	100.0 [NR]	50.0 [NR]
2007-2009	94.9 [14.1]	93.7 [8.5]	83.1 [5.1]	98.0 [12.3]	100.0 [NR]	50.0 [NR]
2008-2010	95.4 [14.0]	93.7 [8.4]	82.6 [4.8]	98.0 [12.7]	100.0 [NR]	50.0 [NR]
2009-2011	95.9 [14.5]	93.7 [8.4]	84.1 [4.9]	98.0 [13.1]	100.0 [NR]	50.0 [NR]
2010-2012	95.4 [14.0]	93.7 [8.2]	83.1 [4.6]	98.0 [12.2]	100.0 [NR]	50.0 [NR]
2011-2013	93.9 [13.9]	93.7 [8.2]	82.6 [4.4]	98.0 [12.2]	100.0 [NR]	50.0 [NR]
2012-2014	94.4 [13.6]	93.7 [8.0]	83.1 [4.6]	98.0 [11.5]	100.0 [NR]	50.0 [NR]
2013-2015	94.9 [13.7]	93.7 [8.1]	84.1 [4.7]	98.0 [11.2]	100.0 [NR]	50.0 [NR]
2014-2016	93.9 [13.5]	94.9 [8.0]	83.1 [4.4]	98.0 [11.0]	100.0 [NR]	50.0 [NR]
2015-2017	94.9 [0.9]	94.9 [0.5]	80.6 [0.2]	98.0 [0.7]	100.0 [NR]	50.0 [NR]
Reduction in % area exceeded from 1996 to 2016	3.6	3.8	9.0	0.0	0.0	0.0
[Reduction in AAE in same period]	[19.6]	[13.6]	[7.1]	[13.7]	[NR]	[NR]

\* Some sites cross the England/Wales or England/Scotland border and have been assigned to these border areas. However, in calculating AAE each 1x1 km square (or part thereof) within each site has been assigned to a single country, so AAE results are calculated for individual countries only.

**Table 3.5: Trends in nutrient nitrogen exceedances for SPAs; percentage of sites (with SRCL) with exceedance of SRCL for at least one feature, and maximum AAE of all sites/features: (a) % of sites with SRCL with exceedance of SRCL for at least one feature, (b)[maximum AAE kg N ha<sup>-1</sup> year<sup>-1</sup>]. NR = Not recorded.**


	England	Wales	Scotland	NI	Eng/Wales*
Number of sites	78	17	145	14	3
Number of sites with SRCL for at least one feature	72	14	124	12	3
1995-1997	97.2 [18.7]	100.0 [18.5]	74.2 [6.5]	83.3 [14.9]	100.0 [NR]
1998-2000	94.4 [17.4]	100.0 [13.4]	69.4 [4.9]	83.3 [8.9]	100.0 [NR]
1999-2001	97.2 [17.8]	100.0 [14.4]	69.4 [5.2]	83.3 [9.1]	100.0 [NR]
2001-2003	95.8 [18.7]	85.7 [15.9]	62.9 [5.2]	83.3 [13.6]	100.0 [NR]
2002-2004	93.1 [17.1]	100.0 [16.2]	66.9 [5.2]	83.3 [11.1]	100.0 [NR]
2003-2005	93.1 [17.2]	100.0 [17.0]	65.3 [4.9]	83.3 [10.7]	100.0 [NR]
2004-2006	93.1 [14.3]	92.9 [13.7]	66.9 [4.8]	83.3 [10.1]	100.0 [NR]
2005-2007	90.3 [14.2]	100.0 [13.7]	68.5 [5.2]	83.3 [11.1]	100.0 [NR]
2006-2008	88.9 [13.6]	100.0 [12.4]	69.4 [4.9]	83.3 [11.1]	66.7 [NR]
2007-2009	91.7 [13.3]	100.0 [11.9]	65.3 [4.0]	83.3 [11.6]	100.0 [NR]
2008-2010	91.7 [13.4]	100.0 [11.6]	62.9 [3.8]	83.3 [12.0]	100.0 [NR]
2009-2011	93.1 [13.8]	100.0 [11.3]	68.5 [4.0]	91.7 [13.3]	100.0 [NR]
2010-2012	90.3 [13.3]	92.9 [11.2]	62.9 [3.7]	83.3 [12.2]	100.0 [NR]
2011-2013	90.3 [12.9]	92.9 [11.3]	64.5 [3.4]	83.3 [11.8]	100.0 [NR]
2012-2014	88.9 [12.8]	78.6 [11.2]	62.9 [3.5]	83.3 [10.4]	66.7 [NR]
2013-2015	88.9 [12.9]	78.6 [11.1]	62.9 [3.7]	83.3 [10.5]	66.7 [NR]
2014-2016	88.9 [12.9]	78.6 [10.9]	61.3 [3.5]	83.3 [10.4]	100.0 [NR]
2015-2017	87.5 [0.9]	78.6 [0.8]	56.5 [0.2]	83.3 [0.7]	100.0 [NR]
Reduction in % area exceeded from 1996 to 2016 [Reduction in AAE in same period]	9.7 [17.8]	21.4 [17.7]	17.7 [6.3]	0.0 [14.2]	0.0 [NR]

\* Some sites that cross the England/Wales have been assigned to this border area. No SPAs cross the England/Scotland border. However, in calculating AAE each 1x1 km square (or part thereof) within each site was assigned to a single country, so AAE results are calculated for individual countries only.


**Table 3.6: Trends in nutrient nitrogen exceedances for SSSIs (ASSIs in Northern Ireland); percentage of sites (with SRCL) with exceedance of SRCL for at least one feature, and maximum AAE of all sites/features: (a) % of sites with SRCL with exceedance of SRCL for at least one feature, (b)[maximum AAE kg N ha<sup>-1</sup> year<sup>-1</sup>]. No SSSIs cross the England/Wales or England/Scotland borders, so all SSSIs have been assigned to a single country.**

	England	Wales	Scotland	NI
Number of sites	4115	1018	1452	291
Number of sites with SRCL for at least one feature	2954	686	938	188
1995-1997	95.2 [20.9]	100.0 [20.7]	90.1 [7.0]	94.1 [16.3]
1998-2000	94.2 [18.1]	98.8 [15.3]	85.0 [5.3]	86.2 [10.9]
1999-2001	94.8 [18.8]	98.8 [16.0]	85.9 [5.7]	87.8 [11.3]
2001-2003	96.4 [20.4]	97.4 [17.6]	82.9 [5.6]	87.8 [15.2]
2002-2004	92.6 [18.8]	98.8 [17.8]	84.3 [5.9]	88.8 [14.0]
2003-2005	93.2 [19.1]	98.8 [18.7]	83.6 [5.7]	88.8 [13.6]
2004-2006	89.4 [16.2]	98.7 [15.7]	83.2 [5.6]	88.3 [12.8]
2005-2007	89.9 [16.1]	99.0 [15.7]	84.6 [5.8]	89.4 [14.1]
2006-2008	88.6 [15.3]	98.8 [14.2]	83.7 [5.3]	92.0 [14.0]
2007-2009	89.5 [14.6]	98.5 [13.7]	81.2 [4.3]	93.1 [14.7]
2008-2010	89.9 [14.6]	98.4 [13.4]	80.3 [4.1]	93.1 [15.1]
2009-2011	90.0 [15.2]	98.5 [13.4]	81.1 [4.4]	93.1 [16.2]
2010-2012	89.1 [14.6]	98.0 [13.0]	80.1 [4.1]	92.6 [14.8]
2011-2013	87.7 [14.4]	98.1 [13.2]	80.2 [4.0]	93.1 [14.5]
2012-2014	87.8 [13.9]	98.0 [12.9]	79.7 [4.1]	92.6 [13.2]
2013-2015	87.5 [14.1]	98.1 [12.8]	80.3 [4.3]	92.6 [13.3]
2014-2016	87.3 [13.6]	98.1 [12.3]	78.9 [3.0]	88.8 [13.1]
2015-2017	87.1 [0.9]	97.7 [0.9]	72.9 [0.2]	87.2 [0.9]
Reduction in % area exceeded from 1996 to 2016 [Reduction in AAE in same period]	8.1 [20.0]	2.3 [19.8]	17.2 [6.8]	6.9 [15.4]

(a) SACs



(b) SPAs



(c) SSSIs




Figure 3.2: Average Accumulated Exceedance (AAE) of nutrient nitrogen critical loads by CBED deposition for 2015-17; maps show the maximum AAE for any feature within each site (other features may have lower or no exceedance).

## Section 4: Critical levels and their exceedances

This section of the report focuses on the trends in exceedance of the ammonia critical levels set to protect (a) lichens and bryophytes; (b) higher plants. The trends are based on rolling 3-year mean ammonia gaseous concentrations, and only cover the period from 2009-11 to 2014-16 since ammonia concentration estimates are not available for earlier periods. For brevity, the three-year means are mainly referred to in this report using the middle year, for example “2010 to 2015” means “2009-11 to 2014-16”. It should be noted that the 1x1 km ammonia concentration data are routinely a year behind the deposition data, due to the timing of availability of the required input data, so the most recent concentration data used for this report are for the period 2014-16.

### Correction December 2019

A mapping error that affected the underlying ammonia concentration data was detected in December 2019, after the original publication of this report in August 2019. The error affected the 2014-2016 mean, but not previous dates. Correcting the error made only a small change in the reported exceedances. At UK and country levels, the biggest change was in Northern Ireland, where the area exceeding the  $1 \mu\text{g m}^{-3}$  limit was originally calculated as 71.3%. After correction, this area was 70.7%. However, it does make more difference at some specific locations. The data on APIS have been corrected and updated (on 13<sup>th</sup> December 2019). The summary data and maps presented here have also been updated to use the correct concentration data.

### 4.1 Critical levels of ammonia

These critical levels are defined as the concentration of ammonia above which direct adverse effects on sensitive vegetation may occur according to present knowledge (CLRTAP 2017). Critical levels are also defined for other pollutants: sulphur dioxide, some N oxides, and ozone. These other pollutants are not considered here; further information on these can be found in CLRTAP (2017).

The critical levels for ammonia were reviewed and updated at an international workshop held in 2006 (UNECE 2007) and approved by the Task Forces of the International Cooperative Programmes (ICPs) of the CLRTAP. Unlike the critical loads, critical levels are only defined for two vegetation types (Table 4.1), rather than a range of habitats. This means that critical levels of ammonia have not been applied to individual habitats or habitat features of designated sites in the UK. The critical level exceedance metrics used are described in Section 4.3.

**Table 4.1: Critical levels of ammonia (CLRTAP, 2014)**

Vegetation type	Critical level $\text{NH}_3$ [ $\mu\text{g m}^{-3}$ ]	Time period
Lichens and bryophytes (including ecosystems where lichens and bryophytes are a key part of the ecosystem integrity)	1	Annual mean concentration
Higher plants (including heathland, semi-natural grassland and forest ground flora)	3*	Annual mean concentration

\*An explicit uncertainty range of 2-4  $\mu\text{g m}^{-3}$  was set for higher plants; this is intended to be useful when applying the critical level in different assessment contexts (e.g. precautionary approach or balance of evidence).

## 4.2 Concentrations of ammonia

The FRAME model (Singles et al. 1998) is used to provide 1x1 km resolution ammonia concentration data for the UK. This relatively high resolution is needed to spatially separate source (agricultural) areas from sink (natural ecosystems) areas (Hallsworth et al. 2010). Modelled ammonia concentrations are calibrated relative to annually averaged measurements from the National Ammonia Monitoring Network, using the median bias to adjust the concentrations. Data from all stations in the monitoring network are used for the calibration, with the exception of one station very close to a point source that was not representative of the surrounding area. The FRAME ammonia concentrations are updated annually using the new emissions data. For critical level exceedance calculations, rolling 3-year mean concentrations are used to reduce inter-year variability.

## 4.3 Calculation of critical levels exceedance

The critical level exceedance metrics calculated for this Trends Report are:

- The percentage land area in England, Wales, Scotland, Northern Ireland and UK where ammonia concentrations exceed the critical levels.
- The percentage area of N-sensitive habitats in England, Wales, Scotland, Northern Ireland and UK where ammonia concentrations exceed the critical levels. The habitat areas are based on the habitat distribution maps used for mapping nutrient N critical loads (Section 1.1).
- The percentage of designated sites (SAC, SPA, SSSI) in England, Wales, Scotland, Northern Ireland and the UK, where ammonia concentrations exceed the critical levels anywhere across a site.

## 4.4 Trends in ammonia critical levels exceedance

This section summarises the results for each of the metrics described above.

### 4.4.1 UK land area with exceedance of ammonia critical levels


Ammonia concentrations exceed the critical level of  $1 \mu\text{g m}^{-3}$  across more than 60% of the UK land area (Table 4.2); this varies spatially with <20% of Scotland, and more than 80% of England and Northern Ireland with ammonia concentrations above  $1 \mu\text{g m}^{-3}$ . The area of the UK with ammonia concentrations above  $1 \mu\text{g m}^{-3}$  has slightly decreased from 63.9% in 2010 to 62.8% in 2015 (Table 4.2).

**Table 4.2: Percentages of the UK land area where ammonia concentrations exceed critical levels.**

Critical level ( $\mu\text{g m}^{-3}$ )	Concentration data years	% land area where ammonia concentrations exceed critical levels				
		England	Wales	Scotland	NI	UK
$1 \mu\text{g m}^{-3}$	2009-2011	88.9	58.9	19.7	85.6	63.9
	2010-2012	88.5	59.0	19.8	86.0	63.7
	2011-2013	86.9	57.6	19.4	86.7	62.7
	2012-2014	84.0	55.1	17.0	85.5	60.1
	2013-2015	85.3	56.3	17.8	87.5	61.3
	2014-2016	87.1	57.5	18.9	89.3	62.8

Change in % area exceeded from 2010 to 2015		-1.8	-1.4	-0.8	+3.8	-1.1
3 $\mu\text{g m}^{-3}$	2009-2011	5.3	0.9	0.1	13.4	3.7
	2010-2012	5.5	0.9	0.2	14.7	3.9
	2011-2013	5.2	0.7	0.2	15.6	3.8
	2012-2014	3.6	0.4	0.1	12.4	2.7
	2013-2015	3.9	0.4	0.1	17.2	3.1
	2014-2016	5.3	0.9	0.1	23.0	4.3
Change in % area exceeded from 2010 to 2015		+0.1	0.0	0.0	+9.6	+0.6

The ammonia critical level of 3  $\mu\text{g m}^{-3}$  is exceeded over a much smaller area of the UK, mostly <5%, with the exception of England and NI where 5.3% and 23.0%, respectively, of the land area was exceeded in 2015 (Table 4.2). The main areas of the UK where this critical level is exceeded are parts of Northern Ireland and west/central England, with smaller areas across other regions of England (Figure 4.1).



Ammonia concentrations ( $\mu\text{g m}^{-3}$ )



$\leq 1$  (Critical levels not exceeded)



1 - 3 (Critical level for lichens and bryophytes exceeded)



$> 3$  (Critical levels for lichens and bryophytes, and higher plants exceeded)

**Figure 4.1: FRAME 1x1 km mean ammonia concentrations for 2014-16.**

#### 4.4.2 Nitrogen-sensitive habitats with exceedance of ammonia critical levels

One quarter of the mapped area of N-sensitive habitats in the UK receive ammonia concentrations above the critical level of  $1 \mu\text{g m}^{-3}$  (Table 4.3). The results vary spatially across the UK depending on (a) the variability in ammonia concentrations (see Figure 4.1), and (b) the distributions of the different N-sensitive habitats. Although 59% of the area of N-sensitive habitats are found in Scotland, as the ammonia concentrations are generally low in this part of the country (Figure 4.1), only 3.5% of the habitat area coincides with ammonia concentrations above  $1 \mu\text{g m}^{-3}$  and no areas with concentrations above  $3 \mu\text{g m}^{-3}$  (Table 4.3). The highest exceedances are seen in England and Northern Ireland, with the ammonia concentrations above  $1 \mu\text{g m}^{-3}$  for ~63-71% of their N-sensitive habitat areas and ~2-7% above  $3 \mu\text{g m}^{-3}$ . The percentage area of N-sensitive habitats in the UK with exceedance of the critical level of  $1 \mu\text{g m}^{-3}$  has decreased marginally from 25.1% in 2010 to 25.0% in 2015. The differences between the years are small and reflect the fluctuations in ammonia concentrations from one year to the next.

The N-sensitive habitats (Table 4.4) with the highest percentage area of exceedance of the ammonia critical level of  $1 \mu\text{g m}^{-3}$  are calcareous grassland (85.9%), unmanaged woodland (80.2%), managed broadleaved woodland (76.5%) and beech woodland (65.5%), which in total equals 14.6% of the total area of N-sensitive habitats. Other habitats have smaller percentage areas where ammonia concentrations are above  $1 \mu\text{g m}^{-3}$  but these results need to be interpreted with care, for example, only 19.0% of acid grassland is exceeded, but as this habitat occupies a large area across the UK, this equates to 2898 km<sup>2</sup>, which is not too different from the 85.9% of calcareous grassland (3072 km<sup>2</sup>).

**Table 4.3: Percentages of the area of nitrogen sensitive habitats in the UK where ammonia concentrations exceed critical levels of  $1 \mu\text{g m}^{-3}$  and  $3 \mu\text{g m}^{-3}$ , by country.**

		England	Wales	Scotland	NI	UK
<i>N-sensitive habitat area (km<sup>2</sup>)</i>		<i>19522</i>	<i>6837</i>	<i>43200</i>	<i>3467</i>	<i>73027</i>
$1 \mu\text{g m}^{-3}$	2009-2011	65.3	28.9	3.6	59.6	25.1
	2010-2012	64.4	29.5	3.6	60.9	25.0
	2011-2013	62.1	28.8	3.6	63.9	24.4
	2012-2014	57.0	26.6	2.9	60.6	22.3
	2013-2015	59.7	28.0	3.2	65.9	23.6
	2014-2016	62.9	29.2	3.5	70.7	25.0
Change in % area exceeded from 2010 to 2015		-2.4	+0.3	-0.1	+11.1	-0.1
$3 \mu\text{g m}^{-3}$	2009-2011	1.6	0.0	0.0	3.1	0.6
	2010-2012	1.7	0.1	0.0	3.6	0.6
	2011-2013	1.6	0.1	0.0	4.0	0.7
	2012-2014	1.1	0.0	0.0	3.2	0.5
	2013-2015	1.2	0.0	0.0	4.7	0.6
	2014-2016	1.7	0.1	0.0	7.2	0.8
Change in % area exceeded from 2010 to 2015		+0.1	+0.1	0.0	+4.1	+0.2

**Table 4.4: Percentages of the area of nitrogen-sensitive habitats where ammonia critical levels of 1 µg m<sup>-3</sup> and 3 µg m<sup>-3</sup> are exceeded in the UK, by habitat.**

		Acid grassland	Calcareous grassland	Dwarf shrub heath	Bog	Montane	Coniferous woodland (managed)	Broadleaved woodland (managed)	Beech woodland (unmanaged)	Acidophilous oak (unmanaged)	Scots Pine (unmanaged)	Other unmanaged woodland	Dune grassland	Saltmarsh
<i>Habitat area (km<sup>2</sup>)</i>		15235	3578	34826	5526	3129	8383	7482	719	1434	204	1761	323	427
1 µg m <sup>-3</sup>	2009-2011	17.9	88.9	6.5	6.6	0.0	18.6	81.3	75.9	43.8	1.4	82.0	14.4	31.6
	2010-2012	18.2	88.1	6.5	6.7	0.0	18.7	80.6	72.9	42.5	1.4	81.8	14.5	31.5
	2011-2013	18.2	85.4	6.5	6.9	0.0	18.7	77.9	66.1	39.2	1.4	79.9	14.4	28.1
	2012-2014	16.4	80.4	5.5	5.8	0.0	16.6	72.7	55.9	35.2	1.2	76.0	11.2	24.5
	2013-2015	17.6	82.9	6.3	6.9	0.0	18.7	74.6	60.0	36.9	1.2	78.4	10.6	27.0
	2014-2016	19.0	85.9	7.2	7.8	0.0	20.4	76.5	65.5	39.2	1.3	80.2	11.9	29.5
Change in % area exceeded from 2010 to 2015		+1.2	-3.1	+0.7	+1.3	0.0	+1.8	-4.8	-10.4	-4.7	-0.1	-1.8	-2.5	-2.1
3 µg m <sup>-3</sup>	2009-2011	0.5	1.7	0.2	0.1	0.0	0.3	2.0	0.6	0.6	0.0	3.0	0.1	0.5
	2010-2012	0.6	1.8	0.2	0.1	0.0	0.4	2.1	0.5	0.7	0.0	3.2	0.1	0.4
	2011-2013	0.6	1.9	0.2	0.1	0.0	0.4	2.1	0.4	0.7	0.0	3.2	0.1	0.3
	2012-2014	0.5	1.4	0.2	0.1	0.0	0.3	1.4	0.3	0.5	0.0	2.2	0.1	0.2
	2013-2015	0.7	1.4	0.2	0.2	0.0	0.3	1.5	0.3	0.5	0.0	2.8	0.0	0.2
	2014-2016	1.0	2.0	0.3	0.3	0.0	0.5	2.1	0.3	0.6	0.0	4.1	0.1	0.2
Change in % area exceeded from 2010 to 2015		+0.5	+0.3	+0.2	+0.2	0.0	+0.2	+0.1	-0.2	-0.1	0.0	+1.0	0.0	-0.3

#### 4.4.3 Designated sites with exceedance of ammonia critical levels

These results show the percentage of sites (SACs, SPAs, SSSIs) where ammonia critical levels are exceeded; a site is counted as exceeded if the ammonia concentration exceeds the respective critical level anywhere across a site. SACs may contain one or more SSSIs, and some SACs and SPAs may overlap one another (Figure 4.2), however in this analysis the sites are all assessed independently.

##### SACs

Approximately 60% of SACs occur in areas of the UK where ammonia concentrations exceed  $1 \mu\text{g m}^{-3}$  (Table 4.5); the lowest number of sites with exceedance is in Scotland where ammonia concentrations are  $< 1 \mu\text{g m}^{-3}$  at 81.6% of sites. The percentage of sites with exceedance of the  $1 \mu\text{g m}^{-3}$  critical level has fallen by 2.3% between 2010 and 2015, but this varies spatially across the UK with an increase seen in Northern Ireland over this time.

The percentage of SACs with ammonia concentrations above  $3 \mu\text{g m}^{-3}$  is small (Table 4.5):  $<15\%$  for all countries in 2015. In Northern Ireland there are increases in the number of sites exceeding this critical level between 2010 and 2015, reflecting the changes seen in the ammonia concentrations over this time period.

**Table 4.5: Percentage of SACs where ammonia concentrations exceed critical levels of  $1 \mu\text{g m}^{-3}$  and  $3 \mu\text{g m}^{-3}$  anywhere across a site.**

		England	Wales	Scotland	NI	UK
<i>Number of sites</i>		<i>[231]</i>	<i>[85]</i>	<i>[234]</i>	<i>[54]</i>	<i>[614*]</i>
$1 \mu\text{g m}^{-3}$	2009-2011	94.4	72.9	20.9	85.2	62.7
	2010-2012	93.5	75.3	19.7	87.0	62.4
	2011-2013	90.0	72.9	20.1	87.0	60.9
	2012-2014	86.6	70.6	17.5	85.2	58.1
	2013-2015	88.3	71.8	18.0	85.2	59.1
	2014-2016	89.6	72.9	18.4	90.7	60.4
Change in % sites exceeded from 2010 to 2015		-4.8	0.0	-2.6	+5.6	-2.3
$3 \mu\text{g m}^{-3}$	2009-2011	10.0	0.0	0.9	3.7	4.9
	2010-2012	10.0	1.2	1.3	3.7	5.1
	2011-2013	13.4	4.7	1.7	5.6	7.5
	2012-2014	9.5	2.4	1.3	3.7	5.4
	2013-2015	7.8	2.4	0.4	5.6	4.4
	2014-2016	8.7	2.4	0.4	13.0	5.7
Change in % sites exceeded from 2010 to 2015		-1.3	+2.4	-0.4	+9.3	+0.8

\*includes 7 sites on the England/Wales border and 3 on the England/Scotland border.

## SPAs

The results show a 12.5% increase since 2010 in the number of sites in Wales exceeding the critical level of  $1 \mu\text{g m}^{-3}$  though it should be noted there are only 16 sites in Wales. There have been decreases since 2010 in the percentage of exceeded sites in England (-2.6%) and Scotland (-2.9%) and no change in Northern Ireland (Table 4.6). Over 90% of the SPAs in England and 12 of the 13 sites in Northern Ireland receive ammonia concentrations above  $1 \mu\text{g m}^{-3}$ .

The critical level of  $3 \mu\text{g m}^{-3}$  is not exceeded for any SPAs in Wales; this is consistent with the fact that <1% of the land area in Wales has ammonia concentrations above  $3 \mu\text{g m}^{-3}$  (Table 4.2). Overall, the percentage of sites with ammonia concentrations above  $3 \mu\text{g m}^{-3}$  is considerably smaller than the number exceeding the critical level of  $1 \mu\text{g m}^{-3}$ . In Northern Ireland, two extra SPAs showed exceedance of the ammonia critical level of  $3 \mu\text{g m}^{-3}$ , representing a tripling since 2010 of the percentage of sites exceeded to 23.1%.

**Table 4.6: Percentage of SPAs with ammonia concentrations exceeding critical levels of  $1 \mu\text{g m}^{-3}$  and  $3 \mu\text{g m}^{-3}$  anywhere across a site.**

		England	Wales	Scotland	NI	UK
<i>[number of sites]</i>		<i>[76]</i>	<i>[16]</i>	<i>[137]</i>	<i>[13]</i>	<i>[245*]</i>
$1 \mu\text{g m}^{-3}$	2009-2011	97.4	37.5	27.7	92.3	54.3
	2010-2012	97.4	43.8	27.0	92.3	54.3
	2011-2013	96.1	37.5	27.0	92.3	53.5
	2012-2014	90.8	43.8	23.4	92.3	50.2
	2013-2015	92.1	50.0	22.6	92.3	50.6
	2014-2016	94.7	50.0	24.8	92.3	52.7
Change in % sites exceeded from 2010 to 2015		-2.6	+12.5	-2.9	0.0	-1.6
$3 \mu\text{g m}^{-3}$	2009-2011	11.8	0.0	1.5	7.7	5.3
	2010-2012	14.5	0.0	2.2	15.4	6.9
	2011-2013	19.7	0.0	2.2	30.8	9.4
	2012-2014	13.2	0.0	2.2	15.4	6.1
	2013-2015	10.5	0.0	0.7	15.4	4.5
	2014-2016	11.8	0.0	1.5	23.1	6.9
Change in % sites exceeded from 2010 to 2015		0.0	0.0	0.0	+15.4	+1.6

\*includes 3 sites on the England/Wales border.

## SSSIs

The percentage of SSSIs in the UK in areas where ammonia concentrations exceed the critical level of  $1 \mu\text{g m}^{-3}$  decreased by 2.9% between 2010 and 2015, to 70.3% (Table 4.6). Over 80% of the sites in England and Northern Ireland are in locations where this critical level is currently exceeded, as well as 62.7% of sites in Wales and 25.9% of sites in Scotland.

There was a small increase in the number of UK SSSIs/ASSIs showing exceedance of the  $3 \mu\text{g m}^{-3}$  Critical Load for ammonia. In particular, it is worth noting that between 2010 and 2015 there was a 6.9% increase in the percentage of SSSIs in Northern Ireland that received ammonia concentrations above the critical level of  $3 \mu\text{g m}^{-3}$ . England and Wales showed a small increase, and Scotland showed no change.

**Table 4.7: Percentage of SSSIs with ammonia concentrations exceeding critical levels of  $1 \mu\text{g m}^{-3}$  and  $3 \mu\text{g m}^{-3}$  anywhere across a site.**

		England	Wales	Scotland	NI	UK
<i>[number of sites]</i>		[4106]	[1014]	[1430]	[289]	[6839]
$1 \mu\text{g m}^{-3}$	2009-2011	89.6	66.5	27.9	83.4	73.0
	2010-2012	89.1	66.3	27.9	84.4	72.7
	2011-2013	87.5	63.8	27.9	85.5	71.4
	2012-2014	84.0	59.7	24.1	83.0	67.8
	2013-2015	84.9	61.6	24.0	84.4	68.7
	2014-2016	86.5	62.7	25.9	86.5	70.3
Change in % sites exceeded from 2010 to 2015		-3.3	-4.1	-2.2	+2.8	-2.9
$3 \mu\text{g m}^{-3}$	2009-2011	4.8	1	0.4	6.2	3.4
	2010-2012	5	0.9	0.5	7.6	3.6
	2011-2013	5.4	0.7	0.7	8	3.9
	2012-2014	3.7	0.4	0.5	6.6	2.7
	2013-2015	3.7	0.7	0.1	8.7	2.7
	2014-2016	4.9	1.8	0.4	13.1	3.9
Change in % sites exceeded from 2010 to 2015		+0.1	+0.8	0.0	+6.9	+0.5




Figure 4.2: Distribution of SACs, SPAs and SSSIs / ASSIs in the UK

## References

- Bobbink, R. and Hettelingh, J. (2011). Review and revision of empirical critical loads and dose-response relationships: Proceedings of an expert workshop, Noordwijkerhout, 23-25 June 2010. Rijksinstituut voor Volksgezondheid en Milieu RIVM.
- Calver, L. (2003). A suggested improved method for the quantification of critical loads of acidity for peat soils. PhD Thesis, *University of York*.
- Calver, L. J., Cresser, M. S. and Smart, R. P. (2004). Tolerance of *Calluna vulgaris* and peatland plant communities to sulphuric acid deposition. *Chemistry and Ecology*. **20**, 4, 309-320.
- CLRTAP (2017). Mapping Manual. Accessed March 2019.  
<https://www.umweltbundesamt.de/en/cce-manual>
- Fowler, D., Cape, J. N., Leith, I. D., Choularton, T. W., Gay, M. J. and Jones, A. (1988). The influence of altitude on rainfall composition at great dun fell. *Atmospheric Environment* (1967). **22**, 7, 1355-1362.  
<http://www.sciencedirect.com/science/article/pii/0004698188901606>
- Fuller, R. M., Smith, G. M., Sanderson, J. M., Hill, R. A., Thomson, A. G., Cox, R., Brown, N. J., Clarke, R. T., Rothery, P. and Gerard, F. F. (2002). Land Cover Map 2000 (1km dominant target class, GB). *NERC Environmental Information Data Centre*. <https://doi.org/10.5285/abff8409-0995-48d2-9303-468e1a9fe3df>
- Gammack, S. M., Smith, C. M. S. and Cresser, M. S. (1995). The approach used for mapping critical loads for ombrotrophic peats in Great Britain. *Proceedings of a Conference on Acid Rain and its Impact: The Critical Loads Debate*. 180-183. Ensis Publishing, London.
- Hall, J., Curtis, C., Dore, T. and Smith, R. (2015). Methods for the calculation of critical loads and their exceedances in the UK. Report to Defra under contract AQ0826. *Centre for Ecology and Hydrology*.  
<http://www.cldm.ceh.ac.uk/publications/uk-status-reports#overlay-context=uk>
- Hallsworth, S., Dore, A. J., Bealey, W. J., Dragosits, U., Vieno, M., Hellsten, S., Tang, Y. S. and Sutton, M. A. (2010). The role of indicator choice in quantifying the threat of atmospheric ammonia to the 'Natura 2000' network. *Environmental Science & Policy*. **13**, 8, 671-687.  
<http://www.sciencedirect.com/science/article/pii/S1462901110001206>
- Henriksen, A. and Posch, M. (2001). Steady-state models for calculating critical loads of acidity for surface waters. *Water, Air and Soil Pollution: Focus*. **1**, 1-2, 375-398.
- Hettelingh, J.-P., Posch, M., De Smet, P. A. M. and Downing, R. J. (1995). The use of critical loads in emission reduction agreements in Europe. *Water, Air, and Soil Pollution*. **85**, 4, 2381-2388.  
<https://doi.org/10.1007/BF01186190>

Hornung, M., Bull, K., Cresser, M., Hall, J., Langan, S., Loveland, P. and Smith, C. (1995). An empirical map of critical loads for soils in Great Britain. *Environmental Pollution*. **90**, 301-310.

Moss, D. and Davies, C. E. (2002). Cross-references between the EUNIS habitat classification and the nomenclature of CORINE Land Cover. *NERC/Centre for Ecology & Hydrology*. 44.

Nilsson, J. and Grennfelt, P. (1988). Critical loads for sulphur and nitrogen. Report 188:15. Copenhagen, Denmark *UNECE/Nordic Council of Ministers*.

Posch, M., de Smet, P. A. M. and Hettelingh, J.-P. (1999). Critical loads and their exceedances in Europe: an overview. In: Calculation and Mapping of Critical Thresholds in Europe: Status Report 1999. M., P., P.A.M., d. S., J.-P., H. and R.J., D. (Eds). *Coordination Centre for Effects, RIVM, Bilthoven, Netherlands*. 3-11. <http://wge-cce.org/>

Posch, M., de Vries, W. and Hettelingh, J.-P. (1995). Critical loads of sulphur and nitrogen. In: Calculation and Mapping of Critical Thresholds in Europe: Status Report 1995. M., P., P.A.M., d. S., J.-P., H. and R.J., D. (Eds). *Coordination Centre for Effects, RIVM, Bilthoven, Netherlands*. 31-41. <http://wge-cce.org/>

Posch, M. and Hettelingh, J.-P. (1997). Remarks on critical load calculations. In: Calculation and Mapping of Critical Thresholds in Europe: Status Report 1997. M., P., P.A.M., d. S., J.-P., H. and R.J., D. (Eds). *Coordination Centre for Effects, RIVM, Bilthoven, Netherlands*. 25-28. <http://wge-cce.org/>

RoTAP (2012). Review of Transboundary Air Pollution: Acidification, Eutrophication, Ground Level Ozone and Heavy Metals in the UK. Contract Report to the Department for Environment, Food and Rural Affairs. *Centre for Ecology and Hydrology*. <http://www.rotap.ceh.ac.uk>

Singles, R., Sutton, M. A. and Weston, K. J. (1998). A multi-layer model to describe the atmospheric transport and deposition of ammonia in Great Britain. *Atmospheric Environment*. **32**, 3, 393-399. <http://www.sciencedirect.com/science/article/pii/S135223109783467X>

Skiba, U. and Cresser, M. (1989). Prediction of long-term effects of rainwater acidity on peat and associated drainage water chemistry in upland areas. *Water Research*. **23**, 12, 1477-1482.

Smith, C. M. S., Cresser, M. S. and Mitchell, R. D. J. (1992). Sensitivity to acid deposition of dystrophic peat in Great Britain. *Ambio*. **22**, 22-26.

Smith, R. I., Fowler, D., Sutton, M. A., Flechard, C. and Coyle, M. (2000). Regional estimation of pollutant gas dry deposition in the UK: model description, sensitivity analyses and outputs. *Atmospheric Environment*. **34**, 22, 3757-3777. <http://www.sciencedirect.com/science/article/pii/S1352231099005178>

SNIFFER (2007). Source attribution and critical loads assessment for Special Areas of Conservation and Special Protection Areas in the UK. Final Report to SNIFFER for Project AQ02. *Centre for Ecology and Hydrology, Edinburgh*. 47.

Sverdrup, H. and De Vries, W. (1994). Calculating critical loads for acidity with the simple mass balance method. *Water, Air, and Soil Pollution*. **72**, 1-4, 143-162.

Sverdrup, H., De Vries, W. and Henriksen, A. (1990). Mapping critical loads: a guidance to the criteria, calculations, data collection and mapping of critical loads. *Nordic Council of Ministers [etc.]*.

UNECE (2007). Report on the Workshop on Atmospheric Ammonia: Detecting Emission Changes and Environmental Impacts. ECE/ED.AIR/WG.5/2007/3. Geneva *United Nations Economic and Social Council (UNECE), Executive body for the Convention of Long-Range Transboundary Air pollution: Working Group on Strategies and Review*. [http://www.ammonia-ws.ceh.ac.uk/documents/ece\\_eb\\_air\\_wg\\_5\\_2007\\_3\\_e.pdf](http://www.ammonia-ws.ceh.ac.uk/documents/ece_eb_air_wg_5_2007_3_e.pdf)

## **Annex: Critical load exceedances by habitat and country**

This Annex contains the summary critical load exceedance statistics for acidity and for nutrient nitrogen by habitat and country, with separate tables for each country.

Acidity results: Tables A1-A8

Nutrient nitrogen results: Tables N1-N8

**Table A1: Acid-sensitive habitat area in England and percentage area of habitats where acidity critical loads are exceeded, by deposition dataset year**

Parameter	Acid-sensitive habitat areas in England and percentage habitat area with exceedance of acidity critical loads:								
	Acid grassland	Dwarf shrub heath	Bog	Montane	Coniferous woodland (managed)	Broadleaved woodland (managed)	Unmanaged woodland	Freshwaters	All habitats
<i>Habitat area (km<sup>2</sup>)</i>	2669	2462	1006	2	1716	5565	2392	1109	18635
1995-1997	96.9	95.7	99.3	99.8	91.6	76.1	74.4	54.2	75.8
1998-2000	95.6	93.8	99.1	99.8	87.5	70.3	62.6	51.8	71.6
1999-2001	95.6	93.5	99.1	99.8	88.9	70.9	63.3	51.6	71.9
2001-2003	95.5	93.8	99.1	100.0	89.3	71.6	64.2	51.4	72.3
2002-2004	95.2	92.7	98.9	99.8	89.2	72.5	64.0	50.4	72.3
2003-2005	94.9	92.6	98.9	99.8	88.6	71.9	62.6	49.2	71.8
2004-2006	94.8	92.0	99.8	99.8	84.5	62.5	48.6	49.5	66.8
2005-2007	94.5	91.8	99.8	99.8	84.0	61.2	47.2	49.7	66.1
2006-2008	93.9	91.0	99.8	99.8	82.6	57.8	44.0	49.6	64.3
2007-2009	93.6	89.9	99.8	99.8	81.6	56.7	43.7	47.5	63.6
2008-2010	93.3	88.9	99.8	99.8	81.3	56.3	43.5	47.6	63.2
2009-2011	93.6	89.3	99.8	99.8	81.8	57.3	44.1	48.2	63.8
2010-2012	93.4	89.1	99.8	99.8	81.4	55.4	42.1	47.5	62.8
2011-2013	93.3	88.3	99.8	99.8	81.1	54.1	40.9	47.8	62.1
2012-2014	93.0	88.2	99.8	99.8	80.4	52.5	38.9	47.3	61.3
2013-2015	92.7	87.5	99.8	99.8	80.2	51.4	37.6	46.9	60.6
2014-2016	92.4	84.7	99.8	99.8	79.3	49.7	35.0	46.7	59.2
2015-2017	90.8	77.7	99.8	99.8	77.7	49.4	36.1	44.4	57.9
Reduction in % area exceeded from 1996 to 2016	6.1	18.0	-0.5	0.0	13.9	26.7	38.3	9.8	17.9

**Table A2: Acidity AAE (in keq ha<sup>-1</sup> year<sup>-1</sup>) by habitat for England by deposition dataset year.**

Year	AAE (keq ha <sup>-1</sup> year <sup>-1</sup> ) by habitat:								
	Acid grassland	Dwarf shrub heath	Bog	Montane	Coniferous woodland (managed)	Broadleaved woodland (managed)	Unmanaged woodland	Freshwaters	All habitats
1995-1997	1.89	1.47	2.09	2.83	1.84	1.29	1.01	1.18	1.33
1998-2000	1.39	1.07	1.63	1.83	1.38	0.99	0.72	0.86	1.00
1999-2001	1.34	1.02	1.56	1.70	1.38	1.02	0.74	0.80	0.98
2001-2003	1.32	1.01	1.53	1.64	1.48	1.15	0.84	0.77	1.04
2002-2004	1.19	0.87	1.33	1.37	1.48	1.05	0.76	0.68	0.94
2003-2005	1.18	0.83	1.30	1.41	1.45	1.05	0.76	0.65	0.93
2004-2006	1.19	0.82	1.33	1.55	1.17	0.72	0.48	0.64	0.77
2005-2007	1.15	0.77	1.28	1.53	1.10	0.71	0.47	0.58	0.74
2006-2008	1.10	0.73	1.24	1.47	1.02	0.62	0.40	0.55	0.68
2007-2009	0.97	0.61	1.08	1.35	0.90	0.59	0.38	0.49	0.62
2008-2010	0.92	0.58	1.02	1.23	0.88	0.58	0.37	0.48	0.59
2009-2011	0.96	0.61	1.07	1.33	0.94	0.61	0.39	0.49	0.62
2010-2012	0.95	0.62	1.09	1.28	0.90	0.56	0.35	0.50	0.60
2011-2013	0.98	0.62	1.10	1.34	0.86	0.52	0.33	0.50	0.59
2012-2014	0.93	0.60	1.08	1.27	0.81	0.47	0.29	0.52	0.55
2013-2015	0.91	0.58	1.07	1.26	0.80	0.46	0.28	0.48	0.54
2014-2016	0.84	0.52	0.99	1.20	0.73	0.42	0.25	0.45	0.49
2015-2017	0.73	0.40	0.83	1.08	0.66	0.42	0.25	0.39	0.44
Reduction in AAE from 1996 to 2016	1.16	1.07	1.26	1.75	1.18	0.87	0.76	0.79	0.89

**Table A3: Acid-sensitive habitat area in Wales and percentage area of habitats where acidity critical loads are exceeded, by deposition dataset year**

Parameter	Acid-sensitive habitat areas in Wales and percentage habitat area with exceedance of acidity critical loads:								
	Acid grassland	Dwarf shrub heath	Bog	Montane	Coniferous woodland (managed)	Broadleaved woodland (managed)	Unmanaged woodland	Freshwaters	All habitats
Habitat area (km <sup>2</sup> )	3143	1078	56	18	1048	790	395	1225	7798
1995-1997	98.9	97.2	98.4	100.0	98.0	86.4	89.4	59.1	90.0
1998-2000	96.5	94.0	97.7	100.0	95.7	74.0	75.5	38.8	83.1
1999-2001	96.6	93.5	97.7	100.0	96.2	75.1	76.5	37.2	83.0
2001-2003	96.0	92.1	97.6	100.0	96.0	76.1	78.2	34.7	82.4
2002-2004	95.9	90.7	97.7	100.0	96.9	77.8	82.2	32.4	82.3
2003-2005	96.1	92.7	97.7	100.0	97.0	79.1	83.3	34.4	83.2
2004-2006	95.3	91.9	100.0	100.0	94.4	73.7	75.1	33.0	81.2
2005-2007	95.2	91.6	100.0	100.0	94.1	73.2	75.0	32.6	81.0
2006-2008	94.5	87.9	99.9	100.0	90.3	70.9	72.9	32.0	79.2
2007-2009	93.7	85.3	98.7	100.0	84.2	70.3	71.8	30.3	77.4
2008-2010	93.0	81.9	98.4	100.0	75.3	69.0	69.8	28.5	74.9
2009-2011	93.0	80.8	98.4	100.0	75.1	69.1	69.5	27.2	74.5
2010-2012	93.1	81.1	98.4	100.0	72.8	67.6	67.8	28.4	74.2
2011-2013	93.2	80.9	98.4	100.0	75.0	67.9	68.3	27.5	74.4
2012-2014	92.7	80.8	97.5	100.0	71.8	66.6	66.3	26.3	73.4
2013-2015	92.3	78.0	94.2	100.0	71.2	67.7	64.9	24.3	72.4
2014-2016	92.0	77.1	95.9	100.0	68.5	64.7	59.3	23.5	71.1
2015-2017	91.6	73.9	96.2	100.0	66.2	62.9	58.1	23.6	70.0
Reduction in % area exceeded from 1996 to 2016	7.3	23.3	2.2	0.0	31.8	23.5	31.3	35.5	20.0

**Table A4: Acidity AAE (in keq ha<sup>-1</sup> year<sup>-1</sup>) by habitat for Wales by deposition dataset year.**

Year	AAE (keq ha <sup>-1</sup> year <sup>-1</sup> ) by habitat:								
	Acid grassland	Dwarf shrub heath	Bog	Montane	Coniferous woodland (managed)	Broadleaved woodland (managed)	Unmanaged woodland	Freshwaters	All habitats
1995-1997	1.56	1.33	1.77	2.38	1.87	1.31	1.30	0.50	1.36
1998-2000	1.07	0.81	1.15	1.99	1.00	0.75	0.71	0.24	0.84
1999-2001	1.04	0.75	1.06	1.70	1.00	0.78	0.74	0.21	0.82
2001-2003	1.00	0.70	1.03	1.58	1.12	0.84	0.83	0.20	0.82
2002-2004	0.92	0.61	0.93	1.42	1.17	0.93	0.90	0.18	0.79
2003-2005	0.96	0.69	1.02	1.48	1.28	0.98	0.96	0.20	0.84
2004-2006	0.92	0.68	0.83	1.42	0.93	0.76	0.72	0.19	0.74
2005-2007	0.91	0.67	0.82	1.37	0.92	0.74	0.70	0.18	0.73
2006-2008	0.80	0.53	0.69	1.25	0.72	0.60	0.55	0.15	0.61
2007-2009	0.72	0.43	0.59	1.17	0.61	0.55	0.50	0.13	0.54
2008-2010	0.67	0.38	0.53	1.12	0.55	0.51	0.46	0.10	0.49
2009-2011	0.65	0.34	0.51	1.14	0.54	0.52	0.47	0.10	0.48
2010-2012	0.66	0.35	0.52	1.13	0.51	0.47	0.42	0.11	0.47
2011-2013	0.65	0.37	0.53	1.13	0.52	0.46	0.40	0.11	0.47
2012-2014	0.64	0.38	0.53	1.06	0.49	0.42	0.37	0.09	0.46
2013-2015	0.62	0.38	0.52	0.97	0.50	0.44	0.38	0.08	0.45
2014-2016	0.60	0.35	0.50	0.92	0.44	0.40	0.35	0.08	0.43
2015-2017	0.56	0.28	0.44	0.84	0.42	0.39	0.33	0.07	0.39
Reduction in AAE from 1996 to 2016	1.00	1.05	1.33	1.54	1.45	0.92	0.97	0.43	0.97

**Table A5: Acid-sensitive habitat area in Scotland and percentage area of habitats where acidity critical loads are exceeded, by deposition dataset year**

Parameter	Acid-sensitive habitat areas in Scotland and percentage habitat area with exceedance of acidity critical loads:								
	Acid grassland	Dwarf shrub heath	Bog	Montane	Coniferous woodland (managed)	Broadleaved woodland (managed)	Unmanaged woodland	Freshwaters	All habitats
Habitat area (km <sup>2</sup> )	8336	20190	3955	3034	5111	1096	1016	5338	48083
1995-1997	88.6	65.2	85.3	95.8	72.1	66.4	54.4	18.4	68.2
1998-2000	77.9	40.6	73.1	91.3	60.2	54.8	40.2	15.3	52.6
1999-2001	77.7	38.8	69.4	93.3	60.3	55.7	42.0	15.3	51.6
2001-2003	68.5	30.1	49.5	82.4	53.6	54.7	40.3	12.9	43.0
2002-2004	69.5	32.3	47.3	89.4	56.3	57.2	44.8	12.8	44.8
2003-2005	69.8	31.3	46.3	92.9	56.3	54.4	44.0	13.2	44.5
2004-2006	74.2	35.9	61.4	96.3	52.0	47.4	37.9	13.5	48.0
2005-2007	73.1	31.4	68.1	94.3	51.1	47.7	34.9	13.0	46.1
2006-2008	68.1	24.3	63.7	85.5	47.3	44.1	30.0	12.2	40.7
2007-2009	59.1	16.0	50.5	71.2	43.4	41.3	25.9	10.6	32.9
2008-2010	56.4	15.9	42.4	69.9	42.8	42.4	26.7	10.2	31.5
2009-2011	60.9	18.7	38.5	71.4	46.5	46.6	30.2	11.1	33.9
2010-2012	58.0	17.4	38.1	65.1	45.3	44.0	28.7	11.0	32.2
2011-2013	58.6	16.5	31.8	62.1	44.0	40.0	27.4	10.9	31.0
2012-2014	57.8	15.8	39.1	62.5	42.9	36.6	25.7	11.0	31.0
2013-2015	58.7	16.4	35.1	64.8	44.6	36.8	26.7	11.0	31.4
2014-2016	55.5	14.8	32.0	61.8	43.1	35.8	24.4	10.9	29.5
2015-2017	46.1	9.4	23.9	43.5	39.1	30.6	18.0	7.8	22.7
Reduction in % area exceeded from 1996 to 2016	42.5	55.8	61.4	52.3	33.0	35.8	36.4	10.6	45.5

**Table A6: Acidity AAE (in keq ha<sup>-1</sup> year<sup>-1</sup>) by habitat for Scotland by deposition dataset year.**

Year	AAE (keq ha <sup>-1</sup> year <sup>-1</sup> ) by habitat:								
	Acid grassland	Dwarf shrub heath	Bog	Montane	Coniferous woodland (managed)	Broadleaved woodland (managed)	Unmanaged woodland	Freshwaters	All habitats
1995-1997	0.81	0.29	0.40	0.80	0.76	0.63	0.42	0.16	0.47
1998-2000	0.54	0.14	0.25	0.56	0.40	0.37	0.24	0.10	0.28
1999-2001	0.51	0.14	0.22	0.58	0.40	0.38	0.26	0.09	0.27
2001-2003	0.41	0.12	0.17	0.60	0.39	0.38	0.26	0.06	0.23
2002-2004	0.43	0.12	0.15	0.64	0.42	0.41	0.29	0.06	0.24
2003-2005	0.43	0.11	0.15	0.61	0.40	0.36	0.26	0.07	0.24
2004-2006	0.45	0.12	0.19	0.66	0.32	0.28	0.19	0.07	0.24
2005-2007	0.40	0.09	0.22	0.52	0.31	0.28	0.18	0.06	0.21
2006-2008	0.33	0.06	0.19	0.38	0.27	0.26	0.15	0.05	0.17
2007-2009	0.25	0.03	0.12	0.28	0.22	0.23	0.12	0.03	0.12
2008-2010	0.24	0.04	0.11	0.28	0.22	0.25	0.13	0.03	0.12
2009-2011	0.30	0.06	0.13	0.31	0.27	0.29	0.16	0.04	0.15
2010-2012	0.29	0.05	0.13	0.26	0.26	0.26	0.14	0.04	0.14
2011-2013	0.29	0.05	0.11	0.24	0.24	0.22	0.12	0.04	0.13
2012-2014	0.28	0.05	0.12	0.24	0.23	0.18	0.11	0.04	0.13
2013-2015	0.32	0.06	0.12	0.26	0.27	0.19	0.12	0.05	0.15
2014-2016	0.30	0.05	0.11	0.22	0.24	0.18	0.11	0.04	0.13
2015-2017	0.21	0.02	0.08	0.13	0.18	0.14	0.07	0.02	0.09
Reduction in AAE from 1996 to 2016	0.60	0.27	0.32	0.67	0.58	0.49	0.35	0.14	0.38

**Table A7: Acid-sensitive habitat area in NI and percentage area of habitats where acidity critical loads are exceeded, by deposition dataset year**

Parameter	Acid-sensitive habitat areas in NI and percentage habitat area with exceedance of acidity critical loads:								
	Acid grassland	Dwarf shrub heath	Bog	Montane	Coniferous woodland (managed)	Broadleaved woodland (managed)	Unmanaged woodland	Freshwaters	All habitats
<i>Habitat area (km<sup>2</sup>)</i>	1189	974	437	0	500	0	208	186	3537
1995-1997	86.1	83.0	84.5	0.0	73.4	0.0	49.9	24.0	76.8
1998-2000	78.9	71.3	81.1	0.0	54.1	0.0	43.3	17.6	67.2
1999-2001	77.7	70.3	80.9	0.0	55.3	0.0	43.7	17.6	66.8
2001-2003	77.6	71.1	79.8	0.0	58.5	0.0	45.6	17.6	67.4
2002-2004	81.5	69.8	82.0	0.0	62.0	0.0	47.6	17.6	69.2
2003-2005	80.0	66.4	80.5	0.0	58.7	0.0	46.8	17.6	67.1
2004-2006	78.9	66.0	96.1	0.0	57.1	0.0	44.7	15.9	68.1
2005-2007	79.9	67.0	93.7	0.0	57.6	0.0	45.9	15.9	68.5
2006-2008	79.9	67.0	95.1	0.0	57.5	0.0	45.7	15.9	68.6
2007-2009	80.3	68.4	96.4	0.0	57.3	0.0	46.3	15.9	69.4
2008-2010	80.3	69.2	96.4	0.0	57.5	0.0	46.6	15.9	69.6
2009-2011	81.1	72.8	94.9	0.0	58.7	0.0	48.9	15.9	71.0
2010-2012	79.0	67.5	92.0	0.0	54.2	0.0	46.8	15.9	67.8
2011-2013	80.0	69.3	95.4	0.0	57.1	0.0	46.4	15.9	69.4
2012-2014	77.8	65.1	96.5	0.0	55.5	0.0	43.5	14.3	67.1
2013-2015	77.8	64.7	95.3	0.0	55.1	0.0	44.3	14.3	66.9
2014-2016	76.5	59.7	92.0	0.0	51.5	0.0	43.5	14.3	64.1
2015-2017	74.5	52.2	91.3	0.0	51.2	0.0	43.3	14.3	61.2
Reduction in % area exceeded from 1996 to 2016	11.6	30.8	-6.8	0.0	22.2	0.0	6.6	9.7	15.6

**Table A8: Acidity AAE (in keq ha<sup>-1</sup> year<sup>-1</sup>) by habitat for NI by deposition dataset year.**

Year	AAE (keq ha <sup>-1</sup> year <sup>-1</sup> ) by habitat:								
	Acid grassland	Dwarf shrub heath	Bog	Montane	Coniferous woodland (managed)	Broadleaved woodland (managed)	Unmanaged woodland	Freshwaters	All habitats
1995-1997	0.86	0.76	0.88	0.00	0.98	0.00	0.66	0.34	0.80
1998-2000	0.57	0.39	0.53	0.00	0.44	0.00	0.49	0.18	0.46
1999-2001	0.55	0.38	0.53	0.00	0.46	0.00	0.50	0.18	0.46
2001-2003	0.56	0.42	0.56	0.00	0.60	0.00	0.63	0.21	0.51
2002-2004	0.54	0.34	0.50	0.00	0.55	0.00	0.66	0.17	0.46
2003-2005	0.50	0.29	0.45	0.00	0.51	0.00	0.63	0.15	0.42
2004-2006	0.49	0.29	0.60	0.00	0.45	0.00	0.57	0.15	0.42
2005-2007	0.52	0.30	0.60	0.00	0.48	0.00	0.64	0.16	0.45
2006-2008	0.51	0.30	0.60	0.00	0.47	0.00	0.64	0.16	0.44
2007-2009	0.53	0.30	0.60	0.00	0.48	0.00	0.67	0.16	0.45
2008-2010	0.55	0.33	0.62	0.00	0.50	0.00	0.70	0.17	0.47
2009-2011	0.62	0.38	0.70	0.00	0.57	0.00	0.78	0.17	0.53
2010-2012	0.53	0.32	0.61	0.00	0.49	0.00	0.68	0.15	0.46
2011-2013	0.54	0.32	0.64	0.00	0.49	0.00	0.63	0.15	0.46
2012-2014	0.46	0.26	0.55	0.00	0.41	0.00	0.54	0.14	0.39
2013-2015	0.46	0.24	0.55	0.00	0.41	0.00	0.54	0.13	0.38
2014-2016	0.43	0.22	0.49	0.00	0.37	0.00	0.54	0.12	0.36
2015-2017	0.40	0.20	0.44	0.00	0.34	0.00	0.55	0.12	0.33
Reduction in AAE from 1996 to 2016	0.46	0.56	0.44	0.00	0.64	0.00	0.11	0.22	0.47

**Table N1: Nutrient-sensitive habitat area in England and percentage area of habitats where nutrient nitrogen critical loads are exceeded, by deposition dataset year.**

Parameter	Nitrogen-sensitive habitat areas in England and percentage habitat area with exceedance of nutrient nitrogen critical loads:													
	Acid grassland	Calcareous grassland	Dwarf shrub heath	Bog	Montane	Coniferous woodland (managed)	Broadleaved woodland (managed)	Beech woodland (unmanaged)	Acidophilous oak (unmanaged)	Scots Pine (unmanaged)	Other unmanaged woodland	Dune grassland	Saltmarsh	All habitats
<i>Habitat area (km<sup>2</sup>)</i>	2620	3312	2466	1007	2	1719	5588	650	601	0	1152	93	312	19522
1995-1997	99.8	99.7	100.0	100.0	100.0	100.0	100.0	100.0	100.0	0.0	100.0	96.7	2.5	98.3
1998-2000	96.2	98.7	99.9	100.0	100.0	100.0	100.0	100.0	100.0	0.0	100.0	83.1	1.3	97.6
1999-2001	96.7	98.8	100.0	100.0	100.0	100.0	100.0	100.0	100.0	0.0	100.0	89.4	2.6	97.7
2001-2003	98.4	98.6	99.6	100.0	100.0	100.0	100.0	100.0	100.0	0.0	100.0	72.0	1.4	97.8
2002-2004	98.4	97.1	99.9	100.0	100.0	100.0	100.0	100.0	100.0	0.0	100.0	72.5	1.5	97.6
2003-2005	98.5	97.0	99.9	100.0	100.0	100.0	100.0	100.0	100.0	0.0	100.0	63.6	1.5	97.5
2004-2006	97.6	93.5	99.8	100.0	100.0	100.0	100.0	100.0	100.0	0.0	100.0	50.4	1.1	96.7
2005-2007	97.5	92.2	99.9	100.0	100.0	100.0	100.0	100.0	100.0	0.0	100.0	50.7	1.1	96.5
2006-2008	97.0	90.4	99.8	100.0	100.0	100.0	100.0	100.0	100.0	0.0	100.0	45.6	1.1	96.1
2007-2009	96.1	92.6	99.9	100.0	100.0	100.0	100.0	100.0	100.0	0.0	100.0	45.7	1.1	96.4
2008-2010	95.3	93.9	99.9	100.0	100.0	100.0	100.0	100.0	100.0	0.0	100.0	58.4	1.1	96.5
2009-2011	97.4	94.8	99.9	100.0	100.0	100.0	100.0	100.0	100.0	0.0	100.0	57.3	1.2	97.0
2010-2012	96.7	93.1	99.9	100.0	100.0	100.0	100.0	100.0	100.0	0.0	100.0	45.8	1.1	96.5
2011-2013	96.6	90.4	99.9	100.0	100.0	100.0	100.0	100.0	100.0	0.0	100.0	40.2	1.1	96.0
2012-2014	94.8	91.3	99.9	100.0	100.0	100.0	100.0	100.0	100.0	0.0	100.0	35.4	1.0	95.9
2013-2015	95.1	90.5	99.9	100.0	100.0	100.0	100.0	100.0	100.0	0.0	100.0	34.1	1.1	95.8
2014-2016	94.4	89.0	99.9	100.0	100.0	100.0	100.0	100.0	100.0	0.0	100.0	29.9	1.1	95.4
2015-2017	89.7	89.1	99.7	100.0	100.0	100.0	100.0	100.0	100.0	0.0	100.0	27.7	1.4	94.8
Reduction in % area exceeded from 1996 to 2016	10.1	10.6	0.3	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	69.0	1.1	3.5

**Table N2: Nutrient nitrogen: AAE (in kg N ha<sup>-1</sup> year<sup>-1</sup>) by habitat for England by deposition dataset year.**

Parameter	AAE (kg N ha-1 year-1) by habitat:													
	Acid grassland	Calcareous grassland	Dwarf shrub heath	Bog	Montane	Coniferous woodland (managed)	Broadleaved woodland (managed)	Beech woodland (unmanaged)	Acidophilous oak (unmanaged)	Scots Pine (unmanaged)	Other unmanaged woodland	Dune grassland	Saltmarsh	All habitats
1995-1997	12.5	7.8	14.4	16.3	24.6	25.5	27.2	22.7	28.5	0.0	27.2	4.2	0.0	19.0
1998-2000	8.9	7.7	11.7	13.5	16.6	21.8	25.1	19.8	25.3	0.0	25.7	3.5	0.1	16.8
1999-2001	9.2	8.0	12.1	14.0	16.7	22.5	26.2	20.6	26.4	0.0	26.7	3.7	0.1	17.4
2001-2003	10.3	9.3	13.2	15.2	17.5	25.3	29.9	23.3	28.9	0.0	30.6	3.1	0.0	19.7
2002-2004	8.8	7.2	11.7	13.0	14.9	24.7	27.7	22.4	27.8	0.0	27.8	1.8	2.3	18.0
2003-2005	9.2	7.2	11.7	13.2	15.7	24.9	28.1	22.8	28.0	0.0	28.3	1.9	2.3	18.2
2004-2006	9.5	6.0	11.6	13.5	17.6	20.5	21.7	15.7	22.4	0.0	21.2	1.1	0.0	14.9
2005-2007	9.4	6.0	11.4	13.4	17.9	20.2	21.8	15.3	22.4	0.0	21.4	1.1	0.1	14.9
2006-2008	9.2	5.5	11.1	13.2	17.6	19.3	20.4	13.9	21.3	0.0	20.0	1.0	0.1	14.1
2007-2009	8.0	5.6	10.1	11.9	16.4	18.2	20.7	14.4	21.3	0.0	20.5	1.1	0.1	13.8
2008-2010	7.8	5.8	10.1	11.5	15.4	18.3	20.9	14.6	21.3	0.0	20.8	1.2	0.1	13.9
2009-2011	8.4	6.2	10.7	12.2	16.6	19.3	21.9	15.2	22.2	0.0	21.7	1.2	0.1	14.6
2010-2012	8.1	5.6	10.5	12.1	16.0	18.5	20.4	13.9	20.8	0.0	20.2	1.1	0.1	13.8
2011-2013	8.3	5.1	10.4	12.1	16.5	17.9	19.5	13.3	20.3	0.0	19.2	0.9	0.0	13.3
2012-2014	7.6	4.9	10.2	12.0	15.7	17.0	18.5	12.4	19.1	0.0	18.2	0.8	0.0	12.6
2013-2015	7.9	4.9	10.4	12.4	16.0	17.3	18.6	12.2	19.2	0.0	18.3	0.6	0.0	12.8
2014-2016	7.5	4.7	10.1	12.0	15.8	16.6	18.0	11.3	18.4	0.0	17.7	0.6	0.0	12.3
2015-2017	6.7	4.6	9.0	10.7	15.2	16.4	18.6	12.0	18.8	0.0	18.5	0.6	0.1	12.2
Reduction in AAE from 1996 to 2016	5.8	3.2	5.4	5.6	9.4	9.1	8.6	10.7	9.7	0.0	8.7	3.6	-0.1	6.8

**Table N3: Nutrient-sensitive habitat area in Wales and percentage area of habitats where nutrient nitrogen critical loads are exceeded, by deposition dataset year.**

Parameter	Nitrogen-sensitive habitat areas in Wales and percentage habitat area with exceedance of nutrient nitrogen critical loads:													
	Acid grassland	Calcareous grassland	Dwarf shrub heath	Bog	Montane	Coniferous woodland (managed)	Broadleaved woodland (managed)	Beech woodland (unmanaged)	Acidophilous oak (unmanaged)	Scots Pine (unmanaged)	Other unmanaged woodland	Dune grassland	Saltmarsh	All habitats
<i>Habitat area (km<sup>2</sup>)</i>	3146	171	1094	56	18	1052	798	65	251	0	81	37	68	6837
1995-1997	98.3	95.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	0.0	100.0	89.7	0.1	98.0
1998-2000	88.4	71.3	99.7	95.8	100.0	100.0	100.0	100.0	100.0	0.0	100.0	31.1	0.2	92.5
1999-2001	85.6	70.3	99.3	95.2	100.0	100.0	100.0	100.0	100.0	0.0	100.0	31.3	0.3	91.1
2001-2003	90.8	73.0	99.4	95.8	100.0	100.0	99.9	100.0	100.0	0.0	99.9	20.9	0.0	93.5
2002-2004	90.4	68.8	99.6	96.0	100.0	100.0	100.0	100.0	100.0	0.0	100.0	24.6	0.1	93.3
2003-2005	91.8	73.5	99.7	96.0	100.0	100.0	100.0	100.0	100.0	0.0	100.0	29.9	0.1	94.1
2004-2006	90.1	73.6	99.4	96.0	100.0	100.0	100.0	100.0	100.0	0.0	99.9	23.7	0.1	93.2
2005-2007	90.7	73.3	99.5	96.0	100.0	100.0	100.0	100.0	100.0	0.0	99.9	24.3	0.1	93.6
2006-2008	89.5	70.1	99.5	96.0	100.0	100.0	99.9	100.0	100.0	0.0	99.9	23.7	0.1	92.9
2007-2009	87.1	69.6	99.4	96.0	100.0	100.0	99.9	100.0	100.0	0.0	99.9	16.2	0.1	91.7
2008-2010	82.8	71.9	98.9	95.4	100.0	100.0	99.9	98.4	100.0	0.0	99.9	16.2	0.0	89.7
2009-2011	82.8	72.5	98.7	95.4	100.0	100.0	99.9	98.4	100.0	0.0	99.9	23.6	0.1	89.8
2010-2012	82.4	71.3	99.0	95.4	100.0	100.0	99.9	98.4	100.0	0.0	99.9	22.2	0.0	89.6
2011-2013	84.0	68.4	99.4	96.0	100.0	100.0	99.9	99.6	100.0	0.0	99.9	15.0	0.0	90.3
2012-2014	82.6	64.4	99.0	95.4	100.0	100.0	99.9	99.2	100.0	0.0	99.8	14.3	0.0	89.4
2013-2015	80.6	64.7	98.6	95.2	100.0	100.0	99.8	99.6	100.0	0.0	99.8	23.9	0.0	88.5
2014-2016	79.9	66.9	98.4	95.4	100.0	100.0	99.6	98.2	100.0	0.0	99.8	25.7	0.0	88.2
2015-2017	79.5	65.9	98.5	95.4	100.0	100.0	99.6	98.6	100.0	-9999.0	99.8	25.6	0.0	88.0
Reduction in % area exceeded from 1996 to 2016	18.8	29.1	1.5	4.6	0.0	0.0	0.4	1.4	0.0	9999.0	0.2	64.1	0.1	10.0

**Table N4: Nutrient nitrogen: AAE (in kg N ha<sup>-1</sup> year<sup>-1</sup>) by habitat for Wales by deposition dataset year.**

Parameter	AAE (kg N ha <sup>-1</sup> year <sup>-1</sup> ) by habitat:													
	Acid grassland	Calcareous grassland	Dwarf shrub heath	Bog	Montane	Coniferous woodland (managed)	Broadleaved woodland (managed)	Beech woodland (unmanaged)	Acidophilous oak (unmanaged)	Scots Pine (unmanaged)	Other unmanaged woodland	Dune grassland	Saltmarsh	All habitats
1995-1997	10.6	6.9	14.8	13.9	21.7	26.1	23.2	22.7	25.6	0.0	24.6	3.0	0.0	15.8
1998-2000	6.2	3.7	9.7	8.2	17.6	17.5	16.3	16.5	18.2	0.0	18.3	0.6	0.0	10.3
1999-2001	6.4	4.3	9.6	7.8	15.2	18.3	17.2	17.6	19.0	0.0	19.8	0.7	0.0	10.6
2001-2003	7.3	4.2	10.6	9.1	16.2	21.6	19.6	19.7	22.3	0.0	22.2	0.5	0.0	12.2
2002-2004	6.8	3.9	10.0	8.4	14.9	22.5	21.3	20.3	23.9	0.0	23.5	0.3	0.0	12.2
2003-2005	7.5	4.2	11.3	9.6	15.9	24.0	22.3	21.5	25.1	0.0	24.3	0.4	0.0	13.2
2004-2006	7.0	3.6	10.9	9.3	15.2	19.0	18.2	17.4	20.7	0.0	19.2	0.4	0.0	11.4
2005-2007	7.0	3.4	11.0	9.4	15.2	19.2	18.1	17.1	20.7	0.0	18.9	0.4	0.0	11.4
2006-2008	5.9	2.5	9.5	8.2	14.3	17.0	16.1	14.9	18.6	0.0	16.8	0.2	0.0	9.9
2007-2009	5.5	2.6	8.8	7.6	13.8	16.3	15.9	14.8	18.3	0.0	17.0	0.2	0.0	9.5
2008-2010	5.2	2.7	8.4	7.2	13.7	15.8	15.7	14.7	17.9	0.0	17.1	0.2	0.0	9.2
2009-2011	5.2	2.9	8.1	7.0	14.1	15.7	16.1	15.2	18.2	0.0	17.6	0.3	0.0	9.2
2010-2012	5.1	2.4	8.1	7.0	13.9	15.1	15.0	14.0	17.2	0.0	16.3	0.2	0.0	8.8
2011-2013	5.1	2.0	8.3	7.3	14.0	15.4	14.7	13.7	17.0	0.0	15.7	0.2	0.0	8.9
2012-2014	5.0	1.9	8.5	7.2	13.1	14.9	13.9	12.9	16.3	0.0	15.1	0.1	0.0	8.6
2013-2015	5.1	2.2	8.8	7.4	12.5	15.2	14.7	13.5	17.1	0.0	15.4	0.4	0.0	8.9
2014-2016	5.1	2.4	8.5	7.4	12.1	14.5	14.3	13.0	16.6	0.0	15.1	0.4	0.0	8.7
2015-2017	5.0	2.5	8.0	7.1	11.9	14.6	14.4	12.9	16.6	0.0	15.3	0.4	0.0	8.6
Reduction in AAE from 1996 to 2016	5.6	4.4	6.8	6.8	9.8	11.5	8.8	9.8	9.0	0.0	9.3	2.6	0.0	7.2

**Table N5: Nutrient-sensitive habitat area in Scotland and percentage area of habitats where nutrient nitrogen critical loads are exceeded, by deposition dataset year.**

Parameter	Nitrogen-sensitive habitat areas in Scotland and percentage habitat area with exceedance of nutrient nitrogen critical loads:													
	Acid grassland	Calcareous grassland	Dwarf shrub heath	Bog	Montane	Coniferous woodland (managed)	Broadleaved woodland (managed)	Beech woodland (unmanaged)	Acidophilous oak (unmanaged)	Scots Pine (unmanaged)	Other unmanaged woodland	Dune grassland	Saltmarsh	All habitats
<i>Habitat area (km<sup>2</sup>)</i>	8283	24	20284	3993	3109	5111	1096	4	581	204	282	184	45	43200
1995-1997	52.4	45.7	50.0	36.7	96.6	92.4	89.1	100.0	97.3	61.1	78.1	53.0	1.2	59.4
1998-2000	40.0	32.4	37.9	25.4	95.7	85.0	82.4	100.0	92.6	38.9	72.4	28.0	1.2	48.9
1999-2001	40.6	33.5	40.6	24.7	97.1	88.4	85.2	100.0	95.3	52.3	74.2	28.5	1.2	50.9
2001-2003	40.0	48.6	36.5	24.2	88.9	84.8	82.4	100.0	90.5	49.7	73.4	31.9	0.0	47.7
2002-2004	42.3	41.2	38.9	24.5	92.6	88.6	87.1	100.0	95.6	66.5	74.2	19.3	0.0	50.2
2003-2005	43.7	25.6	39.8	24.9	90.5	87.2	86.6	100.0	95.4	67.8	74.0	18.2	0.0	50.6
2004-2006	44.2	16.1	44.6	25.8	96.5	84.1	83.1	100.0	89.1	58.0	73.5	18.7	0.0	52.9
2005-2007	42.2	24.0	44.4	37.8	96.2	85.5	82.5	100.0	88.8	52.6	74.0	24.1	0.0	53.6
2006-2008	35.1	24.1	38.6	38.8	95.5	82.8	80.1	100.0	84.6	34.2	72.6	25.6	0.0	49.0
2007-2009	28.8	15.3	31.7	27.2	82.6	78.6	77.7	100.0	74.8	30.7	70.8	23.7	0.0	41.8
2008-2010	29.3	43.3	30.1	25.2	80.9	77.4	77.3	100.0	71.7	30.5	70.4	26.9	0.0	40.7
2009-2011	37.8	51.8	33.1	25.5	82.0	80.9	79.6	100.0	79.1	32.4	71.4	29.9	0.0	44.5
2010-2012	36.4	48.7	29.6	24.3	74.2	78.2	78.1	100.0	69.7	26.2	68.2	29.7	0.0	41.4
2011-2013	37.6	22.9	28.8	21.6	71.0	77.8	77.9	100.0	71.8	24.2	69.9	25.4	0.0	40.7
2012-2014	37.1	23.4	28.8	24.9	74.1	76.5	76.9	100.0	69.5	26.0	69.5	22.3	0.0	40.9
2013-2015	40.5	25.4	30.5	25.2	78.3	77.3	77.4	100.0	71.7	26.4	69.7	23.9	0.0	42.8
2014-2016	38.9	25.4	28.9	23.8	73.1	75.7	76.0	100.0	70.5	23.9	69.3	23.2	0.0	41.0
2015-2017	31.8	14.7	23.0	19.0	61.7	70.8	72.6	100.0	58.0	20.8	63.7	19.1	0.0	34.7
Reduction in % area exceeded from 1996 to 2016	20.6	31.0	27.0	17.7	34.9	21.6	16.5	0.0	39.3	40.3	14.4	33.9	1.2	24.7

**Table N6: Nutrient nitrogen: AAE (in kg N ha<sup>-1</sup> year<sup>-1</sup>) by habitat for Scotland by deposition dataset year.**

Parameter	AAE (kg N ha <sup>-1</sup> year <sup>-1</sup> ) by habitat:													
	Acid grassland	Calcareous grassland	Dwarf shrub heath	Bog	Montane	Coniferous woodland (managed)	Broadleaved woodland (managed)	Beech woodland (unmanaged)	Acidophilous oak (unmanaged)	Scots Pine (unmanaged)	Other unmanaged woodland	Dune grassland	Saltmarsh	All habitats
1995-1997	2.7	1.4	2.4	1.9	5.4	11.7	11.6	14.1	8.6	3.3	10.2	1.8	0.1	4.1
1998-2000	1.6	0.7	1.5	1.1	4.3	7.9	8.8	7.1	6.4	2.0	8.1	0.8	0.1	2.7
1999-2001	1.6	0.9	1.6	1.1	5.0	8.5	9.3	7.5	7.1	2.8	8.6	0.9	0.1	2.9
2001-2003	1.6	1.3	1.7	1.2	5.6	9.1	9.8	9.2	6.9	3.2	9.3	0.8	0.0	3.1
2002-2004	1.7	0.6	1.8	1.1	6.1	9.6	10.3	9.4	8.1	4.0	9.7	0.5	0.0	3.3
2003-2005	1.8	0.5	1.7	1.1	6.0	9.5	9.8	9.5	7.9	3.7	9.5	0.4	0.0	3.3
2004-2006	1.9	0.4	1.8	1.2	6.3	7.9	7.8	7.2	6.2	2.6	7.7	0.5	0.0	3.1
2005-2007	1.7	0.5	1.7	1.3	5.4	7.9	8.3	8.4	5.9	2.3	8.0	0.6	0.0	2.9
2006-2008	1.3	0.4	1.4	1.2	4.2	7.4	8.0	8.4	5.0	1.9	7.6	0.6	0.0	2.5
2007-2009	1.0	0.2	1.0	0.9	3.2	6.7	7.7	8.3	4.4	1.6	7.1	0.6	0.0	2.1
2008-2010	1.1	0.6	1.1	0.9	3.2	6.8	8.2	8.5	4.4	1.7	7.4	0.7	0.0	2.2
2009-2011	1.5	0.9	1.4	1.2	3.5	7.9	9.3	9.7	5.4	1.9	8.3	0.9	0.0	2.6
2010-2012	1.5	0.8	1.3	1.1	2.8	7.5	8.6	9.1	4.8	1.6	7.8	0.8	0.0	2.4
2011-2013	1.4	0.3	1.2	1.1	2.8	7.3	7.7	8.2	4.8	1.5	7.2	0.6	0.0	2.3
2012-2014	1.4	0.2	1.2	1.1	3.0	7.1	6.9	7.2	4.6	1.6	6.6	0.5	0.0	2.3
2013-2015	2.0	0.4	1.5	1.2	3.4	7.9	7.4	7.7	5.0	1.7	7.4	0.6	0.0	2.7
2014-2016	1.9	0.3	1.4	1.2	3.1	7.6	7.3	7.9	4.7	1.5	7.3	0.6	0.0	2.5
2015-2017	1.3	0.3	0.9	0.9	2.1	6.6	6.7	7.5	3.6	1.0	6.7	0.5	0.0	1.9
Reduction in AAE from 1996 to 2016	1.4	1.1	1.5	1.0	3.3	5.1	4.9	6.6	5.0	2.3	3.5	1.3	0.1	2.2

**Table N7: Nutrient-sensitive habitat area in NI and percentage area of habitats where nutrient nitrogen critical loads are exceeded, by deposition dataset year.**

Parameter	Nitrogen-sensitive habitat areas in England and percentage habitat area with exceedance of nutrient nitrogen critical loads:													
	Acid grassland	Calcareous grassland	Dwarf shrub heath	Bog	Montane	Coniferous woodland (managed)	Broadleaved woodland (managed)	Beech woodland (unmanaged)	Acidophilous oak (unmanaged)	Scots Pine (unmanaged)	Other unmanaged woodland	Dune grassland	Saltmarsh	All habitats
<i>Habitat area (km<sup>2</sup>)</i>	1186	70	983	470	0	502	0	0	0	0	246	9	1	3467
1995-1997	84.7	20.2	99.0	99.1	0.0	99.9	0.0	0.0	0.0	0.0	100.0	84.4	0.0	92.6
1998-2000	60.7	23.8	91.9	88.7	0.0	94.8	0.0	0.0	0.0	0.0	96.8	49.7	0.0	80.0
1999-2001	64.4	22.6	92.8	93.6	0.0	96.8	0.0	0.0	0.0	0.0	97.7	49.8	0.0	82.5
2001-2003	73.3	24.2	92.5	92.7	0.0	96.9	0.0	0.0	0.0	0.0	97.9	24.9	16.5	85.4
2002-2004	73.1	24.0	94.2	94.0	0.0	98.3	0.0	0.0	0.0	0.0	99.0	49.7	8.2	86.3
2003-2005	66.3	23.2	93.8	93.7	0.0	98.1	0.0	0.0	0.0	0.0	99.0	49.7	8.2	83.8
2004-2006	69.3	23.2	93.9	94.1	0.0	97.5	0.0	0.0	0.0	0.0	98.2	49.7	11.1	84.8
2005-2007	73.4	23.6	94.0	95.7	0.0	98.0	0.0	0.0	0.0	0.0	98.5	24.9	11.1	86.4
2006-2008	73.5	24.0	94.6	96.4	0.0	98.1	0.0	0.0	0.0	0.0	98.9	24.9	11.1	86.8
2007-2009	78.4	25.8	95.0	96.5	0.0	98.3	0.0	0.0	0.0	0.0	99.0	24.9	11.1	88.7
2008-2010	81.4	27.0	95.0	96.5	0.0	98.3	0.0	0.0	0.0	0.0	99.0	24.9	11.1	89.7
2009-2011	85.7	34.3	95.1	96.6	0.0	98.1	0.0	0.0	0.0	0.0	99.0	49.7	16.5	91.4
2010-2012	80.0	24.8	94.1	94.2	0.0	97.5	0.0	0.0	0.0	0.0	98.4	49.7	11.1	88.5
2011-2013	81.7	27.1	95.1	96.6	0.0	98.3	0.0	0.0	0.0	0.0	99.0	49.7	11.1	89.9
2012-2014	71.7	23.9	95.0	96.7	0.0	98.4	0.0	0.0	0.0	0.0	99.0	24.9	11.1	86.4
2013-2015	74.8	23.6	95.0	96.5	0.0	98.1	0.0	0.0	0.0	0.0	99.0	24.9	11.1	87.4
2014-2016	69.4	23.6	94.1	94.2	0.0	97.5	0.0	0.0	0.0	0.0	98.4	24.9	11.1	84.8
2015-2017	67.9	23.6	93.6	94.1	0.0	97.5	0.0	0.0	0.0	0.0	98.3	22.4	11.1	84.2
Reduction in % area exceeded from 1996 to 2016	16.8	-3.4	5.4	5.0	0.0	2.4	0.0	0.0	0.0	0.0	1.7	62.0	-11.1	8.4

**Table N8: Nutrient nitrogen: AAE (in kg N ha<sup>-1</sup> year<sup>-1</sup>) by habitat for NI by deposition dataset year.**

Parameter	AAE (kg N ha <sup>-1</sup> year <sup>-1</sup> ) by habitat:													
	Acid grassland	Calcareous grassland	Dwarf shrub heath	Bog	Montane	Coniferous woodland (managed)	Broadleaved woodland (managed)	Beech woodland (unmanaged)	Acidophilous oak (unmanaged)	Scots Pine (unmanaged)	Other unmanaged woodland	Dune grassland	Saltmarsh	All habitats
1995-1997	5.9	1.6	11.3	9.5	0.0	19.1	0.0	0.0	0.0	0.0	18.5	3.7	0.0	10.6
1998-2000	3.0	1.4	7.1	5.3	0.0	11.1	0.0	0.0	0.0	0.0	15.2	2.5	0.0	6.5
1999-2001	3.2	1.4	7.3	5.6	0.0	11.7	0.0	0.0	0.0	0.0	15.7	3.0	0.0	6.8
2001-2003	4.4	2.6	9.4	7.3	0.0	15.8	0.0	0.0	0.0	0.0	19.6	3.6	0.3	8.9
2002-2004	4.4	1.6	8.6	6.7	0.0	15.6	0.0	0.0	0.0	0.0	21.4	3.8	0.1	8.7
2003-2005	4.1	1.4	8.2	6.2	0.0	15.1	0.0	0.0	0.0	0.0	21.0	3.5	0.0	8.3
2004-2006	4.0	1.5	8.0	6.3	0.0	13.8	0.0	0.0	0.0	0.0	19.2	3.4	0.2	7.9
2005-2007	4.8	1.7	8.8	6.8	0.0	15.1	0.0	0.0	0.0	0.0	21.5	3.9	0.4	8.8
2006-2008	4.8	1.7	8.8	6.9	0.0	15.2	0.0	0.0	0.0	0.0	21.4	3.8	0.4	8.8
2007-2009	5.3	1.9	9.3	7.3	0.0	15.8	0.0	0.0	0.0	0.0	22.7	4.1	0.5	9.4
2008-2010	5.6	2.0	9.8	7.6	0.0	16.3	0.0	0.0	0.0	0.0	23.4	4.2	0.5	9.8
2009-2011	6.6	2.4	10.9	8.7	0.0	17.7	0.0	0.0	0.0	0.0	25.3	4.3	0.6	10.9
2010-2012	5.4	2.1	9.7	7.5	0.0	15.9	0.0	0.0	0.0	0.0	22.9	3.9	0.4	9.6
2011-2013	5.4	1.9	9.7	7.8	0.0	15.9	0.0	0.0	0.0	0.0	21.8	4.0	0.4	9.5
2012-2014	4.3	1.6	8.5	6.6	0.0	14.4	0.0	0.0	0.0	0.0	19.6	3.8	0.3	8.3
2013-2015	4.3	1.6	8.5	6.7	0.0	14.5	0.0	0.0	0.0	0.0	19.9	3.4	0.2	8.4
2014-2016	4.2	1.6	8.2	6.2	0.0	13.8	0.0	0.0	0.0	0.0	19.9	3.3	0.2	8.1
2015-2017	4.0	1.6	8.0	5.8	0.0	13.5	0.0	0.0	0.0	0.0	20.2	3.2	0.1	7.9
Reduction in AAE from 1996 to 2016	1.9	0.0	3.3	3.7	0.0	5.6	0.0	0.0	0.0	0.0	-1.7	0.5	-0.1	2.7