

Appendices

CONTENTS

Appendix 1	Devolved Administration GHG Inventory Estimation Methodology
Appendix 2	Greenhouse Gas Inventories for England, Scotland, Wales and Northern Ireland, 1990 to 2009, presented in National Communication Reporting Format
Appendix 3	International Shipping and Aviation Emission Estimates for England, Scotland, Wales and Northern Ireland, 1990 to 2009
Appendix 4	Traded and Non-Traded GHG Emission Estimates for England, Scotland, Wales and Northern Ireland in 2009
Appendix 5	End User GHG Emission Estimates for England, Scotland, Wales and Northern Ireland, 1990, 2003 to 2009
Appendix 6	Relationships between source categories as defined by the IPCC and the NC categories used in reporting.
Appendix 7	2010-2011 DA GHGI Improvement Programme – Industry Sector Task.

Appendix 1

Devolved Administration GHGI Estimation Methodology

CONTENTS

1.1	Introduction	3
1.1.1	Reporting Format	3
1.1.2	General Approach	3
1.2	Energy Industries	7
1.2.1	Electricity Production	7
1.2.2	Petroleum Refining	7
1.2.3	Manufacture of Solid Fuels	10
1.2.4	Other Energy Industries	10
1.3	Manufacturing Industries and Construction	10
1.3.1	Iron and Steel	11
1.3.2	Other Industry	11
1.4	Transport	15
1.4.1	Aviation and Navigation	15
1.4.2	Road Transportation	16
1.4.3	Railways	29
1.5	Other Sectors	31
1.5.1	Commercial & Institutional	31
1.5.2	Residential	34
1.5.3	Agriculture, Forestry & Fisheries	35
1.6	Military	35
1.7	Fugitive Emissions from Fuels	35
1.7.1	Coal Mining	35
1.7.2	Solid Fuel Transformation	35
1.7.3	Oil and Natural Gas	35
1.8	Industrial Processes	39
1.8.1	Minerals Industries	39
1.8.2	Chemical Production	39
1.8.3	Metal Production	44
1.8.4	Use of Halocarbons and SF ₆	44

1.9	Agriculture	45
	Planned improvements to the inventory	46
1.10	Land Use, Land Use Change & Forestry (LULUCF)	47
1.10.1:	5A Forest Land	47
1.11	Waste	55
1.11.1	Solid Waste Disposal on Land	55
1.11.2	Waste Water Handling	56
1.11.3	Waste Incineration	56
1.12	Uncertainties	56
1.12.1	Introduction	56
1.12.2	DA GHGI Uncertainty Estimation	57
1.12.3	Trend Uncertainty Analysis	57

1.1 INTRODUCTION

The UK Greenhouse Gas Inventory compiles national estimates of greenhouse gas emissions for submission to the UN Framework Convention on Climate Change under the requirements of the Kyoto Protocol. The most recent version of the inventory, published in April 2011, presents UK greenhouse gas emission estimates for the period 1990 to 2009 (MacCarthy *et al*, 2011).

This report presents separate inventories of greenhouse gas emissions for England, Scotland, Wales and Northern Ireland for the years 1990, 1995 and 1998 to 2009 that are consistent with the 1990 to 2009 UK Greenhouse Gas Inventory.

The six direct greenhouse gases are considered:

- Carbon dioxide (CO₂);
- Methane (CH₄);
- Nitrous oxide (N₂O);
- Hydrofluorocarbons (HFCs);
- Perfluorocarbons (PFCs); and
- Sulphur hexafluoride (SF₆).

1.1.1 Reporting Format

The DA GHG inventories are presented in National Communication reporting format, in order that emissions align with policy analysis requirements of the DA Governments. Within the discussion of inventory compilation methodology, source data and trends, the IPCC sector nomenclature is used, as this enables information to be presented at a much greater level of detail, aligned with specific emission sources. The mapping between National Communication and IPCC sector format reporting is summarised in the table in Appendix 6.

Emissions from the offshore oil and gas industry are included within the UK GHG inventory, and are reported within the DA GHG inventory as "Unallocated" emissions.

The UK Inventory also reports emissions from international marine and aviation bunkers separately, as memo items to the main UK inventory dataset, in line with the reporting requirements of the United Nations Framework Convention on Climate Change (UNFCCC). Emission estimates for these international transport sources are presented separately from the main DA GHG inventory data. For more details, see Appendix 3.

1.1.2 General Approach

The UK Inventory is based on UK statistics for activities producing greenhouse gas emissions. These include fuel consumption, industrial production, agriculture and land use change and forestry. In principle, it would be ideal to obtain a complete set of equivalent statistics for each constituent country to compile each inventory.

Such a set of statistics is not available for all sources and for all constituent countries and hence it is necessary to disaggregate UK emissions into the four constituent countries by an estimation procedure.

For most sources in the UK Inventory, the emission of a pollutant from a source is calculated from the general equation:

$$E = Ae \quad \text{[Equation 1]}$$

where

E	=	Emission of pollutant (tonnes)
A	=	Activity (unit activity)
e	=	Emission Factor (tonnes pollutant/unit activity)

The activity unit may be fuel combustion (tonnes), or production of product (tonnes) or numbers of animals.

A modified equation is used in the compilation of the Devolved Administration GHG inventories:

$$E_i = \frac{d_i A_e}{\sum_{j=1}^5 d_j} \quad [\text{Equation 2}]$$

where

- E_i = Emission (in tonnes) from either England (1), Scotland (2), Wales (3), Northern Ireland (4) or "Unallocated" (5)
- d_i = A driver representing the contribution of the region to UK emissions
- i = 1, 2, 3, 4, 5

The driver, d_i can be any one of:

1. The value of the activity data for the region. [For example, consumption of specific fuels or industrial production figures for the region.];
2. The fraction of the UK activity in the region;
3. The value of a surrogate activity data statistic in the region. Where the required activity is unavailable on a regional basis, a surrogate value may be used. [For example, employment statistics or manufacturing output of a specific product, used as a surrogate for consumption data of a given fuel.]; and
4. In cases where the emissions are derived from a complex model, the driver will be the actual emission for the region calculated from the model.

The modified equation [2] ensures that the sum of the emissions from England, Scotland, Wales and Northern Ireland, plus any "unallocated" (i.e. offshore) emissions, equals the total UK emission reported within the national inventory.

Where the driver is fuel consumption, then the sum of the drivers should add up to the UK consumption. However, in practice this may not be the case if the data are taken from different sources or may be based on the financial rather than the calendar year. The estimation procedure removes such discrepancies.

Thus the compilation of the greenhouse gas inventories for the constituent countries of the UK reduces to the estimation of a set of drivers, each appropriate to emissions from a specific source. In compiling the 2009 inventories, over 180 drivers have been calculated.

Subsequent sections discuss the estimation of the drivers for each source category. Most of the detailed discussion is concentrated on the more complex categories, whilst simpler sources are summarised in Tables A1.1 to A1.10. The IPCC classification is used throughout (IPCC, 1997a), and the following section provides a description of the abbreviations used throughout the Appendix 1 discussion.

1.1.3 Improvements to DA Inventory Datasets

In the derivation of the 1990-2009 DA GHGI datasets, the inventory methods and data sources for several GHG emission source sectors have been revised and improved, including:

- Fuel combustion emissions from the domestic, commercial, public administration and small-scale industrial sectors, using new data from domestic housing surveys, censuses and the Phase II EUETS fuel use and emissions data;
- Coal mining
- Upstream oil & gas exploration and production sources
- Gas network combustion emissions
- Road transport
- Agricultural soils
- Land Use, Land Use Change and Forestry sources

Data and methodological improvements and revisions within the latest DA GHG inventories are summarised in Chapter 7 of the main report, with detailed estimation methods presented in this appendix. Where EUETS data have been used to improve the emission estimates for high emitting source sectors such as power stations, refineries, cement kilns and other industrial combustion sources, the details are provided within Annex 11 of the National Inventory Report 2011 (MacCarthy *et al*, 2011) and within the EUETS improvement programme report at:

http://uk-air.defra.gov.uk/reports/cat07/1106161307_IP_Task_2_EUETS_Issue_1.pdf

Summary of Abbreviations

AEAT	AEA Technology plc
BCA	British Cement Association
BERR	Department for Business Enterprise & Regulatory Reform
BGlass	British Glass
CA	Coal Authority
CAA	Civil Aviation Authority
DAs	Devolved Administrations
DARD	Department of Agriculture and Rural Development (Northern Ireland)
DTI	Department of Trade and Industry (now DECC)
DfT	Department for Transport
DECC	Department for Energy and Climate Change
DETI	Department of Enterprise, Trade and Investment (Northern Ireland)
DETR / DEFRA	Department of Environment, Transport & the Regions / Department for Environment, Food and Rural Affairs
DFPNI	Department of Finance and Personnel, Northern Ireland
DLTR	Department for Local Government, Transport and the Regions
E	England
EA	The Environment Agency of England & Wales
EAF	Electric Arc Furnace
EM	Enviros March
EPER	European Pollutant Emissions Register
EUETS	EU Emission Trading Scheme
IPCC	Intergovernmental Panel on Climate Change
ISR	Inventory of Statutory Releases (NI DoE)
ISSB	Iron and Steel Statistics Bureau
LPG	Liquefied petroleum gas
LRC	London Research Centre
MAFF	Ministry of Agriculture, Fisheries and Food (now DEFRA)
MSW	Municipal Solid Waste
NA	Not Available
NAEI	National Atmospheric Emissions Inventory
NI DoE	Northern Ireland Department of Environment
NIEA	Northern Ireland Environment Agency
NIO	Northern Ireland Office
NO	Not occurring
OFMDFM	Office of the First Minister and the Deputy First Minister (Northern Ireland)
ONS	Office for National Statistics
OPG	Other petroleum gas
PI	Pollution Inventory of the Environment Agency of England & Wales
S	Scotland
SEPA	The Scottish Environment Protection Agency
SPRI	Scottish Pollution Release Inventory
SSF	Solid smokeless fuel
SPRU	Science Policy Research Unit
UKOOA	UK Offshore Operators Association, now called "Oil & Gas UK"
UKPIA	United Kingdom Petroleum Industry Association
WO	Welsh Office
WS	Welsh Statistics

1.2 ENERGY INDUSTRIES

The drivers used for the energy industries are summarised in Table A1.2. This shows the base sources used in the National Atmospheric Emissions Inventory (NAEI) database, which correspond to the IPCC sources. The activity data used in the UK Inventory are shown together with the drivers used in the inventories for the constituent countries for 1990, 1995 and 1998 to 2009. The derivation of drivers sometimes differs between years depending on data availability.

1.2.1 Electricity Production

Emissions are based on fuel consumption data provided by the major power generators in Great Britain and the Northern Ireland Office for 1990 to 1999: Scottish Power (2004), Scottish and Southern Energy (2004), Innogy (2004), PowerTech (2004), AES Drax (2004). From 2000 onwards, emissions data from the Pollution Inventory (Environment Agency, 2010a) the Scottish Pollution Release Inventory (SEPA, 2010a) and the Inventory of Statutory Releases (Northern Ireland Environment Agency, 2010a) has been used to estimate DA emissions. For emissions in 2005 onwards, fuel use and emissions data reported within the EUETS (Environment Agency, 2010b; SEPA, 2010b; Northern Ireland Environment Agency, 2010b) have been used to revise and update the annual fuel emission factors that are applied within the UK GHGI, and the DA GHGI. The emissions data reported via the EUETS are used to estimate DA share of UK emissions, whilst maintaining the emission totals consistent with the UK GHGI data and UK fuel use reported in DUKES. Country-specific electricity generation data (DECC, 2010b) are then used as a comparator against reported emissions, as a quality check for the power station emissions data.

Emissions from plant generating electricity from municipal solid waste combustion are less certain for pre-1999, but all the plant are known to be in England for 1990-98 and so the emissions will correspond to the UK emissions. Since 1999, two plants have been commissioned in Scotland, at Lerwick and Dundee and emissions estimations are based on emissions data reported to the EA and SEPA.

A small number of plants generate heat rather than electricity and these are categorised as 1A4a commercial and institutional. Some generating plant burn poultry litter, or meat and bone meal, and these are all located in England. The distribution of landfill gas and sewage gas generation is assumed to correspond to the distribution of landfill sites and sewage treatment plant (see Waste, Section 1.11).

1.2.2 Petroleum Refining

UKPIA have provided a site-by-site breakdown of UK refining emissions for 1997 and 1999 – 2009 (UKPIA, 2010), presenting the emissions of a range of pollutants from combustion, process and fugitive sources. In addition, UKPIA have advised that refinery throughput did not vary significantly between 1990 and 1997. The EUETS data also provides (from 2008 onwards) a comprehensive scope of refinery emissions broken down by process and fuel, and these data are used to derive emission factors for fuel oil, natural gas and other petroleum gases (OPG) use in refineries within the UK and DA GHG inventories.

Furthermore, in the 1990-2009 GHG inventory cycle, the activity data reported in the EUETS for petroleum coke use in refineries has been used in preference to the reported DUKES fuel use allocation to the sector. Further revisions in future inventory cycles are anticipated for other fuels including fuel oil and OPG, as a result of the EUETS data analysis and consultation with the DECC energy statistics team. Emissions for 1998 are based on carbon dioxide emissions reported in the Pollution Inventory (EA: 1999a).

Table A1.2a: Energy Industries (Base Year – 1990)¹

IPCC Category	NAEI Sources	Activity: Consumption	Fuel	1990
Electricity Production	Power Stations	Coal, oil, natural gas		Consumption data from Power Generators
		Unrefined natural gas		NO
		Sewage gas		Sewage methane recovered
		Landfill gas		As landfill methane
		Orimulsion, MSW, poultry litter and tyres		All plant in England
Petroleum Refining	Refineries	All fuels		UKPIA CO ₂ emission estimates for pre-1997
Manufacture of Solid Fuels	Coke Production	Colliery Methane		All such plant assumed to be in England.
		Coke Oven gas, natural gas		Coal feed to coke ovens, ISSB, WS, DTI
		Coke		Coke breeze consumption, ISSB
		Blast Furnace gas		Coke consumed in blast furnaces, ISSB
	SSF Production	All fuels		Coal feed to SSF plant, DTI, WS
Other Energy Industries	Collieries	All other fuels		Deep mined coal production, data from British Coal Authority
		Coke oven gas		All such plant assumed to be in England.
	Gas Production	Colliery methane		Deep mined coal production, data from British Coal Authority
		LPG and Natural gas		DA share of aggregate data from EUETS installation data for natural gas use from 2005-2009.
	Offshore Own Gas Use / Gas Separation Plant	Unrefined natural gas, LPG, OPG		Extrapolated from 1995 on oil and gas arrivals, DTI
	Nuclear Fuel Prod.	natural gas		All plant in England.

¹ See Section 1.1.3 for abbreviations

Table A1.2b: Energy Industries (1995; 1998 to 2009)

IPCC Category	NAEI Sources	Activity: Consumption	Fuel	Data Sources / Comments
Electricity Production	Power Stations	Coal, oil, natural gas		Emissions data and fuel consumption data from Power Generators; PI, EPER & ISR data from 2000 onwards; EUETS data from 2005 onwards. Emission totals are always held consistent with the UK GHGI and DUKES fuel use totals.
		Unrefined natural gas		Some power facilities have used this fuel since 1995. Data provided by plant operators.
		Sewage gas		Sewage methane recovered
		Landfill gas		As landfill methane
		Orimulsion, MSW, poultry litter		From 1999, some MSW plant now also in Scotland.
Petroleum Refining	Refineries	All fuels		UKPIA CO ₂ emission estimates for pre-1997. Pollution Inventory CO ₂ emission estimates for 1998. UKPIA data for 1999 onwards. EUETS data from 2008 onwards are used to inform fuel-specific emission factors by DA. Deviations from DUKES fuel use allocations have been made for petroleum coke only, using EUETS activity data instead.
Manufacture of Solid Fuels	Coke Production	Colliery Methane		All such plant assumed to be in England.
		Coke oven gas, natural gas		Coal feed to coke ovens, ISSB, WS, DTI and (since 1999) PI data.
		Coke		Coke breeze consumption, ISSB.
		Blast Furnace gas		Coke Consumed in Blast Furnaces, ISSB.
	SSF Production	All fuels		Coal feed to SSF plant, DECC, WS.
Other Energy Industries	Collieries	All other fuels		Deep mined coal production, data from British Coal Authority.
		Coke oven gas		(1995 – current) No such plant operating.
	Gas Production	Colliery methane		Deep mined coal production, data from British Coal Authority.
		LPG and Natural gas		EUETS installation data for natural gas use from 2005-2009. All other years estimated based on the aggregate DA share from the 2005 to 2009 EUETS data.
	Offshore Own Gas Use / Gas Sepn.	Unrefined natural gas, LPG, OPG		(1995 – current) Oil & Gas UK EEMS CO ₂ estimates for terminals, DECC activity data. EUETS data for terminals.
	Nuclear Fuel Prod.	natural gas		(1995 – current) Data not available.

1.2.3 Manufacture of Solid Fuels

This category comprises the production of coke and solid smokeless fuel (SSF). Country-specific data on coke ovens in the iron and steel industry are reported in detail by ISSB (2010). Two coke ovens in England and Wales are not attached to an integrated iron and steel facility, and the consumption of coal by these ovens is estimated from WO (1998) and UK data (DECC: 1991, 2000-2010). The Welsh statistics are only available to 1993, so this data is used as an estimate of the Welsh non-iron and steel coking coal consumption in 1995. For 1998 to 2009, the non-iron and steel coking coal consumption data is apportioned between England and Wales using carbon dioxide emissions for the particular sites reported in the Pollution Inventory (EA: 2010a) and EUETS (EA: 2010b).

The generic driver for coke oven fuel consumption is the regional consumption of coking coal (ISSB, 2010). This driver is also used for natural gas consumption and coke oven gas consumption. Some coke ovens use blast furnace gas as fuel and the availability depends on blast furnace gas capacity (see Industrial Processes). Small amounts of colliery methane are consumed in the manufacture of solid fuels and this was judged to occur entirely in England where coking occurs in close proximity to deep mining. Small amounts of coke breeze are also used, and this has been disaggregated using data on other coke consumption from ISSB.

The estimation of emissions from SSF production is rather uncertain, as limited fuel use data are available from processes across the UK. Moreover, many of these are the new briquetting processes rather than coking processes and produce negligible emissions. For SSF plant operating in England and Wales, it is possible to estimate regional consumption using UK data (DECC, 2010a) and Welsh data (WO, 1998). Welsh data for 1995 has been estimated, whilst all SSF coking plant still operating since 1998 are known to be in England. Thus the driver used is coal consumed by SSF plant.

1.2.4 Other Energy Industries

This category consists of a number of small emissions from collieries, the gas industry, the nuclear fuel industry and a large emission from offshore natural gas use. In the DA inventories, emissions from oil and gas terminals are based on data provided by DECC (2010e). Installation-specific data are only available for post-1995. Emissions for 1990 are extrapolated based on 1995 Oil & Gas UK data and the arrivals of crude oil and natural gas in Scotland and England (DTI, 1991; 1996). The category of "gas separation plant" is assumed to be a subset of the gas used in oil and gas terminals and is treated in the same way as "offshore own gas use", with emissions allocated based on the DECC data on gas consumption in terminals.

Emissions from gas combustion at installations linked to the gas supply network comprise activities at compressor stations, LNG terminals and other above ground installations. The UK GHGI estimates are based on the sector natural gas allocation in DUKES. A large number of the larger sites (compressor sites, LNG terminals) report their fuel use and emissions to the EUETS. Since 2008, the reported fuel use and emissions from these large sites in EUETS exceeds that reported within DUKES and the UK GHGI, indicating that there is a small gas mis-allocation within DUKES. The DA estimates for the sector are based on the aggregated emissions data from the EUETS dataset. The EUETS data are used for each year from 2005 onwards. For 1990-2004, the DA share of the gas use is estimated based on the 2005-2009 EUETS totals. These estimates are uncertain, as the UK data are an under-report, the EUETS data only cover the larger sites on the network and may not be representative of the overall DA split of activity, and the 1990-2004 data are extrapolated from more recent data, assuming that the DA trends follow the UK-wide trend.

Other sources are minor and are covered in Table A1.2.

1.3 MANUFACTURING INDUSTRIES AND CONSTRUCTION

The drivers used to estimate DA-specific fuel consumption from these sectors are summarised in Table A1.3.

1.3.1 Iron and Steel

The ISSB (2010) provides annual reports of the detailed regional consumption of fuel by the steel industry. The consumption of coke by sinter plant is estimated as the non-blast furnace coke consumption (as this is the main other use of coke). The consumption of coke oven gas is distributed as proportional to regional figures for coal feed to coke ovens, whilst the consumption of blast furnace gas is distributed as proportional to regional figures for coke feed to blast furnaces. (The production of these gases is estimated to be proportional to the fuel used as feedstock.)

The ISSB reports the general consumption of coal, fuel oil, gas oil, LPG and natural gas by the primary iron and steel industry. This is a narrower definition than that used by DECC, which includes foundries and finishing plant, and therefore the DECC data used in the UK GHGI is higher than the ISSB data. Nevertheless, the regional ISSB data is used as a surrogate, since the distribution of the wider steel industry is directly linked to that of the primary industry, and the emissions from the secondary plant are considerably lower than the primary plant.

1.3.2 Other Industry

DECC (2010a) reports sub-national sales of solid and liquid fuels, but only as totals for England and Wales (combined), Scotland and Northern Ireland, based on reported sales data from refineries and collieries. These data have previously been used to provide estimates for other industrial fuel use, by difference from consumption data from specific sources (e.g. road transport, heavy industry). In recent years, however, DECC has started to produce more detailed sub-national energy use data (DECC, 2010b), based on local electricity and gas consumption patterns, as part of a project to develop Local Authority carbon dioxide emissions data. These statistics use local electricity and gas use data from the National Grid and the gas supply network operators (formerly Transco). Solid and liquid fuel use is calculated using point source consumption data (for major industrial sites), and a complex modelling process to distribute remaining UK fuel allocations that uses employment and population data, and takes account of smoke control zones and the patterns of gas and electricity consumption.

Table A1.3a: Manufacturing Industry and Construction (Base Year – 1990)¹

IPCC Category	NAEI Sources	Activity: Fuel Consumption	1990
Iron & Steel	Sinter Plant	Coke-breeze	Other coke consumption, ISSB
	Iron & Steel	Blast furnace gas	Coke consumed in blast furnaces, ISSB, WO
		Coke oven gas	Coal feed to coke ovens, ISSB, WS
		Coke	Coke consumed in blast furnaces, ISSB, WO
		Fuel oil, gas oil, LPG, natural gas, coal	Consumption of specified fuel, ISSB
Other Industry	Other Industry	Burning oil, fuel oil, gas oil	Sub-national oil consumption, DECC
		OPG	All such plant are located in Scotland, DECC
		LPG	Sub-national energy statistics, DECC
		Lubricants	Sub-national energy data, DECC, less estimate of road transport use.
		Natural gas	Natural gas consumed, data from Transco
		Colliery Methane	Deep mined coal production, British Coal Authority
		Coal, coke	Sub-national energy statistics, DECC
		Coke oven gas	Coal feed to coke ovens, ISSB, WO, WS
		SSF	Sub-national energy statistics, DECC
		Wood	GDP data.
	Cement	Coal, oil, gas, petrocake, tyres, waste oil	Regional cement capacity, BCA
	Ammonia (combustion)	Natural Gas	All such plant are located in England
	Autogenerators	Coal	All such plant are located in England
		Natural gas	(Data sources exactly as per “Other Industry” above)
	Other-Industry: Off-road	Gas oil, petrol	Industrial employment data (ONS)

1 See Section 1.1.3 for abbreviations

Table A1.3b: Manufacturing Industry and Construction (1995; 1998 to 2009)

IPCC Category	NAEI Sources	Activity: Fuel Consumption	Data Sources / Comments
Iron & Steel	Sinter Plant	Coke-breeze	Other coke consumption, ISSB
	Iron & Steel	Blast furnace gas	Coke Consumed in blast furnaces, ISSB, WO
		Coke oven gas	Coal feed to coke ovens, ISSB, WS
		Coke	Coke consumed in blast Furnaces, ISSB, WO
		Fuel oil, gas oil, LPG, natural gas, coal	Consumption of specified fuel, ISSB
Other Industry	Other Industry	Burning oil, fuel oil, gas oil	Sub-national oil consumption, DECC
		OPG	All such plant are located in Scotland, DECC
		LPG	Sub-national energy statistics, DECC
		Lubricants	Sub-national energy data, DECC, less estimate of road transport use.
		Natural gas	Natural gas consumed, data from Transco (now UK National Grid) & (since 1995) from Phoenix Gas (NI). Sub-national energy statistics (DECC) and AEA point source data, analysed to minimise double-counting.
		Colliery Methane	Deep mined coal production, British Coal Authority
		Coal, coke	Sub-national energy statistics, DECC; Coal consumption, WO, NIO
		Coke oven gas	Coal feed to coke ovens, ISSB, WO, WS
		SSF	Sub-national energy statistics, DECC
		Wood	GDP data.
	Cement	Coal, oil, gas, petrocake, tyres, waste oil	Regional cement capacity, BCA; For 2002 onwards, based on emissions reported to the EUETS, PI, SPRI and ISR (EA, NIEA and SEPA).
	Ammonia (combustion)	Natural Gas	All such plant are located in England
	Autogenerators	Coal	All such plant are located in England
		Natural gas	(Data sources exactly as per "Other Industry" above)
	Other-Industry: Off-road	Gas oil, petrol	Industrial employment data. (ONS)

The DECC sub-national energy statistics are revised and improved each year through targeted sector research aimed at reducing uncertainties in the modelling approach, and are now National Statistics. Previously the regional fuel use in these sectors has been developed using a complex balance approach based on limited source data. The lack of consistent and comprehensive fuel use or fuel sales data from across the DAs (especially for solid and liquid fuels) leads to significant potential errors in the distribution of UK fuel use across the regions; data gaps and inconsistencies have previously been addressed by making “expert judgement” assumptions over the time series.

The DECC sub-national energy statistics are therefore regarded as a more accurate indication of other industrial fuel use for recent years, and have been used to derive estimates for combustion of fuels such as fuel oil, gas oil and coal. These data are based predominantly on analysis of available point source data, supplemented by production and employment surveys. Several industry sectors are now 100% covered under EUETS, such as the cement sector, and hence uncertainties in the DA GHGI estimates are much lower than previously for such sources. To supplement EUETS data, additional information from other pollution inventories (PI, SPRI, ISR) are used to improve the accuracy of the allocation of industrial combustion sources. To reduce the risk of double-counting emissions, the mapping of area sources has been revised to remove the proxy data (i.e. employment or production indices) associated with those major point sources that can be accurately allocated. In future it is hoped that gradually more of the point source data can be researched, assessed, allocated to a specific source and country, and that the uncertainty of the emissions data from this source sector can continue to be reduced. See Chapter 7 for an overview of recent research tasks.

Note that the sub-national energy statistics have only been produced by DECC since 2003, and complete data (i.e. all fuels) are only available up to 2008, with gas and electricity data available up to 2009 within the DECC publication *Energy Trends December 2010*. The sub-national data are used to extrapolate estimates back across the time-series (assuming UK trends across all DAs) in many instances where more detailed data for the earlier years is absent. Hence the emission estimates & trends from solid and liquid fuels within 1A2f “other industry” sources remain amongst the more uncertain estimates within the DA inventories, due to uncertainties from the modelling approach to derive the source activity data and the back-casting of emission estimates for the earlier years.

Liquid Petroleum Gas (LPG) has a number of uses, primarily in sectors such as domestic use and the growing sector of LPG use in road transport applications. Industrial use of LPG has been disaggregated based on DECC sub-national energy statistics (DECC, 2010b) for recent years, maintaining the mass balance approach for the earlier years where complete data are available.

The driver for emissions from lubricant use is based on regional lubricant sales (DECC, 2010a) with England and Wales being disaggregated based on regional manufacturing employment statistics (ONS, 2010a).

DECC (2010c) provides data on natural gas sales to consumers categorised by consumer size and region in Great Britain, excluding consumption by large industrial users and power generators. Consumption data for gas use in Northern Ireland is supplied by Phoenix Natural Gas Ltd (2010) for 1999 onwards, and Firmus Energy (2010). These data sources are used to assess the overall gas use data for each country. Note however, that the DECC data are incomplete due to issues of commercial confidentiality for several large gas using sites, and a series of assumptions are made to estimate the gas use at these “missing” sites. Furthermore, the local authority gas use estimates do not cover a calendar year and are weather-corrected and are therefore not directly consistent with the annual fuel use data by sector that are reported in DUKES, which are used to underpin the UK and DA GHG inventory emission estimates from gas combustion. The overall gas use data for each country are uncertain as a result of these scope and reporting limitations.

The gas use within each economic sub-sector at country-level is then analysed based on the available data from the DECC sub-national energy statistics, supplemented by estimates of major point source gas use derived from analysis of the EUETS and pollution inventory emissions data. Similar to the approach adopted for gas oil, fuel oil and coal, the analysis of point source data enables greater direct allocation of gas use to industry or commercial sectors, reducing allocation uncertainties. Note that the driver determined for “other industry” is also used for “autogenerators”. In Northern Ireland, supplementary information from gas suppliers provides a slightly more detailed breakdown of gas use by end-user sector, and this has been used to revise the allocations between industrial and commercial sectors across recent years.

Drivers for fuel consumption in cement kilns are based on annual regional clinker capacity data for 1990, 1995, 1998-2001 supplied by the British Cement Association (BCA: 2004). These are applied to all fuels, with a correction factor applied to Northern Ireland to account for the absence of natural gas. Where the UK estimate of fuel consumed in cement kilns has been revised for a given year, the regional consumptions have also been revised. From 2002 onwards the emissions data reported to the PI, SPRI and ISR are used to disaggregate UK emissions (EA: 2010a, SEPA: 2010a, NIDoE: 2010a), until 2008 where the EUETS Phase II reporting scope covers all UK cement kiln sites. For 2008 and 2009 therefore, the EUETS data are used to derive the DA estimates, in order that local fuel use patterns and emission factors are fully reflected within the DA inventory data, and to maintain consistency between the EUETS and DA GHGI.

“Autogeneration” refers to electricity generation by industry for its own use. In the case of coal, the key autogenerators are Alcan and Brunner Mond, both of which are located in England. Gas autogeneration is distributed according to the other natural gas “other industry” driver.

1.4 TRANSPORT

The drivers used for transport are summarised in Table A1.4.

1.4.1 Aviation and Navigation

The disaggregation of the domestic aviation emissions uses a database of aircraft movement data from the Civil Aviation Authority, also used in the compilation of the UK GHG inventory.

The CAA database includes details of individual flights (airport origin, destination, fuel type, plane type, engine type), covering both domestic and international flights. Only domestic UK flights have been included in the DA GHG inventory data. Estimates of emissions from take off and landing cycles and aircraft cruise have been calculated. The protocol adopted for disaggregating emissions across DAs is to assign all emissions from a flight to the DA of flight origin.

Changes in the DA GHGI estimates reflect changes made within the 2009 UK GHGI. Here, flights between the UK and overseas territories have been included as domestic aviation. Previous inventories included flights from the UK to overseas territories as international aviation, recorded as a memo item. Details of the aviation methodology can be found in the NIR.

The driver for emissions from aircraft support vehicles is calculated based on aircraft movement data from the UK’s major airports (CAA, 2010).

Emissions from navigation (coastal shipping and fishing) have been revised for all DAs as a consequence of a new methodology used in the UK inventory for estimating coastal and international shipping. The method is centred around a new procedure developed by Entec under contract to Defra for calculating fuel consumption and emissions from shipping activities around UK waters using a bottom-up procedure based on detailed shipping movement data for different vessel types, fuels and journeys (Entec, 2010). Previous emission estimates for coastal and international marine have been based on total deliveries of fuel oil, marine diesel oil and gas oil to marine bunkers and for national navigation given in national energy statistics (DUKES, 2010). This has led to very erratic time series trends in fuel consumption and emissions which bears little resemblance to other activity statistics associated with shipping such as port movement data. The total fuel delivery statistics given in DUKES (marine bunker plus national navigation) are believed to be an accurate representation of the amount of fuel made available for marine consumption, but there is more uncertainty in the ultimate distribution and use of the fuels for domestic and international shipping consumption.

The overall approach can be summarised as follows:

- Fuel consumption and emissions for domestic journeys are taken from the Entec study based on detailed movement data for 2007. Entec provided an uplift to their bottom-up estimates to take account of missing vessel movements

- Fuel consumption and emissions for fishing vessels are taken from the Entec study and reported separately under 1A4ciii
- Entec's estimates for domestic shipping fuel consumption and emissions backcast to 1990 and forecast to 2009 are used. These are derived from applying trends in port movement data as proxies for changes in activities of different types of vessels.
- A reconciliation with fuels data in DUKES is made whereby the difference between the sum of the currently reported fuel deliveries for marine bunkers and national navigation in DUKES (less fuel used for naval shipping) and the fuel consumption estimate for domestic shipping taken from Entec is assigned to international shipping.

From the UK inventory for domestic navigation, the disaggregation of emissions between each constituent country is based on port movement data (DfT: 2010).

[See Appendix 3 for estimates of the DA share of the UK emissions from international shipping and international aviation, which are not directly part of the UK GHGI but are reported as "memo items" in the UK submission to the UNFCCC.]

1.4.2 Road Transportation

Carbon dioxide, methane and nitrous oxide are emitted from the exhaust of all road vehicles with internal combustion engines. Carbon dioxide is the principal product of combustion and emissions are directly related to the fuel efficiency of the vehicle.

Methane is a hydrocarbon emitted as a result of the incomplete combustion of the fuel. Nitrous oxide is a by-product of the combustion process and emitted from partial oxidation of nitrogen present in the air.

All these pollutants are emitted by different amounts from vehicles of similar size running on petrol and diesel fuel. For example, diesel cars tend to be more fuel-efficient than petrol cars of a similar size, so their carbon emissions are lower. None of these pollutants are subject to regulatory type-approval emission limits as are those which have an impact on air quality. However, emissions of GHGs are affected by technologies introduced to reduce emissions of the regulated air quality pollutants. Methane emissions are lower from petrol vehicles fitted with a three-way catalyst, although the reduction in emissions of this pollutant by the catalyst is not as efficient as it is for other hydrocarbons. Measurements also suggest that a three-way catalyst, which is efficient at reducing NO_x emissions, can actually increase emissions of nitrous oxide, formed as a by-product of the catalyst NO_x reduction process, but evidence suggests that this is mainly a problem only for early generation catalyst cars.

Disaggregation of UK emissions across the DAs is based on local data from road traffic surveys run by the UK Department for Transport and the Department for Regional Development in Northern Ireland. Vehicle kilometre figures for different vehicle types and road types are combined with fuel consumption or emission factors. The vehicle kilometre data are also subject to uncertainty, but have shown a consistent growth in traffic across all the regions up to 2007. Traffic levels decreased slightly between 2007 and 2009, except for Northern Ireland.

It is worth noting that the IPCC Reference Manual states that "*the CORINAIR (programme), with a view to the input requirements of atmospheric dispersion models, applies the principle of territoriality (emission allocation according to fuel consumption) whereas the IPCC is bound to the principle of political responsibility (allocation according to fuel sale). For the IPCC, countries with a big disparity between emissions from fuel sales and fuel consumption have the option of estimating true consumption and reporting the emissions from consumption and trade separately.*" (IPCC, 1996).

UK emissions of carbon dioxide from road transport are reported to IPCC on the basis of fuel sales. However, basing road transport emissions on fuel sales in each constituent country of the UK does not provide a representative picture of trends in road transport emissions at regional level, due to issues of cross-border fuel sales (especially between Northern Ireland – Republic of Ireland) and sales data accounting issues within the UK (e.g. "supermarket sales" in Scotland allocated to original point of sale in northern England). Estimates based on fuel consumption calculated from traffic data in each DA are therefore regarded as a more representative approach, and are consistent with the CORINAIR (now EMEP/EEA) guidance.

Table A1.4a: Transport (Base Year – 1990)¹

IPCC Category	NAEI Sources	Activity: Fuel Consumption	1990
Civil Aviation	Domestic cruise; Domestic Take-off & Landing	Aviation Gasoline, Jet Gasoline	CAA database of flight information (CAA, 2009). Fuel consumption: Digest of UK Energy Statistics (1990)
Road Transportation	Road Transport	Petrol, Diesel oil	Road fuel sales, DECC; vehicle km, DfT Traffic data: National Traffic Census, DfT Dept of Regional Development (NI: 1990) Fuel consumption: Digest of UK Energy Statistics (1990)
Railways	Railways	Gas oil	Gas oil consumption back calculated from fuel data from ATOC/ORR using train km data from DfT
Navigation	Coastal shipping	Gas oil, Fuel oil	Back calculated from 2007 estimates by Entec based on detailed shipping movements. Backcasting done from 2007 using trends in port movement data, DfT Maritime Statistics Fuel consumption: Digest of UK Energy Statistics (1990)
Other	Aircraft Support	Gas oil	Regional aircraft movements, DfT Fuel consumption: Digest of UK Energy Statistics (1990)

See Section 1.1.3 for abbreviations

Table A1.4b: Transport (1995; 1998 to 2009)¹

IPCC Category	NAEI Sources	Activity: Fuel Consumption	Data Sources / Comments
Civil Aviation	Domestic cruise; Domestic Take-off & Landing	Aviation Gasoline, Jet Gasoline	CAA database of flight information (CAA, 2010) Fuel consumption: Digest of UK Energy Statistics (1990-2010)
Road Transportation	Road Transport	Petrol, Diesel oil, LPG	Vehicle km, DfT, NI DRD Emission factors: Boulter et al. (2009) COPERT 4 (EEA, 2007) Fuel efficiency: Road Freight Statistics, DfT (1990-2010) Composition of fleet: Vehicle Licensing Statistics Report, DfT (GB) Dept of Regional Development (NI). Traffic data: National Traffic Census, DfT (England, Scotland, Wales: 1990-2010) Dept of Regional Development (NI: 1990-1999), Traffic Census Report (NI: 2000), Vehicle Kilometres of Travel Survey of Northern Ireland Annual Report (NI: 2001), Traffic and Travel Information, DRDNI (NI: 2002- 2010) Fuel consumption: Digest of UK Energy Statistics (1990-2010), Welsh Office fuels data (WO, 1998)
Railways	Railways	Gas oil	Gas oil consumption back calculated from fuel data from ATOC/ORR using train km data from DfT. Gas oil consumption from Translink Fuel consumption: Digest of UK Energy Statistics (1990-2010)
Navigation	Coastal shipping	Gas oil, Fuel oil	Back calculated from 2007 estimates by Entec based on detailed shipping movements. Backcasting and forwardcasting done from 2007 using trends in port movement data, DfT Maritime Statistics Fuel consumption: Digest of UK Energy Statistics (1990-2010)
Other	Aircraft Support	Gas oil	Regional aircraft movements, DfT Fuel consumption: Digest of UK Energy Statistics (1990-2009)

Total emissions from road transport in each region are calculated from the following information:

- Emission factors for different types of vehicles. In the case of carbon emissions, fuel consumption factors can be used because the mass of carbon emitted is proportional to the mass of fuel consumed. Emission factors (g/km) and fuel consumption factors depend on the vehicle type and fuel type (petrol or diesel) and are influenced by the drive cycle or average speeds on the different types of roads;
- Traffic activity, including distance and average speed travelled by each type of vehicle on each type of road;
- Fleet composition in terms of the age of the fleet and the petrol/diesel mix. The age of the fleet determines the proportion of vehicles manufactured in conformity with different exhaust emission regulations (which have been successively tightened over the past 30 years); and
- One of the defining factors for the inventories is the proportion of petrol cars fitted with a three-way catalyst since this became mandatory for all new cars first registered in the UK from around August 1992, in accordance with EC Directive 91/441/EEC. The proportion of cars and vans running on diesel fuel is also an important factor. The sensitivity to the age of the fleet will be much less for the 1990 inventory because there were very few cars then fitted with catalysts and the difference in emissions from cars made to the earlier emission standards was much smaller.

There are a number of improvements made to the 2009 DA road transport inventory. These followed on from the DA Stakeholders Meeting held last year (2010) and the recent improvement programme which assessed regional data to improve road transport emissions estimates (Pang et al., 2011). The improvement programme comprises of two phases: Phase I was a scoping study which assessed the availability of regional data (e.g. vehicle fleet composition, speed and use of alternative fuels), and the likely impact on GHG emissions if adopting such information; Phase II was the actual implementation of the findings from Phase I, which also represents the changes made to the 2009 DA GHGI. The DA Stakeholders Meeting promoted better communication with key statistics and policy leads within the DAs, which led to method improvement, more detailed analysis and understanding of the underlying statistics. The improvements that have been made are discussed further in the following sections.

1.4.2.1 Emission factors

All the emission factors were consistent with those used in the latest UK Greenhouse Gas Emissions Inventory (MacCarthy *et al.*, 2011). Nitrous oxide and methane emissions factors are unchanged and are the same as those used in the last DA GHG inventory compilation; they are taken from the COPERT 4 methodology (EEA, 2007) and emission factors developed by TRL on behalf of DfT (Boulter et al., 2009) respectively. For methane, full emission factor-speed relationships were available for cars and LGVs, whereas for HGVs, buses and motorcycles only single averaged factors for urban, rural and motorway roads were available.

The uncertainties in the CH₄ and N₂O factors can be expected to be quite large. However, the relative differences between emission factors used for different technologies, Euro standards and fuels are likely to reflect realistic trends.

Fuel consumption factors are also unchanged and are based on the fuel consumption-speed relationships for detailed categories of vehicles compiled by TRL on behalf of DfT. They are used in conjunction with fleet-average fuel efficiency and vehicle CO₂ factors from other sources. These include fuel efficiency factors for HGVs and buses from sources in DfT. Further details on fuel consumption factors for other vehicle types can be found in the UK GHGI report for 2009 (MacCarthy *et al.*, 2011).

Tables A1.4.1 to A1.4.3 show the fuel consumption and emission factors used for the inventory broken down by vehicle type, road type and emission standard which the vehicle was compliant with when manufactured and first registered. Tables A1.4.4 and A1.4.5 present the fleet-averaged fuel consumption factors for rigid and articulated HGVs, buses and coaches respectively from 1990-2009 for urban, rural and motorway conditions. For the other vehicle types and pollutants, CH₄ and N₂O,

where the original source of the factors provided them as speed-emission factor equations, emission factors are calculated at average speeds typical of the road types shown in the tables A1.4.1 to A1.4.3. The average speeds used were same as those used in the last DA inventory as described below.

The emission factors shown in Tables A1.4.1-A1.4.3 refer to hot exhaust emissions that is the emissions occurring from the vehicle when the engine and catalyst are at their normal operating temperatures. The excess emissions occurring when the vehicle is started with the engine and catalyst cold was taken into account for calculating N₂O emissions from petrol cars and vans using the methodology given in COPERT 4 (EEA, 2007). Details of the cold start method are given in the latest UK Greenhouse Gas Emissions Inventory (MacCarthy *et al.*, 2011), but essentially it uses mg/km “cold start” emission factors for each Euro standard in combination with the distances travelled with the vehicle not fully warmed up. For the previous DA inventories, cold start distances are based on GB-averaged trip length information (assumed to be the same in each DA region) and ambient temperature. However, DA-specific data on trip lengths have been sought during Phase I of the recent improvement programme. No significant difference in passenger car trip lengths were found for Scotland and Wales compared with the GB average currently assumed in inventories, but trip lengths are shorter in Northern Ireland and this information has been incorporated in the 2009 DA inventory. Data for estimating cold start effects on methane emissions are not available, but the effects are considered to be probably smaller and within the range of uncertainty in the hot exhaust methane emission factors.

1.4.2.2 Age and composition of the fleet

Regional licensing statistics for England, Wales and Scotland between 1994 and 2009 (DfT, 2010a) were interrogated during Phase I of the recent DA improvement programme (Pang *et al.*, 2011). The dataset were provided by DfT (hereafter referred to as DVLA data) from their Driver and Vehicle Licensing Agency database. The analysis showed that there are some regional differences in the composition of fleet, including:

- The proportion of diesel cars in the fleet is similar in England and Scotland, but is consistently slightly higher in Wales.
- Scotland and Wales have a slightly higher proportion of smaller engine-size petrol cars compared with England and the GB average;
- Scotland has a newer petrol car fleet than England and Wales, while Wales appears to have an older diesel car fleet than England and Scotland.
- The van fleet in Scotland is newer than the GB average, while in Wales the van fleet is older

Through Part II of the DA improvement programme, this DA-specific fleet data (petrol/diesel mix, age distribution, car size) for England, Scotland and Wales from the DVLA is introduced to the 2009 DA inventories for the first time.

For Northern Ireland, the proportion of diesel cars is much higher and it also has a newer car fleet than the GB average and this has previously been accounted for in the Northern Ireland inventory. The information is based on vehicle licensing statistics for private and light goods vehicles (PLG) in Northern Ireland Transport Statistics from the Central Statistics and Research Branch of the Department of Regional Development in Northern Ireland (DRDNI, 2010a).

1.4.2.3 Traffic data

The preferred indicators for road transport activity in emission inventories are traffic data in terms of vehicle kilometres travelled per year disaggregated by vehicle and road type. For the UK national inventory (the NAEI), vehicle kilometre data for the road network in Great Britain are provided by DfT for each vehicle type on roads classified as trunk, principal and minor roads in built-up areas (urban) and non-built-up areas (rural) and motorways (DfT, 2010b). These estimates are based on traffic counts from the rotating census and core census surveys.

A consistent time series of vehicle km data for 1993 to 2009 by road type and vehicle type for England, Wales and Scotland was made available by DfT for the first time (DfT, 2010c). Vehicle km data for 1993 was scaled to derive the 1990 values for England, Wales and Scotland, based on the GB trend between 1990 and 1993.

Vehicle kilometre data in Northern Ireland for different road classes and vehicle categories are available from the Traffic and Travel Information 2009: Vehicle Kilometres of Travel Annual Report produced for the Department for Regional Development (DRDNI, 2010b). There have been revisions to vehicle km activity for cars, LGVs and rigid HGVs across the whole time series for Northern Ireland. This followed from clarification with Traffic Information and Control Centre at DRDNI on the vehicle classifications used in the traffic census report which are different to those used by DfT for the rest of the UK. The vehicle km data provided by DRDNI are categorised into five groups, with the first and second categories as 'Car and Vans' and 'Med Commercial'. In the past, 'Car and Vans' vehicle km data were further split into cars and LGVs based on the Northern Ireland's licensing statistics, while 'Med Commercial' had been associated with rigid HGVs. Through clarification with DRDNI, the van component of the 'Car and Vans' category is associated with car-sized vans, while 'Med Commercial' category is associated with medium goods van. This means that LGVs and rigid vkm have been misallocated (and overestimated) in the past. Therefore corrections have been made to the vehicle km allocation methodology. Additional information was also provided by DRDNI which gave a better indication of the split between cars and vans in the vehicle kilometres figures and a better indication of the fuel split between petrol and diesel vehicles within the car and van vehicle kilometre data. This information is based on more detailed licensing data according to vehicle body type and was provided for years 2002-2009 via direct communication with DRDNI (2010c); these figures are believed to give a more accurate picture of the car and van activity data in Northern Ireland for these years than previously used, but confirm the higher proportion of diesel cars in the Northern Ireland fleet compared with the car fleet in GB.

During Phase I of the recent improvement programme, officials at each DA were initially contacted regarding availability of speed data in each DA region. Responses from the DAs show that all DAs collect speed data based on traffic count data, surveys, moving car methods and traffic models. However, it was not possible within the timescales of Phase I improvement programme to obtain and analyse comparable vehicle speed data for each country to determine whether there were any statistically significant differences to average speeds assumed at GB level. Further investigation would therefore be needed on the use of DA-specific speed data, subject to the priorities of the DAs. The vehicle speed data used to calculate emission factors from the emission factor-speed relationships are the same as those used in the UK inventory compilation and applied to all previous years. Average UK speed data for traffic in a number of different areas were taken from the following main sources: Transport Statistics Great Britain (DfT, 2008b) provided averages of speeds in Central, Inner and Outer London surveyed at different times of day during 1990 to 2007. Speeds data from other DfT's publications such as 'Road Statistics 2007: Traffic, Speeds and Congestion' (DfT, 2008c) and 2008 national road traffic and speed forecasts (DfT, 2008d) were used to define speeds in other urban areas, rural roads and motorways. Where new information is not available, previous NAEI assumptions were maintained or road speed limits used for the vehicles expected to observe these on the type of road concerned.

1.4.2.4 Estimation of Emissions of Methane and Nitrous Oxide

Emissions of CH₄ and N₂O from road transport in the regions are calculated by combining the vehicle emission factors, fleet composition data and vehicle kilometre data for the different vehicle, fuel and road types. The emissions from petrol and diesel vehicles in each DA are normalised so that the totals across all DAs equal the UK emissions calculated for the pollutant and fuel type.

1.4.2.5 Estimation of Road Transport Carbon Dioxide Emissions

Road transport has been a very significant and growing source of CO₂ across all of the constituent countries of the UK.

For the purposes of the UK's reporting to the UNFCCC on greenhouse gas emissions under the Kyoto Protocol, the UK is required to use estimation and reporting methodologies that comply with IPCC guidance. The recommended methodology for estimation of CO₂ emissions from road transport sources applies the principle of political responsibility for emissions, whereby fuel sales data are used as the basis for the estimates. In this way, across a group of countries such as the Member States of

the EU, there is no risk of double-counting road transport CO₂ emissions due to the use of different estimation methodologies¹.

Therefore, for the purposes of reporting to the UNFCCC and the determination of progress towards Kyoto Protocol emission reduction targets, the UK uses fuel sales data as the basis for CO₂ emission estimates from road transport in the National Inventory Report. However, for the purposes of compiling the Devolved Administration GHG inventories, the use of regional fuel sales data is problematic due to a couple of key issues:

- **Cross-border fuel sales** - This factor is especially evident in Northern Ireland, where the price differential between fuel in the UK and the Republic of Ireland may have encouraged purchase of fuel from outside of the UK (BERR: Personal Communication, 2004); and
- **Supermarket fuel sales** - Where a supermarket chain purchases its fuel from storage facilities in England and then sells the fuel in other parts of the UK, the emissions from that fuel sold will be incorrectly attributed to England. Although this is known to be a potential source of inconsistency in the reporting of regional fuel sales from supermarkets, it is also likely to be evident across other economic sectors too (BERR: Personal Communication, 2004).

Adopting the IPCC estimation method of using fuel sales data in each DA produces CO₂ emission trends from road transport in Northern Ireland and Scotland that buck the UK trend of increasing emissions with time, contrary to vehicle kilometre data that is collected across the UK.

¹ Note that the UK methodology for estimating emissions of methane and nitrous oxide from road transport sources is based on vehicle kilometre data, in accordance with IPCC guidance.

Table A1.4.1: Fuel Consumption Factors for Road Transport (in g fuel/km)

g fuel /km		Urban	Rural	Motorway
Petrol cars	Pre-Euro 1	66.4	62.8	69.1
	Euro 1	61.4	57.9	64.1
	Euro 2	58.8	55.3	61.5
	Euro 3	55.0	51.4	57.6
	Euro 4	50.8	47.2	53.4
Diesel cars	Pre-Euro 1	60.3	55.0	61.2
	Euro 1	58.5	53.2	59.4
	Euro 2	54.9	49.6	55.8
	Euro 3	50.2	44.9	51.1
	Euro 4	47.7	42.4	48.7
Petrol LGVs	Pre-Euro 1	68.7	64.1	70.0
	Euro 1	63.6	59.0	64.8
	Euro 2	60.9	56.3	62.1
	Euro 3	57.1	52.5	58.3
	Euro 4	52.3	47.7	53.6
Diesel LGV	Pre-Euro 1	61.9	68.4	91.9
	Euro 1	76.7	84.4	110.1
	Euro 2	71.5	77.5	106.0
	Euro 3	63.2	69.8	104.0
	Euro 4	63.2	69.8	104.0
Mopeds, <50cc, 2st	Pre-Euro 1	25.5		
	Euro 1	15.3		
	Euro 2	12.3		
	Euro 3	10.7		
Motorcycles, >50cc, 2st	Pre-Euro 1	27.5	30.2	
	Euro 1	25.3	27.8	
	Euro 2	25.3	27.8	
	Euro 3	25.3	27.8	
Motorcycles, >50cc, 4st	Pre-Euro 1	35.3	35.1	53.9
	Euro 1	33.5	33.2	46.9
	Euro 2	31.6	31.9	49.3
	Euro 3	31.6	31.9	49.3

Table A1.4.2: Methane Emission Factors for Road Transport (in mg/km)

mg CH ₄ /km		Urban	Rural	Motorway
Petrol cars	Pre-Euro 1	73.0	21.8	57.7
	Euro 1	15.0	5.2	20.9
	Euro 2	15.8	9.6	9.7
	Euro 3	5.0	4.1	7.2
	Euro 4	1.3	1.0	1.8
Diesel cars	Pre-Euro 1	12.3	10.2	10.0
	Euro 1	6.1	6.3	6.2
	Euro 2	2.9	1.7	1.2
	Euro 3	1.4	1.1	1.1
	Euro 4	1.0	0.8	0.7
Petrol LGVs	Pre-Euro 1	73.0	21.8	57.7
	Euro 1	15.0	5.2	20.9
	Euro 2	15.8	9.6	9.7
	Euro 3	5.0	4.1	7.2
	Euro 4	1.3	1.0	1.8
Diesel LGV	Pre-Euro 1	11.8	4.0	22.0
	Euro 1	6.7	1.7	5.8
	Euro 2	2.9	1.7	1.2
	Euro 3	2.2	0.6	1.0
	Euro 4	1.5	0.4	0.7
Rigid HGVs	Pre-Euro I	185.5	50.2	43.6
	Euro I	85.0	23.0	20.0
	Euro II	54.4	20.0	18.6
	Euro III	47.6	21.4	18.2
	Euro IV	2.6	1.6	1.2
	Euro V	2.3	1.4	1.1
Artic HGVs	Pre-Euro I	381.8	174.5	152.7
	Euro I	175.0	80.0	70.0
	Euro II	112.0	69.6	65.1
	Euro III	98.0	74.4	63.7
	Euro IV	5.3	5.6	4.2
	Euro V	4.7	5.0	3.8
Buses & coaches	Pre-Euro I	381.8	174.5	152.7
	Euro I	175.0	80.0	70.0
	Euro II	113.8	52.0	45.5
	Euro III	103.3	47.2	41.3
	Euro IV	5.3	5.6	4.2
	Euro V	4.7	5.0	3.8
Mopeds, <50cc, 2st	Pre-Euro 1	219.0		
	Euro 1	43.8		
	Euro 2	24.1		
	Euro 3	19.7		
Motorcycles, >50cc, 2st	Pre-Euro 1	150.0	150.0	
	Euro 1	99.0	106.5	
	Euro 2	30.0	31.5	
	Euro 3	12.0	13.5	
Motorcycles, >50cc, 4st	Pre-Euro 1	200.0	200.0	200.0
	Euro 1	127.9	138.6	148.7
	Euro 2	126.7	93.1	107.1
	Euro 3	76.2	32.6	31.8

Table A1.4.3: N₂O Emission Factors for Road Transport (in mg/km)

mg/km	Standard	Urban	Rural	Motorway
Petrol cars	Pre-Euro 1	10.0	6.5	6.5
	Euro 1	21.3	13.8	6.9
	Euro 2	10.7	3.4	1.8
	Euro 3	1.4	0.6	0.5
	Euro 4	1.8	0.6	0.5
Diesel cars	Pre-Euro 1	0	0	0
	Euro 1	2	4	4
	Euro 2	4	6	6
	Euro 3	9	4	4
	Euro 4	9	4	4
Petrol LGVs	Pre-Euro 1	10.0	6.5	6.5
	Euro 1	22.0	13.8	6.9
	Euro 2	16.3	9.3	5.8
	Euro 3	10.5	4.6	4.6
	Euro 4	0.8	1.3	1.3
Diesel LGV	Pre-Euro 1	0	0	0
	Euro 1	2	4	4
	Euro 2	4	6	6
	Euro 3	9	4	4
	Euro 4	9	4	4
Rigid HGVs	Pre-1988	30	30	30
	88/77/EEC	30	30	30
	Euro I	30	30	30
	Euro II	30	30	30
	Euro III	30	30	30
	Euro IV	30	30	30
Artic HGVs	Pre-1988	30	30	30
	88/77/EEC	30	30	30
	Euro I	30	30	30
	Euro II	30	30	30
	Euro III	30	30	30
	Euro IV	30	30	30
Buses	Pre-1988	30	30	30
	88/77/EEC	30	30	30
	Euro I	30	30	30
	Euro II	30	30	30
	Euro III	30	30	30
	Euro IV	30	30	30
Mopeds, <50cc, 2st	Pre-Euro 1	1		
	Euro 1	1		
	Euro 2	1		
	Euro 3	1		
Motorcycles, >50cc, 2st	Pre-Euro 1	2	2	
	Euro 1	2	2	
	Euro 2	2	2	
	Euro 3	2	2	
Motorcycles, >50cc, 4st	Pre-Euro 1	2	2	2
	Euro 1	2	2	2
	Euro 2	2	2	2
	Euro 3	2	2	2

Table A1.4.4: Fuel Consumption Factors for HGVs (in g fuel/km)

g fuel/km	Rigid HGVs			Artic HGVs		
	urban	rural	m-way	urban	rural	m-way
1990	271	220	228	437	337	344
1991	275	224	231	436	335	343
1992	275	224	232	432	333	341
1993	266	216	223	411	316	323
1994	258	210	218	404	311	318
1995	262	215	222	394	304	311
1996	257	212	219	387	299	306
1997	255	211	218	386	299	306
1998	244	203	210	369	287	293
1999	249	208	216	369	287	293
2000	247	208	215	368	287	294
2001	257	217	225	373	292	298
2002	250	212	219	371	290	296
2003	259	220	227	376	293	300
2004	250	212	219	363	283	289
2005	245	209	216	358	279	286
2006	252	214	221	361	281	287
2007	260	221	228	367	286	293
2008	271	230	237	382	297	304
2009	271	229	237	383	297	304

Table A1.4.5: Average fuel consumption factors for buses and coaches (in g fuel/km) in the fleet based on DfT's BSOG data

g fuel/km	Urban	Rural	Motorway
1990	305	190	216
1991	305	190	216
1992	305	190	216
1993	304	190	216
1994	300	188	214
1995	296	185	212
1996	290	182	209
1997	289	182	210
1998	289	183	212
1999	300	191	222
2000	313	200	234
2001	314	201	236
2002	327	210	247
2003	342	220	259

g fuel/km	Urban	Rural	Motorway
2004	355	228	269
2005	370	237	280
2006	356	228	269
2007	350	224	264
2008	379	242	286
2009	379	242	286

In recent years AEA has moved away from using regional fuel sales data and instead has either directly used regional vehicle km data to estimate road transport carbon dioxide emissions in each DA or has used regional vehicle km data as a means to proportion the total UK road transport carbon dioxide emissions between each DA region. This is believed to provide a more representative assessment of transport emission trends of carbon dioxide within the constituent countries of the UK. The two methods for calculating carbon dioxide emissions based on regional vehicle km data are described in the following sections.

Disaggregation of UK Carbon Dioxide Emissions by DA: Constrained Method:

In this method the sum of the DA inventories for carbon dioxide are constrained to meeting the total of the UK inventory for road transport which for carbon dioxide is derived from UK fuel sales data for petrol and DERV from DECC. The vehicle km data for each region are used to provide an estimated allocation of the total UK road transport emissions across the constituent countries. In constraining to sum to the national totals, this approach is consistent with that adopted across every other source sector in the DA GHG inventories.

However, the criticism of this method is that the presentation of results does not always provide a carbon dioxide emission trend for the DAs that is directly consistent with the vehicle kilometre trend data, as the fluctuations in UK fuel data (from DECC) have a more significant impact on the resultant emission trends.

Direct Calculation of DA Emissions: Unconstrained Method:

In this method, carbon dioxide emissions from constituent countries are derived directly from the regional vehicle km data and are not constrained to the UK totals based on national fuel consumption data. This method removes any year to year fluctuations caused by the normalisation process and enables the emission trends to mirror the smooth trends in vehicle km.

The difference in results between the constrained and unconstrained methods at DA level largely reflects the difference in the results at UK level between bottom-up calculated fuel consumption using vehicle km data and fuel consumption factors and the fuel sales data in DUKES. The reason for a disparity has previously been attributed to cross-border fuel sales ("fuel tourism") although model uncertainty was always emphasised as an additional, and probably a major explanation for the differences.

Any change in the methodologies or the factors used to calculate fuel consumption will affect the magnitude of the difference between calculated fuel consumption at national level and sales figures from DUKES and so, in turn, it will affect the disparity between the DA carbon dioxide emissions from the constrained and unconstrained approaches. The disparity has been changed across the 1990 – 2009 time series. At UK level, the calculated petrol consumption in 1990 is 0.1% higher than petrol sales; in 2009, it is 4% lower. Calculated diesel consumption in 1990 is 2% lower than diesel sales; in 2009, it is 3% higher.

The trend in road transport carbon dioxide emissions for each DA and the UK calculated by the constrained and unconstrained methods across the time series is shown in Table A1.4.6 and Figure A1.4.1. Note that in the table, figures labelled "vkm" refer to the unconstrained method; figures labelled "Fuel sales" refer to the constrained method.

Further details on the fuel consumption vs. fuel sales reconciliation issue and normalisation procedure applied at UK level are given in MacCarthy *et al.*, 2011.

Note that emissions of methane and nitrous oxide both at UK level and for the DAs are calculated directly from vehicle km data and emission factors, with no normalisation to fuel sales data involved.

Table A1.4.6: Comparison between methods of Carbon Dioxide emissions for each DA (kt CO₂)². Vkm refer to the unconstrained method. Fuel sales refer to the constrained method

Methodology	vkm	Fuel Sales	Vkm	Fuel Sales	vkm	Fuel Sales	vkm	Fuel Sales	vkm	Fuel Sales
DA	England		Scotland		Wales		Northern Ireland		UK	
1990	90,786	91,393	8,948	9,011	5,459	5,492	2,943	2,957	108,136	108,854
1995	92,393	92,556	9,025	9,051	5,519	5,521	3,173	3,171	110,110	110,299
1998	96,230	96,748	9,354	9,406	5,725	5,753	3,334	3,349	114,643	115,256
1999	97,578	97,505	9,454	9,441	5,761	5,761	3,470	3,470	116,264	116,176
2000	97,221	96,806	9,361	9,311	5,701	5,687	3,573	3,563	115,856	115,368
2001	98,145	96,723	9,407	9,264	5,735	5,671	3,635	3,592	116,922	115,249
2002	98,806	98,498	9,550	9,514	5,836	5,833	3,762	3,742	117,954	117,588
2003	99,238	97,977	9,687	9,552	5,877	5,829	3,977	3,902	118,779	117,260
2004	99,581	98,957	9,700	9,623	5,946	5,932	3,933	3,895	119,160	118,407
2005	98,461	99,382	9,621	9,692	5,875	5,937	3,921	3,940	117,878	118,951
2006	98,793	99,503	9,827	9,879	5,939	5,991	3,945	3,957	118,504	119,329
2007	99,022	100,314	9,932	10,041	5,984	6,066	4,014	4,048	118,952	120,469
2008	96,870	96,662	9,804	9,754	5,886	5,890	3,923	3,898	116,483	116,204
2009	92,731	92,776	9,415	9,389	5,623	5,639	3,871	3,856	111,640	111,659

1.4.2.6 : Disaggregation of Emissions from LPG fuel Use

All emissions from LPG-fuelled vehicles are disaggregated based on the supply infrastructure that has developed in recent years to provide for this relatively new market. Information on LPG fuel supply stations was obtained from the Energy Saving Trust website, and the number of stations per DA has been used as an activity parameter to distribute UK-based emission figures for LPG consumption across each DA. It is hoped that in future years, actual LPG sales data by DA may become available to provide a more accurate methodology, though it should be noted that consumption of LPG as a transport fuel is still very small in comparison with consumption of petrol and diesel and has been declining from levels reached in 2006.

Emissions of methane and nitrous oxide from LPG consumption have been calculated for the first time based on an estimate of the number of vehicles and distances travelled using this fuel. This improvement applied to all DAs.

² The totals in this table include emissions from Diesel and Petrol use, but do not include the small emissions from LPG and lubricants. These figures are therefore not directly comparable with the road transport emissions presented in Appendix 2, which include emissions from all fuel use.

1.4.2.7 Trends in GHG Emissions from Road Transport

Table A1.4.7 below sets out the carbon dioxide and GHG emissions from 1990 to the latest inventory year (2009) from the two methods of estimating road transport emissions of carbon dioxide.

Table A1.4.7: Emissions of GHGs from UK road transport, according to fuel type and percentage changes from 1990 to the latest inventory year (kt CO₂ equivalent).


Calculation method	GHG	Fuel used	1990	2009	Percentage change 1990-2009
Constrained	Carbon	LPG	-	316.04	
		Petrol and DERV	108,853.53	111,659.21	
		Lubricants	262.77	121.25	
	CH ₄	LPG	-	0.81	
		Petrol and DERV	627.83	87.43	
		LPG	-	2.77	
	N ₂ O	Petrol and DERV	1,170.04	966.55	
		Sum	110,914.16	113,154.07	2.02%
Calculation method	GHG	Fuel used	1990	2009	Percentage change 1990-2009
Unconstrained	Carbon	LPG	-	316.04	
		Petrol and DERV	108,135.90	111,640.48	
		Lubricants	262.77	121.25	
	CH ₄	LPG	-	0.81	
		Petrol and DERV	627.83	87.43	
		LPG	-	2.77	
	N ₂ O	Petrol and DERV	1,170.04	966.55	
		Sum	110,196.54	113,135.33	2.67%

The emissions of methane and nitrous oxide are estimated using vkm data in both of the calculation methods, and the total emissions of these GHGs from the two methods are identical. Carbon emissions of LPG and lubricants burnt in engines are very small relative to emissions from the combustion of petrol and DERV. For convenience, the emissions from LPG and lubricants have not been constrained to fuel sales, and, have been assumed equal in magnitude in both calculation methods in the comparison above. The emissions are quoted to 0.01 kt purely for convenience, to avoid the risk of rounding errors. The number of decimal places used should not be taken as indicative of the accuracy of the estimates.

1.4.3 Railways

Emissions from railway locomotives in Great Britain are disaggregated based on diesel oil consumption data supplied by the Association of Train Operating Companies (ATOC) and the Office of Rail Regulation (ORR). Fuel consumption was estimated for Intercity and regional rail movements on the basis of reported train kilometres travelled and adjusted to align with the ORR estimates of fuel consumption. As no data from ATOC/ORR were available for 2009, fuel consumption was estimated on the basis of trends in train km from 2008. Fuel consumption for rail freight was also estimated from train km data and adjusted to align with ORR estimates of fuel consumption for the rail freight sector. Again, no data from ATOC/ORR were available for rail freight in 2009, so fuel consumption was estimated on the basis of trends in tonne km from 2008. Alignment with data provided by ORR marked an improvement in the methodology and the change was reflected across the time series.

The data from ATOC includes fuel use data for each passenger railway company whose area of operation can in most cases be allocated to one of England, Wales and Scotland. Emissions from railways in Northern Ireland are based on fuel consumption data supplied by Translink. There are no rail freight activities in Northern Ireland.

Figure A1.4.1: Road transport CO₂ emission trends 1990-2009 calculated by constrained and unconstrained methodologies

1.5 OTHER SECTORS

1.5.1 Commercial & Institutional

Emissions estimates for the source categories “public administration” and “miscellaneous and commercial” have previously been based on regional proxy activity data including GVA (as a broad indicator of economic activity across the DAs) or regional employment statistics. Similar to the source categories for small-scale industry and the domestic sector, there is very little detailed solid or liquid fuel use data for these sectors and hence the estimates are subject to greater uncertainty than well-documented sectors (i.e. energy-intensive industries).

The DECC sub-national energy statistics (DECC, 2010b), provide estimates of fuel use by Local Authority for each of these sectors, split by solid fuel types and “oil”. These data are estimates that are based on (i) local electricity and gas meter data, and (ii) modelled estimates of the distribution of solid and liquid fuels using proxy data, concessionary coal data and information on smoke control zones. The estimation methodology has been developed for the latest inventory cycle and now follows a similar method to that described for other industrial combustion (see section 1.3.2). For gas oil, coal and gas, the available point source emissions data and fuel use data from EUETS and the pollution inventories have been analysed to allocate emissions to the DAs. The remaining emissions are allocated to the DAs using the energy modelling approach consistent with the DECC sub-national energy statistics.

National gas sales data for the commercial sector were previously reported by DTI (1992), but for later years (1995 to date) UK National Grid has provided data for regional gas use in the 73-732 MWh range. The UK National Grid source provides the closest data available for commercial and institutional consumers, but the total is lower than UK data reported by DECC (2010a). This data is used to distribute miscellaneous and public service gas use in GB.

Natural gas use data for Northern Ireland are supplied by Phoenix Gas for 1999 onwards (Phoenix Gas, 2010), with a new supplier, Firmus Energy, also providing sales data for 2005 onwards (Firmus Energy, 2010). The commercial consumption is used as an estimate for Northern Ireland miscellaneous and public service gas consumption. A more detailed split of gas use across the domestic, commercial and industrial sectors in Northern Ireland in recent years has been provided by Phoenix Gas, and expert judgement has been used to estimate the overall allocation of gas use to these sectors from the total gas sales data for Northern Ireland.

Consultation with DFPNI has led to the provision of detailed new energy data from public sector energy reports from 2002 to 2009, covering all fuels (including electricity) used in public sector buildings in Northern Ireland. The Public Sector Energy Campaign (PSEC) data have been used to replace previous estimates of fuel use in that sector, for most (but not all) fuels. The data scope covers building energy use and is a close match to the DUKES category description, and therefore the data have been used directly to inform gas and solid fuel use within the public sector in Northern Ireland. The reported gas oil use in the PSEC report is significantly higher than that currently reported for the UK as a whole; in the UK GHGI programme, the limited data on gas oil has been identified as problematic and these data from PSEC should now also be taken into consideration to help inform future gas oil allocations to the public sector. In the current NI inventory, therefore, there is a small under-report in public sector emissions due to this discrepancy. However, for other fuels the use of the PSEC data provides a more accurate estimate of sector emissions and trends.

Stationary combustion by the railway sector is classified as a commercial source. Consumption of burning oil, fuel oil, and coke is relatively insignificant, and has therefore been allocated according to the diesel oil driver used for locomotives. Natural gas consumption for electricity generation refers to the London Underground (Lotts Road power station – closed in 2001).

DECC (2010a) reports a small amount of solid waste (municipal, industrial & hospital) consumption for energy production in the commercial and miscellaneous sectors. Little is known about the distribution of these installations, but the emissions have been distributed using the split derived for MSW incinerators.

Table A1.5a: Other Sectors (Base Year – 1990)¹

IPCC Category	NAEI Sources	Activity: Consumption	Fuel	1990
Commercial & Institutional	Miscellaneous, Public service	Coal		DECC Sub-national energy statistics
		SSF		DECC Sub-national energy statistics
		Natural gas		Commercial Sales, DECC.
		Landfill gas		Landfill methane emissions
		Sewage gas		Sewage methane recovered
		fuel oil, gas oil		DECC Sub-national energy statistics
		MSW		As MSW incinerators
		Burning oil		DECC Sub-national energy statistics
	Railways (Stationary)	fuel oil, burning oil, coal		Sub-national oil consumption, DECC
		Natural gas		Assumed as all England
Residential	Domestic	Wood		Domestic wood mapping grid
		Peat		Domestic peat consumption data, CEH
		SSF, coke, LPG		Sub-national energy statistics (SSF), DECC & Housing Condition Survey data, NI HECA, census data
		Natural gas		Domestic Gas data, DECC
		Burning oil, gas oil,		Sub-national energy statistics (oil), DECC & Housing Condition Survey data, NI HECA, census data
		Coal, anthracite		Sub-national energy statistics (coal, anthracite), DECC & Housing Condition Survey data, NI HECA, census data
		Fuel oil		Regional population, ONS
	House & Garden	DERV, petrol		Regional dwellings, ONS
Agriculture, Forestry & Fishing	Agriculture – stationary combustion	coal, coke, natural gas		Agricultural employment, MAFF
		burning oil, gas oil, fuel oil		DECC Sub-national energy statistics
		straw		Wheat production, MAFF
	Agricultural mobile machinery	Gas oil, petrol		Agricultural off-road mapping grid

1 See Section 1.1.3 for abbreviations

Table A1.5b: Other Sectors (1995; 1998 to 2009)

IPCC Category	NAEI Sources	Activity: Fuel Consumption	Data Sources / Comments
Commercial & Institutional	Miscellaneous, Public service	Coal	DECC Sub-national energy statistics, analysis of point source data and energy modelling data, including EUETS data and PI/SPRI/ISR data. Northern Ireland Public Sector Energy Campaign data (DFPNI, 2010)
		SSF	DECC Sub-national energy statistics
		Natural gas	Natural gas consumed, Transco (now UK National Grid), Phoenix, Firmus, PSEC data (DFPNI, 2010)
		Landfill gas	Landfill methane emissions
		Sewage gas	Sewage methane recovered
		fuel oil, gas oil	DECC Sub-national energy statistics, analysis of point source data and energy modelling data, including EUETS data and PI/SPRI/ISR data, PSEC data (DFPNI, 2010)
		MSW	As MSW incinerators
		Burning oil	DECC Sub-national energy statistics, PSEC data (DFPNI, 2010)
	Railways (Stationary)	fuel oil, burning oil, coal	Regional gas oil consumption, Network Rail (GB) and Translink (NI)
		Natural gas	Assumed as all England
Residential	Domestic	Wood	Domestic wood mapping grid
		Peat	Domestic peat consumption data, CEH
		SSF, coke, LPG	Sub-national energy statistics (SSF), DECC, 2009 mapping grid, using Housing Condition Survey data, NI HECA, DEMScot model, census data
		Natural gas	Domestic Gas sub-national split for GB from DECC, Transco & UK gas network operators, Data from Phoenix Gas and Firmus Energy (NI)
		Burning oil, gas oil,	Sub-national energy statistics (oil) from DECC, 2009 mapping grid using Housing Condition Survey data, NI HECA, DEMScot model, census data
		Coal, anthracite	Sub-national energy statistics (coal, anthracite), DECC, 2009 mapping grid using Housing Condition Survey data, NI HECA, DEMScot model, census data
		Fuel oil	Regional population, ONS
		DERV, petrol	Regional dwellings, ONS
	House & Garden		
Agriculture, Forestry & Fishing	Agriculture stationary combustion	coal, coke, natural gas	Agricultural employment, MAFF/Defra
		burning oil, gas oil, fuel oil	DECC Sub-national energy statistics
		straw	Wheat production, MAFF/Defra
	Agricultural mobile machinery	Gas oil, petrol	Agricultural off-road mapping grid

1.5.2 Residential

The emission estimates for the residential combustion of fuels within the DAs has been revised within the latest inventory cycle. At UK level, activity data have been revised across the time series for peat use and petcoke use in the residential sector (MacCarthy, 2011), with small revisions to natural gas activity data in recent years.

In addition to the UK changes, the AEA emissions mapping team have updated the domestic sector emission maps to include new census data and more detailed housing data for Northern Ireland, integrating further improvements following the revisions to GB domestic sector estimates in the 1990-2008 cycle. Therefore, the domestic DA fuel allocations are not fully consistent with the DECC sub-national energy statistics published in December 2010, as the revisions to mapping grids has occurred since that publication.

The new data on peat use in the domestic sector, provided following research by CEH, greatly reduces the estimates for peat use in the domestic sector, especially in Northern Ireland. The maps and CEH peat data only provide snapshots of analysis for the latest year and where large revisions to previous data are evident, the DA inventory compilation must consider the impacts on time-series consistency for the sector as a whole. In order to enable a sector-wide quality check on the time-series data for the sector, estimates of the energy allocations in the sector across all fuels were derived (including estimated electricity use in each DA in the domestic sector).

Using the time series of population data, the energy use estimates per capita were calculated, in order to review the relative energy intensity per head across the time series for each DA. This is a quality check of limited usefulness given the variability in housing stock, fuel availability (e.g. on gas grid or not) climatic considerations and inter-annual variations of factors such as fuel price and average temperature which will all affect the local energy use in the sector. Nevertheless, the quality check led to some small revisions in the DA trends, but more work is recommended to further develop the domestic sector dataset for each of the DAs, as this is a sector where DA policy levers can have a large impact and currently the evidence base for the energy use in the sector is uncertain.

The domestic sector emission estimates have been extrapolated back from recent mapping grid estimates, drawing on data on primary fuels used in the sector from Housing Condition Surveys in 1996, 2001 and 2006 (HECA NI, 2008), which provide a summary of fuel-switching trends away from solid fuels due to the development of the gas network and use of burning oil in Northern Ireland since the late 1990s. This approach is used for coal, anthracite, gas oil and burning oil. The Scottish Housing Condition Survey, census data and DEMScot model have been considered within new domestic sector energy mapping work for 2009, which also considered more recent (2009) housing post-code level data for Great Britain, retaining some assumptions on the DA share of housing types from the 2001 census. (Personal Communication: Tsagatakis, 2010)

Domestic natural gas consumption data is estimated for GB using the split presented in the DECC sub-national energy statistics for 2005 onwards (DECC, 2010a) whilst data for earlier years draws upon regional data obtained from Transco and other GB gas supply network operators. Domestic gas use in Northern Ireland is based on data from Phoenix Gas (2010) and Firmus Energy (2010). However, the accuracy of the DA domestic gas use estimates is subject to uncertainty, as gas companies use scale of annual consumption to allocate users to the domestic sector, which implies that small-scale commercial users will also be reported in the domestic sector. Furthermore, the sub-national energy statistics are based on meter point data that are weather-corrected and cover the "gas year" (January to January) rather than the calendar year.

The consumption of fuel oil by the domestic sector is a very small amount, and is distributed simply according to population (ONS, 2010a). Domestic use of wood is estimated across the time-series using the latest mapping grid information on wood use (NAEI mapping, 2011). Domestic peat use estimates by DA are provided by the Centre for Ecology & Hydrology (Personal Communication: Thomson, 2010).

1.5.3 Agriculture, Forestry & Fisheries

Emissions from solid fuel use in the agriculture sector are not very significant, whilst regional gas use data in this sector are not available, and hence emissions from these sources are allocated on the basis of regional employment figures from DEFRA (2010a).

Work by AEA (NAEI, 2008) to derive a more detailed split of regional off-road fuel use (i.e. mainly gas oil use in tractors and other mobile machinery) has utilised research to determine the regional distribution of different land uses and farm types (pasture, arable, forestry). These data have been combined with data on the intensity of mobile machinery use by farm type (tractor hours per hectare of arable land, tractor hours per head of livestock), to derive an agricultural off-road mapping grid to estimate geographical distribution of fuel use in the sector. These data have been used to estimate DA GHG emissions from agricultural mobile machinery.

1.6 MILITARY

Emissions from military aircraft and naval vessels are allocated across the DAs based on regional GVA data (ONS, 2010a). Army vehicle emissions are included within road transport data and other army emissions are included within public service categories but are not clearly defined.

1.7 FUGITIVE EMISSIONS FROM FUELS

1.7.1 Coal Mining

Methane emissions arise from coal mining activities. Emissions from operating mines are estimated based on the amounts of deep mined and open cast coal produced. DA inventory estimates are based on regional coal production derived from a number of sources: Coal Authority (2010), BGS (2010), WO (1998), SO (1999), BERR (1996). A small emission occurs from coal storage and transport, which is based on deep mined coal production. Data suggests that only small amounts of coal are transported outside of the region of production and no attempt has been made to allow for this. Hence coal storage and transport emissions are distributed according to deep-mined production (Coal Authority, 2010).

Methane emissions from abandoned coal mines have been updated in a recent research undertaken by WSP (2011) on behalf of DECC. As a result, DA emission allocations from this source have been revised across the time series. The report of this research can be found at:

http://uk-air.defra.gov.uk/reports/cat07/1107080945_1775-ghg-improvement-project-wsp-report.pdf

1.7.2 Solid Fuel Transformation

For coke ovens, three fugitive emissions are estimated:

1. A 'residual' emission of CO₂ which reflects the difference between the carbon input to the coke oven and the carbon content of the coke and coke-oven gas produced.
2. Emissions from the flaring of coke-oven gas.
3. Emissions of methane from the process.

These are disaggregated based on the regional consumption of coking coal discussed in Section 1.2.3. For solid smokeless fuel (SSF) plant, the only fugitive emissions estimated are the 'residual' CO₂ emission and some process methane. The driver used is that for regional consumption of coal by SSF plant (see Section 1.2.3). It is known that some petroleum coke is used in SSF production but the amount is uncertain. The same driver is applied to the petroleum coke consumption.

1.7.3 Oil and Natural Gas

All emissions from the oil & gas exploration and production industry that occur offshore are reported within the DA GHGI data as unallocated. Emissions from onshore oil and gas terminals in England, Wales and Scotland and from a small number of on-shore oil and gas fields, are based on operator reported data. In the 1990-2009 inventory cycle, the allocation of sites in the upstream exploration and production sector has been improved to enable emissions to be estimated separately for oil production sites and gas production sites. This improvement was implemented within the UK GHGI (MacCarthy et

al, 2011). The data management system for the DA GHG inventories has been improved in order to present DA-specific estimates of GHG emissions from the oil production and gas production sectors, leading to an improvement in emissions data detail and transparency.

The estimates of terminal flaring and venting emissions are based on DECC (2010e) EEMS data for 1995, 1998-2009. Data is unavailable for 1990, so these are extrapolated based on flaring volumes for Scottish Terminals and natural gas arrivals to gas terminals in England (DTI, 1991, 1996).

The 2000-2009 UK GHG inventories include a correction to account for flaring on onshore oil and gas fields excluded by the DECC EEMS inventory. Onshore flaring volumes are obtained from DECC sources (DECC, 2010d). Their significance in the UK national GHG inventory is minimal, but the data is more significant for the DA GHG inventories. Wytch Farm, which lies a few miles off the south coast of England, is classified as on-shore for this purpose.

The DECC EEMS inventory data provides data for fugitive emissions of carbon dioxide and methane from terminals for 1998-2009. Methane emissions arise from venting, oil storage and tanker loading and unloading, whilst carbon dioxide emissions arise from venting and processes. A more aggregated set of data for 1995 has been provided by UKOOA (1999), whilst estimates for 1990 have been calculated by extrapolation of data of oil and gas arrivals in England and Scotland (DTI, 1991, 1996) split across the sources and regions based on the 1995 dataset.

UK inventory estimates of emissions of methane due to leakage from the gas transmission system are based on UK National Grid data of leakage from the high-pressure network, Above Ground Installations and the low-pressure networks. Estimates are provided by National Grid (2010) and the other gas network operators: Northern Gas Networks (2010), Scotia Gas Networks (2010), Phoenix Gas (2010) and Wales & West Utilities (2010). Estimates are provided by Local Distribution Zones, enabling direct allocation to each of the constituent countries. The DA GHGI allocation method was reviewed in 2010 and an error was identified and corrected for recent years. In the 1990-2009 UK GHGI cycle, the method to derive the compositional analysis of natural gas provided in the UK network (including the calculation of the average national annual methane and carbon dioxide content of natural gas) was revised and updated (MacCarthy et al, 2011) and small revisions to these data have resulted in small recalculations in the DA leakage estimates, although there has been no revision to the DA disaggregation method.

Table A1.7a: Fugitive Emissions from Fuels (Base Year – 1990)¹

IPCC Category	NAEI Sources	Activity: Fuel Consumption	1990
Coal Mining	Deep mined coal Coal storage & transport	Deep mine coal production	Regional deep mine production, British Coal Authority. CH ₄ from closed coal mines from WYG 2005
	Open cast coal	Open cast mine coal production.	Regional open cast mine production, British Coal Authority CH ₄ from closed coal mines from WYG 2005
Solid Fuel transformation	Coke production	Coke production	Coal feed to coke ovens, ISSB, WS, DECC
	Flaring	Coke oven gas	Coal feed to coke ovens, ISSB, WS, DECC
	SSF production	Coal, Petrocoke	Coal feed to SSF plant, DECC, WS
Oil and gas production	Offshore Oil & Gas	NA	Fugitive emissions from Terminals (extrapolated from 1995)
	Oil Terminal Storage	NA	1998 driver
	Onshore Loading	Oil loaded	1998 driver
Venting & Flaring	Offshore Flaring	Volume gas flared	Flaring at terminals and onshore fields, UKOOA, DECC
	Offshore Venting	NA	Fugitive emissions from Terminals (extrapolated from 1995)
Natural Gas	Gas Leakage	Natural gas leakage	National Grid (Transco), Northern Gas Networks, Scotia Gas Networks, Wales & West Utilities

1 See Section 1.1.3 for abbreviations

Table A1.7b Fugitive Emissions from Fuels (1995; 1998 to 2009)

IPCC Category	NAEI Sources	Activity: Fuel Consumption	Data Sources / Comments
Coal Mining	Deep mined coal	Deep mine coal production	Regional deep mine production, British Coal Authority.
	Coal storage & transport		CH ₄ from closed coal mines from WSL 2011.
	Open cast coal	Open cast mine coal production.	Regional open cast mine production, British Coal Authority
			CH ₄ from closed coal mines from WSL 2011.
Solid Fuel transformation	Coke production	Coke production	Coal feed to coke ovens, ISSB, WS, DECC and (1999-current) PI
	Flaring	Coke oven gas	Coal feed to coke ovens, ISSB, WS, DECC and (1999-current) PI
	SSF production	Coal, Petrocoke	Coal feed to SSF plant, DECC, WS
Oil and gas production	Offshore Oil & Gas	NA	Oil & Gas UK GHG emissions from Terminals, DECC EEMS
	Oil Terminal Storage	NA	Data from storage emissions, DECC EEMS
	Onshore Loading	Oil loaded	Data from loading emissions, DECC EEMS
Venting & Flaring	Flaring	Volume gas flared	Flaring at terminals and onshore fields, Oil & Gas UK, DECC
	Venting	NA	Data from venting emissions, DECC EEMS
Natural Gas	Gas Leakage	Natural gas leakage	National Grid (Transco), Northern Gas Networks, Scotia Gas Networks, Wales & West Utilities, Phoenix Gas

1.8 INDUSTRIAL PROCESSES

These sources report process and fugitive emissions from industrial processes as opposed to the emissions from fuel combustion used to provide energy to these processes. (Table A1.2 covers combustion emissions.) The drivers used for process and fugitive industrial releases are summarised in Table A1.8.

1.8.1 Minerals Industries

Large emissions of carbon dioxide arise from the degradation of limestone used in cement and lime kilns. Cement emissions are estimated from the production of cement clinker, with regional emission estimations based on plant capacity data supplied by the British Cement Association³ (2004) for 1990 to 2001. From 2002 to 2007, the regional split is based on reported emissions from the PI, SPRI and ISR, whereas in 2008 and 2009 all cement sites now report under EUETS and hence the emissions from combustion and process sources by site are derived from EUETS data. Through discussions with environmental regulators it has been determined that lime calcination only occurs in England.

Limestone and dolomite are also used in iron and steel production. Corus suggest that it would be impossible to identify all the different uses of limestone and dolomite in iron and steel making. The major use is in blast furnaces, and so emissions have been disaggregated based upon regional iron production figures (ISSB, 2010).

Limestone, dolomite and soda ash are also used in glass production. Emissions were previously disaggregated using plant capacity and carbon dioxide emissions data from British Glass for 1990, 1995, 1998 and subsequently extrapolated for 1999 and 2000. However, the improvement of data supplied via the Pollution Inventory (Environment Agency, 2010) has enabled more accurate disaggregation for the years 2000 and 2001. Historic data has therefore been revised where appropriate and the Pollution Inventory data now provides a more accurate methodology for regional disaggregation of UK data from 2002 onwards.

The 2009 EUETS dataset contains a much greater coverage of sites and emission sources (combustion and process) from the glass industry, which has led to time-series revisions of DA activity and emissions. Previously the DA allocation of emissions from the glass sector was based on site information on production capacity, but access to fuel use data for 2009 has enabled these assumptions to be over-written with some fuel-specific DA allocations. These revisions have led to improvements to DA GHGI data and also to the DA air quality pollutant inventories for emission estimates in recent years.

The inventory also reports carbon dioxide and methane emissions from Fletton brick production, as introduced in 2000. These bricks are made from Fletton clay which contains a significant amount of naturally occurring carbonaceous material and all such production occurs in England.

1.8.2 Chemical Production

The UK Inventory reports emissions of carbon dioxide from ammonia production; nitrous oxide from adipic acid production and nitrous oxide from nitric acid production. Following the closure of a (nitric acid) fertiliser plant in Belfast in late 2001, all of the nitric acid, ammonia and adipic acid plants are within England. Prior to that, plant capacities for nitric acid production facilities were used to estimate the split in UK chemical production GHG sources.

³ Production capacity data are used for cement emissions as the actual annual production data from cement plant are commercially confidential.

Table A1.8a: Industrial Processes (Base Year – 1990)¹

IPCC Category	NAEI Sources	Activity Data	1990
Cement Production	Cement (decarbonising)	Clinker production	Regional cement production capacity, BCA
Lime Production	Lime (decarbonising)	Limestone consumption	All such plant located in England
Limestone and Dolomite Use	Glass production	Limestone and dolomite consumption	Regional glass production, British Glass
	Blast Furnaces	Limestone and dolomite consumption	Iron production, ISSB
Soda Ash Production and Use	Glass production	Soda Ash Consumption	Regional glass production, British Glass
Mineral Products: Other	Fletton Brick Production	Fletton Brick Production	All such plant located in England
Ammonia Production	Ammonia feedstock	Natural gas feedstock	All such plant located in England
Nitric Acid Production	Nitric Acid Production	Plant capacity	Regional plant capacity
Adipic Acid Production	Adipic Acid Production	Adipic acid made	All such plant located in England
Chemical Industry: Other	Methanol Production	Production of Methanol	All such plant located in England
Chemical Industry: Other	Ethylene Production	Production of Ethylene	Plant capacities
Chemical Industry: Other	Chemical Industry	NA	Extrapolated from PI data
Iron and Steel	Electric Arc Furnace	EAF steel production	Regional EAF production, ISSB
	Flaring	Blast furnace gas	Coke consumed in blast furnaces, ISSB, WO
Aluminium Production	Aluminium production	Primary aluminium produced	Regional aluminium plant capacity, ALCAN
SF ₆ Used in Aluminium and Magnesium Foundries	SF ₆ Cover gas	NA	Regional consumption & sales data, EM industry report 1999

1. See Section 1.1.3 for abbreviations

Table A1.8a: Industrial Processes (Base Year – 1990) (continued)

IPCC Category	NAEI Sources	Activity Data	1990
Halocarbon & SF ₆ By-Product Emissions	Halocarbon Production	NA	All such plant are located in England.
Refrigeration and Air Conditioning	Refrigeration	NA	Regional population, ONS
	Supermarket Refrigeration	NA	Regional GDP, ONS
	Mobile Air conditioning	NA	Vehicle Registration data, AEAT industry report 2003
Foam Blowing	Foams	NA	Regional population, ONS
Fire Extinguishers	Fire fighting	NA	Regional population, ONS
Aerosols	Metered Dose Inhalers	NA	Regional population, ONS
	Aerosols (halocarbons)	NA	Regional population, ONS
Other	Electronics	NA	Regional electronics plant consumption, EM industry report 1999
	Training shoes	NA	Regional population, ONS
	Electrical Insulation	NA	Regional electrical capacity, AEAT industry report 2003

Table A1.8b: Industrial Processes (1995; 1998 to 2009)

IPCC Category	NAEI Sources	Activity Data	Data Sources / Comments
Cement Production	Cement (decarbonising)	Clinker production	Point source data from EUETS (2008 only), and the PI/SPRI/ISR (EA, SEPA & NIEA).
Lime Production	Lime (decarbonising)	Limestone consumption	All such plant located in England.
Limestone and Dolomite Use	Glass production	Limestone and dolomite consumption	Regional glass production, BGlass. EUETS (2009)
	Blast Furnaces	Limestone and dolomite consumption	Iron production, ISSB
Soda Ash Production and Use	Glass production	Soda Ash Consumption	Regional glass production, BGlass. EUETS (2009)
Mineral Products: Other	Fletton Brick Production	Fletton Brick Production	All such plant located in England
Ammonia Production	Ammonia feedstock	Natural gas feedstock	All such plant located in England
Nitric Acid Production	Nitric Acid Production	Plant capacity	Regional plant capacity, PI/SPRI/ISR. Since 2002, all such plant located in England.
Adipic Acid Production	Adipic Acid Production	Adipic acid made	All such plant located in England
Chemical Industry: Other	Methanol Production	Production of Methanol	All such plant located in England
Chemical Industry: Other	Ethylene Production	Production of Ethylene	Plant Capacities, PI/SPRI/ISR
Chemical Industry: Other	Chemical Industry	NA	PI/SPRI/ISR data, or extrapolated from PI data
Iron and Steel	Electric Arc Furnace	EAF steel production	Regional EAF production, ISSB
	Flaring	Blast furnace gas	Coke Consumed in blast furnaces, ISSB, WO
Aluminium Production	Aluminium production	Primary aluminium produced	UK plant production & emissions data, Alcan, Rio-Tinto, EA, SEPA
SF ₆ Used in Aluminium and Magnesium Foundries	SF ₆ Cover gas	NA	Regional consumption & sales data from industry reports compiled by EM & AEAT

Table A1.8b: Industrial Processes (1995; 1998 to 2009) (continued)

IPCC Category	NAEI Sources	Activity Data	Data Sources / Comments
Halocarbon & SF ₆ By-Product Emissions	Halocarbon Production	NA	All such plant are located in England.
Refrigeration and Air Conditioning	Refrigeration	NA	Regional population, ONS
	Supermarket Refrigeration	NA	Regional GDP, ONS
	Mobile Air conditioning	NA	Vehicle Registration data, AEAT industry report 2003
Foam Blowing	Foams	NA	Regional population, ONS
Fire Extinguishers	Fire fighting	NA	Regional population, ONS
Aerosols	Metered Dose Inhalers	NA	Regional population, ONS
	Aerosols (halocarbons)	NA	Regional population, ONS
Other	Electronics	NA	Regional electronics plant consumption, EM industry report 1999 & AEAT industry report 2003
	Training shoes	NA	Regional population, ONS
	Electrical Insulation	NA	Regional electrical capacity, AEAT industry report 2003

The UK inventory reports emissions of methane from methanol production, ethylene production and the other chemical industry. The methanol plant is located in England, whilst ethylene production occurs in England, Scotland and Wales. These emissions are distributed based on data reported in the PI (Environment Agency, 2010a), SPRI (SEPA, 2010a) and plant capacity. Emissions are extrapolated to 1990 and 1995 based on plant capacities.

The emissions from the “other chemical industry” sector are disaggregated to England and Wales based on the site data in the Pollution Inventory. Data on emissions from other chemical processes are not available for Scotland.

1.8.3 Metal Production

In the iron and steel industry, emissions of carbon dioxide arise from electric arc furnaces through the consumption of the graphite anodes. Regional data on steel production from electric arc furnaces is used to determine the regional drivers for this activity (ISSB, 2010).

The flaring of waste blast furnace gas is disaggregated according to the distribution of blast furnaces, using the driver derived for coal consumption by blast furnaces (ISSB, 2010).

Emissions of carbon dioxide from iron and steel making are estimated from a mass balance on the coke consumed in blast furnaces; the blast furnace gas produced; the pig iron produced; the pig iron used in steel making and the crude steel produced. The emissions are distributed using appropriate drivers for each source and sink taken from ISSB (2010). These include regional data on coke consumed in blast furnaces, pig iron production and crude steel production.

The electrolytic process used to produce aluminium results in a carbon dioxide emission as the petroleum coke anode is consumed. Emission estimations are based on plant capacity data provided by Alcan (2004), for years up to 2002. The DA emissions data for 2003 onwards are based on PI and SPRI data (EA, 2010a; SEPA, 2010a). There have been some significant changes in the aluminium industry in recent years, with the closure of the Kinlochleven plant in 2000, and the expansion of the Lynmouth plant, and hence there has been a swing in emissions from this sector from Scotland to England.

The anode baking process within aluminium production also results in emissions of PFCs, and estimates are provided by plant operators (Alcan, 2010).

1.8.4 Use of Halocarbons and SF₆

The UK emissions of halocarbons and sulphur hexafluoride were based on estimates from a model prepared initially by Enviros March (1999). This model has now been updated by AEAT (Haydock et al, 2003), with further updates at UK level for the refrigeration sectors during 2008-9 (Personal Communication: MacCarthy, 2010). For some sources, the emission is equal to the consumption of fluid (e.g. aerosols). For other sources the emissions occur during product manufacture, leakage during product lifetime, and at product disposal (e.g. refrigerators). In these cases emissions are estimated from a time dependent model of the bank of fluid held in products, accounting for unit production and disposal.

Data for HFC emissions from metered dose inhalers in the UK are also taken from the EM & AEAT predictive models. The National Asthma Campaign's National Asthma Audit (1999-2000) concluded that:

“There is little variation in asthma prevalence among children or adults throughout Great Britain.”

Therefore, the regional split of emissions is proportional to population.

Supermarket refrigeration is regarded as sufficiently different from other refrigeration to warrant a separate study. Emissions are based on a market review of the number and size of supermarkets in the regions, combined with discussions with gas manufacturers on the sales into this sector. Discussions with supermarket owners also suggest that regional use could be approximately equated to sales volume, which in turn could be approximated by regional GVA estimates, which have been obtained from ONS (2010a). The DA GVA data are therefore used to estimate the share of refrigeration emissions within the UK.

Air conditioning systems in cars began to use HFC134a from around 1993. Data is supplied by SMMT on regional sales of new cars. Initially, installation of air conditioning was skewed towards company cars, which are broadly distributed according to population.

PFCs and SF₆ are used to cushion the soles of some training shoes. Data have previously been gathered from discussions with Nike. Sales figures for the devolved regions of the UK were not available, and therefore the regional split is made according to population.

Sulphur hexafluoride is used in electrical switchgear within the electricity transmission system. UK estimates are based on discussions with industry sources and summarised within the EM & AEAT model. Regional estimates are determined through consultation with power supply companies (NIE, Scottish Power & Scottish Electric, National Grid) and the Electricity Association.

For aerosols, the split by region is made on the basis of population, although use of these gases often have industrial applications. Making the split using population has the advantage of making the data directly comparable with the figures for the baseline years of 1990 and 1995.

Other sources such as fire extinguishers are very small and are likely to be distributed with the general population.

Emissions of SF₆ cover gas from magnesium production is based on regional sales and consumption data. This stable market is assessed within the AEAT model (2003), with all production located in England & Wales.

Emissions data for regional emissions from semiconductor wafer manufacture are estimated from manufacturing data and consultation with relevant trade associations, and incorporated within the AEAT model (2003).

1.9 AGRICULTURE

Rothamsted Research provides all data and information pertaining to agricultural sources within the Devolved Administration emission inventories.

The UK inventory is disaggregated into England, Scotland, Wales and Northern Ireland, with all default factors and emission factors carried over from the national inventory.

Regional crop areas are obtained from the Defra June Agricultural Census for 1990, 1995, 1998 - 2009 (Defra, 2010), Welsh Assembly Government (2010), Scottish Government (2010) and DARD (2010). Crop production data is taken from Agriculture in the UK and Basic Horticultural Statistics for the UK. The Welsh Assembly Government also provides crop area data in Wales for this inventory.

Fertiliser applications are derived from regional crop areas and average application rates published in the British Survey of Fertiliser Practice for 1990, 1995, 1998 - 2009 (BSFP, 2010), which presents data for England and Wales, Scotland and Great Britain. Application rates in Northern Ireland are assumed to be the same as Scottish applications. In many cases, the sample size used to estimate fertiliser use in Scotland is considered too small to be sufficiently robust and in these cases, the Great Britain data are used. Where application rates are not available for particular crop types, the crop area is amalgamated with a similar crop with a known fertiliser application rate. Where annual applications are not available, fertiliser application for a different year are used.

Livestock numbers are obtained from CEH (Ulrike Dragosits) calculations based on data from Defra, Scottish Government, the Welsh Assembly and DARD Agricultural Census data for 1990, 1995, 1998 - 2009.

Other data such as milk production and fat in milk are obtained from Agriculture in the UK (Defra, 2010).

Dairy cattle weight is from slaughter weight data provided by Sarah Thompson, Defra. A Tier 2 methodology is used for the calculation of the enteric emissions from beef cattle, but a time series of cattle weights are not available, and so a constant weight of 500 kg has been assumed. A country

specific emission factor is used, assuming a weight of 500 kg. A Tier 2 methodology is used for the calculation of the emissions from other cattle but weight is not changed from year to year.

N excretion factors are kept in agreement with the UK NH_3 inventory (Cottril and Smith, ADAS).

As in the national inventory, the area of cultivated histosols (soils of high organic content) is assumed to be equivalent to the area of Eutric Histosols, and is disaggregated according to a percentage split estimated by the Soil Survey and Land Research Centre (personal communication).

Time series of the application of sewage sludge to land was included in the 2009 inventory. Emissions from sewage sludge used as fertiliser are reported under agricultural soils by IPCC. The calculation involves estimating the amount of nitrogen contained per dry matter unit of sludge and applying IPCC emission factors. Data sources for the annual production of sewage sludge (as dry matter) were obtained from Brian Chambers (pers. comm.ADAS) for 1996; from Directive 86/278 for 2004-2006 on the basis of the data contained in the records referred to in Article 10; Sludge used in agriculture (Tonnes dry matter/year); from OFWAT, the Water Commissioner for Scotland and the Northern Ireland regulator, UREGNI for 2008-2009. The amounts for the missing years were derived by interpolation/extrapolation of the available data. The UK follows the IPCC (1997) methodology. This assumes that 20% of the total sludge N applied to soil volatilises as NO_x and NH_3 and therefore does not contribute to N_2O emissions.

A correction to the calculation of the emissions from leaching was introduced. Previously the UK subtracted volatilised N and N_2O emitted from the applied N to calculate the fraction leached. According to the IPCC no subtraction is required for the calculation of the fraction leached, so the UK has now corrected this methodology.

The impact of the changes due to inclusion of sewage sludge and the correction of the calculations in leaching produced an increase in total emissions of N_2O in all years and for example for the year 2008, the increase was 6.4, 6.9, 5.5 and 6.0% for England, Wales, Scotland and Northern Ireland respectively.

There are no regional differences in the parameters currently used to calculate emissions (except for activity data).

Planned improvements to the inventory

UK emission factors are currently under review for:

- EF1, emission factor for direct soil emissions; from a literature review and a field measurement programme.
- EF3, emission factor from manure management systems); from a literature review and a field measurement programme and,
- EF5, nitrogen leaching/runoff factor; from a field measurement programme

The UK aims to improve the spatial disaggregation of the inventory by calculating emissions using parameters specific at DA level.

The UK is improving the link between the NH_3 and GHG inventories, and incorporating NO_x in a study (desk/experimental) which will review the current value of 20% of N lost as NH_3 and NO_x .

Any small differences in national-regional census data are removed by normalising the DA inventories such that the sum of England, Scotland, Wales and Northern Ireland equals the UK emission. (For details of the normalisation procedure, see Section 1.1.2.) In particular, for census years prior to 2001, the supply of data for the constituent countries from different sources (MAFF, DANI and the Scottish Office) was not concurrent with obtaining the UK data and submitting the UK inventory. This lack of synchronicity was not conducive to ensuring that there was good agreement between the sum of country data and the UK figure, and consequently the UK and disaggregated inventory estimates. This problem was rectified in some recent years by the supply of all required land use and livestock data directly from DEFRA statistics. As this is no longer possible, the current protocol is to obtain data from constituent countries prior to submission of the UK inventory, altering UK figures where necessary.

A new spreadsheet will be submitted from the next inventory submission, with structure for potential inclusion of Tier 2 methodology for CH₄ in all livestock categories.

A large programme of improvements to the UK inventory is underway, as a platform of projects to provide country specific data has been funded by Defra. Improvements include analysis of available data from the literature, and specific experimental work to improve the estimates of emissions of N₂O and CH₄.

1.10 LAND USE, LAND USE CHANGE & FORESTRY (LULUCF)

The estimates for the LULUCF sector within the DA emissions inventories are made by the Centre for Ecology and Hydrology. The methods for estimating emissions and removals of greenhouse gases from this sector, data sources and underlying assumptions are described in the latest UK national inventory report for 1990-2009 (MacCarthy et al., 2011).

Detailed information on the data and methods used in the LULUCF inventory is available in the 1990-2009 UK Greenhouse Gas Inventory Report, available on the National Atmospheric Emissions Inventory website <http://naei.defra.gov.uk/>. Chapter 7 and Annex 3.7 contain information on the LULUCF sector, and Chapter 11 contains additional information on the reporting of LULUCF activities for the Kyoto Protocol. Additional information on LULUCF and KP-LULUCF inventory reporting has been made available at <http://ecosystemghg.ceh.ac.uk/>.

The current LULUCF inventory methods use a combination of top-down and bottom-up approaches, based on activity data for each of the Devolved Administrations and the UK as a whole. As a result of this approach, estimates of emissions and removals from LULUCF activities are automatically produced at the DA and UK scale.

Changes between the 2009 and the 2008 inventory

There are differences in the estimated net emissions in the 1990-2009 inventory compared with the previous inventory. There have been updates to methods and activity data this year, together with internal restructuring of the LULUCF calculation methods. Sector 5, which covers LULUCF emissions and removals reporting, now reports a slightly larger source in 1990 compared with the 1990-2008 inventory (3.876 vs. 2.953 Mt CO₂e) and a larger sink from 2000 onwards (-4.005 vs. -1.942 Mt CO₂e in 2008). These differences are due to the inclusion of new activity data for land use change between 2000 and 2007, the reporting of N₂O emissions from land-use conversion to cropland, the reporting of on-site emissions for the 5D Wetlands category and other minor revisions to methods, activity data and emission factors. These changes have been undertaken as part of the development of the inventory, or in response to international review team recommendations. The main changes are described below, with the main differences between inventories in the Devolved Administrations described in their respective sections.

The inclusion of new data on land use change from the Countryside Survey 2007 has made a marked difference to the estimates of net emissions/removals. In previous inventories, rates of land use change were based on the rate of change between 1990 and 1998, rolled forward to the most recent inventory year. With the new dataset, we have been able to recalculate rates of change for 2000 to 2007, and have extrapolated rates for 2008 and 2009 using the most recent rates (the LUC model is set up to use decadal land use change, e.g. 1990-1999 and 2000-2009). Rates of land use change for 1999 are assumed to be the same as those for 1990-1998. Changes to the rates of land use change affect the estimates of soil carbon stock changes from these land use conversions.

1.10.1: 5A Forest Land

Forest Land remaining Forest Land

There are 1168 kha of land that was covered in forest since before 1921: 727 kha in England, 312 kha in Scotland, 123 kha in Wales and 6 kha in Northern Ireland. This forest is assumed to be in overall carbon balance because of its age, and hence has a carbon stock change of zero. The area of forest reported in this category from 1990 to 2009 has been adjusted to take account of forest converted to other land use categories.

Land converted to Forest Land

Carbon stock changes

In the previous versions of the inventory, all land afforested since 1921 was included in Category 5A2 Land Converted to Forest as soil carbon fluxes due to conversion take many decades to reach equilibrium. In the 2009 Inventory, this category has been restructured to use a 20 year transition period for land use converted to forest. The fluxes from conversions prior to the 20 year transition period are still reported, now under the Category 5A1 Forest Remaining Forest. There is therefore no change in reported removals / emissions in the overall 5A Category.

Changes in carbon stocks in forest standing biomass, litter and soils are calculated with the C-Flow forest carbon model (Dewar and Cannell 1992, Cannell and Dewar 1995, Milne et al. 1998). The model is driven by records of annual forest planting in each DA on previously unforested land (available from 1922 onwards). Forests accumulate carbon (removing it from the atmosphere) in their biomass and soils as they grow, but timber harvesting and planting activities disturb this accumulation and result in emissions of carbon to the atmosphere. The net carbon stock change at any one time depends on the balance between these different activities. Forestry management cycles operate over long time scales (50 years+) so the rate of carbon removal in the present day is driven by the rate of forest planting in previous decades.

In 2009 around 59% of the area under forest land use in the UK had been planted since 1921 (401 kha in England, 1028 kha in Scotland, 161 kha in Wales and 82 kha in Northern Ireland). The activity data for annual forest planting (private and state) for the C-Flow model are provided by the Forestry Commission (for England, Scotland and Wales) and the Northern Ireland Department of Agriculture. The data is sub-divided into conifer and broadleaf planting and is time-consistent and reliable. There are differences in trends between conifer and broadleaf planting since 1990.

All DAs have seen a decline in annual conifer planting rates 1990-2009, however, annual broadleaf planting has on average increased in all DAs (Table A1.9).

The C-Flow model also requires input data on the stemwood growth rate and harvesting pattern, which come from standard management and yield tables used by the Forestry Commission (Edwards and Christie 1981). Following work by Milne et al. (1998) DA-specific growth patterns are used for conifers but the same growth pattern is used for all new broadleaf planting in the UK. Milne et al. (1998) have shown that different yield class assumptions for conifer and broadleaf planting have an effect of less than 10% on estimated carbon uptakes for the country as a whole. CEH and Forest Research are undertaking research to characterise forest planting (including species mix) and management in greater spatial and temporal detail. Comparisons of forest census data have already indicated that there is non-standard management (shorter rotations) in England and Wales during the mid-20th century. These variations in management have been incorporated into the forest model but they have only a small impact on net carbon stock change.

The C-Flow method is a Tier 3 modelling approach, so there are no explicit emission factors. Instead parameter values for either conifer or broadleaf trees are required to estimate (i) stemwood, foliage, branch and root masses from the stemwood volume, and (ii) the decomposition rates of litter, soil carbon and wood products. Losses of soil carbon due to the disturbance by planting activity are also considered on a DA-specific basis. From these values and the activity data C-Flow calculates the net changes in the pools of standing trees, litter and soil carbon.

Table A1.9: Forest Planting Rates since 1990

Annual Planting Rate kha a ⁻¹	England	Scotland	Wales	Northern Ireland
Conifer				
1990-1994	0.827	9.536	0.216	1.025
1995-1999	0.547	6.505	0.091	0.498
2000-2004	0.556	3.536	0.084	0.224
2005	0.191	1.865	0.013	0.034

2006	0.082	1.011	0.008	0.043
2007	0.064	2.000	0.037	0.034
2008	0.054	0.744	0.002	0.056
2009	0.007	1.168	0.020	0.020
Broadleaf				
1990-1994	3.828	3.383	0.312	0.238
1995-1999	4.016	4.997	0.420	0.297
2000-2004	4.507	5.180	0.368	0.435
2005	4.523	3.823	0.521	0.321
2006	3.030	2.984	0.452	0.563
2007	2.573	4.579	0.388	0.451
2008	1.999	3.427	0.201	0.496
2009	2.091	2.272	0.089	0.269

The overall uncertainty in emissions/removals from 5A Forest Land is estimated to be 25% but a full analysis of uncertainties is planned for future inventories.

Nitrous Oxide emissions from N fertilization of forests

Information on forest fertilisation was gathered from a search of the relevant literature and discussion with private chartered foresters and the Forestry Commission (Skiba, 2007). In the UK the general recommendation is not to apply fertiliser to forests unless it is absolutely necessary: it is not applied to native woodlands, mature forest stands or replanted forests. The instances where N fertiliser is applied to forests are first rotation (afforestation) forests on "poor" soil, e.g. reclaimed slag heaps, impoverished brown field sites, upland organic soils. In terms of the inventory, this means that N fertilisation is assumed for Settlement converted to Forest land and Grassland converted to Forest Land on organic soils. A Tier 1 approach is used with the amount of N fertiliser calculated using a fixed application rate and the areas of relevant forest planting for each DA from the same datasets used in the CFlow model for 5.A.2. Land converted to Forest land.

An application rate of 150 kg N ha⁻¹ is assumed based on Forestry Commission fertilisation guidelines (Taylor 1991). The guidelines recommend applying fertiliser on a three-year cycle until canopy closure (at c. 10 years), but this is thought to be rather high (Skiba, 2007) and unlikely to occur in reality, so two applications are adopted as a compromise. These applications occur in year 1 and year 4 after planting. As a result, emissions from N fertilisation since 1990 include emissions from forests that were planted before 1990 but received their second dose of fertiliser after 1990. The emission factor for N₂O of applied nitrogen fertiliser has been changed to 1% in line with the application of the IPCC 2006 Guidelines. Emissions of N₂O from N fertilisation of forests have fallen since 1990 due to reduced rates of new forest planting.

Emissions from wildfires on forest land

Estimates of emissions from wildfires on forest land are included in the inventory. The approach is Tier 2, using country-specific activity data and default emission factors. There is no information as to the age and type of forest that is burnt in wildfires, so all wildfire emissions are recorded under 5.A.2.

Estimates of the area burnt in wildfires 1990-2009 are made for Great Britain (data from the Forestry Commission) and Northern Ireland (data from the Forest Service). No data on areas burnt in wildfires has been collected or published since 2004. Activity data for 2005 to 2009 is extrapolated using a Burg regression equation based on the trend and variability of the 1990-2004 dataset. These areas refer only to fire damage in state forests; no information is collected on fire damage in privately owned forests. The area of private-owned forest that was burnt each year was assumed to be in proportion to the percentage of the state forest that was burnt each year. The Great Britain total forest estimate was split between the individual countries in proportion to their forest areas. An estimated 957 ha of forest is burnt on average every year (358 ha in England, 427 ha in Scotland, 91 ha in Wales and 75 ha in Northern Ireland) between 1990 and 2009, although there is high inter-annual variability.

There is no information on the type (conifer or broadleaf) or age of forest that is burnt in wildfires in the UK. Therefore, the amount of biomass burnt is estimated from the mean forest biomass density in each country of the UK, as estimated by the C-Flow model. These densities vary with time due to the different afforestation histories in each country. A combustion efficiency of 0.5 is used with a carbon fraction of dry matter of 0.5 to estimate the total amount of carbon released, and hence emissions of CO₂ and non-CO₂ gases (using the IPCC emission ratios).

The uncertainty for the wildfire activity data is estimated to be 50% for the activity data 1990-2004, but 100% for the 2005 to 2009 values, as these have been extrapolated from previous years. The IPCC default of 70% uncertainty is used for the emission factors.

1.10.2 5B Cropland

Cropland remaining Cropland

Changes in non-forest biomass resulting from yield improvements result in a carbon sink. The activity data on cropland area is DA-specific and the emission factor is fixed at 2% p.a. (Sylvester-Bradley et al. 2002).

The application of lime (limestone, chalk and dolomite) to Cropland (and Grassland) produces emissions of CO₂. Data on the use of limestone, chalk and dolomite for agricultural purposes is reported in the Business Monitor of Mineral Extraction in the UK (Office of National Statistics 2008). Estimates of the individual materials are provided by the British Geological Survey each year as only the totals are published because of commercial confidentiality rules for small quantities. The amount of lime applied is variable from year to year but generally reducing over time. The area that receives lime is calculated from the regional agricultural censuses, the Fertiliser Statistics Report (Agricultural Industries Confederation, 2006) and the British Survey of Fertiliser Practice (BSFP, 2009). This data is DA-specific from 2000 onwards for England, Scotland and Wales and estimated from GB rates before that. There is no specific information on the % area limed in Northern Ireland, which is therefore assumed to be the same as that in England. Fixed emission factors (for limestone/chalk and dolomite) are used to estimate emissions. The uncertainty in the activity data and emission factors are judged to be low and consistent over time.

Past drainage of lowland peatlands in England produces an ongoing change in soil carbon stocks and therefore carbon emissions. The baseline (1990) area of drained peatland is taken as 150,000 ha (Bradley 1997) with an assumption of no further drainage since that time. The annual loss decreases in proportion to the amount of carbon remaining (there are different implied emission factors for “thick” and “thin” peat).

Land converted to Cropland

Emissions and removals in this category result from (i) changes in soil carbon stocks, and (ii) changes in non-forest biomass carbon stocks due to land use change to Cropland. The methodology also applies to land converted to Grassland and to Settlement. Changes in carbon stocks from land use change to Forest land are calculated by the C-Flow model, as described above. Emissions of nitrous oxide (N₂O) from the disturbance associated with land-use conversion to cropland are reported for the first time in this inventory.

Changes in soil carbon stocks due to land use change to Cropland

Land use change results in soil carbon stock change because soil carbon density generally differs under different land uses and the land use change initiates a transition from one density value to another. The change in vegetation cover and management affects the amount of carbon that goes into the soil from biomass decomposition and the initial disturbance of the soil releases carbon to the atmosphere. An increase in soil carbon density results in a removal of CO₂ from the atmosphere, and vice versa. The rate of loss or gain of carbon depends on the type of land use transition. For transitions where carbon is lost, e.g. Grassland conversion to Cropland, the rate is “fast” (50-150 years), but for transitions where carbon is gained the rate is much slower.

The method used for this activity links a land use change matrix to a dynamic model of carbon stock change. Matrices of land use change have been constructed back to 1950 for each DA using country-specific land surveys (MLC 1986, Haines-Young et al 2000, Cooper and McCann 2002).

Areas of land use were assigned to the IPCC GPG categories – Forest land, Cropland, Grassland, Settlements and Other Land. The data available for the UK does not distinguish Wetlands from other

land use types so this category is not used for land use change, although data on peat extraction are now reported under the wetlands category.

Area change data exists up to 2007 (when the most recent Countryside Survey took place) and extrapolated from there to the latest inventory year (2009). Since 1990 an estimated 49.5 kha yr⁻¹ have been converted to Cropland in England, 15.6 kha yr⁻¹ in Scotland, 6.0 kha yr⁻¹ in Wales and 3.5 kha yr⁻¹ in Northern Ireland. This is the first inventory to include extrapolations based upon the 2007 Countryside Survey. Previous inventories were based upon the extrapolations from the 1998 Countryside Survey.

A database of soil carbon density, based on information on soil type, land cover and carbon content, is available at 1km scale for the UK (Bradley et al. 2005): this gives the soil carbon densities under the different land use categories for each UK country. DA-specific times, from the initial to the new soil carbon density following a land use change, are used with the core differential equation to calculate carbon fluxes per unit area (hence the implied emission factors will change over time).

Changes in non-forest biomass stocks resulting from land use change to Cropland

Changes in carbon stocks in non-forest biomass due to land use change are based on the same area matrices used for estimating changes in carbon stocks in soils. The biomass carbon densities for five basic land use types are then weighted by their occurrence in each DA to calculate mean biomass densities for Cropland, Grassland and Settlements. The mean biomass carbon densities for each land type were then weighted by the relative proportions of change occurring between land types, in the same way as the calculations for changes in soil carbon densities. Changes between these equilibrium biomass carbon densities were assumed to happen in a single year.

A Monte Carlo approach is used for both activities to vary the inputs for the core equation, and also give an estimate of uncertainty. The uncertainty in changes in soil carbon from the initial to the final equilibrium value is up to $\pm 11\%$ of the mean value. The uncertainty of the areas of land use change in each land use transition is assumed to be $\pm 30\%$ of the mean. Independent estimates of the uncertainty for each DA are not currently available.

1.10.3 5C Grassland

Grassland remaining Grassland

Emissions resulting from the application of lime are described in the Cropland remaining Cropland section. Peat extraction for horticultural previously reported for England, Scotland and Northern Ireland (it is negligible in Wales) in this category is now reported under Category 5D, Wetlands. Peat extraction for fuel use is reported in the Energy sector of the inventory.

Land converted to Grassland

Methods for estimating changes in non-forest biomass and soils due to land use change to Grassland are described in the Land converted to Cropland section. Since 1990 an estimated 68.1 kha yr⁻¹ have been converted to Grassland in England, 19.2 kha yr⁻¹ in Scotland, 5.0 kha yr⁻¹ in Wales and 5.1 kha yr⁻¹ in Northern Ireland.

When Forest land is converted to Grassland (i.e. deforestation) it is assumed that 60% of the standing biomass is removed as timber products and the remainder is burnt. This burning of forest biomass produces emissions of CO₂, CH₄ and N₂O.

Activity data on deforestation to Grassland in the UK is estimated from Forestry Commission unconditional felling licence data where there is no requirement to restock by replanting or regeneration (see Levy and Milne 2004). The most recent data from the Forestry Commission has been collated: these are estimates for England for 1990-2008 and estimates for GB in 1999-2001.

Estimates for Scotland and Wales are extrapolated from the England estimates using the GB:England ratios in 1999-2001. Areas of deforestation for 2003-2009 are estimated by extrapolation from earlier years. There are no data currently available for Northern Ireland.

Annual rates of deforestation to Grassland were low and stable from 1990 to 1998 (E: 0.16 kha yr⁻¹, S: 0.05 kha yr⁻¹, W: 0.01 kha yr⁻¹) when they increased rapidly to a peak in 2001 (E: 0.8 kha yr⁻¹, S: 0.25 kha yr⁻¹, W: 0.06 kha yr⁻¹). There is anecdotal evidence from the Forestry Commission that this was

due in part to felling of exotic conifers for heathland restoration. The latest Forestry Commission data on unconditional felling licenses gives deforestation rates of 0.59 kha yr⁻¹ in England, 0.199 kha yr⁻¹ in Scotland and of 0.04 kha yr⁻¹ in Wales.

Emissions from biomass burning are estimated using the rates described in the IPCC 1996 guidelines (IPCC 1997a, b, c), based on a mature broadleaf forest stand. Only immediate losses are considered because sites are normally cleared completely for development, leaving no debris to decay. Changes in soil carbon stocks after deforestation are estimated with the other land use transitions.

This method is also used to estimate biomass burning emissions from Forest land converted to Settlements. Deforestation to Cropland in the UK since 1990 is negligible.

1.10.4 5D Wetlands

In previous inventories, data were not included for this category as Wetlands either fell within the Grassland category or open water, which is included in the Other Land category, due to the classifications in the land surveys used to construct the land use change matrix. In this inventory, data on peat extraction is reported within the wetlands category for the first time. The annual volume of peat extraction in Scotland and England is obtained from the Business Monitor of Mineral Extraction in Great Britain (ONS, 2008). Extraction in Northern Ireland is estimated from a 1997 survey (Cruikshank and Tomlinson 1998). The volume extracted is variable in England (650,000 – 1,600,000 m³ yr⁻¹) and Scotland (100,000-700,000 m³ yr⁻¹) but constant in Northern Ireland (628,000 m³ yr⁻¹). Fixed DA-specific emission factors (Cruikshank and Tomlinson 1998) are used to estimate emissions from this activity. Activity data for the area of offsite emissions from the extraction of horticultural peat were reported to be 4.57kha in England, 1.29 kha in Scotland, 0.48 kha in Wales and 1.03 kha and Northern Ireland.

1.10.5 5E Settlements

No changes in carbon stocks are reported for land In Settlements remaining Settlements. Methods for estimating changes in non-forest biomass and soils due to land use change to Settlements are described in the Land converted to Cropland section. Since 1990 an estimated 10.2 kha yr⁻¹ have been converted to Settlements in England, 2.3 kha yr⁻¹ in Scotland, 1.6 kha yr⁻¹ in Wales and 1.6 kha yr⁻¹ in Northern Ireland.

The method used for estimating emissions from biomass burning from Forest land converted to Settlements (deforestation) is the same as that used for Forest land converted to Grassland. The activity data comes from the Ordnance Survey's annual assessment of land use change for map updates (collected on behalf of the Department of Communities and Local Government) (DCLG 2008). Deforestation to Settlements is estimated to be the area that has moved from the forest land-use class to the non-rural land use class (see Levy and Milne 2004 for more details). At present, this data is only available for England for 1990-2005 (smoothed using a five-year moving average); estimates for Scotland and Wales are made by extrapolation from the English rates, assuming that England accounted for 72 per cent of deforestation, based on the distribution of licensed felling between England and the rest of GB in 1999 to 2001. Annual rates of deforestation to Settlements are variable but stable, with an average of 0.35 kha yr⁻¹ for England, 0.11 kha yr⁻¹ for Scotland and 0.03 kha yr⁻¹ for Wales. Deforestation is not currently estimated for Northern Ireland.

The activity data for deforestation to Settlements come from a continuous rolling survey programme and were updated this year. This revision had a knock-on effect on other parts of the land use change matrix but the overall impact on emissions/removals was relatively small.

1.10.6 5F Other Land

No emissions or removals are reported in this category. It is assumed that there are very few areas of land of other types that become bare rock or water bodies, which make up the majority of this type.

1.10.7 5G Other Activities

Changes in stocks of carbon in harvested wood products (HWP) are reported here. The C-Flow model (described in the Forest land section) estimates changes in stocks of HWP produced from the management and harvesting of conifer and broadleaf forests planted since 1922. The activity data used for calculating this activity is the annual forest planting rates. For a given forest stand, carbon enters the HWP pool when thinning is undertaken (depending upon the species thinning first occurs c. 20 years after planting) and when harvesting takes place.

According to this method the total HWP pool from UK forests is presently increasing, driven by historical expansion of the forest area and the resulting history of production harvesting (and thinning). The stock of carbon in HWP (from UK forests planted since 1921) has been increasing since 1990 but this positive stock change rate recently reversed, reflecting a severe dip in new planting during the 1940s. The net carbon stock change in the HWP pool has returned to a positive value (i.e. an increasing sink) in 2007, and is forecast to increase sharply as a result of the harvesting of the extensive conifer forests planted between 1950 and the late 1980s. Different DAs have different patterns depending on their forest planting history.

Ongoing Improvements in Data and Method Planned for the Next Inventory Cycle

Forest Land

The area reported under 5.A.1 Forest Remaining Forest is likely to be revised when the new National Forest Inventory becomes available. National forest statistics do not currently capture forest conversion to other land uses, so a separate adjustment is made to the forest areas and carbon stock changes reported in the inventory. The NFI map due in spring 2011 will establish a new baseline of the total forest area in the UK.

Forest Research has been carrying out an analysis of Forestry Commission GB National Forest Inventory (NFI) data, for a base year of 2005, as well as records maintained in the Forestry Commission Sub-compartment Database (SCDB), for a base year of 2009, to establish the detailed composition of woodlands in England, Scotland and Wales in terms of tree species, growth rates and types of management. The SCDB (which covers slightly less than half of the forest area in Britain) gives detailed, stand-by-stand information about species, growth rate and management prescription. This is being combined with NFI data on the species composition of private woodlands to obtain an overall on the status and management of woodlands in England, Scotland and Wales. Contact has also been made with experts in the Northern Ireland Forest Service with the aim of undertaking a similar exercise for Northern Ireland. At the same time, the Forest Research CARBINE forest carbon accounting model has been significantly upgraded to enable the representation of a much more complete range of tree species and growth rates, as well as many more examples of management prescriptions involving different rotation periods, no thinning, thinning and, where appropriate, management according to continuous cover principles and even no management. New estimates based on this information and the supporting methodology should be available in the summer of 2011.

Forest Research has been seeking to determine a better understanding of the contribution to the carbon balance made by woodlands established before 1921. This has involved analysing the Forestry Commission NFI data for the base year of 2005 in combination with Forestry Commission reports on areas of new planting since 1920. This also involves complex modelling of changes in the age class structure of forest areas, based on species composition, growth rate and management of woodland areas, in particular rotation periods. The aim is to factor out post-1920 woodlands from NFI data to obtain a picture of the age class structure, status and management of pre-1921 woodlands. An initial attempt to demonstrate proof of concept through development and application of an appropriate methodology to woodlands in Wales has been successful. The methodology is now being implemented in detail and applied to analysis of woodlands in England, Scotland and Wales. Contact has also been made with experts in the Northern Ireland Forest Service with the aim of undertaking a similar exercise for Northern Ireland. New estimates based on this analysis should be available in summer 2011.

As described earlier, Forest Research is undertaking a major analysis of woodland areas in terms of species composition, growth rates, age class structure and management, with the aim of producing results by summer 2011. These results will be disaggregated to the level of each country (England, Scotland, Wales and Northern Ireland). At this point it will be appropriate to review the methodology and assess any impacts on reported and projected emissions and removals due to forestry.

Forest Research has been analysing the spatial distribution of afforestation of soils, and the likelihood of associated drainage and fertilisation, with an initial focus on Scotland. The Scottish Soil Map Classification and the Forestry Commission National Forest Inventory (NFI) map for 2001 have been compared for this purpose, enabling woodland areas to be categorised according to coniferous, broadleaf or mixed species composition and association with deep or shallow peat soils and other major soil types (e.g. brown earths, surface water gleys etc.). Extant and historical Forestry Commission guidance on site establishment practices will be a strong indicator of drainage and fertilisation activities for the woodlands on different soil types. The impacts of woodland establishment (on different soil types) on fluxes of CO₂ and non-CO₂ greenhouse gases are also being assessed as

part of this study. The results of the analysis for Scotland should be available by Spring 2011. Work is about to start on a similar analysis for woodlands in Wales with the aim of completing an analysis for England, Scotland and Wales by Autumn 2011. Consideration is being given to extending this study to Northern Ireland.

Improved representation of the fate of forest carbon following harvesting is being achieved by taking more detailed account of woodland composition and management, in particular patterns of thinning and clear felling (on appropriate rotations) as part of the analysis being carried out by Forest Research as described earlier.

Cropland

This category will be internally restructured in the next inventory submission, moving to the IPCC 20-year default period for land remaining in the Land Converted to Cropland category. This will resolve the discrepancies between the areas reported in the CRF tables and those in the Land Use Transition Matrix tables in 1990-1991 and 2008-2009. It is thought that these discrepancies arise because some cropland (and pasture grassland) is in multi-year rotations and these areas have therefore undergone several land use transitions between 1950 and the current inventory year (with the area changes being cumulatively reported in Category 5.B.2, Land Converted to Cropland). Restructuring will allow the total area of Cropland to be made consistent with the cropland area reported in the annual agricultural census or Countryside Survey. Soil carbon stock changes resulting from historical land use change will still be reported in the Cropland remaining Cropland category, and work is planned to resolve the rotational land use change issue.

The weighting used for changes in soil carbon density between different land use types is currently based on the 1990 and 1998 Countryside Survey data. This approach will be updated to include the 2007 Countryside Data in the next inventory submission. No account is currently taken of other carbon stock changes in perennial woody biomass on cropland, for example fruit orchards or crops grown for biofuel production. The area of such crops is currently small (orchards cover 23.7 kha in 2008 or 0.4% of the total croppable area (Defra 2009)). This is an area of potential improvement in the inventory, although not a high priority.

A literature review of non-forest biomass yield improvements will be undertaken and the carbon stock change calculations updated accordingly.

A current development project is examining the scope for the effects of land management policies to be better reflected in the inventory. This has a particular focus on soil carbon. An assessment of the availability and quality of activity data and country-specific emission factors is necessary before this development can be implemented. A wider review of data sources is also in progress: this will be used to assess whether there are additional data sources that could be used in the inventory or contribute to uncertainty estimates. The review will also assess other models that could potentially be used, although a full methods comparison is beyond the intended scope of the review.

Grassland

This category will be internally restructured in the next inventory submission, moving to the IPCC 20-year default period for land remaining in the Land converted to Grassland category. This will allow the reporting of areas to be made more consistent with other international reporting, in a similar manner to the planned changes to reporting cropland as outlined above.

Input activity data for deforestation remain a problem but outputs from the National Forest Inventory and map may help to resolve this (see the Forest Land section for further information). The initial focus involves the comparison of snapshot woodland area maps from the Forestry Commission's NIWT (1995-1999) and the new NFI map. In principle, it should be possible to derive provisional deforestation estimates from such a comparison, however technical issues concerning the methods used to prepare the two maps are still being addressed. The publication of the new NFI map has been put back until mid-2011. The intention is that eventually deforestation estimates should be obtained directly from periodic National Forest Inventories (NFIs), as results from these become available.

There is currently no national activity data for wildfires and controlled burning on non-forest land use types. There is a new Vegetation Fire Monitoring Standard in development which may produce useful data in the future but this will only cover England at first. We will investigate the potential for extrapolating burnt areas from the number of fire incidents, for which statistics are available.

Wetlands

Work is in progress to develop the time series of areas undergoing peat extraction and post-extraction restoration, as suitable data is known to be available in Scotland and Wales. It may be possible to develop a Tier 2 methodology for this category that separates peatlands being converted for peat extraction from those already in production. The method combining site locations with Google Earth imagery could be extended to Northern Ireland if a similar register of commercial extraction sites is available. Further work on volume conversion factors for off-site emissions from horticultural peat has been undertaken and will be included in the next inventory submission. Additional work on uncertainty analysis in this category is required. The area of land converted to permanently Flooded Land is assumed to be small in the UK but activity data will be sought. We also plan to start reporting areas of open water in this category (rather than as part of the Other Land category). It is assumed that there are no emissions associated with such areas.

Settlements

This category will be internally restructured in the next inventory submission, moving to the IPCC 20-year default period for land remaining in the Land converted to Settlement category.

Input activity data for deforestation remain a problem but outputs from the National Forest Inventory and map may help to resolve this (see the Forest Land section and for further information).

Other Land

We have started reporting areas in the CRF Table for this category. The planned restructuring of the Cropland, Grassland and Settlement categories in the next inventory submission will improve consistency in area reporting. The intention is to start reporting areas of open water in the Wetlands category.

Other Activities

The emission factors and activity data for harvested wood products will be kept under review. A watching brief on the development of methods of HWP estimation has been maintained (Matthews 2009). It is too early to proceed with full implementation of any specific method and preparation of estimates until methodologies have been agreed by the IPCC.

1.11 WASTE

1.11.1 Solid Waste Disposal on Land

In the UK Inventory, emissions are estimated based on a model of methane production in landfill sites. The generation of methane is assumed to follow a first order model with different decay rates for different types of waste. The model requires data on waste disposals and waste composition from 1945 onwards.

The UK GHGI waste model has been revised and updated during the 1990-2009 inventory cycle by Eunomia Research & Consulting. This update incorporated a range of DA-specific data on waste composition, MSW arisings, and the report can be found at:

<http://decc.gov.uk/publications/DirectoryListing.aspx?tags=40>

The AEA team provided a suite of DA-specific waste sector datasets that have been incorporated into the landfill model revision funded by Defra during the 1990-2009 UK GHGI cycle. Consultation with waste sector experts during 2010 led to the collation of new information from recent waste survey and compositional studies including: MSW composition (Scotland WRAP study, Wales WRAP study), public sector waste arisings survey (Wales), Commercial & Industrial & Agricultural waste fate studies (Wales), Northern Ireland waste compositional study and Commercial & Industrial waste sector report.

The revised model retains much of the design and functionality of the previous GHGI waste model, with no adjustment made for different proportions of waste disposal to landfill in each DA. Due to a lack of detailed local data, it has also had to be assumed that the degree of methane recovery is the same in each region.

In the 1990-2009 DA GHGI compilation, data from the www.WasteDataFlow.org website have been used to estimate DA landfill activity. The Waste Data Flow website is used by Local Authorities to track progress towards waste sector objectives. Reports from the Waste Data Flow organisation have been reviewed, and these provide a more detailed split of waste disposal options undertaken in the DAs, with ultimate fate of municipal waste recorded against numerous options, primarily: recycling, landfill and incineration. Hence this source of data (which has limited data back to 1999) has been used to derive regional drivers for emissions of methane from landfill waste. Note that this method is still limited in its accuracy, as the lack of detailed local data necessitates that average UK waste composition and average UK methane recovery is still assumed within each DA.

1.11.2 Waste Water Handling

Emissions from waste-water handling are based on population statistics for the UK. These are taken from the Office of National Statistics (ONS, 2009a) and assume that the application of different sludge treatment and disposal options is uniform across the UK. There has been some improvement to the UK GHGI methodology for this source during the 1990-2009 cycle, but insufficiently complete DA-specific data is currently available. Improvements to this source estimation method will be considered during the next inventory cycle but is dependent on accessing new data from water companies, notably those in England and Wales.

1.11.3 Waste Incineration

The UK Inventory reports emissions from the incineration of sewage sludge, municipal solid waste and some chemical waste. Regional estimates are based on DEFRA (2010a) which reports data for the amount sewage incinerated for Scotland, Northern Ireland and England & Wales.

Emission drivers from MSW Incinerators for 1990-1995 are based on capacity data for individual incinerators taken from RCEP (1993). It is assumed there were no significant changes between 1990 and 1995. Estimates for recent years are based on plant capacity data (Patel, 2000). All of the larger MSW incinerator plant have been re-fitted during the late 1990s to generate electricity and are therefore reported as power stations in the regional inventories. A handful of smaller waste incinerators (municipal, industrial and clinical) are used for district heating and are reported as commercial or miscellaneous. The disaggregation of these smaller heat-generating plant is based on the same driver as for larger MSW incinerators, as there is no specific source of information that provides a more satisfactory estimation of the regional split. The total consumption of these incinerators is reported within the Digest of UK Energy Statistics (DECC, 2010a).

Emissions from clinical waste incineration are allocated to the regions based on a set of plant capacity data for 1998. Emissions data from chemical waste incineration are available for England and Wales only, based on data taken from the Pollution Inventory (Environment Agency, 2010a), and these data are used for the DA estimates also. Some chemical waste incineration takes place in Scotland but no emissions data are available, and hence the emissions contributed from this source are currently omitted from the Scottish inventory data. No chemical incinerators have been identified in Northern Ireland.

1.12 UNCERTAINTIES

1.12.1 Introduction

The uncertainties in the UK Inventory are estimated using a Monte Carlo simulation. Eggleston et al (1998) and Salway et al (2001) describe this in detail. The method involves estimating the uncertainties in the activity data and the emission factors for all the emission source categories and then using a Monte Carlo simulation package to calculate the uncertainty in the emission totals. In order to apply a similar approach to the DA GHG inventories, it is necessary to estimate uncertainties for the DA activity data (i.e. fuel consumption, production data). The same emission factors are used in the DA inventories as in the UK Inventory, so their uncertainties are known. In the UK Inventory uncertainties in the activity data for fuel use are estimated on the basis of the statistical differences

between fuel supply and demand data reported in the energy statistics. However, such data is not available for the DA-specific activity data used. Moreover, for some sources, no direct activity data is available at all, and it has been necessary to distribute the UK data using surrogate data (e.g. employment statistics). In such cases, it is impossible to say whether the surrogate statistics are an accurate indicator of fuel consumption.

1.12.2 DA GHGI Uncertainty Estimation

The uncertainties in the DA GHG inventories are also estimated using a Monte Carlo simulation. In order to simplify the calculations, the source categories are broader than those used in the UK GHG Inventory simulation. In the DA inventory simulation, the combustion categories are effectively the total consumption of a particular fuel. This contrasts with the UK simulation where there is a further disaggregation into sectoral categories (e.g. power stations, refineries). The rationale for this is that it is more practicable to estimate the uncertainty in the total consumption of a fuel in a region than to attempt to estimate uncertainties in diverse sectors where in some cases surrogates have been used.

For each of the broad source categories, an estimate of the activity uncertainty has been made for Scotland, Wales, Northern Ireland, and Unallocated emissions. The approach adopted was to estimate a factor to scale the UK uncertainty with, based on knowledge of the relative uncertainties of the DA estimates to each other, and to the UK total.

The factors used to weight the uncertainties for each of the DAs have been reviewed as part of the model update. The DA inventories have evolved since the uncertainty model was set up, and so some of the assumptions that were previously used are now no longer relevant. For example, different methods used to be used for fuel sales data in Northern Ireland and therefore the Northern Ireland estimate was assumed to be less uncertain than the other DAs. Now that DECC regional energy statistics are used across all four DAs, this difference in uncertainty is no longer assumed.

It is important to note that the uncertainties in the inventories for the UK, England, Scotland, Wales, Northern Ireland, Unallocated are inter-dependent, because:-

UK Emissions = [England + Scotland + Wales + Northern Ireland + Unallocated]

Therefore uncertainties from the UK, Scotland, Wales, Northern Ireland, and Unallocated emissions are estimated using the model, and uncertainty in the English emissions is calculated.

In many of the non-combustion sources (e.g. industrial processes, coal mines) the overall uncertainty is dominated by the emission factor and the uncertainty in the activity data is not a determining factor. Therefore, there is unlikely to be any significant variation in uncertainties between DAs. In these cases, a low uncertainty for the activity data (say 1%) may be assumed for each DA, whilst the UK uncertainty for the emission factor is applied.

For sources where the UK total is made up as a sum of the DA totals, such as agriculture or LULUCF, no additional uncertainty is introduced in the process of deriving the UK split, and therefore the UK uncertainty parameters are applied directly to the DA estimates.

In the case of halocarbons and SF₆ emissions it is not considered feasible to attempt to assume varying uncertainties across the DAs. The F-gas uncertainties have been considered at a sector level. This means that the uncertainty for each sector emission at UK level has been applied to the DA estimates, so that the overall uncertainty for each of the F-gases reflects the mix of sources that are the most significant for each of the DAs, and their relative uncertainties.

1.12.3 Trend Uncertainty Analysis

The DA uncertainty model has been extended to provide estimates of the uncertainty in the emissions trend. The model uses, where possible, the same principles as the UK uncertainty model, however, these estimates are currently indicative since it will take further improvement work to refine the estimates fully.

In order to estimate the uncertainty on the trend, it was necessary to make an estimate of the uncertainty in the base year (1990 for CO₂, CH₄ and N₂O, and 1995 for the F-gases). This estimate is made for the UK Inventory, as part of the analysis presented in the National Inventory Report (MacCarthy *et al.*, 2011). Therefore, it was possible to make the DA uncertainty estimates using the

method described above in conjunction with the UK estimates for the base year. The DA weighting factors were reviewed as part of this process, because changes to the source data and methods used across the time series mean that it is not always appropriate to apply the same weighting factor in the latest year as in the base year.

In addition to the estimation of the uncertainty in each year, it was also necessary to consider correlations between sources across years. The UK uncertainty model considers correlations in the estimates of

- N₂O from agriculture;
- CH₄ from landfills;
- CH₄ from leakage from the gas distribution network; and
- N₂O from waste water treatment.

These correlations have been replicated in the DA model. The uncertainty in the trend is particularly sensitive to the correlation in emissions from agricultural soils.

The uncertainty in the trend between 2008 and 2009 has been considered. However, the correlations between activity data and emission factors between adjacent years are not currently well understood, and it has therefore not been possible to produce a meaningful estimate at this time.